



Revista de Osteoporosis y Metabolismo Mineral

ISSN: 1889-836X

ISSN: 2173-2345

Sociedad Española de Investigaciones Óseas y
Metabolismo Mineral

Martínez Arias, L; Panizo García, S; Carrillo López, N; Barrio Vázquez, S; Quirós
González, I; Román García, P; Mora Valenciano, I; Miguel Fernández, D; Añón
Álvarez, E; Fernández Martín, JL; Ruiz Torres, MP; Cannata Andía, JB; Naves Díaz, M
Efecto de la enzima antioxidante catalasa en la calcificación vascular y desmineralización ósea
Revista de Osteoporosis y Metabolismo Mineral, vol. 9, no. 1, 2017, January-March, pp. 13-19
Sociedad Española de Investigaciones Óseas y Metabolismo Mineral

DOI: <https://doi.org/10.4321/S1889-836X2017000100003>

Available in: <https://www.redalyc.org/articulo.oa?id=360955391003>

- How to cite
- Complete issue
- More information about this article
- Journal's webpage in redalyc.org

redalyc.org

Scientific Information System Redalyc

Network of Scientific Journals from Latin America and the Caribbean, Spain and
Portugal

Project academic non-profit, developed under the open access initiative

Martínez Arias L¹, Panizo García S¹, Carrillo López N¹, Barrio Vázquez S¹, Quirós González I³, Román García P³, Mora Valenciano I⁴, Miguel Fernández D², Añón Álvarez E², Fernández Martín JL¹, Ruiz Torres MP⁴, Cannata Andía JB^{1,5}, Naves Díaz M¹

1 Servicio de Metabolismo Óseo y Mineral - Instituto Reina Sofía de Investigación Nefrológica - REDinREN del ISCIII - Hospital Universitario Central de Asturias - Oviedo (España)

2 Laboratorio de Medicina - Hospital Universitario Central de Asturias - Oviedo (España)

3 Universidad/Instituto Sanger - Cambridge (Reino Unido)

4 Departamento de Biología de Sistemas - Universidad de Alcalá de Henares - REDinREN del ISCIII - Alcalá de Henares - Madrid (España)

5 Departamento de Medicina - Universidad de Oviedo - Oviedo (España)

Effects of the catalase antioxidant enzyme in vascular calcification and bone demineralization

DOI: <http://dx.doi.org/10.4321/S1889-836X2017000100003>

Correspondence: Manuel Naves Díaz - Servicio de Metabolismo Óseo y Mineral - Instituto Reina Sofía de Investigación Nefrológica - REDinREN del ISCIII - Hospital Universitario Central de Asturias - Edificio FINBA, 1ª planta - Avenida de Roma, s/n - 33011 Oviedo (Spain)
e-mail: manuel@hca.es

Date of receipt: 23/05/2016

Date of acceptance: 12/07/2016

Work awarded a scholarship Research Molecular Biology Bone FEIOMM 2012.

Summary

Objetives: Assess the role of the catalase antioxidant enzyme in the vascular calcification process associated with chronic renal failure (CRF) and its effect on bone mass.

Material and methods: Wild type C57/BL6J mice (WT) and transgenic mice (TG) were used, that overexpress the catalase enzyme, to which CRF was induced. Control WT and TG mice were used in simulated intervention. After 16 weeks, the mice were sacrificed, with serum samples taken for biochemical markers as well as residual pieces of kidney, aorta and tibias. An *in vitro* model of primary culture of smooth vascular muscle cells (SVMC) taken from the WT and TG aorta which underwent eight days of 3 mM phosphorus and 2 mM calcium calcifying medium.

Results: A significant increase in Runx2 gene expression, calcium renal deposit and bone structure deterioration at trabecular level was only detected in WT mice with CRF. This was not observed in TG mice with CRF.

Only in the case of WT mice SVMC, did added calcification medium raise calcium levels, proteic Runx2 expression and the reactive oxygen species of mitochondria with low catalase enzyme.

Conclusions: Calcifying catalase over-expression was observed in both *in vivo* and *in vitro*, with *in vivo* showing that this reduction was accompanied by an improvement in bone parameters under study.

Key words: vascular calcification, bone, antioxidants, oxidative stress, catalase, μ CT, chronic renal failure.

Introduction

Cardiovascular disease currently represents the leading cause of death in the developed world. It is expected to continue rising in the coming decades due to our aging population. One factor that contributes to cardiovascular risk is oxidative stress¹. Different stimuli are associated with the development of cardiovascular disease including macrophage activation, hyperglycemia, LDL oxidation and even angiotensin II which exert their harmful effects, at least partially, through the local synthesis of reactive oxygen species²⁻⁵.

On the other hand, there is evidence of a positive relationship between oxidative stress and vascular calcifications⁶. Vascular smooth muscle cells (VSMC) subjected to oxidative stress increase the activity of alkaline phosphatase and calcium deposition, indicating their transdifferentiation toward cells capable of mineralizing (osteoblast-like/chondrocyte)⁷. In VSMC in primary culture of mouse aorta, the oxidative stress induced by hydrogen peroxide or glucose oxidase promotes calcification and overexpression of Runx2 (Cbfa1), transcription factor related bone osteogenic differentiation, mediated AKT⁸. Interestingly, these same stimuli have the opposite effect on osteoblast precursor cells, demonstrating the importance of hydrogen peroxide or oxygenated water (H₂O₂) in the process of differentiation of mesenchymal cells lineage⁹.

The main antioxidant enzymes involved in the catalytic removal of hydrogen peroxide is catalase, glutathione peroxidases and thyroidin peroxidase. Of these, the catalase is the most efficient in removing the hydrogen peroxide enzyme.

One of the factors affecting aging is the progressive accumulation of oxidative damage. This damage may be due to exposure to normal intracellular oxidative stress or, in pathological situations, an increase in this stress due to inflammation or other causes can actually cause accelerated aging. Therefore, our objective was to evaluate the role played by overexpression of catalase on the process of vascular calcification associated with moderate kidney disease and its effect on bone mass¹⁰. For this purpose, a transgenic mouse model overexpressing catalase enzyme, subjected to chronic renal failure was used.

Material and methods

Experimental model *in vivo*

Establishment of chronic renal failure (CRF)

C57/BL6J wild (WT) and C57/BL6J transgenic (TG) overexpressing the antioxidant enzyme catalase were used. To induce CRF, mice underwent 3 months old to a first intervention, which involved the lateral opening of the animal on the right side where the two poles of the kidney is cauterized. Isoflurane anesthesia was used (1-2%) by inhalation. A week after the first operation, the animal underwent a second intervention, which consisted of opening the left side and complete removal of the kidney. After 16 weeks of the last operation, the animals were sacrificed by exsanguination, anesthetized with CO₂. In slaughter animal serum he was

obtained to analyze and general biochemical markers of bone metabolism: urea, calcium, phosphorus, Ca-P product, iPTH and FGF23. the remaining piece of kidney, aorta and tibias were also extracted.

The left tibia, which was preserved in ethanol 70°, was analyzed by computerized microtomography (μCT) with a SkyScan 1174, Bruker μCT (Kontich, Belgium) equipment. The 2D and 3D morphometric analysis was carried out using the CTAn software. The region of interest (ROI) was defined manually in each sample. For the trabecular region, 150 cuts were selected and threshold levels used in the gray scale between 78 and 250. The morphometric analyses were based on internal plug-ins CTAN in 2D and 3D. Morphometric parameters were measured trabecular bone volume (BV/TV,%), trabecular spacing (TbSp, μm), trabecular number (ToNb, mm⁻¹) and trabecular porosity (TbPo, μm).

All studies were approved and authorized by the Committee on Animal Experimentation of the University of Oviedo.

In vitro experimental model

Primary cultures of aortic VSMCs from C57/BL6J WT and TG were used. To do this, the aortas of animals chopped and put explants to grow in culture dishes coated with fibronectin to promote adhesion.

Cells were cultured in DMEM medium supplemented with fetal bovine serum at 10%. When cells reached 60-70% confluency was replaced with DMEM-F12 supplemented medium with 0.1% bovine albumin (control medium) and calcifying medium supplemented medium consisting of control phosphorus and calcium at concentrations of 3 mM and 2 mM, respectively. Cells were incubated under these conditions for 8 days.

The basal activity of catalase was measured in VSMC from WT and TG mice using the commercial kit "catalase assay kit" (Cayman Chemical, 707002), following the protocol established by the manufacturer.

Levels of reactive oxygen species in VSMC cultured WT and TG mice with control medium and calcifying were measured with a fluorochrome specific mitochondria, dihydrorhodamine 123 (DHR 123).

Proteins of cell cultures were extracted with a buffer composition 50 mM Tris-HCl, 150 mM NaCl, 1% NP-40, sodium deoxycholate 0.5%, 1.0 mM EDTA and 0.1% SDS inhibitor proteases. the sample was sonicated in cold 10 minutes to prevent the breakdown of proteins and centrifuged at 10,000 rpm for 5 minutes at 4°C proteins of the supernatant was collected, quantified by the Bio-Rad DC method and stored at -80.

20 μg of protein were electrophoresed on acrylamide gels of 0.75 mm thickness under denaturing conditions (SDS-PAGE), for identification using known molecular weight markers (Molecular Weight Markers Rainbow™, GE Healthcare, United Kingdom). Proteins were transferred to a nitrocellu-

lose membrane (Hybond Amersham™ 0.45 µm PVDF, GE Healthcare, UK). The transfer was carried out in the cold for one hour at 100 v. After the same, the membrane was blocked for one hour with 5% milk in phosphate buffered saline (PBS). Subsequently, membranes were kept overnight with the primary antibody (monoclonal antibody of catalase in 1: 5000 dilution in BSA, Runx2 1: 500 in BSA against glyceraldehyde 3-phosphate dehydrogenase (GAPDH) diluted 1: 5000 in BSA, all from Santa Cruz Biotechnology, Inc., Santa Cruz, California, USA). The next day, the membrane was washed three times at intervals of 10 minutes each with wash solution composed of PBS and Tween-20 (Sigma-Aldrich), and subsequently incubated with antibody or goat against rabbit. The membrane was washed with the washing solution described above and the protein was detected by ECL Western Clarity™ Substrate (BIO-RAD, USA) commercial kit. The relative quantification of the intensity of the bands obtained in the Western blotting was performed with Image Lab™ Software and Molecular Imager scanner ChemiDoc™ XRS+ (both from BIO-RAD, USA).

Common techniques employed

qRT-PCR

The analysis by qRT-PCR was performed with the cDNA obtained from different experimental conditions using the High Capacity Reverse Transcription Kit (Applied Biosystems) and phenol RNA extraction. For normalization of results constitutive expression of genes (rRNA Runx2 and GAPDH) and $\Delta\Delta CT$ relative quantification method was used.

Study of vascular calcification

After addition of 0.6 N HCl for 24 hours to extract the cell or tissue calcium, the calcium content was quantified by ortho-cresolphthalein method complexone. Calcium levels were relativized to total protein content.

Statistical analysis

For the statistical analysis of the results the SPSS 17.0 software was used. Comparison of the treatment groups was performed using chi-square in the case of categorical variables and by Student's t test for numerical variables.

Results

a) In vivo studies:

Biochemical markers

The CRF in WT and TG mice showed increased levels of urea, Ca-P product, and iPTH FGF23. In the case of FGF23 increase it was much higher (10 times) in WT mice with CRF regarding their Sham control.

Moderate CRF effect on vascular calcification and bone level changes

Although the C57/BL6J mouse strain used for the calcification studies did not allow us to obtain vascular calcification in the aortas (Table 1), a significant increase in gene expression of Runx2, osteo-

blast differentiation marker, was observed only in the WT group with CRF (Figure 1). This increase was not observed in the TG animals with CRF. The effect of CRF on the calcification process was studied in other soft tissue such as the kidney, with an increase of calcium content in CRF mice with a higher increase in WT than in TG mice (Table 1).

Vascular level changes were also observed at bone level. In WT animals with CKD deterioration of bone structure it was about its Sham control group with a statistically significant decrease in trabecular bone volume (Figure 2A) and the number of trabeculae (Figure 2C), and increased trabecular separation (Figure 2B) and trabecular porosity (Figure 2D). The group TG mice with CRF did not differ with respect to their Sham control group at trabecular level.

b) In vitro studies:

Effect of overexpression of catalase in the hydrogen peroxide purification and Runx2 expression in VSMCs

The VSMCs from the TG mice had increased activity and protein expression of the catalase enzyme at baseline compared with those of WT mice (Figure 3 A and B).

When calcifying medium was added to VSMC WT mice there was increased calcium content. This effect was not observed in VSMC TG mice (Figure 4A). In parallel, the protein expression of Runx2 was studied in VSMC of WT and TG mice with calcifying medium for 8 days and an increase in Runx2 expression in VSMC of WT mice was observed which was not observed in TG (Figure 4B).

Study of the levels of markers of oxidative stress and protein levels of catalase

An increased level of fluorescence of the probe DHR123 was observed in the VSMC of WT mice cultured with calcifying medium, indicating an increase in mitochondrial oxygen reactive species. This effect was not observed in VSMC of TG mice, but there was a decrease of reactive oxygen species (Figure 5). Lower protein expression was also observed of the enzyme catalase by culturing the VSMC of WT mice for 8 days with calcifying medium relative to those from TG mice (Figure 6).

Discussion

In this work we have been able to confirm that overexpression of catalase antioxidant enzyme in a mouse model protects the process of vascular calcification and bone deterioration. In the VSMC of these same mice, a decrease was observed in the levels of reactive oxygen species, but also osteogenic proteins such as Runx2.

At the biochemical level, increased FGF23 was particularly noticeable in WT animals with CRF regarding the Sham group (10 times). This effect was less marked in the TG with CRF whose increase was twice that of the control group (Sham). Some authors have postulated the role of FGF23 as a calcification inducer^{11,12}. It has even been associated with high levels of renal mortality in patients¹³.

Table 1. Values of biochemical WT and TG mice with and without CRF


	WT Sham (n=7)	WT CRF (n=9)	TG Sham (n=10)	TG CRF (n=10)
Urea (mg/dL)	47±3	83±13*	41±4	94±19*
Ca-P product (mg ² /dL ²)	69±7	81±12*	60±14	74±13*
iPTH (pg/mL)	316 (0.834)	3,941 (3,649-4,499)*	338 (35-1,836)	3,411 (1,204-3,868)*
FGF23 (pg/mL)	111 (102-125)	1,103 (773-1,143)*	224 (204-268)	437 (153-988)
Calcium aorta (µg/mg)	92±7	95±6	82±11	93±5
Calcium kidney (µg/mg)	3.0 (1.9-4.0)	562.8 (200.2-636.3)*	0 (0-3.3)	20.7 (3.7-123.8)*

WT: wild mouse; TG: transgenic mouse; CRF: chronic renal failure. *p<0.05 with respect to their respective Sham.

On the other hand, it inhibits FGF23 expression of CYP27B1 gene encoding the alpha 1-hydroxylase suppressing renal calcitriol synthesis from its precursor 25-hydroxyvitamin D₃¹⁴. Furthermore, FGF23 activates CYP24 gene expression that encodes 24 hydroxylase, the enzyme that hydrolyzes and inactivates calcitriol¹⁵. This contributes to the decline in the synthesis of calcitriol, but also to degradation which leads to decreased levels of vitamin D, a factor that could induce vascular calcification as has been shown in epidemiological studies^{16,17}.

The calcium content in the aorta of TG mice with CRF was similar to WT mice with CRF. While this may seem paradoxical and contradictory to other observed results, Giachelli et al. have confirmed the absence of calcification in the aorta in the same strain of mice used for transgenic generation¹⁸. However, the protective effect of the overexpression of catalase preventing calcium accumulation was observed in other soft tissue such as the kidney.

Figure 1. Levels of gene expression of Runx2 (relative units) in aorta in different groups. WT: wild mouse; TG: transgenic mouse; CRF: chronic renal failure



*p<0.05 with respect to their respective Sham.

Moreover, high levels of FGF23 have been associated with suppression of osteoblast differentiation and mineralization of bone matrix *in vitro*¹⁹. This could perhaps explain the negative effect of CRF on bone deterioration in WT mice and to a much lesser extent in the TG mice. Recent studies by our group have shown bone loss in a rat model with CRF and high levels of phosphorus, PTH and FGF23. In the same study, gene silencing studies have confirmed that FGF23 only has a direct effect activating Dkk1²⁰, inhibitor of Wnt pathway which is involved in pathway inactivation. Thus, high levels of FGF23, as observed in our experimental model in WT mice with CRF, could have contributed to the decrease in bone mineral density through inactivation of the Wnt pathway.

The increase in reactive oxygen species contributes to increased osteogenic protein, being a stimulus for the start of the calcification process^{7,8}. This effect has been observed in VSMC of WT mice subjected to calcifying stimuli. However, in VSMC of TG mice subjected to a calcifying stimulus, not only did the expression of reactive oxygen species not increase but it decreased, as with the protein expression of Runx2, osteogenic protein and early marker process calcification. In fact, the importance of hydrogen peroxide has noted as a second messenger involved in intracellular signaling²¹ regulated by oxidative stress. A drop of hydrogen peroxide by catalase overexpression contributes to lower oxidative stress and decreased vascular calcification process through inhibition of Runx2.

In view of the results obtained, studies are needed to ascertain the mechanisms by which decreased oxidative stress confers an advantage at both vascular and bone level. The reduction of the inflammatory process, maintaining protein levels of renal Klotho, the main molecule involved in aging, or regulation of the Wnt pathway may help explain the different behavior, so further research into the field is required.

Figure 2. Changes to bone level; WT: wild mouse; TG: transgenic mouse; CRF: chronic renal failure. A) Trabecular bone volume (BV/TV) in different groups. * $p < 0.05$ relative to its respective Sham. B) Trabecular separation (TbSp) in the different groups. * $p < 0.001$ relative to its respective Sham. C) Number of trabeculae (TbN) in the different groups. * $p < 0.05$ relative to its respective Sham. D) Porosity trabecular (PoTb) in the different groups. * $p < 0.05$ relative to its respective Sham




Figure 3. A) Basal Activity of catalase enzyme in WT and TG VSMC mice. B) Protein expression in relative units of catalase enzyme in VSMC WT and TG mice by Western blotting; WT: wild mouse; TG: transgenic mouse. * $p < 0.001$ compared to WT group




Figure 4. Effect of the addition of calcifying medium (Ca and P) in VSMC WT and TG mice on: A) The calcium content for 8 days. B) The protein expression of Runx2 for 8 days (the values of the densitometry vs GAPDH) are shown by Western Blot. WT: wild mouse; TG: transgenic mouse. * $p < 0.05$ vs control




Figure 5. Effect of addition of calcifying medium (Ca and P) on the levels of reactive oxygen species in mitochondrial VSMC WT and TG mice. UR: relative units DHR123 fluorescence probe. WT: wild mouse; TG: transgenic mouse. * $p < 0.05$ compared to control





Figure 6. Effect of the addition of calcifying medium (Ca and P) in VSMC WT and TG mice on protein expression (relative units) of catalase at 8 days of culture by Western blotting. WT: wild mouse; TG: transgenic mouse



In summary, the overexpression of catalase enzyme reduced the calcification process both *in vivo* and *in vitro*, showing *in vivo* that this decline was accompanied by an improvement in bone parameters studied.

Declaration of interest: The authors declare no conflicts of interest.

Acknowledgements: This work has been possible thanks to funding obtained by the FEIOMM 2012 scholarship to promote research in the area of MARROW MOLECULAR BIOLOGY. This work has also been partially funded with help from the National R+D+I 2008-2011 State Plan for R & D 2013-2016 Carlos III Health Institute (ISCIII) - European Regional Development Fund PI11/00667 and PI13/00014, Plan for Science, Technology and Innovation 2013-2017 of the Principality of Asturias (GRUPIN14-028), Foundation for the Promotion in Asturias of Applied Scientific Research and Technology (FICYT), Reina Sofía Research Institute nephrology, Renal Foundation Iñigo Alvarez de Toledo, Renal Research Network-REDinREN ISCIII (RD06/0016/1013, RD12/0021/1023 and RD16/0009), by the Metabolic Research Development Company Asturiana.

Bibliography

1. Stocker R, Keaney JJ. Role of oxidative modifications in atherosclerosis. *Physiol Rev.* 2004;84:1381-478.
2. Rajagopalan S, Meng XP, Ramasamy S, Harrison DG, Galis ZS. Reactive oxygen species produced by macrophage-derived foam cells regulate the activity of vascular matrix metalloproteinases in vitro. Implications for atherosclerotic plaque stability. *J Clin Invest.* 1996;98:2572-9.
3. Ushio-Fukai M, Alexander RW, Akers M, Yin Q, Fujio Y, Walsh K, et al. Reactive oxygen species mediate the activation of Akt/Protein kinase B by angiotensin II in vascular smooth muscle cells. *J Biol Chem.* 1999;274:22699-704.
4. Uemura S, Matsushita H, Li W, Glassford AJ, Asagami T, Lee KH, et al. Diabetes mellitus enhances vascular matrix metalloproteinase activity: role of oxidative stress. *Circ Res.* 2001;88:1291-8.
5. Zmijewski JW, Moellering DR, Le Goffe C, Landar A, Ramachandran A, Darley-Usmar VM. Oxidized LDL induces mitochondrially associated reactive oxygen/nitrogen species formation in endothelial cells. *Am J Physiol Heart Circ Physiol.* 2005;285:252-61.
6. Sutra T, Morena M, Bargnoux AS, Caporiccio B, Canaud B, Cristol JP. Superoxide production: a calcifying cell signalling event in osteoblastic differentiation of vascular smooth muscle cells exposed to calcification media. *Free Radic Res.* 2008;42:789-97.
7. Mody N, Parhami F, Sarafian TA, Demer LL. Oxidative stress modulates osteoblastic differentiation of vascular and bone cells. *Free Radic Biol Med.* 2001;31:509-19.
8. Byon CH, Javed A, Dai Q, Kappes JC, Clemens TL, Darley-Usmar VM, et al. Oxidative stress induces vascular calcification through modulation of the osteogenic transcription factor Runx2 by AKT signalling. *J Biol Chem.* 2008;283:15319-27.
9. Wang L, Zhang YG, Wang XM, Ma LF, Zhang YM. Narigin products human adipose-derived mesenchymal stem cells against hydrogen peroxide-induced inhibition of osteogenic differentiation. *Chem Biol Interact.* 2015;242:255-61.
10. Roman-Garcia P, Carrillo-Lopez N, Fernandez-Martin JL, Naves-Diaz M, Ruiz-Torres MP, Cannata-Andia JB. High phosphorus diet induces vascular calcification, a related decrease in bone mass and changes in the aortic gene expression. *Bone.* 2010;46:121-8.
11. Fang Y, Ginsberg C, Sugatani T, Monier-Faugere MC, Malluche H, Hruska KA. Early chronic kidney disease-mineral bone disorder stimulates vascular calcification. *Kidney Int.* 2014;85:142-50.
12. Jimbo R, Kawakami-Mori F, Mu S, Hirohama D, Majtan B, Shimizu Y, et al. Fibroblast growth factor 23 accelerates phosphate-induced vascular calcification in the absence of Klotho deficiency. *Kidney Int.* 2014;85:1103-11.
13. Gutiérrez OM, Mannstadt M, Isakova T, Rauh-Hain JA, Tamez H, Shah A, et al. Fibroblast growth factor 23 and mortality among patients undergoing hemodialysis. *N Engl J Med.* 2008;359:584-92.
14. Shimada T, Hasegawa H, Yamazaki Y, Muto T, Hino R, Takeuchi Y, et al. FGF-23 is a potent regulator of vitamin D metabolism and phosphate homeostasis. *J Bone Miner Res.* 2004;19:429-35.
15. Liu S, Tang W, Zhou J, Stubbs JR, Luo Q, Pi M, et al. Fibroblast growth factor 23 is a counter-regulatory phosphaturic hormone for vitamin D. *J Am Soc Nephrol.* 2006;17:1305-15.
16. Dobnig H, Pilz S, Scharnagl H, Renner W, Seelhorst U, Wellnitz B, et al. Independent association of low serum 25-hydroxyvitamin D and 1, 25-dihydroxyvitamin D levels with all-cause and cardiovascular mortality. *Arch Intern Med.* 2008;168:1340-9.
17. Naves-Díaz M, Cabezas-Rodríguez I, Barrio-Vázquez S, Fernández E, Díaz-López JB, Cannata-Andía JB. Low calcidiol levels and risk of progression of aortic calcification. *Osteoporos Int.* 2012;23:1177-82.
18. Lau WL, Linnes M, Chu EY, Foster BL, Bartley BA,

- Somerman MJ, et al. High phosphate feeding promotes mineral and bone abnormalities in mice with chronic kidney disease. *Nephrol Dial Transplant*. 2013;28:62-9.
19. Wang H, Yoshiko Y, Yamamoto R, Minamizaki T, Kozai K, Tanne K, et al. Overexpression of fibroblast growth factor 23 suppresses osteoblast differentiation and matrix mineralization in vitro. *J Bone Miner Res*. 2008;23:939-48.
20. Carrillo-López N, Panizo S, Alonso-Montes C, Román-García P, Rodríguez I, Martínez-Salgado C, et al. Direct inhibition of osteoblastic Wnt pathway by fibroblast growth factor 23 contributes to bone loss in chronic kidney disease. *Kidney Int*. 2016;90:77-89.
21. Cozzolino M, Brenna I, Ciceri P, Volpi E, Cusi D, Brancaccio D. Vascular calcification in chronic kidney disease: a changing scenario. *J Nephrol*. 2011;24(Suppl 18):S3-S10.