



Revista de Osteoporosis y Metabolismo Mineral

ISSN: 1889-836X

ISSN: 2173-2345

Sociedad Española de Investigaciones Óseas y
Metabolismo Mineral

Roca Ruiz, LJ; González López, MC

Valoración de la densidad de masa ósea en la indicación quirúrgica. Nueva herramienta

Revista de Osteoporosis y Metabolismo Mineral, vol. 12, núm. 1, 2020, pp. 32-33

Sociedad Española de Investigaciones Óseas y Metabolismo Mineral

DOI: <https://doi.org/10.4321/S1889-836X2020000100006>

Disponible en: <https://www.redalyc.org/articulo.oa?id=360963260006>

- Cómo citar el artículo
- Número completo
- Más información del artículo
- Página de la revista en redalyc.org

UAEH
redalyc.org

Sistema de Información Científica Redalyc

Red de Revistas Científicas de América Latina y el Caribe, España y Portugal

Proyecto académico sin fines de lucro, desarrollado bajo la iniciativa de acceso
abierto

Assessment of bone mass density in the surgical indication. New tool

DOI: <http://dx.doi.org/10.4321/S1889-836X2020000100006>

Roca Ruiz LJ¹, González López MC²

1 Shoulder Unit

2 Traumatology Unit

Orthopedic and Traumatology Surgery. Virgen Macarena. University Hospital. Seville (Spain)

The proximal humerus fracture represents 5 to 8% of all fractures and is twice as frequent in women as in men. These fractures occur mainly in patients with bone fragility. They are among the most frequent along with hip and distal radius fractures in patients older than 65 years¹⁻⁴, thus presenting a multidisciplinary challenge. Since proximal humerus fractures have been considered fragility fractures, the role of general and local bone mineral density is increasingly gaining attention in the literature⁵⁻⁸.

The influence of local bone mineral density on the functional outcome of the treatment of proximal humerus fractures is controversial. Classically, it has not been sufficiently addressed in the literature. However, the most recent studies show that osteoporosis can negatively affect surgical treatment and subsequent consolidation of fractures of the proximal humerus. That is why bone quality should be part of the preoperative evaluation^{6,9}.

Barnett and Nordin first reported the determination of cortical thickness as a predictor of skeletal mineralization in 1960¹⁰. Since then, measurements of the cortical thickness of the femoral shaft and metacarpals have been widely used to estimate osteoporotic changes in bone. However, cortical thickness of the distal humerus has been shown to be an even more reliable predictor for detecting generalized osteoporosis than that of femoral or metacarpal cortical osteoporosis¹¹.

The use of a simple measurement to determine the bone quality of the proximal humerus could help in making surgical decisions, allowing the indication of the most appropriate technique. For example, it may be possible to predict the safety of screw fixation in bone¹¹.

The Tingart measurement¹¹ is the most frequently used method to measure bone quality in AP x-rays of the shoulder. However, in patients presenting a proximal humerus fracture, the reference points required for the Tingart measurement are often involved in the fracture. In addition, measurement errors must be corrected by x-ray magnification, and there is not always a reference to perform it.

Recently, another index that relates cortical thickness to bone quality is increasing in the literature: the deltoid tuberosity index (DTI). The necessary measurements for it are made immediately above the upper end of the deltoid tuberosity. At that level, the outer cortical edges become parallel; the DTI is equal to the relationship between the external cortical diameter and the internal endosteal diameter. When this ratio is less than 1.4, there will be low bone mineral density in the proximal humerus⁹.

Unlike what happens with the Tingart index, the location of the precise measurements to calculate the DTI are far from the fracture lines. Furthermore, the deltoid tuberosity generally appears well defined in AP x rays, possibly due to the antalgic position that is normally adopted, with the arm in internal rotation⁹.

In their study, Spross et al.⁹ found that the correlation between radiographic measurements and local bone mineral density was strong for the DTI and moderate for the Tingart measurement. Likewise, inter-observer reproducibility was higher in DTI.

Thus, we consider DTI to be a reliable, simple, and applicable tool to assess local bone quality in the proximal humerus. Furthermore, its use has better clinical applicability in patients with proximal humerus fractures than the Tingart index, since sometimes the fracture lines reach the reference points of this measurement.

In this way, Spross et al.¹² have generated a comprehensive algorithm as a treatment guide for FHP, where the demands and biology of the patient are prioritized, being a useful tool for decision-making, achieving a low rate of complications and revisions.

We thus believe that a comprehensive patient assessment, with its different facets, weighing each one in its proper measure, will bring us closer to reality. Hence, considering this global vision of the patient, not limiting ourselves solely and exclusively to the fracture, will make the difference between being good or achieving excellence.



Bibliography

1. Fjalestad T, Iversen P, Hole MØ, Smedsrud M, Madsen JE. Clinical investigation for displaced proximal humeral fractures in the elderly: a randomized study of two surgical treatments: reverse total prosthetic replacement versus angular stable plate Philos (The DELPHI-trial). *BMC Musculoskeletal Disord.* 2014;15:323.
2. Court Brown CM, McQueen MM. The relationship between fractures and increasing age with reference to the proximal humerus. *Curr Orthop.* 2002;16:213-22.
3. Bengner U, Johnell O, Redlund-Johnell I. Changes in the incidence of fracture of the upper end of the humerus during a 30 year period: a study of 2125 fractures. *Clin Orthop.* 1988;231:179-82.
4. Mather J, MacDermid JC, Faber KJ, Athwal GS. Proximal humerus cortical bone thickness correlates with bone mineral density and can clinically rule out osteoporosis. *J Shoulder Elbow Surg.* 2013;22(6):732-8.
5. Jung SW, Shim SB, Kim HM, Lee JH, Lim HS. Factors that influence reduction loss in proximal humerus fracture surgery. *J Orthop Trauma.* 2015;29:276-82.
6. Hertel RW. Fractures of the proximal humerus in osteoporotic bone. *Osteoporos Int.* 2005;16(Suppl. 2):S65-72.
7. Nho SJ, Brophy RH, Barker JU, Cornell CN, MacGillivray JD. Management of proximal humeral fractures based on current literature. *J Bone Joint Surg.* 2007;89:44-58.
8. Mazzucchelli RA, Enny K, Zdravkovic V, Erhardt JB, Jost B, Spross C. The influence of local bone quality on fracture pattern in proximal humerus fractures. *Injury.* 2018;492:359-63.
9. Spross C, Kaestle N, Benninger E, Fornaro J, Erhardt J, Zdravkovic V, et al. Deltoïd Tuberosity Index: a simple radiographic tool to assess local bone quality in proximal humerus fractures. *Clin Orthop Relat Res.* 2015;473:3038-45.
10. Barnett E, Nordin BEC. The radiological diagnosis of osteoporosis: a new approach. *Clin Radiol.* 1960;11:166-9.
11. Tingart MJ, Apreleva M, von Stechow D, Zurakowski D, Warner JJ. The cortical thickness of the proximal humeral diaphysis predicts bone mineral density of the proximal humerus. *J Bone Joint Surg Br.* 2003;85(4):611-7.
12. Spross C, Meester J, Mazzucchelli RA, Puskás GJ, Zdravkovic V, Jost B. Evidence-based algorithm to treat patients with proximal humerus fractures-a prospective study with early clinical and overall performance results. *J Shoulder Elbow Surg.* 2019;28(6):1022-32.