

**Revista Internacional de
Contaminación Ambiental**

Revista Internacional de Contaminación Ambiental

ISSN: 0188-4999

claudio.amescua@atmosfera.unam.mx

Universidad Nacional Autónoma de México

México

Luis-Villaseñor, Irasema E.; Zamudio-Armenta, Olga O.; Voltolina, Domenico; Rochin-Arenas, Jesús A.; Gómez-Gil, Bruno; Audelo-Naranjo, Juan M.; Flores-Higuera, Francisco A.

BACTERIAL COMMUNITIES OF THE OYSTERS *Crassostrea*
cortezensis AND *C. sikamea* OF COSPITA BAY, SINALOA, MEXICO

Revista Internacional de Contaminación Ambiental, vol. 34, no. 2, 2018, May-July, pp. 203-213

Universidad Nacional Autónoma de México

México

DOI: <https://doi.org/10.20937/RICA.2018.34.02.02>

Available in: <https://www.redalyc.org/articulo.oa?id=37056657002>

- How to cite
- Complete issue
- More information about this article
- Journal's webpage in redalyc.org

UNAM
redalyc.org

Scientific Information System Redalyc

Network of Scientific Journals from Latin America and the Caribbean, Spain and
Portugal

Project academic non-profit, developed under the open access initiative

**BACTERIAL COMMUNITIES OF THE OYSTERS *Crassostrea corteziensis*
AND *C. sikamea* OF COSPITA BAY, SINALOA, MEXICO**

Irasema E. LUIS-VILLASEÑOR^{1*}, Olga O. ZAMUDIO-ARMENTA¹, Domenico VOLTOLINA²,
Jesús A. ROCHIN-ARENAS¹, Bruno GÓMEZ-GIL³, Juan M. AUDELO-NARANJO¹ y
Francisco A. FLORES-HIGUERA¹

¹ Universidad Autónoma de Sinaloa, Facultad de Ciencias del Mar, Paseo Claussen s/n, Mazatlán, Sinaloa, México

² Centro de Investigaciones Biológicas del Noroeste, Apartado 1132, Mazatlán, Sinaloa, México

³ Centro de Investigación en Alimentación y Desarrollo, Unidad Mazatlán, Apartado 711, Mazatlán, Sinaloa, México

*Author for correspondence; irasemaluis@uas.edu.mx

(Received December 2016; accepted September 2017)

Key words: bacteria, cultured oysters, wild oysters

ABSTRACT

This work aimed to quantify the bacterial loads and determine the taxonomic composition of the microbial communities of oysters *Crassostrea corteziensis* and *C. sikamea*, as well as the surrounding waters of Cospita Bay, using two sets of samples (M1 and M4) obtained in the dry season and two (M2 and M3) in the rainy season. Bacterial loads were quantified with the viable count technique and out of the 320 strains isolated for their different morphologies, 84 were identified with molecular techniques to the species level. These species pertained to the families Vibrionaceae, Bacillaceae, Brucellaceae, Micrococcaceae, Pseudoalteromonaceae, Rhodobacteraceae, Shewanellaceae and Staphylococcaceae. The higher concentrations of total bacteria were in the rainy season (samples M2 and M3), when families Vibrionaceae and Brucellaceae dominated. The highest *Vibrio* spp. and *Pseudomonas* spp. concentrations were in winter (samples M4). Probably because of its poor health conditions, the Kumamoto oyster *C. sikamea* had the highest diversity of bacterial species and the highest number of known pathogens for aquatic organisms.

Palabras clave: bacterias, ostiones cultivados, ostiones silvestres

RESUMEN

En la presente investigación se cuantificó la carga bacteriana y se determinó la composición de la microbiota de los ostiones *Crassostrea corteziensis* y *C. sikamea*, así como del agua circundante de Bahía Cospita, usando dos muestras obtenidas durante el periodo de estiaje (muestras M1 y M4) y dos obtenidas en el periodo de lluvias (muestras M2 y M3). La carga bacteriana se cuantificó con la técnica de cuenta viable y se aislaron con criterios morfológicos 320 cepas, de las cuales de 84 se identificaron con técnicas moleculares hasta el nivel de especie y fueron agrupadas en las familias Vibrionaceae, Bacillaceae, Brucellaceae, Micrococcaceae,

Pseudoalteromonaceae, Rhodobactereceae, Shewanellaceae y Staphylococcaceae. La mayor concentración de bacterias totales se presentó en la temporada de lluvias (muestreos M2 y M3), predominando las familias Vibrionaceae y Brucellaceae. La mayor concentración de *Vibrio* spp. y *Pseudomonas* spp. se presentó en invierno (muestreo M4). El ostión Kumamoto (*C. sikamea*) presentó una mayor diversidad de especies y el mayor número de bacterias patógenas para organismos acuáticos, que es un probable reflejo de su estado de salud deficiente.

INTRODUCTION

The yearly growth rate of bivalve culture in Mexico is close to 1.2 % (CONAPESCA 2013), and has been limited by frequent mortalities caused by infectious diseases of different origin, such as the phenomenon known as summer mortality syndrome. This has been associated with the presence and virulence of pathogenic bacteria, or of herpesvirus OsHV-1 (Samain and McCombie 2008, Cáceres-Martínez and Vázquez-Yeomans 2013, Petton et al. 2015). Most of these diseases are related to factors that favor pathogens growth, such as excessive densities or poor feeding conditions (Trabal-Fernández et al. 2014).

Other factors generally associated to mortality outbreaks are high temperatures, low oxygen availability or excessive energetic expenditures, mainly associated to spawning (Cheney et al. 2000, Li et al. 2007, de Decker and Saulnier 2011). There are several studies on the composition and seasonal variations of the bacterial communities associated to bivalve molluscs (Llanos et al. 2002, Romanenko et al. 2008). Some of these studies stress the role of several species of the genus *Vibrio*, which may affect larvae, juveniles and adults of several aquatic organisms, including oysters, mussels, clams and scallops (Paillard et al. 2004, Beaz-Hidalgo et al. 2010).

In particular, *V. splendidus* and *V. aestuarianus* have been associated to the summer mortalities that affect oyster production worldwide (Gay et al. 2004, Garnier et al. 2007, 2008, Labreuche et al. 2010). These and several other *Vibrio* species, in particular *V. parahaemolyticus*, are also important vectors of food-borne diseases for marine shellfish consumers (Daniels et al. 2000, McLaughlin et al. 2005, García-Lázaro et al. 2010, Kokashvili et al. 2015).

This indicates the importance of gaining better knowledge on the composition and seasonal variations of the bacterial communities associated to cultured and wild populations of commercially important bivalves.

The cultivable *Vibrio* flora harbored in healthy oysters is a bivalve-specific community with densities ranging from 10^2 to 10^4 colony forming units (CFU)/

mg of tissues, and higher loads may be expected at higher temperatures (Saulnier et al. 2010). However, if bacterial concentrations are too high, they may induce negative stress, with consequent high mortalities and low condition indices (Harekrishna et al. 2014). Additionally there is a general lack of information on the structure and persistence of the bacterial community present in wild and cultivated oysters of commercial interest, such as *C. corteziensis* and *C. sikamea*.

This study aimed to determine the concentration and taxonomy of culturable heterotrophic bacteria present in the oysters *Crassostrea corteziensis* and *C. sikamea* cultured in Cospita Bay, Sinaloa, NW Mexico, and of the bacterioflora suspended in the surrounding water.

MATERIALS AND METHODS

Cospita Bay lies within coordinates 24° 04' 30" to 24° 05' 65" N and 107° 08' 42" to 107° 08' 06" W. Samples were obtained at quarterly intervals from May 2014 to January 2015. Two sets were from the dry period (M1: dry-hot, May 7th 2014; and M4: dry-cold, January 7th, 2015), and two from the warm-rainy season (M2 and M3, August 29th and October 1st, 2014). Samples of *C. corteziensis* and *C. sikamea* (30 of each species) were donated by a local commercial oyster farm. Wild *C. corteziensis* (30, from roots of different mangroves) and several water samples (mixed in equal parts to give one-L composite sample) were also obtained in sterile containers from areas surrounding each sampling point. Surface salinity, temperature and oxygen concentration were determined in situ in the central area of the oyster farm using a refractometer and an YSI 55 dissolved oxygen meter.

Oyster and water samples were transported in ice boxes (< 7 °C) to the laboratory, where 100 µL of the original water sample and of its 1:10 dilution were spread plated in triplicate on Zobell medium for total culturable marine bacteria (2216), on cetrimide

Pseudomonas medium and on 2.5 % NaCl-added TCBS for *Vibrio*-like bacteria detection. DIFCO media were used for all isolation and purification work.

Oysters were thoroughly washed and brushed under running tap water, rinsed with sterile distilled water (DW) and each sample of 30 oysters was divided into five groups of six oysters chosen at random. The oysters of each group were shucked and the liquor and soft parts were homogenized for 90 s in 450 mL of 3 % saline solution using a sterile food blender. The resulting suspension was used to prepare four serial dilutions (10^{-2} to 10^{-5}), which were spread-plated (100 μ L) in the same growth media used for water samples. All plates were read after 48 h at 30 ± 2 °C and the results served to calculate bacteria concentrations in seawater (CFU/mL) and in the oysters' soft tissues (CFU/g, wet weight).

Colonies with different morphologies, spread-plated on the respective growth medium, were purified with the cross streak technique. Five-six colonies were preserved in 1.5 mL Eppendorf vials with 1 mL of 96 % ethanol, and used later for their identification using molecular techniques. The original purified colonies were considered pure strains, assigned an identification key and preserved in glycerol at -80 °C in the laboratory's bacterial cultures collection.

For species identification, the DNA of each ethanol-preserved strain was extracted with the Wizard® Genomic DNA Purification Kit (lot A1120, Promega). After adjusting concentration to 50 ng/ μ L, the ribosomal 16S gene (16S ARNr) was amplified using end-point polymerase chain reaction (PCR) and universal primers Forward 27f.1 (AGR GTT TGA TCM TGG CTC AG) and Reverse 1492R2 (GGT TAC CTT GTT ACG ACT T). The amplification program was: 16S: 94 °C/2' \rightarrow 35 cycles (94 °C/1' \rightarrow 56 °C/1' \rightarrow 72 °C/1') \rightarrow 72 °C/5' \rightarrow 4 °C/ ∞ .

The products obtained were sequenced at Macro-gen (Republic of Korea), and the sequences obtained

were used for species identification with the public databases BLAST (Altschul 1997) and EzTaxon (Chun et al. 2007).

The sequences obtained in FASTA format were aligned with program ClustalW (Thompson et al. 1994). The alignments were exported to version 5.0 of the program MEGA and in order to obtain the molecular phylogenetic tree of each strain (Tamura et al. 2011). Using the neighbor-joining test (NJ) (Saitou and Nei 1987) with the p-distance method and 500 bootstrap repetitions was used to that end, considering transitions and transversions. The bacterium *Klebsiella pneumonia* served as outgroup.

All bacterial concentrations, transformed to the respective R1 rank values (Conover 2012), were normal and homoscedastic (Kolmogorov-Smirnov and Bartlett's tests). Therefore, the mean concentrations of the bacteria present in the water samples, determined with each of the three growth media in the four sampling dates, were compared using one-way ANOVA tests, separating the different means with Tuckey's tests. Those of the bacteria present in wild and cultured oysters were compared with two-ways ANOVA and Tuckey's tests. All statistical tests were performed with SigmaPlot 11.0 software. In all cases, the significance level was $\alpha = 0.05$ (Zar 1996).

RESULTS

Temperature and salinity values ranged from 32.3 to 23.2 °C and 35.0 to 23.2 ppt, respectively. In both cases, differences were significant. Oxygen concentrations were significantly higher in the first two sampling dates than in the remaining samplings (8.09 and 7.46, compared to 4.87 and 5.35 mg/L). In all cases differences were not clearly related to climatic events (**Table I**).

TABLE I. MEAN (\pm SD) TOTAL MARINE AND *Vibrio*-LIKE BACTERIA CONCENTRATIONS IN 10^3 COLONY FORMING UNITS/mL (CFU/mL), AND MEAN TEMPERATURE, SALINITY AND DISSOLVED OXYGEN OF THE WATER SAMPLES OBTAINED IN COSPITA BAY ON THE FOUR SAMPLING DATES

Sampling	Temp. (°C)	Salinity (ppt)	O ₂ (mg/L)	CFU/mL	
				Total*	<i>Vibrio</i>
(M1) Dry-warm	27.5 \pm 0.3 ^{ab}	35.0 \pm 1.0 ^a	8.09 \pm 1.8 ^a	0.87 \pm 0.25 ^a	0.31 \pm 0.18 ^{ab}
(M2) Wet-warm	32.3 \pm 0.2 ^a	23.2 \pm 1.5 ^b	7.46 \pm 2.1 ^a	3.92 \pm 2.87 ^a	0.13 \pm 0.12 ^b
(M3) Wet-warm	28.2 \pm 0.5 ^{ab}	33.5 \pm 0.7 ^a	4.87 \pm 0.9 ^b	2.36 \pm 1.75 ^a	0.14 \pm 0.13 ^{ab}
(M4) Dry-cold	23.2 \pm 0.3 ^b	33.3 \pm 0.9 ^a	5.35 \pm 1.1 ^b	8.77 \pm 0.10 ^a	0.87 \pm 0.35 ^a

* Non parametric test. Different letters indicate significant differences between data in the same column ($p = 0.05$, $a \geq ab \geq b$ and $a > b$)

The concentrations of culturable marine bacteria ranged from 875 to 3920 CFU/mL, and there were no differences between sampling dates. Those of *Vibrio*-like bacteria varied from 127.50 to 875 CFU/mL. These were significantly different, and were determined in M2 and M4 samples, respectively (**Table I**). The cetrimide medium gave negative results in all water samples.

The concentrations of total marine bacteria tended to be higher in the samples of the rainy than in the dry season, although with overlapping values in dry-cold samples of *C. corteziensis* and dry-warm samples of *C. sikamea*, respectively (**Table II**). The tendency was similar in the case of *Vibrio*-like bacteria. For *C. corteziensis* differences between rainy and dry seasons were significant in all cases, while for *C. sikamea* the lowest and highest values were observed in the samples of the dry-warm and dry-cold season, respectively (**Table II**).

Pseudomonas showed the same species-related trends. The lowest and highest mean annual values were recorded in the dry-warm and dry-cold samples in *C. sikamea*, while in *C. corteziensis* mean concentrations were higher in the rainy than in the dry season samples (**Table II**).

In the available samples of the dry period (M1 and M4), bacterial loads were significantly lower in cultured than in wild oysters. In these, marine and *Vibrio*-like bacteria were significantly higher in the dry-cold, rather than in the dry-warm samples. In cultured oysters, there were no differences between the mean concentrations of *Pseudomonas* and marine bacteria in samples obtained in the warm or in the cold period while that of *Vibrio*-like bacteria was significantly higher in the dry-cold samples (**Table III**).

TABLE III. MEAN (\pm SD) BACTERIAL CONCENTRATIONS (10^3 CFU/g) OF CULTURED AND WILD *Crassostrea corteziensis* OF COSPITA BAY IN THE SAMPLES OF THE DRY SEASON M1 (WARM) AND M4 (COLD)

	Cultured	Wild
<i>Marine bacteria</i>		
M1 Dry-warm	1.86 ± 0.62^c	34.20 ± 18.84^b
M4 Dry-cold	3.82 ± 0.50^c	1416.66 ± 381.88^a
<i>Vibrio</i>		
M1 Dry-warm	0.08 ± 0.08^d	1.05 ± 0.74^b
M4 Dry-cold	0.51 ± 0.24^c	40.99 ± 0.00^a
<i>Pseudomonas</i>		
M1 Dry-warm	0.04 ± 0.02^b	0.57 ± 0.16^a
M4 Dry-cold	0.06 ± 0.06^b	2.88 ± 3.72^a

Different letters indicate significant differences ($p = 0.05$, $a > b > c > d$)

CFU: colony forming units

Bacteria identification

The number of strains isolated for their morphologies was notably lower in the dry (cold + warm) (M1 + M4 = 95) than in the rainy season (M2 + M3 = 225) samplings. Among these, molecular techniques allowed identification of 84 sequences with 90-100 % similarity to those of taxonomic entities available in public databases (**Fig. 1**).

The families identified in M2 and M3 samplings in both oyster species were Vibrionaceae, Bacillaceae, Brucellaceae, Micrococcaceae, Pseudoalteromonaceae, Rhodobactereaceae and Staphylococcaceae, and the most frequent was Vibrionaceae (76 and 71 % of all sequences observed in M2 and M3,

TABLE II. MEAN (\pm SD) BACTERIAL CONCENTRATIONS (IN 10^3 CFU/mL) OF *Crassostrea corteziensis* AND *C. sikamea* DETERMINED ON THE FOUR SAMPLING DATES M1 TO M4 IN COSPITA BAY

	M1	M2	M3	M4
<i>Marine bacteria</i>				
<i>C. corteziensis</i>	1.86 ± 0.62^b	7.20 ± 6.88^{ab}	31.88 ± 17.60^a	3.82 ± 0.50^b
<i>C. sikamea</i>	0.71 ± 0.36^b	20.60 ± 12.86^a	18.93 ± 14.11^a	9.99 ± 1.00^{ab}
<i>Vibrio</i>				
<i>C. corteziensis</i>	0.15 ± 0.05^c	4.15 ± 3.27^b	12.35 ± 11.45^b	0.51 ± 0.24^c
<i>C. sikamea</i>	0.06 ± 0.03^c	14.88 ± 11.44^b	13.86 ± 9.03^{ab}	27.33 ± 2.71^a
<i>Pseudomonas</i>				
<i>C. corteziensis</i>	0.06 ± 0.01^{bc}	0.30 ± 0.15^b	0.07 ± 0.01^b	0.03 ± 0.02^c
<i>C. sikamea</i>	0.02 ± 0.01^c	0.38 ± 0.20^b	0.18 ± 0.14^{abc}	2.28 ± 0.23^a

Different letters indicate significant differences between data in the same row ($p = 0.05$, $a > b$)

CFU: colony forming units




Fig. 1. DNA 16S-based phylogenetic tree of the 84 bacterial strains isolated in Cospita Bay oyster and water samples

respectively), followed by Bacillaceae (8.16 %) and Brucellaceae (25.71 %) in M2 and M3, respectively (Fig. 2).

The number of bacteria species observed exclusively in samples of one oyster species was higher for *C. sikamea* (seven: *Vibrio neptunius*, *V. harveyi*, *V. diabolicus*, *V. parahaemolyticus*, *V. ponticus*, *V. alfacensis* and *Bacillus firmus*), than for *C. corteziensis* (four: *Vibrio sinaloensis*, *Bacillus* sp., *B. clausii* and *Micrococcus yunnanensis*). Seven species (families Xanthomonadaceae, Rhodobacteraceae, Staphylococcaceae, Vibrionaceae and Bacillaceae) were found only in water samples (*Lysobacter spongiicola*, *Donghicola* sp., *D. eburneus*, *Bacillus* sp., *Staphylococcus cohnii*, *S. warnei* and *V. hepatarius*) (Table IV).

DISCUSSION

Silva-Neta et al. (2015) detected higher bacterial concentrations at the end of the dry season, which corresponds to our M1 sampling, while the general tendency to higher bacteria concentrations in oyster tissues in the rainy season, and the correspondingly lower values observed in *C. corteziensis* during the dry season, seem to agree with the direct relationship with water temperature described by Deepanjali et al. (2005) and Aagesen and Häse (2014) for *V. parahaemolyticus* concentrations, or for aerobic heterotrophs and *Vibrio*-like bacteria determined by Fay et al. (2012).

When they were collected, concentrations of heterotrophic bacteria were higher in wild than in




Fig. 2. Bacterial families identified in wild and cultured oysters and in Cospita Bay water samples during the wet-warm season (M2 and M3 samples)

cultured oysters, possibly because of the high levels of organic matter and bacterial concentrations common in their environment (Mahasneh 2001). An additional explanation is the stress caused by tide-dependent exposures (Bishop and Peterson 2006, Farcy et al. 2009), which might have caused an increase in the internal bacterial load. The absence of wild organisms during the rainy period was due to mortalities related to high energy expenditure caused by early summer spawning (Frías-Espericueta et al. 1997) or to stress induced by high temperatures and low salinities. Additionally, summer conditions seem to favor the parasite *Perkinsus marinus*, whose presence and high incidence was detected in the area object of this study (Cáceres-Martínez et al. 2012, Cáceres-Martínez and Vázquez-Yeomans 2013).

Summer mortalities with similar etiologies have been documented in different geographic areas for several oyster species such as *C. hongkongensis* (Wang et al. 2016), *C. gigas* (Li et al. 2007), *C. virginica* (Soniat et al. 2006, Heilmayer et al. 2008) and *C. ariakensis* (Kelly et al. 2011). In all these cases, increasing temperatures would seem to be the driving factor, either causing an increase of energy losses or favoring bacterial or parasite establishment and growth.

Although its presence and abundance should be confirmed with long-term regular observations, the dominance of Vibrionaceae in both oyster species in the warm rainy period coincides with the indication that *Vibrio* spp. are the most abundant bacteria in oysters during summer months (Pujalte et al. 1999, Wang et al. 2016). Among the species of *Vibrio* detected in *C. sikamea*, some are known pathogens for invertebrates (*V. parahaemolyticus*, *V. hepatarius*, *V. brasiliensis*, *V. neptunius* and *V. owensii* [Ruangan and Kitao 1991, Thompson et al. 2003, Cano-Gómez et al. 2010, Rivera-Posada et al. 2011, Gómez-Gil et al. 2012]), while others, such as *V. xuii*, *V. ponticus*, *V. alfacensis* and *V. harveyi* may affect vertebrates (Pujalte et al. 2003, Thompson et al. 2003, Xie et al. 2007). Among these pathogens, *V. neptunius* was isolated in cultures of diseased *Nodipecten nodosus* larvae (Thompson et al. 2003), and it may cause high mortalities in mussel larvae (Kesarodi-Watson et al. 2009) and in oyster hatcheries (Prado et al. 2005).

Known pathogens for molluscs were not detected in *C. corteziensis*. However, the bacteria *V. sinaloensis*, *Micrococcus yunnanensis*, *Bacillus clausii* and *V. natriegens*, which were detected only in this oyster, are known pathogens of *Lutjanus guttatus*, *Litopenaeus vannamei* and *Acanthaster planci* (Gómez-Gil et al. 2008, Flores-Miranda et al. 2011, Rivera-Posada et al. 2011).

Of all known pathogenic *Vibrio* of major medical interest listed by Daniels and Shafaie (2000), only *V. parahaemolyticus* was present in our oyster samples. Among those isolated from surface waters, the coagulase-negative *Staphylococcus warneri* and *S. cohnii* are of mild clinical interest as possible sources of nosocomial infections (Martínez and Máttar 2006, Fariña et al. 2013, Soldera et al. 2013), while *Lysobacter spongiicola* pertains to a genus of potential importance for its production of several extracellular enzymes (de Bruijn et al. 2015), and because several *Lysobacter* species have been suggested for biological plant disease control (Hashizume et al. 2004, Li et al. 2008).

ACKNOWLEDGMENTS

Supported by projects Programa de Fomento y Apoyo a Proyectos de Investigación, PROFAPI 2013/212 and Consejo Nacional de Ciencia y Tecnología, Apoyos a Infraestructura Científica, INFR-2015/251448. D.M. Cruz-Zamudio and G. Solano-Cota helped with the field and analytical work.

TABLE IV. BACTERIAL STRAINS (M) ISOLATED FROM *Crassostrea corteziensis*, *C. sikamea* AND COSPITA BAY WATER SAMPLES

Strain	Identification	<i>C. corteziensis</i>	<i>C. sikamea</i>	water
182	<i>Vibrio</i> sp.	M2		
183	<i>Vibrio</i> sp.	M2		
187	<i>Vibrio mediterranei</i>		M2	
190	<i>Vibrio</i> sp.	M2		
192	<i>Vibrio mediterranei</i>		M2	
195	<i>Vibrio sinaloensis</i>	M2		
196	<i>Vibrio neptunius</i>		M2	
197	<i>Vibrio harveyi</i>		M2	
200	<i>Vibrio brasiliensis</i>	M2		
201	<i>Vibrio</i> sp.		M2	
203	<i>Vibrio hepatarius</i>	M2		
206	<i>Vibrio rotiferianus</i>		M2	
207	<i>Vibrio rotiferianus</i>	M2		
208	<i>P. flavipulchra</i>	M2		
209	<i>Lysobacter spongiicola</i>			M2
210	<i>Donghicola</i> sp.			M2
211	<i>Bacillus firmus</i>		M2	
212	<i>Vibrio owensii</i>		M2	
213	<i>Vibrio hepatarius</i>		M2	
214	<i>Bacillus</i> sp.			M2
218	<i>Vibrio diabolicus</i>		M2	
220	<i>Vibrio owensii</i>	M2		
222	<i>Donghicola eburneus</i>			M2
225	<i>Bacillus</i> sp.	M2		
226	<i>Vibrio harveyi</i>		M2	
227	<i>P. flavipulchra</i>	M2		
228	<i>Vibrio brasiliensis</i>	M2		
229	<i>Vibrio</i> sp.	M2		
230	<i>Vibrio communis</i>	M2		
231	<i>Vibrio brasiliensis</i>	M2		
241	<i>Vibrio harveyi</i>		M2	
242	<i>Vibrio parahaemolyticus</i>		M2	
245	<i>Vibrio communis</i>		M2	
247	<i>Staphylococcus cohnii</i>			M2
248	<i>Vibrio hepatarius</i>			M2
250	<i>Vibrio owensii</i>	M2		
263	<i>Micrococcus yunnanensis</i>	M2		
267	<i>Bacillus clausii</i>	M2		
300	<i>Vibrio rotiferianus</i>	M2		
302	<i>Vibrio rotiferianus</i>		M2	
303	<i>Vibrio rotiferianus</i>		M2	
305	<i>Vibrio rotiferianus</i>	M2		
311	<i>Vibrio rotiferianus</i>			M2
315	<i>Vibrio ponticus</i>		M2	
318	<i>Vibrio rotiferianus</i>	M2		
322	<i>Vibrio communis</i>	M2		
327	<i>Vibrio rotiferianus</i>	M2		
328	<i>Vibrio rotiferianus</i>	M2		
330	<i>Staphylococcus warneri</i>			M2
337	<i>Vibrio rotiferianus</i>		M3	
341	<i>Vibrio alfacensis</i>		M3	
350	<i>Vibrio rotiferianus</i>	M3		
351	<i>P. flavipulchra</i>	M3		
353	<i>Vibrio mediterranei</i>		M3	
359	<i>Vibrio brasiliensis</i>		M3	

TABLE IV. BACTERIAL STRAINS (M) ISOLATED FROM *Crassostrea corteziensis*, *C. sikamea* AND COSPITA BAY WATER SAMPLES

Strain	Identification	<i>C. corteziensis</i>	<i>C. sikamea</i>	water
360	<i>Vibrio mediterranei</i>	M3		
372	<i>Vibrio rotiferianus</i>	M3		
391	<i>Vibrio mediterranei</i>	M3		
394	<i>Vibrio mediterranei</i>		M3	
401	<i>Vibrio brasiliensis</i>		M3	
407	<i>Vibrio communis</i>	M3		
410	<i>Vibrio</i> sp.		M3	
414	<i>Vibrio owensii</i>		M3	
427	<i>Vibrio communis</i>		M3	
437	<i>Vibrio xuii</i>		M3	
457	<i>Vibrio natriegens</i>	M3		
467	<i>Vibrio communis</i>	M3		
468	<i>Ochrobactrum cytisi</i>	M3		
473	<i>Ochrobactrum cytisi</i>	M3		
474	<i>Ochrobactrum cytisi</i>	M3		
475	<i>Vibrio owensii</i>	M3		
479	<i>Vibrio rotiferianus</i>	M3		
481	<i>Vibrio diabolicus</i>		M3	
494	<i>Ochrobactrum cytisi</i>	M3		
500	<i>Vibrio natriegens</i>	M3		
507	<i>Ochrobactrum cytisi</i>		M3	
508	<i>Vibrio diabolicus</i>		M3	
514	<i>Vibrio rotiferianus</i>	M3		
522	<i>Ochrobactrum cytisi</i>		M3	
533	<i>Ochrobactrum cytisi</i>	M3		
561	<i>Vibrio communis</i>	M3		
570	<i>Vibrio owensii</i>		M3	
574	<i>Ochrobactrum cytisi</i>	M3		
576	<i>Ochrobactrum cytisi</i>	M3		

REFERENCES

- Aagesen A.M. and Häse C.C. (2014). Seasonal effects of heat shock on bacterial populations, including artificial *Vibrio parahaemolyticus* exposure, in the Pacific oyster, *Crassostrea gigas*. Food Microbiol. 38, 93-103. DOI: 10.1016/j.fm.2013.08.008
- Altschul S.F. (1997). Evaluating the statistical significance of multiple distinct local alignments. In: Theoretical and computational methods in genome research (S. Suhai, Ed.). Plenum, New York, USA, pp. 1-14.
- Beaz-Hidalgo R., Balboa S., Romalde J. and Figueras M. (2010). Diversity and pathogenicity of *Vibrio* species in cultured bivalve molluscs. Environ. Microbiol. Rep. 2 (1), 34-43. DOI: 10.1111/j.1758-2229.2010.00135.x
- Bishop N.J. and Peterson C.H. (2006). Direct effects of physical stress can be counteracted by indirect benefits oyster growth on a tidal elevation gradient. Oecologia 147, 426-433. DOI: 10.1007/s00442-005-0273-3
- Cáceres-Martínez J., García-Ortega M., Vásquez-Yeomans R., Pineda-García T.J., Stokes N.A. and Carnegie R.B. (2012). Natural and cultured populations of the mangrove oyster *Saccostrea palmula* from Sinaloa, Mexico, infected by *Perkinsus marinus*. J. Invertebr. Pathol. 110, 321-325. DOI: 10.1016/j.jip.2012.03.019
- Cáceres-Martínez J. and Vásquez-Yeomans R. (2013). Enfermedades, parásitos y episodios de mortalidad de ostiones de importancia comercial en México y sus implicaciones para la producción. Ciencia Pesquera 21, 5-48.
- Cano-Gómez A., Goulden E.F., Owens L. and Høj L. (2010). *Vibrio owensii* sp. nov., isolated from cultured crustaceans in Australia. FEMS Microbiol. Lett. 302 (2), 175-181. DOI: 10.1111/j.1574-6968.2009.01850.x
- Cheney D.P., MacDonald B.F. and Elston R.A. (2000). Summer mortality of Pacific oyster, *Crassostrea gigas* (Thunberg): Initial findings on multiple environmental stressors in Puget Sound, Washington, 1998. J. Shellfish Res. 19, 353-359.
- Chun J., Lee J.H., Jung Y., Kim M., Kim S., Kim B.K. and Lim Y.W. (2007). EzTaxon: A webbased tool for the identification of prokaryotes based on 16S ribosomal

- RNA gene sequences. *Int. J. Syst. Evol. Microbiol.* 57, 2259-2261. DOI: 10.1099/ijs.0.64915-0
- CONAPESCA (2013). Anuario estadístico de acuicultura y pesca 2013. Secretaría de Agricultura, Ganadería, Desarrollo Rural, Pesca y Alimentación, Comisión Nacional de Acuicultura y Pesca. Mazatlán, Mexico, 299 pp.
- Conover W.J. (2012). The rank transformation-an easy and intuitive way to connect many nonparametric methods to their parametric counterparts for seamless teaching introductory statistics courses. *WIREs Comp. Stat.* 4, 432-438. DOI: 10.1002/wics.1216
- Daniels N.A. and Shafaie A. (2000). A review of pathogenic *Vibrio* infections for clinicians. *Infect. Med.* 17 (10), 665-685
- Daniels N.A., MacKinnon L., Bishop R., Altekruze S., Ray B., Hammond R.M., Thompson S., Wilson S., Bean N.H., Griffin P.M. and Slutsker L. (2000). *Vibrio parahaemolyticus* infections in the United States, 1973-1998. *J. Infect. Dis.* 181, 1661-1666. DOI: 10.1086/315459
- De Bruijn I., Cheng X., de Jager V., Gómez Expósito R., Watrous J., Patel N., Postma, J., Dorrestein P.C., Kobayashi D. and Raaijmakers J.M. (2015). Comparative genomics and metabolic profiling of the genus *Lysobacter*. *BMC Genomics* 16, 991. DOI: 10.1186/s12864-015-2191-z
- Deepanjali A., Kumar H.S., Karunasagar I. and Karunasagar I. (2005). Seasonal variation in abundance of total and pathogenic *Vibrio parahaemolyticus* bacteria in oysters along the southwest coast of India. *App. Environ. Microbiol.* 71, 3575-3580. DOI: 10.1128/AEM.71.7.3575-3580.2005
- De Decker S. and Saulnier D. (2011). Vibriosis induced by experimental cohabitation in *Crassostrea gigas*: Evidence of early infection and down-expression of immune related genes. *Fish Shellfish Immunol.* 30, 691-699. DOI: 10.1016/j.fsi.2010.12.017
- Farcy E., Voiseux C., Lebel J.-M. and Fiévet B. (2009). Transcription genes in the Pacific oyster exposed to acute thermal stress. *Cell Stress Chaperones* 14, 371-380. DOI: 10.1007/s12192-008-0091-8
- Fariña N., Carpinelli L., Samudio M., Guillén R., Laspina F., Sanabria R., Abente S., Rodas L., González P. and de Kaspar H.M. (2013). *Staphylococcus* coagulasa-negativa clínicamente significativos. Especies más frecuentes y factores de virulencia. *Rev. Chil. Infectol.* 30, 480-488. DOI: 10.4067/S0716-10182013000500003
- Fay P.J., Richards G.P. and Ozbay G. (2012). Water quality parameters and total aerobic bacterial and Vibrionaceae loads in eastern oysters (*Crassostrea virginica*) from oyster-gardening sites. *Arch. Environ. Contam. Toxicol.* 62, 628-637. DOI: 10.1007/s00244-011-9736-1
- Flores-Miranda M.C., Luna-González A., Campa-Córdova Á.I., González-Ocampo H.A., Fierro-Coronado J.A. and Partida-Arangure B.O. (2011). Microbial immunostimulants reduce mortality in whiteleg shrimp (*Litopenaeus vannamei*) challenged with *Vibrio sinoaloensis* strains. *Aquaculture* 320, 51-55. DOI: 10.1016/j.aquaculture.2011.08.005
- Frias-Espicueta M.G., Páez-Osuna F. and Osuna-López J.I. (1997). Seasonal changes in the gonadal state of the oysters *Crassostrea iridescens* and *Crassostrea corteziensis* (Filibranchia: Ostreidae) in the Northwest coast of Mexico. *Rev. Biol. Trop.* 45, 1061-1065.
- García-Lázaro M., Almodóvar-Pulido M.C., Rivero A. and Torre-Cisneros J. (2010). Cólera y otras infecciones del género *Vibrio*. *Medicine* 11 (56), 3317-3321. DOI: 10.1016/S0304-5412(14)70776-0
- Garnier M., Labreuche Y., García C., Robert M. and Nicolas J.L. (2007). Evidence for the involvement of pathogenic bacteria in summer mortalities of the Pacific oyster *Crassostrea gigas*. *Microb. Ecol.* 53 (2), 187-196. DOI: 10.1007/s00248-006-9061-9
- Garnier M., Labreuche Y. and Nicolas J.L. (2008). Molecular and phenotypic characterization of *Vibrio aestuarianus* subsp. *francensis* subsp. nov., a pathogen of the oyster *Crassostrea gigas*. *Syst. Appl. Microbiol.* 31 (5), 358-365. DOI: 10.1016/j.syapm.2008.06.003
- Gay M., Renault T., Pons A.M. and Le Roux F. (2004). Two *Vibrio splendidus* related strains collaborate to kill *Crassostrea gigas*: taxonomy and host alterations. *Dis. Aquat. Org.* 62, 65-74. DOI: 10.3354/dao062065
- Gómez-Gil B., Fajer-Ávila E., Pascual J., Macián M.C., Pujalte M.J., Garay E. and Roque A. (2008). *Vibrio sinaloensis* sp. nov., isolated from the spotted rose snapper, *Lutjanus guttatus* Steindachner, 1869. *Int. J. Syst. Evol. Microbiol.* 58(7), 1621-1624. DOI: 10.1099/ijs.0.65719-0
- Gómez-Gil B., Roque A., Chimento L., Moreira A.P.B., Lang E. and Thompson F.L. (2012). *Vibrio alfacensis* sp. nov., isolated from marine organisms. *Int. J. Syst. Evol. Microbiol.* 62 (12), 2955-2961. DOI: 10.1099/ijs.0.033191-0
- Harekrishna J., Keshab C., Mondal, Maity C., Das A., Halder S.K. and Mitra A. (2014). Effect of microbial load on the condition index of the edible oyster, *Saccostrea cucullata* in the Sundarbans. *India. Int. J. Aquat. Biol.* 2, 238-245.
- Hashizume H., Hirose S., Sawa R., Muraoka Y., Ikeda D., Naganawa H. and Igarashi M. (2004). Tripropeptins, novel antimicrobial agents produced by *Lysobacter* sp. *J. Antibiot. (Tokyo)* 57, 52-58.
- Heilmayer O., Digialleonardo J., Qian L. and Roesijadi G. (2008). Stress tolerance of a subtropical *Crassostrea*

- virginica* population to the combined effects of temperature and salinity. *Est. Coast. Shelf Sci.* 79, 179-185. DOI: 10.1016/J.ECSS.2008.03.022
- Kelly C.J., Laramore S.E., Scarpa J. and Newell R.I.E. (2011). Seasonal comparison of physiological adaptation and growth of suminoe (*Crassostrea ariakensis*) and eastern (*Crassostrea virginica*) oysters. *J. Shellfish Res.* 30, 737-749. DOI: 10.2983/035.030.0314
- Kesarcodi-Watson A., Kaspar H., Lategan M.J. and Gibson L. (2009). Two pathogens of greenshell mussel larvae, *Perna canaliculus*: *Vibrio splendidus* and a *V. coralliilyticus/neptunius*-like isolate. *J. Fish Dis.* 32 (6), 499-507. DOI: 10.1111/j.1365-2761.2009.01006.x
- Kokashvili T., Whitehouse C.A., Tskhvediani A., Grim C.J., Elbakidze T., Mitaishvili N., Janelidze N., Jaiani E., Haley B.J., Lashkhi N., Huq A., Colwell R.R. and Tediashvili M. (2015). Occurrence and diversity of clinically important *Vibrio* species in the aquatic environment of Georgia. *Front. Pub. Health* 3, 232. DOI: 10.3389/fpubh.2015.00232
- Labreuche Y., Le-Roux F., Henry J., Zatylny C., Huvet A., Lambert C., Soudant P., Mazel D. and Nicolas J.L. (2010). *Vibrio aestuarianus* zinc metalloprotease causes lethality in the Pacific oyster *Crassostrea gigas* and impairs the host cellular immune defenses. *Fish Shellfish Immunol.* 29, 753-758. DOI: 10.1016/j.fsi.2010.07.007
- Li Y., Qin J.G., Abbott C.A., Li X. and Benkendorff K. (2007). Synergistic impacts of heat shock and spawning on the physiology and immune health of *Crassostrea gigas*: An explanation for summer mortality in Pacific oyster. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 293 (6), 2353-2362. DOI: 10.1152/ajpregu.00463.2007
- Li S., Jochum C.C., Yu F., Zaleta-Rivera K., Du L., Harris S.D. and Yuen G.Y. (2008). An antibiotic complex from *Lysobacter enzymogenes* strain C3: Antimicrobial activity and role in plant disease control. *Phytopathology* 98, 695-701. DOI: 10.1094/PHYTO-98-6-0695
- Llanos J., Cid M., Navarro S., Dinamarca A. and García-Tello P. (2002). Atypical bacteria accompanying the scallop *Argopecten purpuratus*. *Invest. Mar.* 30(2), 57-59. DOI: 10.4067/S0717-71782002000200005.
- Mahasneh A.M. (2001). Seasonal incidence of some heterotrophic aerobic marine bacteria in an *Avicennia marina* habitat along a Qatari Coast. *J. Biol. Sci.* 1(7), 666-670. DOI: 10.3923/jbs.2001.666.670
- Martínez P. and Máttar S. (2006). Posible aislamiento clínico de *Staphylococcus cohnii* resistente a vancomicina: reporte de caso. *Infectio* 10 (3), 175-177.
- McLaughlin J.B., DePaola A., Bopp C.A., Martinek K.A., Napolilli N.P., Allison C.G., Murray S.L., Thompson E.C., Bird M.M. and Middaugh J.P. (2005). Outbreak of *Vibrio parahaemolyticus* gastroenteritis associated with Alaskan oysters. *N. Engl. J. Med.* 353, 1463-1470. DOI: 10.1056/NEJMoa051594
- Paillard C., LeRoux F. and Borrego J.J. (2004). Bacterial disease in marine bivalves, a review of recent studies: trends and evolution. *Aquat. Living Res.* 17, 477-498. DOI: 10.1051/alr:2004054
- Petton B., Bruto M., James A., Labreuche Y., Alunno-Bruscia M. and Leroux F. (2015). *Crassostrea gigas* mortality in France: The usual suspect, a herpesvirus, may not be the killer in this polymicrobial opportunistic disease. *Front. Microbiol.* 6, 686. DOI: 10.3389/fmicb.2015.00686
- Prado S., Romalde J.L., Montes J. and Barja J.L. (2005). Pathogenic bacteria isolated from disease outbreaks in shellfish hatcheries. First description of *Vibrio neptunius* as an oyster pathogen. *Dis. Aquat. Org.* 67(3), 209-215. DOI: 10.3354/dao067209
- Pujalte M.J., Ortigosa M., Macián M.C. and Garay E. (1999). Aerobic and facultative anaerobic heterotrophic bacteria associated to Mediterranean oysters and seawater. *Int. Microbiol.* 2, 259-266.
- Pujalte M.J., Sitjà-Bobadilla A., Macián M.C., Belloch C., Alvarez-Pellitero P., Pérez-Sánchez J., Uruburu F. and Garay E. (2003). Virulence and molecular typing of *Vibrio harveyi* strains isolated from cultured dentex, gilthead sea bream and European sea bass. *Syst. Appl. Microbiol.* 26(2), 284-292. DOI: 10.1078/072320203322346146
- Rivera-Posada J.A., Pratchett M., Cano-Gómez A., Arango-Gómez J.D. and Owens L. (2011). Refined identification of *Vibrio* bacterial flora from *Acanthaster planci* based on biochemical profiling and analysis of housekeeping genes. *Dis. Aquat. Org.* 96 (2), 113-123. DOI: 10.3354/dao02389
- Romanenko L.A., Uchino M., Kalinovskaya N.I. and Mikhailova V.V. (2008). Isolation, phylogenetic analysis and screening of marine mollusc-associated bacteria for antimicrobial, hemolytic and surface activities. *Microbiol. Res.* 163, 633-644. DOI: 10.1016/j.micres.2006.10.001
- Ruangpan L. and Kitao T. (1991). *Vibrio* bacteria isolated from black tiger shrimp, *Penaeus monodon* Fabricius. *J. Fish Dis.* 14 (3), 383-388. DOI: 10.1111/j.1365-2761.1991.tb00836.x
- Saitou N. and Nei M. (1987). The neighbor-joining method: A new method for reconstructing phylogenetic trees. *Mol. Biol. Evol.* 4, 406-425.
- Samain, J.F. and McCombie H. (Eds.) (2008). Summer mortality Pacific oyster *Crassostrea gigas*: The Morést Project. Éditions Quae, Versailles, France, 379 pp.
- Saulnier D., Decker S.D., Haffner P., Cobret L., Maeva R., Garcia C. (2010). A large-scale epidemiological study to identify bacteria pathogenic to Pacific oyster

- Crassostrea gigas* and correlation between virulence and metalloprotease-like activity. Microb. Ecol. 59, 787-798. DOI: 10.1007/s00248-009-9620-y
- Silva-Neta M.T., Maciel B.M., Lopes A.T.S., Marques E.L.S., Rezende R.P., and Boehs G. (2015). Microbiological quality and bacterial diversity of the tropical oyster *Crassostrea rhizophorae* in a monitored farming system and from natural stocks. Genet. Mol. Res. 14, 15754-15768. DOI: 10.4238/2015.December.1.27
- Soldera J., Nedel W.L., Cardoso P.R. and d'Azevedo P.A. (2013). Bacteremia due to *Staphylococcus cohnii* sp. *urealyticus* caused by infected pressure ulcer: case report and review of the literature. Sao Paulo Med. J. 131 (1), 59-61. DOI: 10.1590/S1516-31802013000100010
- Soniat T.M., Klinck J.M., Powell E.N. and Hoffmann E.E. (2006). Understanding the success and failure of oyster populations: Climatic cycles and *Perkinsus marinus*. J. Shellfish Res. 25, 83-93. DOI: 10.2983/0730-8000(2006)25[83:utsafo]2.0.co;2
- Tamura K., Peterson D., Peterson N., Stecher G., Nei M., and Kumar S. (2011). MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. Mol. Biol. Evol. 28 (10), 2731-2739. DOI: 10.1093/molbev/msr121
- Thompson F.L., Li Y., Gomez-Gil B., Thompson C.C., Hoste B., Vandemeulebroecke K., Rupp G.S., Pereira A., De Bem M.M., Sorgeloos P. and Swings J. (2003). *Vibrio neptunius* sp. nov., *Vibrio brasiliensis* sp. nov. and *Vibrio xuii* sp. nov., isolated from the marine aquaculture environment (bivalves, fish, rotifers and shrimps). Int. J. Syst. Evol. Microbiol. 53, 245-252. DOI: 10.1099/ijs.0.02447-0
- Thompson J.D., Higgins D.G. and Gibson T.J. (1994). CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. Nucleic Acids Res. 22 (22), 4673-4680.
- Trabal-Fernández N., Mazón-Suástegui J.M., Vázquez-Juárez R., Ascencio-Valle F. and Romero J. (2014). Changes in the composition and diversity of the bacterial microbiota associated with oyster (*Crassostrea corteziensis*, *Crassostrea gigas* and *Crassostrea sikamea*) during commercial production. FEMS Microbiol. Ecol. 88, 69-88. DOI: 10.1111/1574-6941.12270
- Wang R., He J. and Wang J. (2016). Heterotrophic bacterial abundance and diversity in the farming environment and guts of the oyster *Crassostrea hongkongensis*. J. Shellfish Res. 35, 343-350. DOI: 10.2983/035.035.0208
- Xie Z.Y., Hu C.Q., Zhang L.P., Chen C., Ren C.H. and Shen Q. (2007). Identification and pathogenicity of *Vibrio ponticus* affecting cultured Japanese sea bass, *Lateolabrax japonicus* (Cuvier in Cuvier and Valenciennes). Lett. Appl. Microbiol. 45 (1), 62-67. DOI: 10.1111/j.1472-765X.2007.02141.x
- Zar J.H. (1996). Biostatistical analysis. Prentice-Hall, Englewood Cliffs, New Jersey, USA, 620 pp.