



Brazilian Journal of Cardiovascular Surgery

ISSN: 0102-7638

ISSN: 1678-9741

Sociedade Brasileira de Cirurgia Cardiovascular

Li, Zeshu; Liu, Luqi

Patency of Individual and Sequential Coronary Artery Bypass
in Patients with Ischemic Heart Disease: A Meta-analysis

Brazilian Journal of Cardiovascular Surgery, vol. 34, no. 4, 2019, pp. 420-427

Sociedade Brasileira de Cirurgia Cardiovascular

DOI: 10.21470/1678-9741-2018-0284

Available in: <http://www.redalyc.org/articulo.oa?id=398966769009>

- How to cite
- Complete issue
- More information about this article
- Journal's webpage in redalyc.org

UABM
redalyc.org

Scientific Information System Redalyc

Network of Scientific Journals from Latin America and the Caribbean, Spain and
Portugal

Project academic non-profit, developed under the open access initiative

Patency of Individual and Sequential Coronary Artery Bypass in Patients with Ischemic Heart Disease: A Meta-analysis

Zeshu Li^{1,2}, MD; Luqi Liu², MD



DOI: 10.21470/1678-9741-2018-0284

Abstract

Objective: To evaluate the patency of individual and sequential coronary artery bypass in patients with ischemic heart disease.

Methods: We searched PubMed, Cochrane Library, Excerpta Medica Database, and ClinicalTrials.gov databases for controlled trials. Endpoints included graft patency, anastomosis patency, occluded rates in left anterior descending (LAD) system and right coronary artery (RCA) system, in-hospital mortality, and follow-up mortality. Pooled risk ratios (RRs) and standardized mean difference (SMD) were used to assess the relative data.

Results: Nine cohorts, including 7100 patients and 1440 grafts under individual or sequential coronary artery bypass. There were no significant differences between individual and sequential coronary artery bypass in the graft patency (RR=0.96;

95% CI=0.91-1.02; $P=0.16$; $I^2=87\%$), anastomosis patency (RR=0.95; 95% CI=0.91-1.00; $P=0.05$; $I^2=70\%$), occluded rate in LAD system (RR=1.03; 95% CI=0.92-1.16; $P=0.58$; $I^2=37\%$), occluded rate in RCA system (RR=1.36; 95% CI=0.72-2.57; $P=0.35$; $I^2=95\%$), in-hospital mortality (RR=1.57; 95% CI=0.92-2.69; $P=0.10$; $I^2=0\%$), and follow-up mortality (RR=0.96; 95% CI=0.36-2.53; $P=0.93$; $I^2=0\%$).

Conclusion: No significant differences on clinical data were observed regarding anastomosis patency, occluded rate in LAD system, occluded rate in RCA system, in-hospital mortality, and follow-up mortality, indicating that the patency of individual and the patency of sequential coronary artery bypass are similar to each other.

Keywords: Coronary Artery Bypass. Coronary Artery Disease. Myocardial Ischemia. Hospital Mortality. Meta-Analysis as Topic.

Abbreviations, acronyms & symbols

CABG = Coronary artery bypass grafting
 CIs = Confidence intervals
 CI = Control
 CT = Computed tomography
 EMBASE = Excerpta Medica database
 LAD = Left anterior descending
 NOS = Newcastle Ottawa Scale

PDA = Posterior descending artery
 PRISMA = Preferred Reporting Items for Systematic Review and Meta-analysis
 RCA = Right coronary artery
 RRs = Risk ratios
 SMD = Standardized mean difference

INTRODUCTION

Ischemic heart disease is currently the leading cause of death worldwide and will account for 14.2% of all deaths by 2030. Also, it is a major contributor to societal costs of cardiac disease^[1]. Coronary artery bypass grafting (CABG) is one of the common surgeries for cardiac patients, which is the best treatment for advanced ischemic heart disease^[2]. The sequential

grafting technique in CABG was introduced by Flemming et al.^[3] in the 1970s. Since then, different methods of anastomosis such as individual or sequential grafts have been used. However, the efficacy of these methods is controversial.

Our meta-analysis was undertaken to analyze the efficacy of individual and sequential grafts used in patients with ischemic heart disease and under CABG.

¹Department of Thoracic and Cardiovascular Surgery, Shandong Provincial PKUcare Luzhong Hospital, Zibo, Shandong, People's Republic of China.

²Department of Cardiac Surgery, Shandong Provincial Qianfoshan Hospital, Shandong University, Jinan, Shandong, People's Republic of China.

This study was carried out at the Shandong Provincial Qianfoshan Hospital, Shandong University, Jinan, Shandong, People's Republic of China.

Correspondence Address:

Luqi Liu

<https://orcid.org/0000-0002-5185-7608>

Department of Cardiac Surgery, Shandong Provincial Qianfoshan Hospital, Shandong University

No. 16766, Jingshi Road, Jinan, Shandong, People's Republic of China.

Zip Code: 250014

Email: luqizh@yeah.net

Article received on September 14th, 2018.

Article accepted on January 31st, 2019.

METHODS

Using the keywords “coronary artery bypass grafting”, “individual graft”, “sequential graft”, and “ischemic heart disease”, we searched PubMed, Cochrane Library, Excerpta Medica database (EMBASE), and ClinicalTrials.gov databases and got data from inception to February 25, 2018. The search was restricted to studies with humans and had no restrictions in language. In addition, references from randomized trials and relevant reviews that were not identified in the database search were hand-searched.

The following inclusion criteria were applied: (1) patients with ischemic heart disease and under CABG; (2) cohort trials that compared the efficacy of individual and sequential coronary artery bypass; and (3) clinical outcomes reported, such as patency rate, blood flow, and the incidence of death. Reviews, meta-analyses, and observational studies were excluded. The meta-analysis was conducted according to the Preferred Reporting Items for Systematic Review and Meta-analysis (PRISMA) guidelines^[4].

Two investigators independently extracted data from the relevant sources. Authors were contacted when data were incomplete or unclear and conflicts were resolved by discussion. Baseline demographic and quality characteristics (sample size, age, sex, community, and follow-up) of patients were collected from the eligible studies. The patency rate of graft and anastomosis, blood flow, and mortality were recorded. Newcastle Ottawa Scale (NOS) was used to assess the quality of included literatures based on recommendations from a non-randomized methodological level^[5].

Statistical Analysis

Binary classification variables of the clinical endpoints were measured using the risk ratios (RRs) with 95% confidence intervals (CIs). The continuous variables of the clinical endpoints were determined by standardized mean difference (SMD) with 95% CIs. Two-sided P -values < 0.05 were considered statistically significant. Heterogeneity was assessed by the Cochran Q test and I^2 statistic, and Cochran's $P < 0.10$ and $I^2 > 50$ were considered indicative of significant heterogeneity. Pooled analyses were conducted using a fixed effect model, whereas a random effect model was used if there was significant heterogeneity. Publication biases were assessed by funnel plot analysis and Egger's test. Data analysis was conducted using the RevMan 5.3 software (Nordic Cochrane Centre, Cochrane Collaboration, 2013), and sensitivity analysis was performed by the Stata 11.0 software (StataCorp, College Station, Texas, USA).

RESULTS

Data Search Results

We identified nine trials^[6-14] out of 632 records that satisfied our inclusion criteria, as shown in the selection procedure depicted in Figure 1. A total of 7100 patients and 3060 grafts under individual coronary artery bypass and 7380 grafts under sequential coronary artery bypass were included. Baseline characteristics and quality assessment according to NOS is

presented in Table 1. All clinical trials included in our study were middle to high quality cohort studies, with seven to nine NOS scores.

Graft Patency

Seven clinical studies reported the results of graft patency. The analysis of the graft patency rate includes 2374 out of 2739 grafts from the individual coronary artery bypass group and 6803 out of 7210 from the sequential coronary artery bypass group. There are no significant differences between these groups (RR=0.96; 95% CI=0.91-1.02; $P=0.16$; $I^2=87\%$) on the patency of grafts, as shown in Figure 2.

Anastomosis Patency

There were six clinical studies showing the results of anastomosis patency. The analysis of anastomosis patency includes 1100 out of 1400 anastomosis from the individual coronary artery bypass group and 1875 out of 2214 from the sequential coronary artery bypass group. There are no significant differences between these groups (RR=0.95; 95% CI=0.91-1.00; $P=0.05$; $I^2=70\%$) in the patency of anastomosis (Figure 3).

Occluded Rate in Left Anterior Descending (LAD) System

Three clinical studies reported the occluded rate in LAD system. The analysis found the occluded rate in LAD system in 145 out of 235 LAD system anastomoses from the individual coronary artery bypass group and 213 out of 349 from the sequential coronary artery bypass group. There are no significant differences between these groups (RR=1.03; 95% CI=0.92-1.16; $P=0.58$; $I^2=37\%$) in the occluded rate in LAD system, as demonstrated in Figure 4.

Occluded Rate in Right Coronary Artery (RCA) System

The occluded rate in RCA system was reported in three clinical studies. The analysis found the occluded rate in RCA system in 185 out of 403 RCA system anastomoses from the individual coronary artery bypass group and 122 out of 327 from the sequential coronary artery bypass group. There are no significant differences between these groups (RR=1.36; 95% CI=0.72-2.57; $P=0.35$; $I^2=95\%$) in the occluded rate in RCA system, as shown in Figure 5.

In-hospital Mortality

There were three clinical studies reporting the in-hospital mortality rates. The analysis of n of hospital mortality shows 16 of 664 patients from the individual coronary artery bypass group and 71 of 3765 from the sequential coronary artery bypass group. However, there are no significant differences between these groups (RR=1.57; 95% CI=0.92-2.69; $P=0.10$; $I^2=0\%$) in the in-hospital mortality rates (Figure 6).

Follow-up Mortality

There were two clinical studies reporting the follow-up mortality rates. The analysis of n of hospital mortality shows eight




Fig. 1 – Flowchart showing the progress of data selection.

out of 290 patients from the individual coronary artery bypass group and eight out of 275 patients from the sequential coronary artery bypass group. There are no significant differences between these groups ($RR=0.96$; $95\% \text{ CI}=0.36-2.53$; $P=0.93$; $I^2=0\%$) in the follow-up mortality rates (Figure 7).

Sensitivity and Publication Bias Analyses

Sensitivity analysis was conducted by excluding each individual study. It was found that the study of Christenson^[6], in 1998, resulted in a significantly different result, as shown

in Figure 8. A similar meta-analysis outcome was obtained, which demonstrated that our conclusion is stable, and this heterogeneity is not affected by the combined results. No significant evidence of publication bias was obtained using the Begg's test in the study endpoints, as shown in Table 2.

DISCUSSION

CABG has become the gold standard for the treatment of coronary artery disease involving multiple vessels, and it consists of on-pump CABG and off-pump CABG. In the early 1980s, two

Table 1. Baseline characteristics and quality assessment.

References	Community	Comparability			Assessment method	Follow-up time	Quality assessment (NOS)				
		Age	Male (%)	Other factors			Rate (%)	Selection	Comparability	Outcome	Total
Takazawa et al. ^[13] , 2015	Saitama International Medical Center in Japan	71±8	63.4	–	Angiography	14.7±17.5	–	4	1	2	7
Fukui et al. ^[8] , 2012	Sakakibara Heart Institute, Tokyo, Japan	67.2±10.4/67.0±10.7	76.5/85.2	Body surface area	Angiography	12.1(2–21)	50.9	4	2	1	7
Samano et al. ^[12] , 2017	Orebro University Hospital, Örebro, Sweden.	75.6±8.5	74.6/81	Body mass index, hypertension	Angiography	72	93.8	4	2	3	9
Ji et al. ^[10] , 2017	Zhongshan Hospital Fudan University, China	63.6±8.5/62.9±9.4	87.5/90	Smoking history, diabetes mellitus	Computed tomographic angiography	27.0±7.3	–	4	2	2	9
Gao et al. ^[9] , 2010	Patients operated on by a single surgeon	63.6±10.3	89	–	Angiography	26.4±23.6	–	4	2	1	7
Kim et al. ^[11] , 2011	Asan Medical Center	63.7±8.3/62.9±8.3	69.6/69	Hypertension, diabetes mellitus	Dual-source CT	14.8	–	4	2	2	8
Farsak et al. ^[7] , 2003	–	55.2±9.3	87	Atherosclerotic risk factors	Angiography	55.4±17.6	–	3	2	2	7
Vural et al. ^[14] , 2001	YuĖksek İhtisas Hospital in Turkey	49±8	89	Atherosclerotic risk factors	Angiography	69.6	–	4	2	2	8
Christenson et al. ^[6] , 1998	–	58.2±9.2	81	Hypertension, hyperlipidemia, diabetes	Angiography	76	99.10%	3	2	3	8

CT=computed tomography; NOS=Newcastle Ottawa Scale

surgeons published their extensive series of off-pump CABG in patients who received grafts in the LAD and the main RCA, but with more limited and difficult grafting of coronary arteries on the posterior and lateral walls^[15,16]. On-pump CABG provides a motionless operative field, but it can be associated with a number of complications, such as myocardial ischemic injury, strokes, coagulation, and inflammatory responses^[17,18]. To the

present, it has been reported that the advantage of sequential coronary artery bypass technology is that it can save grafts, reduce proximal anastomosis, shorten the operation time, provide a more complete vascularization, and have a satisfactory long-term patency rate^[19,20]. It is more accurate to determine the direction and length of the bridge between anastomoses. There is a study showing that the proximal obstruction of the sequential




Fig. 2 – Forest plot of graft patency. CI=confidence interval




Fig. 3 – Forest plot of anastomosis patency. CI=confidence interval




Fig. 4 – Forest plot of occluded rate in LAD system. CI=confidence interval; LAD=left anterior descending




Fig. 5 – Forest plot of occluded rate in RCA system. CI=confidence interval; RCA=right coronary artery




Fig. 6 – Forest plot of in-hospital mortality. CI=confidence interval




Fig. 7 – Forest plot of follow-up mortality. CI=confidence interval




Fig. 8 – Sensitivity analysis of excluding each individual study. CI=control

Table 2. Publication bias of the Begg's test.

Endpoints	P-value
Graft patency	1.00
Anastomosis patency	0.133
Occluded rate in LAD system	0.296
Occluded rate in RCA system	0.296
In-hospital mortality	0.296

LAD=left anterior descending; RCA=right coronary artery

bridge leads to the reduction of blood flow in multiple coronary arteries, resulting in a large area of myocardial infarction, and endangers the patient's life^[21]. It is also considered that the distal end of the proximal end of the sequential bridge plays an important role in collateral circulation, and patients rarely have myocardial infarction^[22,23]. In addition, some professors and their teams have found out that a sequential bridge can reduce the blood flow resistance of bridges, which reduces the mismatch of vascular resistance and increases the long-term patency rate.

The coronary circulation can be divided into left-dominant, right-dominant, and co-dominant systems. In a left-dominant system, the posterior descending artery (PDA) is supplied by the circumflex artery. In a right-dominant system, the PDA is supplied by the RCA^[24]. In this meta-analysis, we included nine trials with a total of 7100 patients and 1440 grafts under individual or sequential coronary artery bypass. We found out that the individual and sequential coronary artery bypass associated show no significant differences in the graft patency, anastomosis patency, occluded rate in LAD system, occluded rate in RCA system, in-hospital mortality, and follow-up mortality. In a previous study, the patency of sequential coronary artery bypass was lower than of individual coronary artery bypass. In a meta-analysis of RCTs on almost 16,900 patients, it was found no difference in 30-day mortality^[25]. Kowalewski et al.^[26] found out that 19,000 patients demonstrated no significant difference in short-term mortality. Another investigation in a recent meta-analysis of RCTs indicated no difference in patients with over a six-months follow-up (RR, 1.02; 95% CI: 0.86-1.22; $P=0.81$)^[27,28]. In the present study, we included the newest clinical trials and compared the individual and sequential coronary artery bypass groups regarding grafts and anastomosis. Also, we performed sensitivity and publication bias analyses, which demonstrates that our analysis is stable and has no publish bias.

Limitation

Nevertheless, there are some limitations in this meta-analysis. Firstly, the graft used in coronary artery bypass is not unified. Saphenous vein grafts and internal thoracic artery are both included in this study, which might have affected the reliability of the results. Secondly, several of the included clinical trials are cohort studies, instead of randomized clinical trials, which reduces the level of evidence. In addition, the generally

different designs and characteristics of each trial might have also caused heterogeneity. Therefore, more rigorous, large-sample, international trials are needed to further confirm the results.

CONCLUSION

In conclusion, no significant difference on clinical data were observed regarding anastomosis patency, occluded rate in LAD system, occluded rate in RCA system, in-hospital mortality, and follow-up mortality. The patency of individual and the patency of sequential coronary artery bypass are similar to each other.

No financial support.

No conflict of interest.

Authors' roles & responsibilities

ZL	Substantial contributions to the conception or design of the work; or the acquisition, analysis, or interpretation of data for the work; final approval of the version to be published
LL	Substantial contributions to the conception or design of the work; or the acquisition, analysis, or interpretation of data for the work; final approval of the version to be published

REFERENCES

1. The top 10 causes of death [Internet]. World Health Statistics: Geneva; 24 May 2018 [cited 2019 May 15]. Available from: <http://www.who.int/mediacentre/factsheets/fs310/en/>.
2. Harskamp RE, Alexander JH, Ferguson TB Jr, Hager R, Mack MJ, Englum B, et al. Frequency and Predictors of Internal Mammary Artery Graft Failure and Subsequent Clinical Outcomes: Insights From the Project of Ex-vivo Vein Graft Engineering via Transfection (PREVENT) IV Trial. *Circulation*. 2016;133(2):131-8. doi: 10.1161/CIRCULATIONAHA.115.015549.
3. Flemma RJ, Johnson WD, Lepley D Jr. Triple aorto-coronary vein bypass as treatment for coronary insufficiency. *Arch Surg*. 1971;103(1):82-3.
4. Walther S, Schuetz GM, Hamm B, Dewey M. [Quality of reporting of systematic reviews and meta-analyses: PRISMA (Preferred Reporting Items for Systematic reviews and Meta-Analyses)]. *Rofo*. 2011 Dec;183(12):1106-10. doi:10.1055/s-0031-1281809. German.
5. Wells GA, Shea B, O'Connell D, Peterson J, Welch V, Losos M, et al. The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses [Internet]. Ottawa (CA): Ottawa Hospital Research Institute; c2019 [cited 2019 May 15]. Available from: http://www.ohri.ca/programs/clinical_epidemiology/oxford.htm.
6. Christenson JT, Simonet F, Schmuziger M. Sequential vein bypass grafting: tactics and long-term results. *Cardiovasc Surg*. 1998;6(4):389-97. doi:10.1016/S0967-2109(98)00024-6.
7. Farsak B, Tokmakoglu H, Kandemir O, Günaydin S, Aydin H, Yorgancioglu C, et al. Angiographic assessment of sequential and individual coronary artery bypass grafting. *J Card Surg*. 2003;18(6):524-9; discussion 30-31. doi:10.1046/j.0886-0440.2003.02063.x.

8. Fukui T, Tabata M, Morita S, Takanashi S. Sequential free right internal thoracic artery grafting for multivessel coronary artery bypass grafting. *J Thorac Cardiovasc Surg.* 2012;144(4):824-9. doi:10.1016/j.jtcvs.2011.11.021.
9. Gao C, Wang M, Wang G, Xiao C, Wu Y, Li B, et al. Patency of sequential and individual saphenous vein grafts after off-pump coronary artery bypass grafting. *J Card Surg.* 2010;25(6):633-7. doi:10.1111/j.1540-8191.2010.01145.x.
10. Ji Q, Shi Y, Xia L, Ma R, Shen J, Lai H, et al. Revascularization of Left Coronary System Using a Skeletonized Left Internal Mammary Artery-Sequential vs. Separate Grafting. *Circ J.* 2017;82(1):102-9. doi:10.1253/circj.CJ-17-0223.
11. Kim HJ, Lee TY, Kim JB, Cho WC, Jung SH, Chung CH, et al. The impact of sequential versus single anastomoses on flow characteristics and mid-term patency of saphenous vein grafts in coronary bypass grafting. *J Thorac Cardiovasc Surg.* 2011 Mar;141(3):750-4. doi:10.1016/j.jtcvs.2010.05.037.
12. Samano N, Geijer H, Bodin L, Arbeus M, Mannion JD, Dashwood M, et al. The no-touch saphenous vein graft in elderly coronary bypass patients with multiple comorbidities is a promising conduit to substitute the left internal thoracic artery. *J Thorac Cardiovasc Surg.* 2017 Aug;154(2):457-466.e3. doi:10.1016/j.jtcvs.2017.03.048.
13. Takazawa A, Nakajima H, Iguchi A, Tabata M, Morita K, Koike H, et al. Impacts of intraoperative flow on graft patency of sequential and individual saphenous vein grafts. *Innovations (Phila).* 2015;10(2):85-9. doi:10.1097/IMI.0000000000000140.
14. Vural KM, Sener E, Tasdemir O. Long-term patency of sequential and individual saphenous vein coronary bypass grafts. *Eur J Cardiothorac Surg.* 2001;19(2):140-4. doi:10.1016/S1010-7940(00)00629-1.
15. Buffolo E, Andrade JC, Succi J, Leão LE, Gallucci C. Direct myocardial revascularization without cardiopulmonary bypass. *Thorac Cardiovasc Surg.* 1985;33(1):26-9. doi:10.1055/s-2007-1014076.
16. Gaudino M, Angelini GD, Antoniadis C, Bakaeen F, Benedetto U, Calafiore AM, et al. Off-Pump Coronary Artery Bypass Grafting: 30 Years of Debate. *J Am Heart Assoc.* 2018 Aug 21;7(16):e009934. doi:10.1161/JAHA.118.009934.
17. King N. Is there a long-term survival benefit with on pump coronary artery bypass grafting? *Ann Transl Med.* 2017;5(24):498. doi:10.21037/atm.2017.10.10.
18. Puskas JD, Williams WH, O'Donnell R, Patterson RE, Sigman SR, Smith AS, et al. Off-pump and on-pump coronary artery bypass grafting are associated with similar graft patency, myocardial ischemia, and freedom from reintervention: long-term follow-up of a randomized trial. *Ann Thorac Surg.* 2011;91(6):1836-42; discussion 1842-3. doi:10.1016/j.athoracsur.2010.12.043.
19. Al-Ruzzeh S, George S, Bustami M, Nakamura K, Khan S, Yacoub M, et al. The early clinical and angiographic outcome of sequential coronary artery bypass grafting with the off-pump technique. *J Thorac Cardiovasc Surg.* 2002;123(3):525-30. doi:10.1067/mtc.2002.119059.
20. McNamara JJ, Bjerke HS, Chung GK, Dang CR. Blood flow in sequential vein grafts. *Circulation.* 1979;60(2 Pt 2):33-8.
21. Kieser TM, FitzGibbon GM, Keon WJ. Sequential coronary bypass grafts. Long-term follow-up. *J Thorac Cardiovasc Surg.* 1986;91(5):767-72.
22. Christenson JT, Schmuziger M. Sequential Venous Bypass Grafts: Results 10 Years Later. *Ann Thorac Surg.* 1997;63(2):371-6. doi:10.1016/S0003-4975(96)01059-4.
23. Meeter K, Veldkamp R, Tijssen JG, van Herwerden LL, Bos E. Clinical outcome of single versus sequential grafts in coronary bypass operations at ten years' follow-up. *J Thorac Cardiovasc Surg.* 1991;101(6):1076-81.
24. Dong L, Kang YK, An XG. Short-Term and Mid-Term Clinical Outcomes Following Hybrid Coronary Revascularization Versus Off-Pump Coronary Artery Bypass: A Meta-Analysis. *Arq Bras Cardiol.* 2018;110(4):321-30. doi:10.5935/abc.20180044. English, Portuguese.
25. Fattouch K, Guccione F, Dioguardi P, Sampognaro R, Corrado E, Caruso M, et al. Off-pump versus on-pump myocardial revascularization in patients with ST-segment elevation myocardial infarction: a randomized trial. *J Thorac Cardiovasc Surg.* 2009;137(3):650-6; discussion 656-7. doi:10.1016/j.jtcvs.2008.11.033.
26. Deppe AC, Arbash W, Kuhn EW, Slottosch I, Scherner M, Liakopoulos OJ, et al. Current evidence of coronary artery bypass grafting off-pump versus on-pump: a systematic review with meta-analysis of over 16,900 patients investigated in randomized controlled trials†. *Eur J Cardiothorac Surg.* 2016;49(4):1031-41; discussion 1041. doi:10.1093/ejcts/ezv268.
27. Sá MP, Ferraz PE, Escobar RR, Oliveira Nunes E, Lustosa P, Vasconcelos FP, et al. Patency of skeletonized versus pedicled internal thoracic artery in coronary bypass graft surgery: a systematic review, meta-analysis and meta-regression. *Int J Surg.* 2014;12(7):666-72. doi:10.1016/j.ijssu.2014.05.071.
28. Luo T, Ni Y. Short-term and Long-term Postoperative Safety of Off-Pump versus On-Pump Coronary Artery Bypass Grafting for Coronary Heart Disease: A Meta-analysis for Randomized Controlled Trials. *Thorac Cardiovasc Surg* 2015;63(4):319-27. doi:10.1055/s-0035-1544232.

