



Brazilian Journal of Cardiovascular Surgery

ISSN: 0102-7638

ISSN: 1678-9741

Sociedade Brasileira de Cirurgia Cardiovascular

Petersen, Johannes; Holst, Theresa; Pecha, Simon;
Reichenspurner, Hermann; Girdauskas, Evaldas
Rare Presentation of Sinus of Valsalva Aneurysm Treated by Aortic Valve Reimplantation Technique
Brazilian Journal of Cardiovascular Surgery, vol. 37, no. 6, 2022, pp. 949-951
Sociedade Brasileira de Cirurgia Cardiovascular

DOI: <https://doi.org/10.21470/1678-9741-2020-0672>

Available in: <https://www.redalyc.org/articulo.oa?id=398973636024>

- ▶ How to cite
- ▶ Complete issue
- ▶ More information about this article
- ▶ Journal's webpage in redalyc.org

The logo for Redalyc.org, featuring the text 'Redalyc.org' with a stylized red and black graphic element.

Scientific Information System Redalyc

Network of Scientific Journals from Latin America and the Caribbean, Spain and Portugal

Project academic non-profit, developed under the open access initiative

Rare Presentation of Sinus of Valsalva Aneurysm Treated by Aortic Valve Reimplantation Technique

Johannes Petersen¹, MD; Theresa Holst¹, MD; Simon Pecha¹, MD; Hermann Reichenspurner¹, MD, PhD; Evaldas Girდაuskas¹, MD

DOI: 10.21470/1678-9741-2020-0672

ABSTRACT

Sinus of Valsalva aneurysm is a rare cardiac abnormality which can be acquired or of congenital origin. A spontaneous rupture into the right atrium is possible and, if not adequately treated, may result in a progressive heart failure due to the left-to-right intracardiac shunt. If ruptured sinus of Valsalva aneurysm is diagnosed, surgical repair is indicated, and different surgical techniques have been reported. If

concomitant aortic regurgitation is present, aortic valve replacement is usually performed. Herein, we describe an uncommon clinical presentation of a ruptured sinus of Valsalva aneurysm which has been corrected by aortic valve reimplantation.

Keywords: Sinus of Valsalva. Aortic Aneurysm. Heart Atria. Heart Failure. Replantation.

Abbreviations, acronyms & symbols

CT	= Computed tomography
TEE	= Transesophageal echocardiography

INTRODUCTION

Sinus of Valsalva aneurysm is a rare cardiac abnormality which can be of congenital origin (*e.g.*, connective tissue disorder) or acquired, caused by an inflammatory disease (*e.g.*, syphilis, endocarditis)^[1]. Most commonly, the aneurysm is located either in the right or the non-coronary sinus while the left coronary sinus is rarely involved^[1]. During the time course, a spontaneous rupture of sinus of Valsalva aneurysm into the right atrium is possible and, if not adequately treated, may result in a progressive heart failure due to the left-to-right intracardiac shunt or even in a sudden cardiac death^[1]. Depending on the size of perforation, clinical presentation may vary between an asymptomatic heart murmur, mild dyspnea, chest pain^[2], or even signs of acute heart failure^[1].

Diagnosis is confirmed with transesophageal echocardiography (TEE), contrast-enhanced computed tomography (CT), or magnetic resonance imaging^[3]. If ruptured sinus of Valsalva aneurysm is diagnosed, surgical repair is indicated and different surgical techniques have been reported, depending on the type of the aneurysm^[4]. If concomitant aortic regurgitation is present, aortic valve replacement is usually performed^[5]. Herein, we describe an uncommon clinical presentation of a ruptured sinus of Valsalva aneurysm which has been corrected by means of aortic valve reimplantation and simultaneous aortic valve repair.

TECHNIQUE

A 57-year-old male patient was referred to our hospital for catheter-based ablation due to paroxysmal atrial fibrillation. The patient reported intermittent palpitations, dizziness, and a reduced quality of life. Otherwise, the patient was healthy and had no previously diagnosed connective tissue disorder. During the preprocedural diagnostic work-up, a TEE was performed and revealed an ascending aortic aneurysm of 43 mm and a concomitant sinus of Valsalva aneurysm of 20 × 20 mm protruding

¹Department of Cardiovascular Surgery, University Heart & Vascular Center Hamburg, Hamburg, Germany.

This study was carried out at the Department of Cardiovascular Surgery, University Heart & Vascular Center Hamburg, Hamburg, Germany.

Correspondence Address:

Johannes Petersen

 <https://orcid.org/0000-0002-3415-3385>

Department of Cardiovascular Surgery, University Heart & Vascular Center Hamburg Martinistrasse, 52, Hamburg, Germany - Zip Code: 20251

E-mail: joh.petersen@uke.de

Article received on December 3rd, 2020.

Article accepted on December 8th, 2020.

into the right atrium (Figure 1A) associated with an aorto-right atrial shunt (Figure 1B). Right atrial dimensions were significantly enlarged. Mild aortic valve regurgitation was present and left ventricular systolic function was preserved. Subsequent cardiac CT scan confirmed the diagnosis of perforated sinus of Valsalva aneurysm with contrast medium shunting into the right atrium (Figure 1C). Intraoperatively, ruptured sinus of Valsalva aneurysm originating from the non-coronary and the right coronary sinus was present (Figure 2A). Native tricuspid aortic valve was highly asymmetric and right coronary cusp showed geometric height

of only 15 mm in combination with a reduced effective height of 4 mm. (Figure 2A). The geometric height of the non- and the left coronary cusp was 22 mm and 18 mm, respectively. Despite this complex asymmetric anatomy, a valve-sparing procedure was planned, taking into account a good quality of the native cusps. First, the aorto-right atrial shunt was closed, and the right atrial roof restored using a bovine pericardial patch (40 × 20 mm) (Figure 2B). Next, aortic sinus tissue was completely resected preserving only both coronary buttons. During the reimplantation, the three aortic valve commissures were

asymmetrically attached into the 28-mm Gelweave™ Valsalva Graft to mimic the original valve orientation and to achieve sufficient coaptation of aortic cusps. In addition, central plication suture of the right coronary cusp was performed to reach the effective cusp height of 8 mm. Furthermore, pulmonary vein isolation was performed with bipolar radiofrequency and the left atrial appendage was closed using an AtriClip® LAA Exclusion System. After weaning from cardiopulmonary bypass, the TEE showed a normal aortic valve function with only trace residual aortic regurgitation, no residual left-to-right cardiac shunt, and complete closure of the left atrial appendage. The patient recovered uneventfully after the surgery and was discharged in stable sinus rhythm and only trace aortic regurgitation with a mean transvalvular gradient of 13 mmHg.




Fig. 1 - Preoperative transesophageal echocardiography (A, B) and contrast-enhanced computed tomography imaging (C) showing the sinus of Valsalva aneurysm with an aorto-right atrial shunt (arrows).




Fig. 2 - Intraoperative findings of the perforated sinus of Valsalva aneurysm with an aorto-right atrial shunt (arrow) and the asymmetric tricuspid aortic valve configuration (A) and the closure of the right atrial shunt with bovine pericardial patch (B) before aortic valve reimplantation was performed.

DISCUSSION

This case represents an aortic valve sparing root procedure which aims to maintain the geometry of aortic valve considering the marked asymmetry of aortic root. The challenge of reimplantation procedure in this scenario is to keep the asymmetric configuration of aortic root due to extremely dilated non-coronary sinus and large non-coronary cusp of the aortic valve. Aortic valve annulus was almost non-existent below the perforated sinus of Valsalva aneurysm in the non-coronary sinus and was therefore fixed from inside to outside using the standard pledgeted subannular anchoring

sutures. In this specific case, the defect was in the right atrial groove well above the tricuspid valve plane. Therefore, the right atrium was not separately opened.

CONCLUSION

Despite an uncommon clinical presentation, a ruptured sinus of Valsalva aneurysm can be detected by comprehensive preoperative work-up. Further, an aortic valve reimplantation is possible even in a complex asymmetric anatomy if the geometry of the aortic valve is respected.

REFERENCES

1. Feldman DN, Roman MJ. Aneurysms of the sinuses of Valsalva. *Cardiology*. 2006;106(2):73-81. doi:10.1159/000092635.
2. Sanna GD, Talanas G, Denurra C, Ferrandu P, Bullitta L, Terrosu P. An unusual cause of acute chest pain: rupture of the noncoronary sinus of Valsalva into the right atrium. *Am J Emerg Med*. 2016;34(10):2052.e1-3. doi:10.1016/j.ajem.2016.03.014.
3. Alozie A, Kische S, Kaminski A, Ince H. Aorto-right atrial shunt due to perforated sinus of Valsalva aneurysm in a young athlete: a rare congenital defect confirmed by various non-invasive imaging modalities. *Eur Heart J*. 2012;33(1):138. doi:10.1093/eurheartj/ehr230.
4. Guo HW, Xiong H, Xu JP, Wang XQ, Hu SS. A new and simple classification for sinus of Valsalva aneurysms and the corresponding surgical procedure. *Eur J Cardiothorac Surg*. 2013;43(6):1188-93. doi:10.1093/ejcts/ezs673.
5. Lin Y, Yin K, Wang Y, Guo C, Tian Z, Xie Q, et al. Sinus of Valsalva aneurysms with concomitant aortic insufficiency: how should the aortic valve be managed? *Interact Cardiovasc Thorac Surg*. 2018;26(2):210-5. doi:10.1093/icvts/ivx302.

No financial support.

No conflict of interest.

Authors' roles & responsibilities

JP	Substantial contributions to the conception or design of the work; or the acquisition, analysis, or interpretation of data for the work; drafting the work or revising it critically for important intellectual content; final approval of the version to be published
TH	Substantial contributions to the conception or design of the work; or the acquisition, analysis, or interpretation of data for the work; drafting the work or revising it critically for important intellectual content; final approval of the version to be published
SP	Substantial contributions to the conception or design of the work; or the acquisition, analysis, or interpretation of data for the work; drafting the work or revising it critically for important intellectual content; final approval of the version to be published
HR	Substantial contributions to the conception or design of the work; or the acquisition, analysis, or interpretation of data for the work; drafting the work or revising it critically for important intellectual content; final approval of the version to be published
EG	Substantial contributions to the conception or design of the work; or the acquisition, analysis, or interpretation of data for the work; drafting the work or revising it critically for important intellectual content; final approval of the version to be published



This is an open-access article distributed under the terms of the Creative Commons Attribution License.