



Brazilian Journal of Cardiovascular Surgery

ISSN: 0102-7638

ISSN: 1678-9741

Sociedade Brasileira de Cirurgia Cardiovascular

Hou, Jian; Huang, Xiaolin; Shang, Liquan; Chen,
Guangxian; Wu, Huawei; Wu, Zhongkai; Huang, Suiqing
Reduced Angiotensin II Levels Are Correlated with Better
Cardiac Function and Prognosis in Valvular Heart Disease
Brazilian Journal of Cardiovascular Surgery, vol. 38, no. 1, 2023, pp. 104-109
Sociedade Brasileira de Cirurgia Cardiovascular

DOI: <https://doi.org/10.21470/1678-9741-2021-0364>

Available in: <https://www.redalyc.org/articulo.oa?id=398974859014>

- How to cite
- Complete issue
- More information about this article
- Journal's webpage in [redalyc.org](https://www.redalyc.org)

The logo for Redalyc, featuring the word 'redalyc' in a stylized font with a red heart shape integrated into the letter 'a', followed by '.org'.

Scientific Information System Redalyc

Network of Scientific Journals from Latin America and the Caribbean, Spain and
Portugal

Project academic non-profit, developed under the open access initiative

Reduced Angiopietin Factor 2 Levels Are Correlated with Better Cardiac Function and Prognosis in Valvular Heart Disease

Jian Hou^{1,2*}, PhD; Xiaolin Huang^{1,2*}, MD; Liqun Shang^{1,2*}, MD; Guangxian Chen^{1,2}, MD; Huawei Wu³, BSc; Zhongkai Wu^{1,2}, MD, PhD; Suiqing Huang^{1,2}, MD

DOI: 10.21470/1678-9741-2021-0364

ABSTRACT

Introduction: There are few circulating biomarkers for valvular heart disease. Angiopietin (Ang) 1, Ang2, and vascular endothelial growth factor are important inflammation-associated cytokines. The aim of this study was to investigate the clinical significance and association of Ang1, Ang2, and vascular endothelial growth factor in valvular heart disease.

Methods: This is a retrospective study; a total of 62 individuals (valvular heart disease patients [n=42] and healthy controls [n=20]) were included. Plasma levels of Ang1, Ang2, and vascular endothelial growth factor were detected by enzyme-linked immunosorbent assays. We retrospectively collected the baseline characteristics and short-term outcomes; logistic regression was performed to identify predictor for short-term mortality.

Results: Ang2 was significantly decreased in the valvular heart disease group compared with the healthy control group ($P=0.023$), while no significant difference was observed in the Ang1 and vascular endothelial

growth factor levels. The Ang2 level of New York Heart Association (NYHA) I/II patients — but not NYHA III/IV patients — was significantly decreased compared with that of healthy control individuals (NYHA I/II: $P=0.017$; NYHA III/IV: $P=0.485$). Univariable logistic regression analysis indicated that Ang2 was a significant independent predictor for short-term mortality (odds ratio 18.75, $P=0.033$, 95% confidence interval 8.08–102.33). Ang1 was negatively correlated with Ang2 ($P=0.032$, Pearson's correlation coefficient = -0.317) and was positively correlated with vascular endothelial growth factor ($P=0.019$, Pearson's correlation coefficient = 0.359).

Conclusion: Ang2 might serve as a therapeutic and prognostic target for valvular heart disease.

Keywords: Valvular Heart Disease. Angiopietin 1. Angiopietin 2. Heart Valve Diseases. Inflammation. Cytokines.

Abbreviations, Acronyms & Symbols			
Ang	= Angiopietin	NYHA	= New York Heart Association
CI	= Confidence interval	OR	= Odds ratio
CPB	= Cardiopulmonary bypass	PLT	= Platelet
EF	= Ejection fraction	SD	= Standard deviation
ELISA	= Enzyme-linked immunosorbent assay	VEGF	= Vascular endothelial growth factor
ICU	= Intensive care unit	VHD	= Valvular heart disease
NT-pro-BNP	= N-terminal pro B-type natriuretic peptide	WBC	= White blood cell

¹Department of Cardiac Surgery, The First Affiliated Hospital of Sun Yat-sen University, Guangzhou, Guangdong Province, People's Republic of China.

²NHC Key Laboratory of Assisted Circulation, Sun Yat-sen University, Guangzhou, Guangdong Province, People's Republic of China.

³Department of Neurobiology, Physiology and Behavior, College of Biological Sciences, University of California, Davis, California, United States of America.

*Jian Hou, Xiaolin Huang, and Liqun Shang contributed equally to this work.

This study was carried out at the Department of Cardiac Surgery, The First Affiliated Hospital of Sun Yat-sen University, Guangzhou, Guangdong Province, People's Republic of China.

Correspondence Address:

Zhongkai Wu

 <https://orcid.org/0000-0003-1184-1182>

Department of Cardiac Surgery, The First Affiliated Hospital of Sun Yat-sen University

58 Zhongshan Rd. No.2, Guangzhou, Guangdong Province, People's Republic of China - Zip Code: 510080

E-mail: wuzhk@mail.sysu.edu.cn

Article received on June 25th, 2021.
Article accepted on November 2nd, 2021.

INTRODUCTION

Valvular heart disease (VHD) is an important cause of morbidity and mortality, with a prevalence that is predicted to increase dramatically because of the increase in life expectancy in middle-income and high-income nations. Despite the wider appreciation of the emerging importance of VHD, the mechanisms underlying its development are poorly understood, and the role of circulating biomarkers in guiding clinical management in individual patients is relatively unexplored.

Angiopoietin (Ang) 1 and 2, secreted 70-kDa glycoproteins and well-known members of the Ang-Tie pathway, have been demonstrated to participate in a number of cardiovascular diseases^[1-3]. Ang2 mediates vascular leakage and inflammation, while Ang1 has antipermeability, anti-inflammatory, and cardioprotective effects^[4-7]. Higher plasma Ang2 levels were predictive of myocardial infarction^[8,9] and stroke recurrence^[10] and were independent of traditional risk factors. In acute myocardial infarction, the extent of myocardial damage correlated with serum Ang2 and Ang2/Ang1^[11], indicating that Ang1 and Ang2 are potential biomarkers of the severity of cardiac disease.

Vascular endothelial growth factor (VEGF) has been shown to act as a proinflammatory cytokine by inducing the expression of adhesion molecules that cause leukocytes to bind endothelial cells^[12,13]. Moreover, increased VEGF levels were associated with an imbalance of Ang1 and Ang2^[14], suggesting the complicated connection of these inflammation-associated cytokines. However, investigations on the correlation of VEGF and Ang have mostly focused on coagulation, and few have been related to inflammation.

Given that VHD is characterized by an ongoing inflammatory cellular response, we hypothesized that Ang1, Ang2, and VEGF play important roles in the progression and prognosis of VHD. To explore this hypothesis, plasma levels of Ang1, Ang2, and VEGF in VHD patients and healthy individuals were measured.

METHODS

Participants' Clinical Information

Forty-two subjects with VHD were recruited between March 2018 and March 2019. All patients underwent a valve replacement procedure and recovered well. Twenty subjects without VHD and asymptomatic cardiovascular events were recruited as a healthy control group.

The exclusion criteria included age < 18 years old, recent-onset acute cardiovascular or cerebrovascular events, trauma, or surgery, coronary artery heart disease, pregnancy, acute or chronic infection, autoimmune disease, malignant tumor, and pulmonary, hepatic, or renal dysfunction. Clinical data of the patients were reviewed and collected. The study was approved by the clinical research and experimental animal ethics committee ([2017]157), and all patients enrolled in this study provided informed consent prior to participating in the study.

Blood Sample Collection and Detection

Blood samples from VHD patients and healthy controls were obtained from forearm veins using 6.0-mL vacutainer tubes (BD, Plymouth, United Kingdom) containing ethylenediamine tetraacetic acid. After centrifugation at 2,500 rpm for 15 minutes, plasma samples were collected and stored at -80°C within one day for further enzyme-linked immunosorbent assays (ELISAs) detection. The plasma levels of Ang1, Ang2, and VEGF were determined using ELISAs according to the instructions from the manufacturer (R&D Systems, Minneapolis, Minnesota, United States of America).

Statistical Analysis

Normality was tested by Shapiro-Wilk test. Continuous variables with normal distribution were expressed as mean±standard deviation and compared by Student's *t*-test or one-way analysis of variance. Categorical variables were expressed as percentages (number and %) and compared by Chi-squared test. All potential predictors for short-term (within 30 days after surgery) mortality were studied using univariable logistic regression analysis. Correlations were determined using Pearson's correlation method. For all tests, *P*<0.05 was considered statistically significant.

Statistical analysis was performed using GraphPad Prism Software (Version 8, La Jolla, California, United States of America) and IBM Corp. Released 2013, IBM SPSS Statistics for Windows, version 22.0, Armonk, NY: IBM Corp.

RESULTS

Patients' Characteristics

Table 1 shows the demographic data of the patients enrolled in this study. Patients with VHD consisted of 22 men and 20 women (age: 55.65±1.71 years). We retrieved information, including sex, age, New York Heart Association (NYHA) classifications, ejection fraction (EF), N-terminal pro B-type natriuretic peptide level, creatinine, uric acid, white blood cell count, neutrophils, neutrophil percentage, neutrophil-to-lymphocyte ratio, platelet, cardiopulmonary bypass duration, aortic cross-clamping duration, length of tracheal intubation, length of intensive care unit stay, and length of hospital stay from patients' hospital charts.

Concentration of Ang1, Ang2, and VEGF in VHD Patients with Different NYHA Classifications

Ang1, Ang2, and VEGF levels were detected in the plasma samples of VHD patients and the healthy control group (Tables 2 and 3). Compared with the healthy control group,

Table 1. Clinical characteristics of VHD patients (n=42).

Variables	n (%) or mean±SD
Age (years)	55.65±1.71
Male/female (% male)	22/20 (52.38%)
NYHA	
I	2 (4.76%)
II	28 (66.67%)
III	10 (23.81%)
IV	2 (4.76%)
EF (before surgery)	63.41±1.54
EF (after surgery)	56.98±1.70
NT-pro-BNP (before surgery)	1412.10±402.41
Creatinine	89.37±4.87
Uric acid	446.00±20.87
WBC count (109/L)	7.00±0.29
Neutrophil (109/L)	4.07±0.24
Neutrophilic percentage	0.57±0.01
Neutrophil-to-lymphocyte ratio	2.07±0.14
PLT	224.34±9.05
CPB duration (min)	146.81±6.71
Aortic cross-clamping duration (min)	84.93±4.03
Length of tracheal intubation (hour)	33.13±15.52
Length of ICU stay (day)	2.70±0.65
Length of hospital stay (day)	29.74±1.73

The results are expressed as n (%) or mean±standard deviation (SD)

CPB=cardiopulmonary bypass; EF=ejection fraction; ICU=intensive care unit; NT-pro-BNP=N-terminal pro B-type natriuretic peptide; NYHA=New York Heart Association; PLT=platelet; VHD=valvular heart disease; WBC=white blood cell

Table 2. Ang1, Ang2, and VEGF levels in control group and VHD patients.

Parameter	Control (n=20)	Total VHD (n=42)	P-value
Ang1 (pg/mL)	2842.96±769.51	6147.63±1894.10	0.112
Ang2 (pg/mL)	1736.70±95.69	1365.05±98.48	0.023
VEGF (pg/mL)	109.58±17.92	110.86±14.63	0.956

The results are expressed as means±standard deviation. Differences in the data between the two groups were determined by Student's *t*-test

Ang=angiopoietin; VEGF=vascular endothelial growth factor; VHD=valvular heart disease

Ang2 was significantly decreased in the VHD group (control = 1736.70±95.69 pg/mL, VHD = 1365.05±98.48 pg/mL, *P*=0.023), while no significant difference was observed in the Ang1 and VEGF levels. Further, differences in plasma Ang1 and Ang2 and VEGF levels among patients with NYHA I/II classes, NYHA III/

IV classes, and controls were analyzed. Interestingly, only the Ang2 expression level of NYHA I/II patients was significantly decreased compared with that of healthy control individuals, but the same was not true for NYHA III/IV patients (NYHA I/II: 1317.01±120.36 pg/mL vs. healthy control group: *P*=0.017; NYHA

III/IV: 1567.81 ± 189.27 pg/mL vs. healthy control group: $P=0.485$), suggesting that decreased Ang2 might be associated with better cardiac function in VHD.

Univariable Logistic Regression

Univariable analysis (Table 4) showed significant differences between short-term survivors and non-survivors in the following factor: Ang2 (odds ratio 18.75, $P=0.033$, 95% confidence interval 8.08–102.33).

Correlation of Ang1, Ang2, and VEGF Levels

Given the complex connection of Ang1, Ang2, and VEGF^[14], the correlation of Ang1, Ang2, and VEGF levels was also analyzed and is shown in Table 5. Ang1 levels were negatively correlated with Ang2 levels ($P=0.032$, Pearson's correlation coefficient = -0.317) and positively correlated with VEGF levels ($P=0.019$, Pearson's correlation coefficient = 0.359).

DISCUSSION

In the present study, we found that the plasma level of Ang2 was significantly lower in all VHD patients and VHD patients with NYHA I/II classes — but not in VHD patients with NYHA III/IV classes — than in healthy controls. It may suggest an association of decreased Ang2 levels and better cardiac function and prognosis in VHD. The Ang1 expression level was negatively associated with Ang2 but positively associated with VEGF.

The Ang-Tie axis, one of the most important switch signaling pathways for angiogenesis, has been documented in a wide range of cardiovascular diseases and inflammatory diseases^[1,15]. Among the pathway members, the best characterized are Ang1 and Ang2. Ang2 mediates vascular leakage^[4] and inflammation^[5], while Ang1 has antipermeability^[4], anti-inflammatory^[6], and cardioprotective effects^[7] because of their opposite pathophysiological effects in response to binding to the Tie2 receptor^[16]. Circulating Ang2 levels and the Ang2/Ang1 ratio could serve as suitable biomarkers of inflammatory

Table 3. Ang1, Ang2, and VEGF levels in control group and VHD patients.

Parameter	Control (n=20)	VHD with NYHA I/II (n=30)	VHD with NYHA III/IV (n=12)	P-value ^a	P-value ^b
Ang1 (pg/mL)	2842.96±769.51	7841.771±2927.14	3602.88±769.35	0.151	0.857
Ang2 (pg/mL)	1736.70±95.69	1317.01±120.36	1567.81±189.27	0.017	0.485
VEGF (pg/mL)	109.58±17.92	121.02±20.87	88.34±44.70	0.686	0.531

The results are expressed as means±standard deviation. Statistical difference between the groups was evaluated using one-way analysis of variance

P-value^a: NYHA I/II vs. control

P-value^b: NYHA III/IV vs. control

Ang=angiotensin; NYHA=New York Heart Association; VEGF=vascular endothelial growth factor; VHD=valvular heart disease

Table 4. Univariable logistic regression.

	Univariable analysis		
	OR	95% CI	P-value
EF (before surgery) (%)	1.005	(1.001, 1.014)	0.427
EF (after surgery) (%)	1.007	(1.001, 1.016)	0.238
Angiotensin 1 (pg/mL)	1.087	(1.008, 1.193)	0.113
Angiotensin 2 (pg/mL)	18.75	(8.08, 102.33)	0.033
Vascular endothelial growth factor (pg/mL)	1.014	(0.994, 1.031)	0.146
CPB duration (min)	1.001	(0.999, 1.005)	0.135
Aortic cross-clamping duration (min)	1.007	(0.989, 1.022)	0.097

Logistic regressions were used to analyze the predictors of 30-day mortality after surgery

CI=confidence interval; CPB=cardiopulmonary bypass; EF=ejection fraction; OR=odds ratio

Table 5. Correlations among plasma levels of Ang1, Ang2, and VEGF in VHD patients.

Variables	Ang1	Ang2	VEGF
Ang1			
Pearson's correlation coefficient	1.000		
P-value	--		
Ang2			
Pearson's correlation coefficient	-0.317	1.000	
P-value	0.034	--	
VEGF			
Pearson's correlation coefficient	0.359	0.177	1.000
P-value	0.019	0.285	--

Correlations between Ang1, Ang2, and VEGF expression levels were assessed with the parametric Pearson's rank correlation test.

P-value < 0.05 was considered statistically significant

Ang=angiopoietin; VEGF=vascular endothelial growth factor; VHD=valvular heart disease

disease. Circulating Ang2 levels were correlated with markers of endothelial cell activation and 28-day mortality in patients with severe sepsis^[15]. In acute lung injury patients, the Ang2/Ang1 ratio was an independent predictor of mortality^[17]. Higher Ang2 and Ang2/Ang1 were associated with poor cardiac function in acute myocardial infarction patients^[18]. Our results are consistent with previous results. In the present study, lower Ang2 levels were observed in the VHD patients with lower NYHA classes and higher EFs (Table 3). As Ang2 is an important inflammation-associated cytokine^[19,20], it may suggest that lower Ang2 levels might prevent VHD progression and improve prognosis by regulating inflammation levels. The association of Ang2/Ang1 ratio and VHD patient characteristics was not observed in the present study (data not shown); however, a significantly negative correlation of Ang1 and Ang2 was observed (Table 5), which might have been due to the small sample size of our study. Furthermore, Ang1 and Ang2 have been shown to interact with VEGF to induce an inflammatory response^[21]. Our data showed the positive association of Ang1 and VEGF (Table 5), indicating that a synergistic effect of Ang1 and VEGF might exist in VHD, which may regulate the inflammation level. Taken together, these results suggest that Ang2 could affect the disease progression and prognosis of VHD by regulating the inflammatory level via the interaction of Ang1 and VEGF.

In summary, a lower Ang2 level was associated with a better NYHA class in VHD. Univariable logistic regression analysis result indicated that Ang2 was a significant independent predictor for short-term mortality. Furthermore, the association of Ang1, Ang2, and VEGF was found in VHD. Thus, Ang2 might affect the disease progression and prognosis of VHD, which may involve the interaction of Ang1 and VEGF.

Limitations

As a single-center retrospective study, this study has all the limitations of a retrospective observational study. Thus, as an observational study, cause-effect relationships could not be

established. Further prospective studies are necessary to confirm our present findings. Also, the sample size of our study was small. The changes in plasma levels of Ang1, Ang2, and VEGF after valve replacement surgery were not evaluated. Future studies could examine whether the expression of these proteins changes after surgery and how the levels correlate with the outcomes of valve replacement.

CONCLUSION

Ang2 might serve as a therapeutic and prognostic target for VHD.

Financial support: This study was funded by the National Key R&D Program of China (x 2017YFC1105000) and the National Natural Science Foundation of China (81570039, 81770319, 81900294, 82070297).

No conflict of interest.

Authors' Roles & Responsibilities

JH	Substantial contributions to the conception and design of the work; and the analysis of data for the work; final approval of the version to be published
XH	Substantial contributions to the conception and design of the work; final approval of the version to be published
LS	Substantial contributions to the conception and design of the work; final approval of the version to be published
GC	Substantial contributions to the interpretation of data for the work; final approval of the version to be published
HW	Substantial contributions to the analysis of data for the work; final approval of the version to be published
ZW	Substantial contributions to the analysis of data for the work; final approval of the version to be published
SH	Substantial contributions to the interpretation of data for the work; final approval of the version to be published

REFERENCES

- Eklund L, Kangas J, Saharinen P. Angiopoietin-Tie signalling in the cardiovascular and lymphatic systems. *Clin Sci (Lond)*. 2017;131(1):87-103. doi:10.1042/CS20160129.
- Pang S, Tao Z, Min X, Zhou C, Pan D, Cao Z, et al. Correlation between the serum platelet-derived growth factor, angiopoietin-1, and severity of coronary heart disease. *Cardiol Res Pract*. 2020;2020:3602608. doi:10.1155/2020/3602608.
- Chen SM, Li D, Xing X, Li ZP. Higher serum angiopoietin 2 levels are independently associated with coronary microvascular dysfunction in patients with angina in the absence of obstructive coronary artery disease. *Chin Med J (Engl)*. 2020;133(14):1662-8. doi:10.1097/CM9.0000000000000812.
- van der Heijden M, van Nieuw Amerongen GP, van Bezu J, Paul MA, Groeneveld AB, et al. Opposing effects of the angiopoietins on the thrombin-induced permeability of human pulmonary microvascular endothelial cells. *PLoS One*. 2011;6(8):e23448. doi:10.1371/journal.pone.0023448.
- Fiedler U, Reiss Y, Scharpfenecker M, Grunow V, Koidl S, Thurston G, et al. Angiopoietin-2 sensitizes endothelial cells to TNF- α and has a crucial role in the induction of inflammation. *Nat Med*. 2006;12(2):235-9. doi:10.1038/nm1351.
- Gu H, Cui M, Bai Y, Chen F, Ma K, Zhou C, et al. Angiopoietin-1/Tie2 signaling pathway inhibits lipopolysaccharide-induced activation of RAW264.7 macrophage cells. *Biochem Biophys Res Commun*. 2010;392(2):178-82. doi:10.1016/j.bbrc.2010.01.009.
- Dallabrida SM, Ismail N, Oberle JR, Himes BE, Rupnick MA. Angiopoietin-1 promotes cardiac and skeletal myocyte survival through integrins. *Circ Res*. 2005;96(4):e8-24. doi:10.1161/01.RES.0000158285.57191.60.
- Patel JV, Lim HS, Varughese GI, Hughes EA, Lip GY. Angiopoietin-2 levels as a biomarker of cardiovascular risk in patients with hypertension. *Ann Med*. 2008;40(3):215-22. doi:10.1080/07853890701779586.
- Iribarren C, Phelps BH, Darbinian JA, McCluskey ER, Quesenberry CP, Hytopoulos E, et al. Circulating angiopoietins-1 and -2, angiopoietin receptor Tie-2 and vascular endothelial growth factor-A as biomarkers of acute myocardial infarction: a prospective nested case-control study. *BMC Cardiovasc Disord*. 2011;11:31. doi:10.1186/1471-2261-11-31.
- Chen J, Yu H, Sun K, Song W, Bai Y, Yang T, et al. Promoter variant of angiopoietin-2 and plasma angiopoietin-2 are associated with risk of stroke recurrence in lacunar infarct patients. *Biochem Biophys Res Commun*. 2010;398(2):212-6. doi:10.1016/j.bbrc.2010.06.062.
- Chen S, Guo L, Cui M, Sun L, Mi L. Dynamic changes in serum angiopoietin-1, angiopoietin-2, and angiopoietin-2/angiopoietin-1 ratio in acute myocardial infarction patients treated with primary percutaneous coronary intervention. *Biomarkers*. 2012;17(5):441-6. doi:10.3109/1354750X.2012.684152.
- Clauss M, Gerlach M, Gerlach H, Brett J, Wang F, Familletti PC, et al. Vascular permeability factor: a tumor-derived polypeptide that induces endothelial cell and monocyte procoagulant activity, and promotes monocyte migration. *J Exp Med*. 1990;172(6):1535-45. doi:10.1084/jem.172.6.1535.
- Blum S, Issbrücker K, Willuweit A, Hehlhans S, Lucerna M, Mechtcheriakova D, et al. An inhibitory role of the phosphatidylinositol 3-kinase-signaling pathway in vascular endothelial growth factor-induced tissue factor expression. *J Biol Chem*. 2001;276(36):33428-34. doi:10.1074/jbc.M105474200.
- Choudhury A, Freestone B, Patel J, Lip GY. Relationship of soluble CD40 ligand to vascular endothelial growth factor, angiopoietins, and tissue factor in atrial fibrillation: a link among platelet activation, angiogenesis, and thrombosis? *Chest*. 2007;132(6):1913-9. doi:10.1378/chest.07-1565.
- Ricciuto DR, dos Santos CC, Hawkes M, Toltl LJ, Conroy AL, Rajwans N, et al. Angiopoietin-1 and angiopoietin-2 as clinically informative prognostic biomarkers of morbidity and mortality in severe sepsis. *Crit Care Med*. 2011;39(4):702-10. doi:10.1097/CCM.0b013e318206d285.
- Maisonpierre PC, Suri C, Jones PF, Bartunkova S, Wiegand SJ, Radziejewski C, et al. Angiopoietin-2, a natural antagonist for Tie2 that disrupts in vivo angiogenesis. *Science*. 1997;277(5322):55-60. doi:10.1126/science.277.5322.55.
- Ong T, McClintock DE, Kallet RH, Ware LB, Matthay MA, Liu KD. Ratio of angiopoietin-2 to angiopoietin-1 as a predictor of mortality in acute lung injury patients. *Crit Care Med*. 2010;38(9):1845-51. doi:10.1097/CCM.0b013e3181eaa5bf.
- Chen S, Guo L, Chen B, Sun L, Cui M. Association of serum angiopoietin-1, angiopoietin-2 and angiopoietin-2 to angiopoietin-1 ratio with heart failure in patients with acute myocardial infarction. *Exp Ther Med*. 2013;5(3):937-41. doi:10.3892/etm.2013.893.
- Akwii RG, Sajib MS, Zahra FT, Mikelis CM. Role of Angiopoietin-2 in vascular physiology and pathophysiology. *Cells*. 2019;8(5):471. doi:10.3390/cells8050471.
- Li Z, Korhonen EA, Merlini A, Strauss J, Wihuri E, Nurmi H, et al. Angiopoietin-2 blockade ameliorates autoimmune neuroinflammation by inhibiting leukocyte recruitment into the CNS. *J Clin Invest*. 2020;130(4):1977-90. doi:10.1172/JCI130308.
- Sinnathamby T, Yun J, Clavet-Lanthier M $\acute{E}$, Cheong C, Sirois MG. VEGF and angiopoietins promote inflammatory cell recruitment and mature blood vessel formation in murine sponge/Matrigel model. *J Cell Biochem*. 2015;116(1):45-57. doi:10.1002/jcb.24941.



This is an open-access article distributed under the terms of the Creative Commons Attribution License.