

Brazilian Journal of Cardiovascular Surgery

ISSN: 0102-7638 ISSN: 1678-9741

Sociedade Brasileira de Cirurgia Cardiovascular

Jose, Jes; Magoon, Rohan; Kohli, Jasvinder Kaur; Kashav, Ramesh Deep Sternal Wound Infection Following OPCAB: Delving Deeper into the Predisposition! Brazilian Journal of Cardiovascular Surgery, vol. 38, no. 1, 2023, p. 213 Sociedade Brasileira de Cirurgia Cardiovascular

DOI: https://doi.org/10.21470/1678-9741-2021-0547

Available in: https://www.redalyc.org/articulo.oa?id=398974859033



Complete issue

More information about this article

Journal's webpage in redalyc.org



Scientific Information System Redalyc

Network of Scientific Journals from Latin America and the Caribbean, Spain and Portugal

Project academic non-profit, developed under the open access initiative

# Deep Sternal Wound Infection Following OPCAB: Delving Deeper into the Predisposition!

DOI: 10.21470/1678-9741-2021-0547

Dear Editor,

Predisposition to deep sternal wound infection (DSWI) following off-pump coronary artery bypass (OPCAB) grafting surgery classifies as an area of particular research interest. Given the fact that a sound evaluation of the risk factors for DSWI mandates a comprehensive approach, we wish to highlight a few important concerns pertaining to the Engineev et al.<sup>[1]</sup> study recently published in the *Brazilian Journal of Cardiovascular Surgery*.

Interestingly, the index analysis does not outline diabetes mellitus as a preoperative risk factor for DSWI (30.2% in DSWI and 26.2% in non DSWI group, P=0.5)<sup>[1]</sup> albeit the lack of data on perioperative glycaemic control deserves attention. Appropriate to the context, the Mayo Clinic research group delineate as high as 30% increase in adverse outcomes, including infective complications for every 20 mg/dL rise in mean intraoperative glucose levels<sup>[2]</sup>. Moreover, specific literature linking glycaemic fluctuations with infective complications continues to accumulate over the past decade<sup>[3,4]</sup>. Järvelä et al.<sup>[3]</sup> found a significantly heightened rate of postoperative infections in their cardiac surgical cohort manifesting repeated hyperglycaemia (39.7% incidence) as opposed to normoglycaemic or those with single hyperglycaemic episode (12.1% and 8.2%, respectively, P=0.019).

Furnary et al.<sup>[4]</sup> reveal the independent DSWI predictive ability of post-cardiac surgery hyperglycaemia in the Portland Diabetic Project, wherein the subset with 48-hour mean blood glucose levels >200 mg/dL demonstrated a 2.2 times elevated risk of DSWI. Concomitantly, there is convincing evidence to suggest that perioperative glucose control with insulin infusion management protocols considerably attenuate the DSWI

dynamic research area in a more holistic manner.

Correspondence Address:

### Jes Jose<sup>1</sup>, MD

<sup>1</sup>Department of Cardiac Anesthesiology, Sri Jayadeva Institute of Cardiovascular Sciences and Research, Bengaluru, India

incidence<sup>[4-6]</sup>. Alongside the absence of perioperative glucose

data, Enginoev et al.[1] fail to describe the glucose management

In addition, the authors could have elaborated whether or

not any of the study participants were receiving preoperative

corticosteroids<sup>[1,7]</sup>. Herein, a comparative account of the preoperative leucocytic counts of the DSWI and non-DSWI

groups could also have added incremental value<sup>[1,8]</sup>. As much

as we laud the endeavours of Engineev et al.[1], the points of

perioperative relevance elucidated by us and the authors'

explanation would probably assist readers to comprehend this

regime employed in their retrospective study[1].

### Rohan Magoon<sup>2</sup>, DM, MD

https://orcid.org/0000-0003-4633-8851

E-mail: rohanmagoon21@gmail.com

<sup>2</sup>Department of Cardiac Anaesthesia, Atal Bihari Vajpayee Institute of Medical Sciences (ABVIMS) and Dr. Ram Manohar Lohia Hospital, Baba Kharak Singh Marg, New Delhi, India.

## Jasvinder Kaur Kohli<sup>2</sup>, MD

<sup>2</sup>Department of Cardiac Anaesthesia, Atal Bihari Vajpayee Institute of Medical Sciences (ABVIMS) and Dr. Ram Manohar Lohia Hospital, Baba Kharak Singh Marg, New Delhi, India.

### Ramesh Kashav<sup>2</sup>, MD

<sup>2</sup>Department of Cardiac Anaesthesia, Atal Bihari Vajpayee Institute of Medical Sciences (ABVIMS) and Dr. Ram Manohar Lohia Hospital, Baba Kharak Singh Marg, New Delhi India

# **REFERENCES**

- Enginoev S, Rad AA, Ekimov S, Kondrat'ev D, Magomedov G, et al. Risk Factors for Deep Sternal Wound Infection after Off-Pump Coronary Artery Bypass Grafting: a Case-Control Study [ahead of print] Braz J Cardiovasc Surg 2021 DOI: 10.21470/1678-9741-2020-0444
- 2. Gandhi GY, Nuttall GA, Abel MD, Mullany CJ, Schaff HV, et al. Intraoperative hyperglycemia and perioperative outcomes in cardiac surgery patients. Mayo Clin Proc 2005; 80: 862-866
- Järvelä KM, Khan NK, Loisa EL, Sutinen JA, Laurikka JO, Khan JA. Hyperglycemic Episodes Are Associated With Postoperative Infections After Cardiac Surgery. Scand J Surg 2018;107:138-144
- 4. Furnary AP, Wu Y, Bookin SO. Effect of hyperglycemia and continuous intravenous insulin infusions on outcomes of cardiac

- surgical procedures: the Portland Diabetic Project. Endocr Pract 2004;10 Suppl 2:21-33
- Kramer R, Groom R, Weldner D, Gallant P, Heyl B, et al. Glycemic control and reduction of deep sternal wound infection rates: a multidisciplinary approach. Arch Surg 2008; 143:451-456
- Cotogni P, Barbero C, Rinaldi M. Deep sternal wound infection after cardiac surgery: Evidences and controversies. World J Crit Care Med 2015; 4:265-273
- 7. Magoon R, Choudhury A, Sahoo S, Malik V. Steroids for adult cardiac surgery: The debate echoes on. J Anaesthesiol Clin Pharmacol 2019; 35:560-562
- 8. Dey S, Kashav R, Kohli JK, Magoon, R., ItiShri, et al. Systemic Immune-Inflammation Index Predicts Poor Outcome After Elective Off-Pump CABG: A Retrospective, Single-Center Study. J Cardiothorac Vasc Anesth 2021; 35:2397-2404



This is an open-access article distributed under the terms of the Creative Commons Attribution License.