



Brazilian Journal of Cardiovascular Surgery

ISSN: 0102-7638

ISSN: 1678-9741

Sociedade Brasileira de Cirurgia Cardiovascular

Gasperi, Ricardo de; Bodanese, Luiz Carlos; Guaragna, João Carlos
Vieira da Costa; Wagner, Mario Bernardes; Albuquerque, Luciano Cabral
Proposed Risk Score in Patients with Aortic Stenosis Submitted to Valve Replacement Surgery
Brazilian Journal of Cardiovascular Surgery, vol. 38, no. 2, 2023, March-April, pp. 219-226
Sociedade Brasileira de Cirurgia Cardiovascular

DOI: <https://doi.org/10.21470/1678-9741-2022-0254>

Available in: <https://www.redalyc.org/articulo.oa?id=398974952003>

- ▶ [How to cite](#)
- ▶ [Complete issue](#)
- ▶ [More information about this article](#)
- ▶ [Journal's webpage in redalyc.org](#)

The logo for Redalyc.org, featuring the text 'redalyc.org' in a stylized font with a red heart icon to its left.

Scientific Information System Redalyc

Network of Scientific Journals from Latin America and the Caribbean, Spain and Portugal

Project academic non-profit, developed under the open access initiative

Proposed Risk Score in Patients with Aortic Stenosis Submitted to Valve Replacement Surgery

Ricardo de Gasperi¹, PhD; Luiz Carlos Bodanese², PhD; João Carlos Vieira da Costa Guaragna^{2,3}, PhD; Mario Bernardes Wagner⁴, PhD; Luciano Cabral Albuquerque⁵, MD; PhD

¹Department of Interventional Cardiology, Associação Dr. Bartholomeu Tacchini, Bento Gonçalves, Rio Grande do Sul, Brazil.

²Department of Cardiology, Hospital São Lucas, Faculdade de Medicina, Pontifícia Universidade Católica do Rio Grande do Sul, Porto Alegre, Rio Grande do Sul, Brazil.

³Department of Cardiology, Hospital Divina, Porto Alegre, Rio Grande do Sul, Brazil.

⁴Department of Postgraduate Program Stricto Sensu in Medicine and Health Sciences, Faculdade de Medicina, Pontifícia Universidade Católica do Rio Grande do Sul, Porto Alegre, Rio Grande do Sul, Brazil.

⁵Department of Cardiac Surgery, Hospital São Lucas, Pontifícia Universidade Católica do Rio Grande do Sul, Porto Alegre, Rio Grande do Sul, Brazil.

This study was carried out at the Hospital São Lucas, Pontifícia Universidade Católica do Rio Grande do Sul, Porto Alegre, Rio Grande do Sul, Brazil.

ABSTRACT

Introduction: Due to Brazilian population aging, prevalence of aortic stenosis, and limited number of scores in literature, it is essential to develop risk scores adapted to our reality and created in the specific context of this disease.

Methods: This is an observational historical cohort study with analysis of 802 aortic stenosis patients who underwent valve replacement at Hospital São Lucas, Pontifícia Universidade Católica do Rio Grande do Sul, from 1996 to 2018. With the aid of logistic regression, a weighted risk score was constructed based on the magnitude of the coefficients β of the logistic equation. Two performance statistics were obtained: area under the receiver operating characteristic curve and the chi-square (χ^2) of Hosmer-Lemeshow goodness-of-fit with Pearson's correlation coefficient between the observed events and predicted as a model calibration estimate.

Results: The risk predictors that composed the score were valve replacement surgery combined with coronary artery bypass grafting, prior renal failure, New York Heart Association class III/IV heart failure, age > 70 years, and ejection fraction < 50%. The receiver operating characteristic curve area was 0.77 (95% confidence interval: 0.72-0.82); regarding the model calibration estimated between observed/predicted mortality, Hosmer-Lemeshow test $\chi^2 = 3,70$ ($P=0.594$) and Pearson's coefficient $r = 0.98$ ($P<0.001$).

Conclusion: We propose the creation of a simple score, adapted to the Brazilian reality, with good performance and which can be validated in other institutions.

Keywords: Risk Factors. Aortic Valve Stenosis. Heart Valves. Coronary Artery Bypass. Aging.

Abbreviations, Acronyms & Symbols

AUC	= Area under the curve
AVR	= Aortic valve replacement
CABG	= Coronary artery bypass grafting
CI	= Confidence interval
COPD	= Chronic obstructive pulmonary disease
EF	= Ejection fraction
EuroSCORE	= European System for Cardiac Operative Risk Evaluation
NYHA	= New York Heart Association
OR	= Odds ratio
PUCRS	= Pontifícia Universidade Católica do Rio Grande do Sul
ROC	= Receiver operating characteristic
STS	= Society of Thoracic Surgeons
TAVI	= Transcatheter aortic valve implantation

INTRODUCTION

Aortic stenosis presents a growing prevalence as a consequence of life expectancy increasing and natural population aging, having as a main cause the aortic calcification^[1,2]. It is estimated a prevalence of 0.2% among adults and 2.8% in patients > 75 years of age^[1]. Historically, the treatment of choice is aortic valve replacement surgery, but transcatheter aortic valve implantation (TAVI) has expanded its indications according to the latest guidelines, based mainly on the classification of surgical risk^[2-4]. The importance of estimating the surgical risk in aortic stenosis patients, who are candidates for the intervention, is mandatory, because it suggests the risk of death and it also implies the kind of intervention to be performed.

The surgical risk scores most widely used and mentioned by the guidelines are the European System for Cardiac Operative Risk Evaluation (EuroSCORE) and the Society of Thoracic Surgeons (STS) Score^[5-10]. EuroSCORE is a death score based mostly on European

Correspondence Address:

Ricardo de Gasperi

 <https://orcid.org/0000-0002-1982-4523>

Associação Dr. Bartholomeu Tacchini

Rua José Mario Mônaco, 358, Bairro Centro, Bento Gonçalves, RS, Brazil

Zip Code: 95700-068

E-mail: ricardohemodinamica@hotmail.com

Article received on July 6th, 2022.
Article accepted on August 3rd, 2022.

patients and formed by different types of cardiac surgeries^[8-10]. On the other hand, STS Score is an American morbidity and mortality score, made of three big cohorts: coronary artery bypass grafting (CABG), valve surgeries, and valve surgeries combined with CABG^[5-7]. In the Brazilian reality and in the context of valve surgery, there is the Guaragna Score, a risk of death score created in a single center and already validated in other services^[11]. There are several predictors that add mortality risk to cardiac surgery, according to previous studies, such as: advanced age, female gender, diabetes, renal failure, stroke, pulmonary hypertension, advanced functional class of heart failure, endocarditis, hypertension, chronic obstructive pulmonary disease (COPD), atrial fibrillation, previous cardiac surgery, urgent and emergency surgeries, patients critical condition, and degree of ventricular dysfunction^[5-9,11-16]. Due to population aging, prevalence of aortic stenosis, and limited number of scores in literature, especially in the Brazilian reality, is extremely important to develop risk scores adapted to our reality and, mainly, created in the specific scope of this pathology.

METHODS

Study Design

This is an observational study of a historical cohort based on the database of the postoperative cardiac surgery unit at Hospital São Lucas, Pontifícia Universidade Católica do Rio Grande do Sul (PUCRS), in accordance with the principles established in the Declaration of Helsinki and approved by the Research Ethics Committee of Faculdade de Medicina – PUCRS, under the registration number 2,796,970.

Population

Between January 1996 and July 2018, 6,658 patients underwent cardiac surgery at Hospital São Lucas, PUCRS. Of these, 802 patients aged > 18 years and with aortic stenosis who underwent aortic valve replacement alone or in combination with CABG were included in the analysis. Exclusion criteria were patients undergoing aortic valve replacement combined with another valve approach, aortic approaches or associated myectomy, and emergency/urgent surgeries (all 20 cases excluded by urgency were due to acute coronary syndrome) (Figure 1).

Analyzed Variables

The variables initially tested in the statistical analyses were: age, gender, heart failure according to functional class by the New York Heart Association (NYHA) classification, presence of atrial fibrillation, stroke, diabetes mellitus, hypertension, previous heart surgery, chronic obstructive pulmonary disease (diagnosis through clinical data, imaging methods, spirometry, or use of continuous medications), pulmonary arterial hypertension (pulmonary artery systolic pressure > 30 mmHg), current or recent endocarditis (last 60 days), obesity (classified as body mass index ≥ 30 kg/m²), ejection fraction (EF) measured by echocardiography, previous chronic renal failure (creatinine ≥ 1.5 mg/dL), hemodialysis, and isolated or concomitant aortic surgery with CABG.

Outcome

The analyzed outcome was death, being considered during the intraoperative period and throughout the hospitalization period.




Fig. 1 - Diagram of patients included in the analysis (n=802).

Statistical Analysis

Continuous variables were described as mean and standard deviation and compared using Student’s *t*-test, and categorical variables were described by counts and percentages, being compared by the chi-square test. The initial consideration of the variables followed a hierarchical model based on biological plausibility and previous information from the literature regarding to the relevance and strength of the associations of these potential risk factors with the occurrence of death. Once these variables were listed, multiple logistic regression was performed in a forward stepwise selection process, keeping in the model all variables with a significance level of *P*<0.05. The variable EF < 50% was also kept in the model because it is described in literature as having a strong association with the outcome. Then, a weighted risk score was constructed based on the magnitude of the β coefficients of the logistic equation. The coefficients were transformed into odds ratios (OR - exp [β]) and rounded to a whole number through the truncation process. Two performance statistics were obtained: area under the receiver operating characteristic (ROC) curve and Hosmer-Lemeshow chi-square (χ²) goodness-of-fit test with the Pearson’s correlation coefficient between the events observed and those predicted by the model as calibration estimate. The resulting logistic model followed the formula below and, unlike the score, presents direct estimates of the probability of occurrence of the outcome.

$$P(\text{events}) = 1 / 1 + \exp (- (\beta_0 + \beta_1x_1 + \dots + \beta_k x_k))$$

Data were processed and analyzed using the IBM Corp. Released 2013, IBM SPSS Statistics for Windows, version 22.0, Armonk, NY: IBM Corp.

RESULTS

In the study sample of 802 patients, 39.9% were female, with a mean (± standard deviation) age of 62.9 (±13.8) years and ranging from 18 to 91 years, finding a total death rate of 10.5%. The mortality of patients undergoing valve replacement alone was 5.9%, while in patients undergoing valve replacement associated with CABG (28.3% of the sample) it was 22.0%.

Table 1 shows all the variables studied with the univariate calculation analysis. After performing multiple logistic regression of variables, the following predictors obtained statistical significance for the construction of the score: aortic valve surgery combined with CABG, previous renal failure, NYHA class III/IV heart failure, and age > 70 years. The variable EF < 50% reached borderline values for statistical significance (OR 1.66, 95% confidence interval [CI] 0.96 – 2.86, *P*=0.07), and as a strong predictor associated with death in this group of patients, according to literature, it was included in the score composition (Table 2).

At the final risk score, aortic valve surgery combined with CABG received 3 points, previous renal failure received 2 points, heart failure class III/IV received 2 points, and age > 70 years and EF < 50% received 1 point each (Table 3).

Table 1. Univariate analysis (n=802).

Variables	Deaths n=84 (%)	Survivors n=718 (%)	OR	95% CI	P-value
Age > 70 years	51 (60.7)	253 (35.2)	2.84	1.79-4.52	< 0.001
Female	36 (42.9)	284 (39.6)	1.15	0.73-1.81	0.560
NYHA III/IV	48 (57.1)	240 (33.4)	2.66	1.68-4.20	< 0.001
Previous cardiac surgery	0 (0.0)	8 (1.1)	-	-	0.182
Atrial fibrillation	7 (8.3)	52 (7.2)	1.16	0.51-2.65	0.722
Stroke	6 (7.1)	21 (2.9)	2.55	1.00-6.52	0.071
Diabetes mellitus	25 (29.8)	112 (15.6)	2.29	1.38-3.81	0.002
Hypertension	59 (70.2)	392 (54.6)	1.96	1.20-3.20	0.005
COPD	18 (21.4)	81 (11.3)	2.14	1.21-3.79	0.013
Pulmonary hypertension	1 (1.2)	3 (0.4)	2.87	0.29-27.92	0.409
Endocarditis	1 (1.2)	3 (0.4)	2.87	0.29-27.91	0.409
Obesity	14 (16.7)	71 (9.9)	1.82	0.98-3.40	0.073
EF < 50%	28 (33.3)	125 (17.4)	2.37	1.45-3.88	0.001
Renal failure	20 (23.8)	52 (7.2)	4.00	2.25-7.12	< 0.001
Hemodialysis	2 (2.4)	1 (0.1)	17.49	1.57-194.96	0.019
CABG associated	50 (50.9)	177 (24.7)	4.49	2.81-7.17	< 0.001

CABG=coronary artery bypass grafting; CI=confidence interval; COPD=chronic obstructive pulmonary disease; EF=ejection fraction; NYHA=New York Heart Association; OR=odds ratio

Table 2. Logistic regression (n=802).

Variables	Coefficient β	OR	95% CI	P-value
CABG associated	1.35	3.84	2.35-6.28	< 0.001
Renal failure	0.88	2.42	1.29-4.54	0.006
NYHA III/IV	0.80	2.23	1.35-3.67	0.002
Age > 70 years	0.62	1.86	1.13-3.06	0.014
EF < 50%	0.51	1.66	0.96-2.86	0.070

CABG=coronary artery bypass grafting; CI=confidence interval; EF=ejection fraction; NYHA=New York Heart Association; OR=odds ratio

Table 3. Multivariate risk score (n=802).

Variables	Score
CABG associated	3
Renal failure	2
NYHA III/IV	2
Age > 70 years	1
EF < 50%	1

CABG=coronary artery bypass grafting; EF=ejection fraction; NYHA=New York Heart Association

Risk of death according to the risk score and classification (additive score) were divided into four groups — low, medium, high, and very high (Table 4).

The risk model had an accuracy measured by the area under the ROC curve of 0.77 (95% CI 0.72 – 0.82) and, therefore, had good discriminatory ability. There was also a good correlation between predicted and observed mortality: $r = 0.98$ ($P < 0.001$) with $\chi^2 = 3.70$ ($P = 0.594$) (Hosmer-Lemeshow test) (Figures 2 and 3).

DISCUSSION

This study proposed the creation of a risk score for death in patients undergoing aortic valve replacement by including five variables: aortic valve surgery combined with CABG, renal failure, NYHA class III/IV heart failure, age > 70 years, and EF < 50%.

The overall mortality obtained was 10.5%, whereas in isolated aortic valve surgery, it was 5.9%. Compared to literature data, especially in relation to European and North American data, it was reported a high overall mortality rate. When it was compared with STS Score cohort that involves only isolated valve surgery, it was observed a mortality of 3.4%, and 5.6% in the cohort associating aortic valve surgery with CABG^[5,6]. New York State cohort describes a 3.3% mortality for isolated aortic valve replacement and 7.1% for aortic valve replacement associated with CABG^[15]. The United Kingdom data from the Ambler Score cohort shows a mortality rate of 4.9% for aortic valve replacement and 7.9% for combined surgery with CABG^[14]. When national cohorts are analyzed, some numbers closer to this study's results can be seen. Garofallo et al.^[17] presented data from a tertiary center in the same city, also involving patients' interventions from the public and private healthcare system, and reported 8.6% mortality for valve surgery and 20% when associated with CABG. Ribeiro et al.^[18], reviewing data from > 100,000 surgeries performed in Brazil between 2000-2003, described a mortality of

8.9% for valve surgeries and 16.5% for associated surgeries. Bueno et al.^[19] presented data from the 1990s showing mortality for isolated aortic surgery of 8% and 21% for aortic surgery associated with CABG. The Brazilian registry of adult Patients undergoing cArdiovascular Surgery (or BYPASS), a recent Brazilian registry, organized by the Brazilian Society of Cardiovascular Surgery analyzing data on valve surgeries from 920 patients (80% from the public and 20% from the private healthcare system of 17 different institutions in the country), found a mortality in aortic valve replacement isolated of 5.1% and 14.7% in aortic valve replacement associated with CABG. In this analysis, when only interventions in aortic degenerative disease were evaluated, the mortality was 7.8%^[20]. It is worth emphasizing that this analysis included data from more than twenty years of interventions, representing nearly three decades and covering different stages of cardiac surgery within PUCRS hospital. Analyzing the same period, another national reference center in cardiology reported mortality in valve surgery ranging from 7.47% to 13.96%^[21].

According to the studies, it was observed that in aortic valve replacement surgery combined with CABG compared to isolated valve replacement, there is an increased risk of death. Based on the abovementioned national studies, an increase of at least two to three times in the risk of death for combined surgery can be seen^[17,18,20]. Regarding to STS Score and the United Kingdom cohorts, it was observed a less than twofold increase in associated surgery mortality compared to isolated aortic valve, and in New York State cohort, it was twice as high^[5,6,14]. It was found in this study that combined surgery was responsible for adding 3 points (OR 3.84) to the risk of death, being the main score risk predictor. Patients with only this variable would receive 3 points, being moderate risk with an estimated mortality of 6.6%. Since mortality in valve surgery associated with CABG in study cohort was much higher, most patients undergoing this intervention presented

Table 4. Risk of death according to the score (n=802).

Score	Sample	Deaths		Risk category
		n	%	
0	227	4	1.8	Low
1 to 3	362	24	6.6	Medium
4 and 5	127	25	19.7	High
6 to 9	86	31	36.0	Very high




Fig. 2 - Area under the receiver operating characteristic curve in detecting the occurrence of death (n=802). Area under the curve (AUC)=0.77 (95% confidence interval [CI] 0.72- 0.82).




Fig. 3 - Distribution of points in the predicted and observed model (n=802). Person's coefficient $r=0.98$ ($P<0.001$) and chi-square Hosmer-Lemeshow (χ^2 HL) = 3.70 ($P=0.594$).

an association of risk variables. It is highlighted that most of the national services, as well as our service, mostly assisted patients are from the public healthcare system — patients who often have serious socioeconomic problems, difficulties in optimal use of drug treatment, less access to medical care, as well as being in more advanced stages of the disease when undergoing surgical intervention^[17].

Renal failure is another common risk factor across all different risk scores^[5-9,11-16]. In this study, it was defined as a creatinine ≥ 1.5 mg/dL and it was responsible for adding 2 points to the mortality risk. There is a wide variation in the definition of renal failure within heart surgery cohorts. It is known that small increases in creatinine above normal levels and a slight decrease in the glomerular filtration rate are already associated with a worse prognosis in cardiac surgery^[22,23]. In a previous study published by this hospital group, two different risk levels were aggregated — the first, when creatinine was between 1.5 and 2.49 mg/dL, and a second, with a higher risk level, when creatinine was ≥ 2.5 mg/dl or on chronic dialysis^[11]. In the cohort study, there were only three patients on chronic dialysis, compromising the analysis of this variable as being an isolated risk predictor.

In a review by Tjang et al.^[24], NYHA class III/IV heart failure was the most mentioned risk factor among the predictors of mortality in patients undergoing aortic valve replacement. Functional class in heart failure is a strictly clinical factor, with simple assessment and easy applicability, supporting the idea that patients with aortic stenosis should undergo valve replacement before their clinical deterioration. In the study score, the advanced functional class III/IV added 2 points to the increased risk of in-hospital death, and 36% of patients were in this class. Regarding to the STS Score cohorts, there was a division of risks into two groups: first, a lower risk group involving functional classes I to III, and a second higher risk group including only functional class IV^[5,6]. The Ambler Score^[14] and EuroSCORE^[10] do not include the functional class as a risk predictor, only the degree of ventricular dysfunction. In EuroSCORE II, the functional class was incorporated, being divided into II, III, and IV according to NYHA^[9].

Age is a risk predictor mentioned by all risk scores in literature, and the cutoff point, which increases the risk of surgical mortality, varies among them^[5-9,11-16]. EuroSCORE determines an increased risk for mortality in patients > 60 years of age and adds an increase in risk every five years above this cutoff point^[8]. STS Score^[5-7]

determines an increase in risk over the age of 50, and the New York State cohort^[15] shows an increase in risk every decade over the age of 55 years. Although the surgical risk increases with advanced age compared to younger patients, the benefit in relation to medical treatment is substantially advantageous. Octogenarian surgery candidates who, for their own reasons, choose not to undergo the intervention have a mortality increase greater than ten times compared to patients who are intervened. This shows that advanced age by itself is not a contraindication to valve replacement surgery^[25]. It is also true that less invasive approaches such as TAVI and hybrid procedures with stents associated with TAVI must be considered within a general context, including age as well. The latest American Guideline (American Heart Association/American College of Cardiology) on valvulopathy places TAVI superior over valve replacement surgery in patients over 80 years of age, regardless of surgical risk^[4], and The Brazilian Guideline on Valvular Heart Disease^[2] considers TAVI implantation as a class IA indication in patients over 70 years of age, even at low surgical risk.

Ventricular dysfunction is a marker of severity in the valve surgery. EuroSCORE established two risk groups for the outcome death in cardiac surgery according to EF — EF values between 30 and 50% with an OR of 1.5 and EF < 30% with an OR of 2.5^[10]. In EuroSCORE II, the division of EF into more categories was prioritized, with EF 20-29% and < 20% being added^[9]. STS Score in the cohort of valve surgery establishes an OR of 1.09 for each decrease of 10 EF units below 50, regardless of the kind of valve surgery^[5]. Ambler score showed an OR for death of 1.2 for EF of 30-50% and an OR of 1.99 for EF < 30% when compared to EF > 50%^[14]. Guaragna et al.^[11] found an OR of 2.1 for the variable EF ≤ 45 in patients undergoing valve surgery. Specifically in patients with aortic stenosis, the German Aortic Valve Score established an OR for mortality of 1.96 in patients with EF between 30 and 50% and OR of 2.96 for EF < 30%^[12]. In this study, in the multivariate analysis, the variable EF < 50% reached borderline values in relation to its statistical significance (OR 1.66, 95% CI 0.96 – 2.86, P=0.07) and, due to its strong association described in literature, it was included in the final score, receiving 1 point. Only twenty-four patients underwent a replacement valve surgery with EF < 30%.

The surgical risk was divided into four groups, ranging from low to very high. It was emphasized that, in this sample, any of these predictors alone places the patient at medium risk and adds an important increase in the risk of death compared to the low-risk group where none of these predictors is present.

The model had an accuracy measured by the area under the ROC curve of 0.77 (95% CI 0.72 – 0.82) and, therefore, had a good discriminatory ability. There was also a good correlation between predicted and observed mortality — $r = 0.98$ ($P < 0.001$) with $\chi^2 = 3.70$ ($P = 0.594$) (Hosmer–Lemeshow test). Table 5 summarizes the main scores in literature, showing that this risk score, created from a specific group of patients with aortic stenosis, presents a superimposable accuracy.

Limitations

Related to the study limitations. Firstly, patients in need of urgent surgery were excluded from the analysis. The proposal was to create a score focused on the pathology of aortic stenosis, and the twenty cases of urgent surgery were performed on an emergency basis due to acute coronary syndrome. Secondly, variables such as previous hemodialysis, current or recent endocarditis, pulmonary hypertension, and previous cardiac surgery included a very low absolute number of patients in the sample and limited a better analysis of these variables as risk predictors. Thirdly, the risk model was created by analyzing data obtained from a single center without an internal validation group. According to literature, they tend to show lower results when applied to institutions other than where the scores were created. Thus, it is understood that this study has internal validity, however, the study researchers consider it is important a future external validation of this model in other institutions.

Regarding to practical implications of these findings, the identification of risk factors and the creation of a death risk score allow to accurately estimate the surgical risk within the institution, monitor the care quality, and implement measures for service qualification. The model has a good statistical performance and, therefore, has adequate capacity to be tested and validated in other institutions.

Table 5. Scores' accuracy comparative.

Score	AUC ROC
Gasperi	0.77
Guaragna Score ^[11]	0.83
Kotting (German Aortic Valve Score I) ^[12]	0.80
Nashef (EuroSCORE II) ^[9]	0.80
Mejía (Inscor) ^[16]	0.79
Ambler ^[14]	0.77
Roques (EuroSCORE) ^[8]	0.76
O'Brien (STS Score – AVR valve) ^[5]	0.76
Hannan ^[15]	0.76
Shahian (STS Score – AVR + CABG) ^[6]	0.74
Schiller (German Aortic Valve Score II) ^[13]	0.74

AUC=area under the curve; AVR=aortic valve replacement; CABG=coronary artery bypass grafting; EuroSCORE=European System for Cardiac Operative Risk Evaluation; ROC=receiver operating characteristic; STS=Society of Thoracic Surgeons

CONCLUSION

In this study, it is proposed an in-hospital death risk score for patients undergoing aortic valve replacement using five variables: valve replacement surgery combined with CABG, previous renal failure, presence of NYHA class III/IV heart failure, age > 70 years, and EF < 50%. Also, this is a simple score, with good statistical performance, and adapted to the Brazilian reality.

No financial support.

No conflict of interest.

Authors' Roles & Responsibilities

RG	Substantial contributions to the conception or design of the work; or the acquisition, analysis, or interpretation of data for the work; drafting the work or revising it critically for important intellectual content; agreement to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved; final approval of the version to be published
LCB	Substantial contributions to the conception or design of the work; or the acquisition, analysis, or interpretation of data for the work; drafting the work or revising it critically for important intellectual content; agreement to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved; final approval of the version to be published
JCVCG	Substantial contributions to the conception or design of the work; or the acquisition, analysis, or interpretation of data for the work; drafting the work or revising it critically for important intellectual content; agreement to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved; final approval of the version to be published
MBW	Substantial contributions to the conception or design of the work; or the acquisition, analysis, or interpretation of data for the work; agreement to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved; final approval of the version to be published
LCA	Drafting the work or revising it critically for important intellectual content; agreement to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved; final approval of the version to be published

REFERENCES

1. Otto CM, Prendergast B. Aortic-valve stenosis—from patients at risk to severe valve obstruction. *N Engl J Med.* 2014;371(8):744-56. doi:10.1056/NEJMra1313875.
2. Tarasoutchi F, Montera MW, Ramos AIO, Sampaio RO, Rosa VEE, Accorsi TAD, et al. Update of the Brazilian guidelines for valvular heart disease - 2020. *Arq Bras Cardiol.* 2020;115(4):720-75. doi:10.36660/abc.20201047.
3. Baumgartner H, Falk V, Bax JJ, De Bonis M, Hamm C, Holm PJ, et al. 2017 ESC/EACTS guidelines for the management of valvular heart disease. *Eur Heart J.* 2017;38(36):2739-91. doi:10.1093/eurheartj/ehx391.
4. Otto CM, Nishimura RA, Bonow RO, Carabello BA, Erwin JP 3rd, Gentile F, et al. 2020 ACC/AHA guideline for the management of patients with

valvular heart disease: a report of the American college of cardiology/ American heart association joint committee on clinical practice guidelines. *Circulation.* 2021;143(5):e72-e227. Erratum in: *Circulation.* 2021;143(5):e229. doi:10.1161/CIR.0000000000000923.

5. O'Brien SM, Shahian DM, Filardo G, Ferraris VA, Haan CK, Rich JB, et al. The society of thoracic surgeons 2008 cardiac surgery risk models: part 2--isolated valve surgery. *Ann Thorac Surg.* 2009;88(1 Suppl):S23-42. doi:10.1016/j.athoracsur.2009.05.056.
6. Shahian DM, O'Brien SM, Filardo G, Ferraris VA, Haan CK, Rich JB, et al. The society of thoracic surgeons 2008 cardiac surgery risk models: part 3--valve plus coronary artery bypass grafting surgery. *Ann Thorac Surg.* 2009;88(1 Suppl):S43-62. doi:10.1016/j.athoracsur.2009.05.055.
7. Shahian DM, O'Brien SM, Filardo G, Ferraris VA, Haan CK, Rich JB, et al. The society of thoracic surgeons 2008 cardiac surgery risk models: part 1--coronary artery bypass grafting surgery. *Ann Thorac Surg.* 2009;88(1 Suppl):S2-22. doi:10.1016/j.athoracsur.2009.05.053.
8. Roques F, Michel P, Goldstone AR, Nashef SA. The logistic EuroSCORE. *Eur Heart J.* 2003;24(9):881-2. doi:10.1016/s0195-668x(02)00799-6.
9. Nashef SA, Roques F, Sharples LD, Nilsson J, Smith C, Goldstone AR, et al. EuroSCORE II. *Eur J Cardiothorac Surg.* 2012;41(4):734-44; discussion 744-5. doi:10.1093/ejcts/ezs043.
10. Nashef SA, Roques F, Michel P, Gauducheau E, Lemeshow S, Salamon R. European system for cardiac operative risk evaluation (EuroSCORE). *Eur J Cardiothorac Surg.* 1999;16(1):9-13. doi:10.1016/s1010-7940(99)00134-7.
11. Guaragna JC, Bodanese LC, Bueno FL, Goldani MA. Proposta de escore de risco pré-operatório para pacientes candidatos à cirurgia cardíaca valvar. *Arq Bras Cardiol.* 2010;94(4):541-8. doi:10.1590/s0066-782x2010005000026.
12. Kötting J, Schiller W, Beckmann A, Schäfer E, Döbler K, Hamm C, et al. German aortic valve score: a new scoring system for prediction of mortality related to aortic valve procedures in adults. *Eur J Cardiothorac Surg.* 2013;43(5):971-7. doi:10.1093/ejcts/ezt114.
13. Schiller W, Barnewold L, Kazmaier T, Beckmann A, Masseli F, Welz A, et al. The German aortic valve score II. *Eur J Cardiothorac Surg.* 2017;52(5):881-7. doi:10.1093/ejcts/ezx282.
14. Ambler G, Omar RZ, Royston P, Kinsman R, Keogh BE, Taylor KM. Generic, simple risk stratification model for heart valve surgery. *Circulation.* 2005;112(2):224-31. doi:10.1161/CIRCULATIONAHA.104.515049.
15. Hannan EL, Racz MJ, Jones RH, Gold JP, Ryan TJ, Hafner JP, et al. Predictors of mortality for patients undergoing cardiac valve replacements in New York state. *Ann Thorac Surg.* 2000;70(4):1212-8. doi:10.1016/s0003-4975(00)01968-8.
16. Mejía OA, Lisboa LA, Puig LB, Moreira LF, Dallan LA, Pomerantzeff PM, et al. InsCor: a simple and accurate method for risk assessment in heart surgery. *Arq Bras Cardiol.* 2013;100(3):246-54. doi:10.5935/abc.20130043.
17. Garofallo SB, Machado DP, Rodrigues CG, Bordim O Jr, Kalil RA, Portal VL. Applicability of two international risk scores in cardiac surgery in a reference center in Brazil. *Arq Bras Cardiol.* 2014;102(6):539-48. doi:10.5935/abc.20140064.
18. Ribeiro AL, Gagliardi SP, Nogueira JL, Silveira LM, Colosimo EA, Lopes do Nascimento CA. Mortality related to cardiac surgery in Brazil, 2000-2003. *J Thorac Cardiovasc Surg.* 2006;131(4):907-9. doi:10.1016/j.jtcvs.2005.11.022.
19. Bueno RM, Neto VÂ, Melo R. Fatores de risco em operações valvares: análise de 412 casos. *Braz J Cardiovasc Surg.* 1997;12(4):348-58. doi:10.1590/S0102-76381997000400007.
20. Zilli AC, Guizilini S, Rocco IS, Santo JADE, Berwanger O, Kalil RAK, et al. Valve heart surgery in Brazil - the BYPASS registry analysis. *Braz J Cardiovasc Surg.* 2020;35(1):82-90. doi:10.21470/1678-9741-2019-0408.
21. Mejia OAV, Lisboa LAF, Caneo LF, Arita ET, Brandão CMA, Dias RR, et al. Analysis of >100,000 cardiovascular surgeries performed at the heart institute and a new era of outcomes. *Arq Bras Cardiol.* 2020;114(4):603-612. doi:10.36660/abc.20190736.
22. Machado MN, Nakazone MA, Maia LN. Acute kidney injury based on

- KDIGO (kidney disease improving global outcomes) criteria in patients with elevated baseline serum creatinine undergoing cardiac surgery. *Rev Bras Cir Cardiovasc.* 2014;29(3):299-307. doi:10.5935/1678-9741.20140049.
23. Anderson RJ, O'Brien M, MaWhinney S, VillaNueva CB, Moritz TE, Sethi GK, et al. Mild renal failure is associated with adverse outcome after cardiac valve surgery. *Am J Kidney Dis.* 2000;35(6):1127-34. doi:10.1016/s0272-6386(00)70050-3.
 24. Tjang YS, van Hees Y, Körfer R, Grobbee DE, van der Heijden GJ. Predictors of mortality after aortic valve replacement. *Eur J Cardiothorac Surg.* 2007;32(3):469-74. doi:10.1016/j.ejcts.2007.06.012.
 25. Kojodjojo P, Gohil N, Barker D, Youssefi P, Salukhe TV, Choong A, et al. Outcomes of elderly patients aged 80 and over with symptomatic, severe aortic stenosis: impact of patient's choice of refusing aortic valve replacement on survival. *QJM.* 2008;101(7):567-73. doi:10.1093/qjmed/hcn052.

