

The logo for the journal Therya, featuring the word "Therya" in a serif font, underlined, and enclosed in a rounded rectangular border.

Therya

ISSN: 2007-3364

Asociación Mexicana de Mastozoología A. C.

Gutiérrez-Molina, Rigoberto; Cruz-Romero, Anabel; Romero-Salas, Dora;
Ballados-González, Gerardo; Jiménez-Hernández, José A.; Ibarra-
Priego, Nelly; Serna-Lagunes, Ricardo; Sánchez-Montes, Sokani
Molecular evidence for the presence of *Leptospira borgpetersenii*
in synanthropic rodents in the Nautla region, Veracruz, México

Therya, vol. 10, no. 2, 2019, pp. 171-174
Asociación Mexicana de Mastozoología A. C.

DOI: <https://doi.org/10.12933/therya-19-723>

Available in: <https://www.redalyc.org/articulo.oa?id=402362668012>

- How to cite
- Complete issue
- More information about this article
- Journal's webpage in redalyc.org

UDEM The logo for redalyc.org, featuring the text "redalyc.org" in a stylized font with a red dot above the "i".

Scientific Information System Redalyc

Network of Scientific Journals from Latin America and the Caribbean, Spain and
Portugal

Project academic non-profit, developed under the open access initiative

Molecular evidence for the presence of *Leptospira borgpetersenii* in synanthropic rodents in the Nautla region, Veracruz, México

RIGOBERTO GUTIÉRREZ-MOLINA¹, ANABEL CRUZ-ROMERO¹, DORA ROMERO-SALAS¹, GERARDO BALLADOS-GONZÁLEZ¹, JOSÉ A. JIMÉNEZ-HERNÁNDEZ¹, NELLY IBARRA-PRIEGO¹, RICARDO SERNA-LAGUNES² AND SOKANI SÁNCHEZ-MONTES^{3*}

¹ Facultad de Medicina Veterinaria y Zootecnia, región Veracruz. Universidad Veracruzana. Miguel Ángel de Quevedo s/n, CP. 91710. Veracruz, México. Email: iori_luna25@hotmail.com (RGM), anabcruz@uv.mx (ACR), dromero@uv.mx (DRS), balladosgerardo506@hotmail.com (GGBG), mvzjimenez.850114@gmail.com (JAJH), nellydeljesus@yahoo.com.mx (NIP).

² Facultad de Ciencias Biológicas y Agropecuarias, región Orizaba-Córdoba. Universidad Veracruzana. Carretera Peñuelas Amatlán Kilómetro 177, CP. 94500. Veracruz, México. Email: rserna@uv.mx (RSL).

³ Centro de Medicina Tropical, Unidad de Medicina Experimental, Facultad de Medicina. Universidad Nacional Autónoma de México. Dr. Balmis 148, CP. 06726, Ciudad de México. México. Email: sok10108@gmail.com (SSM).

*Corresponding author

The genus *Leptospira* encompasses ten species of spirochetes capable of infecting mammals, particularly rodents. In México, studies focused on the detection of *Leptospira* sp. in rodents are scarce, all of them restricted to three states of the Gulf of México. For this reason, this work aimed to identify the diversity of *Leptospira* species associated with synanthropic rodents in Veracruz, a state where leptospirosis is endemic. Rodents were sampled with Sherman traps placed in 10 Production Units across the Nautla region. Animals were euthanized and their kidneys removed. Subsequently, a 474-bp segment of the outer membrane protein LipL32, present in all pathogenic species, was amplified and sequenced. Sequences were compared vs. reference using the BLAST algorithm: a phylogenetic reconstruction was carried out using the Maximum Likelihood method. In addition, the prevalence of infection in each Production Unit was estimated. Twenty eight rodents of a single species (*Mus musculus*) were caught. *Leptospira* DNA was detected in 17 samples (62.9 %, CI_{95%} 42.3 to 80.6) from seven localities in the Nautla region. The sequences recovered exhibited 99-100% identity to each other and 99 % identity with *Leptospira borgpetersenii* sequences deposited in GenBank. This study confirms the presence of *L. borgpetersenii* in rodents, particularly in *M. musculus*, in México. This study increases the inventory of pathogenic leptospires for the state of Veracruz to three species.

El género *Leptospira* engloba 10 especies de espiroquetas capaces de infectar mamíferos, particularmente roedores. En México se han realizado escasos estudios para la detección de *Leptospira* sp. en roedores, todos ellos restringidos a tres estados del Golfo de México. Por tal motivo el objetivo del presente trabajo fue identificar la diversidad de leptospirosis en roedores sinantrópicos de Veracruz, un estado endémico de leptospirosis. Para la colecta de roedores, se colocaron trampas tipo Sherman en 10 unidades de producción de la región Nautla. Los animales se sacrificaron y se obtuvieron los riñones. Posteriormente se amplificó y secuenció un segmento de 474 pb de la proteína exterior de membrana LipL32 presente en las leptospirosis patógenas. Posteriormente se compararon las secuencias con las de referencia mediante el uso del algoritmo BLAST y se realizó una reconstrucción filogenética mediante el método de Máxima Verosimilitud. Adicionalmente se obtuvieron las prevalencias de la infección por unidad de producción. Se colectaron 28 roedores de una única especie (*Mus musculus*). Se detectó la presencia de ADN de *Leptospira* en 17 muestras (62.9 %; IC_{95%} 42.3 a 80.6) procedentes de siete localidades de la región Nautla. Las secuencias recuperadas exhibieron una similitud del 99-100 % entre sí y una identidad del 99 % con secuencias de referencia de *Leptospira borgpetersenii* depositadas en GenBank. Este estudio confirma la presencia de *L. borgpetersenii* en roedores y en particular con *M. musculus* en México. Este estudio incrementa a tres especies el inventario de leptospirosis patógenas para el estado de Veracruz.

Keywords: México; pathogens; small mammals; spirochetes.

© 2019 Asociación Mexicana de Mastozoología, www.mastozoologiamexicana.org

Introduction

The genus *Leptospira* encompasses a set of thin bacteria with hook-shaped tips that inhabit lentic habitats worldwide, particularly wetlands (Levett 2015). Previously, the existence of two species was recorded: a saprophytic species thriving in aquatic environments (*L. biflexa*) and a pathogenic one causing a zoonosis called leptospirosis (*L. interrogans*; Levett 2015). Today, 35 species are acknowledged, classified into three groups based on their ability to infect vertebrate hosts: non-pathogenic (e. g., *L. biflexa*, *L. idonoi*, and *L. meyeri*), facultative pathogenic (e. g., *L. broomii*, *L. fainei*, and *L. wolffii*) and pathogenic (e. g., *L. alexanderi*, *L. borgpetersenii*, *L. interrogans*, and *L. kirschneri*; Bourhy et al. 2014; Thibeaux et al. 2018).

Ten pathogenic species are currently known, which have been reported in more than 160 species of wild and domestic mammals around the world (Ballados-González et al. 2018). In particular, rodents play a role as reservoirs; many murine species exhibit chronic infections and continually release bacteria in urine, a phenomenon called leptospiuria. Members of the family Muridae serve as the main reservoirs of *Leptospira* sp., and are deemed responsible for spreading the infection to livestock, pets, and humans (da Silva et al. 2010; Colombo et al. 2018).

In México, few studies have been conducted to detect *Leptospira* sp. in rodents, mostly in the states of Tamaulipas, Campeche, and Yucatán (Méndez et al. 2013; Espinosa-Martínez et al. 2015; Torres-Castro et al. 2016; Panti-May et

al. 2017; Torres-Castro et al. 2018). These studies provide molecular evidence of the presence of *L. interrogans* and *L. kirschneri* in various species of rodents of the families Cricetidae, Heteromyidae, and Muridae. However, there are other states of México where human and bovine leptospirosis is a major public health issue. Specifically, the state of Veracruz records an incidence of human leptospirosis of five cases per 100,000 inhabitants, and seroprevalence in livestock ranging between 60 % and 80 % (Moles-Cervantes et al. 2002; Álvarez et al. 2005; Sánchez-Montes et al. 2015; Zárate-Martínez et al. 2015). However, there is no evidence on the *Leptospira* species maintained in rodent populations in the region. The aim of the present work was to determine the presence and diversity of *Leptospira* sp. in synanthropic rodents in the Nautla region, Veracruz, México.

Materials and Methods

This study was conducted in well-defined and delimited areas that have facilities, machinery, and equipment to carry out livestock activities, named Livestock Production Units (LPU), in the municipalities of San Rafael, Nautla, Martínez de la Torre, Vega de Alatorre, and Misantla, in the Nautla region, Veracruz (Figure 1). This region is located in the center-northern portion of the State of Veracruz. It is bordered by the Totonac region to the north, the Capital and Mountain regions to the south, the Gulf of México to the east, and the state of Puebla to the west. This region covers an area of 3,329 km², with 86.1 % dedicated to farming activities; in turn, 43.2 % is covered by pastures for cattle raising.

Rodents were sampled from November 2016 to May 2017 on Production Units across the Nautla region, Veracruz. In each Production Unit, Sherman® traps were placed (8×9×23 cm), using an oat-vanilla mixture as bait attractant. Forty traps were placed per Production Unit, strategically distributed in areas with high probability of occurrence of rodents such as warehouses, farmyards, or inside households. Traps were placed in the afternoon and reviewed the next morning (before 7 am) during two trapping nights

in each locality. The rodents captured were removed from traps, identified, and processed for kidney sample collection following biosafety standards, under collection license FAUT-0250 granted by the Secretariat of Environment and Natural Resources (Semarnat). Animals were euthanized according to the protocol established by NOM-033-SAG/ZOO-2014 using ketamine (Wildlife Pharmaceuticals México SA de CV 04930, México) as anesthetic agent, followed by cervical dislocation. Each rodent was placed in supine position and the absence of reflexes (corneal and podal) was confirmed before dissection to remove the kidneys. Kidney samples were placed in containers with 70 % alcohol and kept at 4 °C until processing.

DNA extraction was carried out in each sample separately, using 500 µl of a 10 % solution of the resin Chelex 100 added with 20 µl of proteinase K; then samples were incubated at 56 °C for two hours (Ballados-González et al. 2018). Afterward, samples were centrifuged at 15,000 rpm for 15 minutes; the supernatant was transferred to new tubes and stored at -20 °C.

Once the sample was obtained, a 474-bp segment of the outer membrane protein *LipL32*, present in the genome of the pathogenic *Leptospira* species, was amplified using the oligonucleotides (ATCTCCGTTGCACTCTTTGC) and *LipL32* reverse (GTCCGCCTACACACCCTTTAC; Vital-Brazil et al. 2010). The reaction mixture consisted of 12.5 µl of a 2X solution of GoTaq® Green Master Mix (Promega Corporation, Madison, WI, USA), 1 µl of each oligonucleotide (2µM each), 6.5 µl of DNase-free water, and 4 µl of DNA (200-300 ng) to make a final volume of 25 µl (Espinosa-Martínez et al. 2015; Ballados-González et al. 2018). Amplicons were visualized on agarose gels using 2 % TAE buffer at 85 V for 45 min.

Positive PCR products were sent to the Biology Institute at Universidad Nacional Autónoma de México for sequencing. The resulting sequences were compared with those deposited in GenBank to determine the similarity between them using the BLAST tool.

Global alignments were carried out between the sequences produced in this study and some representative pathogenic leptospires deposited in GenBank, with the algorithm Clustal W using the software MEGA 6.0 (Tamura et al. 2013). The Tamura's three-parameter model of nucleotide substitution (T92) was selected based on the lowest Bayesian Information Criterion (BIC) score (2,497,948). In addition, a phylogenetic reconstruction was conducted using the maximum likelihood (ML) approach; the support of the topology was validated with 10,000 bootstrap replicates, also in MEGA 6.0.

Results

A total of 28 specimens of *Mus musculus* were collected. None of the collected animals showed signs of disease at the time of collection, nor gross evidence of renal impairment. *Leptospira* DNA was detected in 17 of the 28 samples analyzed (62.9 %; CI_{95%} 42.3 to 80.59). Positive samples




Figure 1. Map of the location of the Production Units sampled in the Nautla region, Veracruz, Mexico. The municipalities that make up the Nautla region are highlighted in green. The localities sampled are marked with blue circles.

were obtained from animals collected in the localities of La Providencia, La Esperanza, Santa Julia, El Pozón, El Laurel, El Cabellal, and Tres Marías (Table 1).

The localities with the highest number of positive mice were Santa Julia (municipality of Misantla), and El Cabellal (municipality of San Rafael), with four positive specimens each (Table 1). Of the 17 positive PCR products, 12 sequences of 460 to 470 base pairs were recovered. The recovered sequences exhibited 99 % identity between them (457/460 bp), and 99 % identity (458/460 bp) with sequences of *L. borgpetersenii* deposited in GenBank. In addition, the phylogenetic analysis encompassed the sequences observed in this study along with the reference sequence of *L. borgpetersenii* in a monophyletic group with a support value of 100 (Figure 2). The sequences generated in this study were deposited in GenBank with access numbers MK568973–MK568984.

Discussion

This work is the first approximation to the study of *Leptospira* in rodents in the northern part of the state of Veracruz. Besides, it represents the first molecular confirmation of the presence of *L. borgpetersenii* in rodents, particularly in *Mus musculus*, in México (Espinosa-Martínez et al. 2015; Torres-Castro et al. 2016, 2018; Panti-May et al. 2017). The reference serovar of *L. borgpetersenii* isolated from *M. musculus* is Ballum, which was detected in the past century in Europe (Yager et al. 1953). Since then, multiple serological and molecular studies have shown that this serovar is widely distributed across *M. musculus* populations worldwide (da Silva et al. 2010; Matsui et al. 2015; Colombo et al. 2018). In México, studies conducted in the state of Durango have reported titers of antibodies to the *L. borgpetersenii* Ballum serovar in domestic animals such as pigs and donkeys (Alvarado-Esquivel et al. 2018; Cruz-Romero et al. 2018). Thus, it is reasonable to assume that domestic animals can be exposed to bacteria through rodents and food or water sources contaminated with rodent urine in farmyards or livestock handling and processing areas.

Table 1. Location of the Production Units sampled in the Nautla region, Veracruz, Mexico. RC = Rodents collected; PR = Positive rodents; % = Prevalence

Locality	Municipality	Latitude	Longitude	RC	PR	%
La Providencia	Vega de Alatorre	19° 56' 0.47"	-96° 32' 59.02"	2	1	50
El Ciervo	Vega de Alatorre	20° 02' 00.31"	-96° 45' 18.97"	1	0	0
La Esperanza	Vega de Alatorre	19° 59' 53.26"	-96° 44' 13.48"	2	1	50
Santa Julia	Misantla	19° 52' 00.26"	-96° 48' 38.30"	5	4	80
El Pozón	Misantla	19° 55' 03.14"	-96° 51' 09.71"	5	2	40
El Laurel	Vega de Alatorre	20° 07' 04.62"	-96° 40' 53.18"	1	1	100
El Paraíso	San Rafael	20° 13' 51.00"	-96° 48' 39.09"	3	0	0
El Cabellal	San Rafael	20° 15' 15.24"	-96° 59' 36.23"	4	4	100
Santa Elena	Martínez de la Torre	20° 13' 55.49"	-97° 01' 23.89"	4	3	75
Arroyo Frio	Martínez de la Torre	19° 58' 43.10"	-97° 00' 25.23"	1	1	100
Total				28	17	60.7




Figure 2. Phylogenetic tree of some pathogenic *Leptospira* species, obtained using the maximum likelihood (ML) approach and the Tamura's three-parameter model (TN93). The sequences recovered from rodents (*Mus musculus*) sampled in this study are marked with red diamonds.

The presence of two additional species of pathogenic leptospires (*L. weilli* and *L. noguchii*) has been confirmed in the state of Veracruz, both in kidney tissue samples of the hematophagous bat *Desmodus rotundus* and the frugivorous bat *Artibeus jamaicensis* (Ballados-González et al. 2018). Therefore, this study increases the inventory of pathogenic leptospires of Veracruz to three species. These findings in rodents suggest the existence of multiple transmission cycles in both wild and anthropic environments, which should be evaluated carefully.

Literature cited

- ALVARADO-ESQUIVEL, C., A. CRUZ-ROMERO, D. ROMERO-SALAS, A. ALVARADO-FÉLIX, M. AGUILAR-DOMÍNGUEZ, J. L. OCHOA-VALENCIA, D. ALVARADO-ESQUIVEL, J. HERNÁNDEZ-TINOCO, J. A. ZAMARRIPA-BARBOZA, AND L. F. SÁNCHEZ-ANGUIANO. 2018. Apparently high *Leptospira* antibody seropositivity in donkeys for slaughter in three municipalities in Durango, México. *Journal of Veterinary Diagnostic Investigation* 30:929–932.
- ALVAREZ, M. A., L. P. MOLES-CERVANTES, D. G. ROSAS, C. N. VÁSQUEZ, AND F. S. GARCÍA. 2005. Retrospective seroprevalence study of bovine leptospirosis in México considering the ecological regions. *Revista Cubana de Medicina Tropical* 57:28–31.
- BALLADOS-GONZÁLEZ, G. G., S. SÁNCHEZ-MONTES, D. ROMERO-SALAS, P. COLUNGA-SALAS, R. GUTIÉRREZ-MOLINA, L. LEÓN-PANIAGUA, I. BECKER, M. L. MÉNDEZ-OJEDA, C. BARRIENTOS-SALCEDO, R. SERNA-LAGUNES, AND A. CRUZ-ROMERO. 2018. Detection of pathogenic *Leptospira* species associated with phyllostomid bats (Mammalia: Chiroptera) from Veracruz, México. *Transboundary and Emerging Diseases* 65:773–781.
- BOURHY, P., L. COLLET, S. BRISSE, AND M. PICARDEAU. 2014. *Leptospira mayottensis* sp. nov., a pathogenic species of genus *Leptospira* isolated from humans. *International Journal of Systematic and Evolutionary Microbiology* 64:4061–4067.

- COLOMBO, V. C., I. GAMIETEA, S. G. LOFFLER, B. F. BRIHUEGA, AND P. M. BELDOMENICO. 2018. New host species for *Leptospira borgpetersenii* and *Leptospira interrogans* serovar Copenhageni. *Veterinary Microbiology* 215:90–92.
- CRUZ-ROMERO, A., C. ALVARADO-ESQUIVEL, D. ROMERO-SALAS, A. O. ALVARADO-FÉLIX, S. SÁNCHEZ-MONTES, J. HERNÁNDEZ-TINOCO, AND L. F. SÁNCHEZ-ANGUIANO. 2018. Seroepidemiology of *Leptospira* infection in backyard pigs in Durango State, México. *European Journal of Microbiology and Immunology* 8:87–90.
- DA SILVA, E. F., S. R. FÉLIX, G. M. CERQUEIRA, M. Q. FAGUNDES, A. C. NETO, A. A. GRASSMANN, M. G. AMARAL, T. GALLINA, AND O. A. DELLAGOSTIN. 2010. Preliminary characterization of *Mus musculus*-derived pathogenic strains of *Leptospira borgpetersenii* serogroup Ballum in a hamster model. *American Journal of Tropical Medicine and Hygiene* 83:336–337.
- ESPINOSA-MARTÍNEZ, D. V., D. S. SÁNCHEZ-MONTES, L. LEÓN-PANIAGUA, C. A. RÍOS-MUÑOZ, M. BERZUNZA-CRUZ, AND I. BECKER. 2015. New wildlife hosts of *Leptospira interrogans* in Campeche, México. *Revista do Instituto de Medicina Tropical de São Paulo* 57:181–183.
- LEVETT, P. N. 2015. Systematics of leptospiaceae. *Current Top Microbiology and Immunology* 387:11–20.
- MATSUI, M., L. ROCHE, M. E. SOUPÉ-GILBERT, M. ROUDIER, V. MONIQUET, AND C. GOARANT. 2015. Experimental hamster infection with a strain of *Leptospira borgpetersenii* Ballum isolated from a reservoir mouse in New Caledonia. *American Journal of Tropical Medicine and Hygiene* 92:982–985.
- MÉNDEZ, C., L. BENAVIDES, A. ESQUIVEL, A. ALDAMA, J. TORRES, D. GAVALDÓN, P. MELÉNDEZ, AND L. MOLES. 2013. Pesquisa serológica de *Leptospira* en roedores silvestres, bovinos, equinos y caninos en el noreste de México. *Revista de Salud Animal* 35:25–32.
- MOLES-CERVANTES, L. P., M. A. CISNEROS-PUEBLA, D. G. ROSAS, N. R. SERRANÍA, AND J. I. TORRES-BARRANCA. 2002. Serological study of bovine leptospirosis in México. *Revista Cubana de Medicina Tropical* 54:24–27.
- PANTI-MAY, J. A., R. R. C. DE ANDRADE, Y. GURUBEL-GONZÁLEZ, E. PALOMO-ARJONA, L. SODÁ-TAMAYO, J. MEZA-SULÚ, M. RAMÍREZ-SIERRA, E. DUMONTEIL, V. M. VIDAL-MARTÍNEZ, C. MACHAÍN-WILLIAMS, D. DE OLIVEIRA, M. G. REIS, M. A. TORRES-CASTRO, M. R. ROBLES, S. F. HERNÁNDEZ-BETANCOURT, AND F. COSTA. 2017. A survey of zoonotic pathogens carried by house mouse and black rat populations in Yucatan, México. *Epidemiological Infections* 145:2287–2295.
- SÁNCHEZ-MONTES, S., D. V. ESPINOSA-MARTÍNEZ, C. A. RÍOS-MUÑOZ, M. BERZUNZA-CRUZ, AND I. BECKER. 2015. Leptospirosis in México: Epidemiology and Potential Distribution of Human Cases. *PLoS One* 10:e0133720.
- TAMURA, K., G. STECHER, D. PETERSON, A. FILIPSKI, AND S. KUMAR. 2013. Mega6: Molecular Evolutionary Genetics Analysis Version 6.0. *Molecular Biology and Evolution* 30:2725–2729.
- THIBEAUX, R., D. GIRAULT, E. BIERQUE, M. E. SOUPÉ-GILBERT, A. RETTINGER, A. DOUYÈRE, M. MEYER, G. IRAOLA, M. PICARDEAU, AND C. GOARANT. 2018. Biodiversity of Environmental *Leptospira*: Improving Identification and Revisiting the Diagnosis. *Frontiers in Microbiology* 1:816.
- TORRES-CASTRO, M., L. GUILLERMO-CORDERO, S. HERNÁNDEZ-BETANCOURT, E. GUTIÉRREZ-RUIZ, P. AGUDELO-FLÓREZ, R. PELÁEZ-SÁNCHEZ, J. F. ZAVALA-CASTRO, AND F. I. PUERTO. 2016. First histopathological study in kidneys of rodents naturally infected with *Leptospira* pathogenic species from Yucatan, México. *Asian Pacific Journal of Tropical Medicine* 9:145–147.
- TORRES-CASTRO, M., B. CRUZ-CAMARGO, R. MEDINA-PINTO, B. REYES-HERNÁNDEZ, C. MOGUEL-LEHMER, R. MEDINA, J. ORTIZ-ESQUIVEL, W. ARCILA-FUENTES, A. LÓPEZ-ÁVILA, H. NO-PECH, A. PANTI-MAY, I. RODRÍGUEZ-VIVAS, AND F. I. PUERTO. 2018. Molecular detection of pathogenic *Leptospira* in synanthropic and wild rodents captured in Yucatán, México. *Biomedica* 38:51–58.
- VITAL-BRAZIL, J. M., I. T. BALASSIANO, F. S. OLIVEIRA, A. D. COSTA, L. HILLEN, AND M. M. PEREIRA. 2010. Multiplex PCR-based detection of *Leptospira* in environmental water samples obtained from a slum settlement. *Memorias do Instituto Oswaldo Cruz* 105:353–355.
- YAGER, R. H., W. S. GOCHENOUR, A. D. ALEXANDER, AND P. W. WETMORE. 1953. Natural occurrence of *Leptospira ballum* in rural house mice and in an opossum. *Proceedings of the Society for Experimental Biology and Medicine* 84:589–590.
- ZÁRATE-MARTÍNEZ, J. P., J. V. ROSETE-FERNÁNDEZ, A. RÍOS-UTRERA, F. T. BARRADAS-PIÑA, AND S. OLAZARÁN-JENKINS. 2015. Prevalencia de Leptospirosis y su relación con la tasa de gestación en bovinos de la zona centro de Veracruz. *Nova Scientia* 7:202–217.

Associated editor: Jesús Fernández

Submitted: November 27, 2018; Reviewed: January 29, 2019;

Accepted: March 21, 2019; Published on line: May 12, 2019.