



Veterinaria México OA

ISSN: 0301-5092

Universidad Nacional Autónoma de México, Facultad de
Medicina Veterinaria y Zootecnia

Iša, Pavel; Paulin, Luis Felipe; Hernandez, Jesus;
Montalvo-Corral, Maricela; López, Susana; Arias, Carlos F
Phylogenetic analysis of avian influenza viruses from migratory waterfowl at Pacific wetlands
Veterinaria México OA, vol. 9, e702, 2022
Universidad Nacional Autónoma de México, Facultad de Medicina Veterinaria y Zootecnia

DOI: <https://doi.org/10.22201/fmvz.24486760e.2022.702>

Available in: <https://www.redalyc.org/articulo.oa?id=42375500003>

- How to cite
- Complete issue
- More information about this article
- Journal's webpage in redalyc.org

UNAM redalyc.org

Scientific Information System Redalyc
Network of Scientific Journals from Latin America and the Caribbean, Spain and
Portugal

Project academic non-profit, developed under the open access initiative

Phylogenetic analysis of avian influenza viruses from migratory waterfowl at Pacific wetlands

Pavel Iša^{1*}

 0000-0002-3175-0993

Luis Felipe Paulin^{1,2}

 0000-0003-2567-3773

Jesus Hernandez³

 0000-0002-5131-3600

Maricela Montalvo-Corral³

 0000-0002-0070-7490

Susana López¹

 0000-0001-6336-9209

Carlos F. Arias¹

 0000-0003-3130-4501

¹ Universidad Nacional Autónoma de México.
Instituto de Biotecnología,
Departamento de Genética del Desarrollo
y Fisiología Molecular,
Morelos-Cuernavaca, México.

² Baylor College of Medicine,
Human Genome Sequencing Center,
Houston, TX, USA.

³ Centro de Investigación en Alimentación
y Desarrollo, AC.
Hermosillo, México

***Corresponding author:**

Email address:

pavel.isa@ibt.unam.mx

Abstract

The surveillance of influenza A viruses (IAV) in migratory waterfowl is of great importance, given the possibility of transmission to susceptible hosts and introduction of viral subtypes, including highly pathogenic strains, to new geographic regions. IAV were previously identified from migratory birds in two wetlands located in Mexico during two wintering seasons. Using next-generation sequencing, thirteen samples of RNA from the field sampled cloacal swabs were subjected to whole genome sequencing. Seven whole IAV genomes were obtained, while remaining six samples provided partial results. Eight different hemagglutinin (HA) and neuraminidase (NA) combinations were identified in analyzed samples, while the analysis of the remaining genes in seven whole genome samples demonstrated four different combinations of viral segments, suggesting a diversity of circulating IAV. Phylogenetic analysis of samples whose whole genomes were sequenced showed that they are closely related to North American IAV, with some segments (NP, H6 and H11) related to the Eurasian virus. The use of high-throughput technologies enabled genetic characterization of field samples and contributed to fostering our knowledge of IAV genomics and virus circulation in Mexico, a lowly surveyed but biological and epidemiological relevant region, thus highlighting the need for future studies in Mexican wetlands.

Keywords: avian influenza viruses; migratory waterfowl; phylogenetic analysis; Pacific wetlands; next-generation sequencing.

Submitted: 2019-04-01

Accepted: 2021-11-10

Published: 2022-08-31

Additional information and declarations
can be found on page 15

© Copyright 2022
Pavel Iša et al.

open access 



Distributed under Creative Commons CC-BY 4.0

Cite this as: Iša P, Paulin LF, Hernández J, Montalvo-Corral M, López S, Arias CF. Phylogenetic analysis of avian influenza viruses from migratory waterfowl at Pacific wetlands. *Veterinaria México OA*. 2022;9. doi: 10.22201/fmvz.24486760e.2022.702.

Study contribution

In this study we characterized whole genome of several influenza A virus isolates obtained from wild birds. The phylogenetic comparison of all individual influenza A viruses segments sequenced in this work showed that they mostly clustered with other isolates from North America, with some being related to viruses circulating in Asia. Our observations suggest that migratory birds, which come to wetlands in Sonora, Mexico, co-migrate along flight corridors with bird populations from different breeding sites in North America, possibly Alaska and Canada, where major flyways may overlap with Asian flyways, allowing the interchange of genetic material and the increase in viral diversity.

Introduction

Migration is one of the important aspects of bird behavior, with billions of specimens migrating thousands of kilometers across the globe. These journeys generally follow a predominantly north to south direction, and link Arctic sites with temperate and tropical areas. Influenza A viruses circulate in many animal species, including humans, pigs, horses, dogs and a wide range of wild and domestic birds.^(1, 2) The virus reservoir in nature is thought to be formed principally by wild aquatic birds, mainly from Anseriformes (particularly ducks, geese and swan) and Charadriiformes (particularly gulls, terns and waders) orders, which contribute to wide-spread of avian influenza A viruses (IAV) during their seasonal migrations on their migratory routes.^(2, 3) Eight migratory flyways have been identified globally, with some of them connecting the high arctic north with the Caribbean, up to south Tierra de Fuego, through the Pacific sea coast.^(2, 4-6) The most important North American flyways identified are Atlantic, Mississippi, Central, and Pacific, where migratory birds stop over wintering grounds along the route. Millions of aquatic birds congregate in their breeding grounds in the northern hemisphere around the world, allowing free transmission of influenza viruses and exchange of viral segments during co-infections. The fact that IAV does not cause disease in wild birds [they are mostly low pathogenic avian influenza, (LPAI)], but are excreted in large quantities in feces, helps their spread. Excreted viruses remain viable during several days, and different viral types have been observed in environmental reservoirs at one point.⁽⁷⁾ Differences in set of gene segments (assortment of viral segments) among continental population of LPAI viruses have been described, and intercontinental reassortment of viral segments has also been found in migratory birds.^(4-6, 8, 9) Highly pathogenic avian influenza (HPAI) strains (H5N2, H5N1, H7N3, H9N2, H7N9) circulate widely and were transferred to caged birds.⁽¹⁰⁻¹¹⁾ Recent surveillance study of IAV in aquatic overwintering birds in Guatemala showed a high virus diversity (showing 52 different genomic sets in 68 analyzed isolates), and demonstrated evidence of introduction of a Eurasian viral PA segment into North Americas IAV population.⁽¹²⁾ In fact, diverse HPAI of Asian-origin have recently been reported in North America.⁽¹³⁾

There is an increased interest in monitoring the migration and spread of IAV strains through the globe. However, the coverage of wild bird sampling and genetic analysis of IAV is not always complete. For example, data available for Mexico are scarce. A previous eco-epidemiological study demonstrated IAV circulation in migratory birds in the Northwestern Mexican wetlands and partial characterization

of hemagglutinin genes of some viruses was performed; however, at that time we could not isolate and characterize whole samples.^(14, 15) For this reason, in this study we took advantage of high-throughput generation sequencing, to achieve full characterization and perform in-depth phylogenetic analysis of the whole genome of several IAV sampled from wild birds in the Pacific flyway migratory grounds in Mexico.

Materials and methods

Ethical statement

Samples were collected *post mortem* from birds captured during official surveillance activities conducted by the Mexican Department of Environment and Natural Resources (Semarnat). At the time of sampling (2007–2009), no further authorization was required.

Collection of samples, RNA extraction and library preparations

Thirteen influenza A virus-positive cloacal swabs, as tested for matrix gene by One-Step qRT–PCR, were collected from opportunistic sampling of 1262 hunter-killed migratory birds as part of a local surveillance program in Sonora, Mexico during the 2007–2008 and 2008–2009 seasons. The collection was realized in Laguna Moroncarit, Huatabampo (26°41'56.4"N 109°31'45.7"W) and EsteroTobari, Cd. Obregon (27°00'43.6"N 109°54'13.1"W).⁽¹⁴⁾ Samples were stored at -80°C with minimum handling until extraction (3 and 4 years reaspectively).

Before extracting viral nucleic acids, the samples were incubated with Turbo DNase (Ambion) to remove external or host DNA. Total RNA for each sample was extracted individually with the PureLink Viral DNA/RNA kit (Invitrogen), using linear acrylamide as a carrier. Individual whole influenza A genome amplification was performed by multi-segment RT-PCR using a set of universal primers, MB-Tuni-12 (5' ACGCGTGATCAGCAAAAGCAGG 3') and MBTuni-13 (5' ACGCGTGATCAGTAGAAACAAGG 3') with the SuperScript III one-step RT-PCR platinum Taq HiFi kit (Invitrogen), as described previously.⁽¹⁶⁾ All RT-PCR products were purified and concentrated by DNA clean & concentrator kit (Zymo) and visualized in 1.5% agarose gels.

Library preparation and sequencing

PCR products from whole genome RT-PCR were used as input material for preparation of sequencing libraries. DNA was sheared by treatment with NEBNext Fragmentase (New England Biolabs) for 15 min at 37 °C. Sheared DNA was purified and concentrated by DNA clean & concentrator kit (ZYMO) and 50 to 100 ng were used for the preparation of the libraries using Illumina's Genomic DNA Sample Preparation kit with Multiplex Sample Preparation Oligokit, as described by the manufacturer.

High-throughput sequencing was performed at the next-generation sequencing core facility located at the Instituto de Biotecnología, UNAM, in Cuernavaca, Morelos. Libraries of approximately 350 nucleotides were used to generate se-

quencing clusters, followed by 45 or 66 cycles of single base pair extensions in the Genome Analyzer Ix sequencer (Illumina, San Diego, CA), which was followed by a multiplex barcode acquisition. Computational analysis was performed using the computational cluster at the Instituto de Biotecnología, UNAM. Image analysis was carried out using the Genome Analyzer Pipeline Version 1.4.

All reads were quality filtered, and aligned to the reference genome generated separately for each sample. The number of filtered and influenza specific reads of nine sequenced samples is shown in [Table S1](#). To choose the appropriate reference genome to map the reads, the reads were first mapped to influenza A strains used in Paulin et al.⁽¹⁷⁾ and the segments with most of the reads mapped were selected for reference genomes. The alignment was performed by SMALT version 0.7.0.1⁽¹⁸⁾ and a consensus sequence was called by SAMtools version 0.1.18. To allow for sequence variability, two separate alignment rounds were performed. In the first round, 15 and 20 mismatches were allowed with 46 or 65 nucleotide long reads, respectively, while during the second alignment round only 5 mismatches were allowed using as a reference the consensus sequence previously generated during the first mapping round. Reads that did not match the reference genome were not considered for further analysis.

Phylogenetic analysis

Phylogenetic analysis was performed separately for each influenza A segment, and in the case of the HA and NA segments, for each identified subtype. Included sequences were selected from four regions reflecting continents where avian species used in this study are observed: Europe (EU), North America (NA), South America (SA), and Asia (AS). Sequences from avian, human, equine and swine sources were selected from NCBI Influenza Virus Resource, and ones with identical nucleotide coding sequences were collapsed. Additionally, sequences with more than 20 consecutive N's were eliminated.

The resulting sequences were clustered into arbitrary groups for each segment/subtype to remove highly similar (but not identical) sequences with cd-hit.⁽¹⁹⁾ For the case of the hemagglutinin and neuraminidase, all the clusters were done at 95 % of sequence identity for a median of 54 sequences per subtype for all samples included, with two outliers: N1 and N2 with 178 and 298 sequences, respectively.

For the rest of the segments, the clusters were constructed at 92 % of sequence identity, having a median of 90 sequences per tree (min = 54, mean = 85, max = 104). The number of sequences used is shown in [Table S2](#). For the selected sequences, the name was changed as follows: the continent of origin, abbreviated source, accession number, gi, species of origin, virus name and number, year, segment number, segment name and for the case of hemagglutinin and neuraminidase, the subtype as shown in the following example: NA_AV_GU051078_269822049_A_semipalmatedsandpiper_Delaware_2144_2000__3_PA. Finally, each field was separated by an underscore instead of a slash to avoid issues with the software used.

Each data set was aligned in nucleotides with the multiple sequence comparison by log-expectation (MUSCLE) algorithm⁽²⁰⁾ using the default parameters.

The presence of mismatches or possible frame shifts was checked after translation to protein. The substitution model was calculated for each alignment with IQTree, in model mode.⁽²¹⁾ The phylogenetic trees were constructed with the IQTree algorithm, using the model selected in the previous step, using the ultrafast bootstrap option with 10 000 bootstraps.⁽²²⁾ In the phylogenetic trees, only the geographical area is shown. The strains sequenced in this study are marked with a black dot next to their name. The bootstrap value for each split is shown in a circle, which the color intensity code indicates its value as a percentage.

The isolates sequenced in this work were grouped into clades after visual inspection of phylogenetic trees. The clades are well defined and the split that separates them has an ultra-fast bootstrap close to 100 percent.

Results and discussion

In total, thirteen avian influenza-positive samples were obtained from 7 green winged teals (*Anas crecca carolinensis*), 1 gadwall (*Anas strepera*), 1 American wigeon (*Anas americana*), 3 northern shovelers (*Anas clypeata*) and 1 redhead (*Aythya americana*). Samples were collected from two migratory avian wintering grounds in Sonora, Mexico; 3 were collected during 2007-2008 wintering season and 10 from the 2008-2009 season.⁽¹⁴⁾ Both locations (Laguna Moroncarit, Huatabampo and Estero Tobari, Cd. Obregon) are among the most important wetlands for avian migration on the Mexican Pacific coast. Given that in previous studies only partial identification of genes was achieved,^(14, 15) high-throughput sequencing was used to improve our data. Using of high-throughput sequencing technologies has shown usefulness for avian influenza virus complete genomic characterization from field samples, otherwise limited using traditional virological approaches.⁽²³⁾

A whole genome sequence with >100× coverage was obtained from 7 samples, which were used in the following analysis, while the remaining six samples generated only partial sequences, allowing only a description of influenza virus subtypes (Table 1). The samples from 2007-2008 season were identified as H6N1 (A/Green-winged teal/Mexico-Sonora/266/2008), and H5N2 (A/Gadwall/Mexico-Sonora/150/2008; A/Redhead/Mexico-Sonora/408/2008) (Table 1).

The samples from the 2008–2009 season were identified as H5N3 (A/Northern shoveler/Mexico-Sonora/796/2008; A/Northern shoveler/Mexico-Sonora/797/2008); A/Green winged teal/Mexico-Sonora/827/2008; A/Green winged teal/Mexico-Sonora/840/2008), H6N1 (A/Northern shoveler/Mexico-Sonora/738/2008), H6N5 (A/Green winged teal/Mexico-Sonora/829/2009), H9N2 (A/American wigeon/Mexico-Sonora/769/2008), H10N3 (A/Green winged teal/Mexico-Sonora/1132/2009), H10N7 (A/Green winged teal/Mexico-Sonora/1116/2009) and H11N3 (A/Green winged teal/Mexico-Sonora/701/2008).

It is of interest that out of the 7 isolates characterized from green winged teal; six different hemagglutinin and neuraminidase combinations were found (H5N3, H6N1, H6N5, H10N3, H10N7, H11N3), with only H6N1 being shared by two birds (Table 1). The H5N3 subtype was found in two Northern shovelers, collected at Estero Tobari, and in Laguna Moroncarit in two Green winged teals. The H5N2 combination was also found more than once, however, in different avian species (Table 1).

Table 1. Avian influenza A from waterfowl analyzed in this study

Sample	Species	Sex	Date of collection	WG ¹	Subtype ²
A/gadwall/Sonora/150/07	Gadwall	F	15/12/2007	ND ³	H5N2
A/green winged teal/Sonora/266/08	Green winged teal	M	19/01/2008	H6N1	H6
A/redhead/Sonora/408/08	Redhead	M	12/02/2008	ND	H5N2
A/green winged teal/Sonora/701/08	Green winged teal	F	13/12/2008	H11N3	
A/northern shoveler/Sonora/738/08	Northern Shoveler	M	13/12/2008	H6N1	H6
A/American wigeon/Sonora/769/08	American Wigeon	F	14/12/2008	H9N2	H9
A/northern shoveler/Sonora/796/08	Northern Shoveler	F	14/12/2008	ND	H5N3
A/northern shoveler/Sonora/797/08	Northern Shoveler	F	14/12/2008	H5N3	H5
A/green winged teal/Sonora/827/09	Green winged teal	F	10/01/2009	ND	H5N3
A/green winged teal/Sonora/829/09	Green winged teal	F	10/01/2009	H6N5	
A/green winged teal/Sonora/840/09	Green winged teal	F	10/01/2009	ND	H5N3
A/green winged teal/Sonora/1116/09	Green winged teal	M	08/02/2009	H10N7	
A/green winged teal/Sonora/1132/09	Green winged teal	M	08/02/2009	ND	H10N3

¹ WG. The whole genome was determined in this study.

² Subtype as determined previously (Montalvo-Corral and Hernandez 2010) or in this study.

³ Not determine the whole genome.

Whole genome sequencing allows analysis of all influenza segments, which could have important implications in influenza pathogenesis. While PB2 and PB1 gene segments from the field samples sequenced in this work clustered together, mostly with other viruses isolated from North American territories (Figures S1 and S2), the PA gene segments clustered into two separate groups (Figure 1), with both groups localized close to the North American isolates. Phylogenetic analysis of genomic segments corresponding to nucleoprotein, matrix and nonstructural proteins showed similar results as those observed for the polymerase gene segments. While matrix genes formed one group, close to other samples from North America (Figure S3), one Mexican sample (A/ Green winged teal/Mexico-Sonora/1116/2009 H10N7) grouped separately in its NS gene segment (Figure 2), and two samples, (A/American wigeon/Mexico-Sonora/769/2008 H9N2; A/ Green winged teal/Mexico-Sonora/1116/2009 H10N7) grouped separately from the remaining Mexican samples in the NP gene segment (Figure 3). NS segment, which groups separately, is closely related to other North American samples, while the NP segments were close to isolates from Asia. These observations confirm the complexity of influenza virus inter-continental circulation.




Figure 1. The phylogenetic tree for the PA gene segment of avian influenza viruses was analyzed in this work. For used sequences, the name was changed as follows: continent of origin, source, gi and accession number, virus name, segment number and segment name. In phylogenetic trees, following area code was used: Europe (EU), North America (NA), South America (SA), Asia (AS) and Mexico-this work (MX). The bootstrap value for each split is shown in a circle with the intensity color code indicating its value as a percentage. The color code of the continent of origin is as follows: green North America (NA), blue South America (SA), brown Europe (EU) purple Asia (AS), red sample sequences in this work (MX).



Figure 2. The phylogenetic tree for the NS gene segment of avian influenza viruses was analyzed in this work. For the selected sequences, the name was changed as follows: continent of origin, source, gi and accession number, virus name, segment number and segment name. In phylogenetic trees, following area code was used: Europe (EU), North America (NA), South America (SA), Asia (AS) and Mexico-this work (MX). The bootstrap value for each split is shown in a circle with the intensity color code indicating its value as a percentage. The color code of the continent of origin is as follows: green North America (NA), blue South America (SA), brown Europe (EU) purple Asia (AS), red sample sequences in this work (MX).




Figure 3. The phylogenetic tree for the NP gene segment of avian influenza viruses was analyzed in this work. For used sequences the name was changed as follows: continent of origin, source, gi and accession number, virus name, segment number, segment name and for the case of hemagglutinin and neuraminidase the subtype. In phylogenetic trees, following area code was used: Europe (EU), North America (NA), South America (SA), Asia (AS) and Mexico-this work (MX). The bootstrap value for each split is shown in a circle with the intensity color code indicating its value as a percentage. The color code of the continent of origin is as follows: green North America (NA), blue South America (SA), brown Europe (EU) purple Asia (AS), red sample sequences in this work (MX).

Hemagglutinin and neuraminidase gene segments

From the seven wild bird IAV-positive samples sequenced, five different hemagglutinin types were found (H5, H6, H9, H10, and H11), with the HA6 type being the only present more than once. Phylogenetic analyses of hemagglutinin gene segments showed that they are closely related to other IAV from North America (Figures S4–S6). The exceptions were H6 and H11 segments, which were grouped closer to isolates from Asia (Figure 4 and 5). Interestingly, the 3 Mexican HA6 samples found in this study were grouped together despite being isolated in two different influenza seasons, from two distinct locations. Sample A/Green-winged teal/Mexico-Sonora/266/2008 H6N1, was collected during season 2007–2008, in Estero Tobari, and samples A/Northern shoveler/Mexico-Sonora/738/2008 H6N1, and A/Green-winged teal/Mexico-Sonora/829/2009 H6N5, were collected during 2008–2009 season, the first from Estero Tobari and the second sample from Laguna Moroncarit (Figure 4). However, sample A/Green winged teal/Mexico-Sonora/701/2008 localized close to the Asian sample described in 2016.

Analysis of the H5 segment characterized in this work showed that sample A/Northern shoveler/Mexico-Sonora/797/2008 H5N3, separated from the previously characterized Mexican H5 samples isolated in 1994–1997 period (Figure S4). This genetic distance might be explained by differences in space-time genetic changes and possible origins, because samples from 1994–1997 were isolated from farm chickens and belonged to an HPAI virus. AIV H5 subtype characterized in this work is low pathogenic avian influenza (LPAI) virus, lacks of hemagglutinin pathogenic markers. The occurrence of the H5 subtype is not as rare as previously thought in wild birds. A large-scale surveillance conducted in the USA between 2006 and 2011 found H5 subtype highly represented among IAV isolates⁽²⁴⁾ and HPAIV H5 was recently found in wild birds in Canada and USA.⁽²⁵⁾ In our study, which included few samples, 6 of 13 IAVs were subtyped as H5 isolates (Table 1). Alternatively, recent data from ducks overwintering in Guatemala showed that only 3 samples from 68 characterized isolates were H5.⁽¹²⁾

With respect to neuraminidase genome segments, five different NA types were identified (N1, N2, N3, N5, and N7), with N1 and N3 types being present in two samples (Table 1), and all are closely related to other North American isolates (Figures S7–S11).

To analyze virus diversity, specific combinations of viral segments in each sample were determined. For this, each segment was classified within the corresponding clade as described in the Materials and Methods. Among the seven fully sequenced isolates, 4 different sets of genome segments were observed (Table 2). First is formed by three viruses (A/greenwingedteal/Sonora/266/08, A/greenwingedteal/Sonora/701/08, and A/northernshoveler/Sonora/738/08), which have all segments in main groups, the second constellation contained two viruses (A/northernshoveler/Sonora/797/08 and A/greenwingedteal/Sonora/829/09), which differ in 116 nucleotides in the PA segment, with the last two having just one example, A/american wigeon/Sonora/769/08, with the NP segment different from the rest, and lastly most divergent sample A/green winged teal/Sonora/1116/09, with PA, NP and NS segments distinct (Table 2). When H and N genes were added for comparison, 6 different sets of genome segments are observed (Table 2).




Figure 4. The phylogenetic tree for the H6 gene segment of avian influenza viruses was analyzed in this work. For used sequences the name was changed as follows: continent of origin, source, gi and accession number, virus name, segment number, segment name and for the case of hemagglutinin and neuraminidase the subtype. In phylogenetic trees, following area code was used: Europe (EU), North America (NA), South America (SA), Asia (AS) and Mexico-this work (MX). The bootstrap value for each split is shown in a circle with the intensity color code indicating its value as a percentage. The color code of the continent of origin is as follows: green North America (NA), blue South America (SA), brown Europe (EU) purple Asia (AS), red sample sequences in this work (MX).




Figure 5. The phylogenetic tree for the H11 gene segment of avian influenza viruses was analyzed in this work. For used sequences the name was changed as follows: continent of origin, source, gi and accession number, virus name, segment number, segment name and for the case of hemagglutinin and neuraminidase the subtype. In phylogenetic trees, following area code was used: Europe (EU), North America (NA), South America (SA), Asia (AS) and Mexico-this work (MX). The bootstrap value for each split is shown in a circle with the intensity color code indicating its value as a percentage. The color code of the continent of origin is as follows: green North America (NA), blue South America (SA), brown Europe (EU) purple Asia (AS), red sample sequences in this work (MX).

Table 2. Sets of genomic segments of avian influenza viruses identified in waterfowl species from Pacific wetlands

Sample	PB2	PB1	PA	NP	M	NS	HANA
A/green winged teal/Sonora/266/08	1	1	1	1	1	1	H6N1
A/green winged teal/Sonora/701/08	1	1	1	1	1	1	H11N3
A/northern shoveler/Sonora/738/08	1	1	1	1	1	1	H6N1
A/american wigeon/Sonora/769/08	1	1	1	2	1	1	H9N2
A/northern shoveler/Sonora/797/08	1	1	2	1	1	1	H5N3
A/green winged teal/Sonora/829/09	1	1	2	1	1	1	H6N5
A/green winged teal/Sonora/1116/09	1	1	2	2	1	2	H10N7

When analyzing all 13 IAV isolates characterized (seven complete whole genomes and six partial genomes that identify the HA and NA subtypes), eight different HA and NA combinations were found, confirming the subtype diversity present in the North American Pacific flyway. This is true especially considering that we could analyze samples from only five different avian species. These results again suggest that there is diversity of influenza viruses present in the wintering sites analyzed. Similar high genomic variability was observed in Guatemala, where out of 68 influenza isolates characterized, 19 different subtype combinations were found (nine HA subtypes and seven NA subtypes).⁽¹²⁾ The Mississippi migratory flyway also showed a high level of genomic variability, with 46 unique HA and NA combinations observed in 297 isolates.⁽²⁶⁾ However, in Kazakhstan, lower variability of HA and NA combinations was observed during a 7 year span (from 2002 to 2009).⁽²⁷⁾

Despite the large dataset of influenza genome sequences now available, knowledge of complete avian influenza virus sequences from wild birds from Mexico and Latin America are very limited.⁽²⁸⁾ Thus, the phylogenetic comparison of all individual IAV segments sequenced in this work showed that they mostly clustered with other isolates from North America. The only exceptions were some of the segments of samples A/Green-winged teal/Mexico-Sonora/266/2008; A/Northern shoveler/Mexico-Sonora/738/2008; and A/Green winged teal/Mexico-Sonora/829/2009 (H6 gene), A/Green winged teal/Mexico-Sonora/701/2008 (H11 segment) and A/American wigeon/Mexico-Sonora/769/2008 and A/Green winged teal/Mexico-Sonora/1116/2009 (NP genomic segment), which appear related to viruses circulating in Asia. These observations support our previous data and show new findings, suggesting that migratory birds, which come to wetlands in Sonora, Mexico, co-migrate along flight corridors with bird populations from different breeding and stop over sites in northern locations of North America, possibly Alaska and Canada, where major flyways may overlap with Asian flyways, allowing the interchange of genetic material and the increase of viral diversity.

Conclusions

Previously we have described isolation and partial characterization of influenza virus isolated from migratory birds in two wetlands in Mexico. In this work using whole genome sequencing we have identified eight different HA and NA combinations

in thirteen analyzed samples, while the analysis of seven whole genome samples demonstrated four different combinations of viral segments, suggesting a diversity of circulating IAV. Phylogenetic analysis of samples whose whole genomes were sequenced showed that they are all LPAIV and closely related to North American IAV, with some segments (NP, H6 and H11) being related to strains circulating in Asia. The use of high-throughput technologies enabled genetic characterization of field samples and contributed to fostering our knowledge of IAV genomics and virus circulation in Mexico, a lowly surveyed but biological and epidemiological relevant region, and highlights the need for future studies in Mexican wetlands.

Data availability

Sequences are publicly available at the National Center for Biotechnology Information (NCBI: <https://www.ncbi.nlm.nih.gov>), accession numbers KY575169-KY575224, and KY593192-KY593208.

Acknowledgments

We thank Dr. Ricardo Grande and MC Veronica Jiménez, in the next-generation sequencing core facility of UNAM for their technical support. The computer analysis was performed using the cluster of the Instituto de Biología, UNAM.

Funding statement

This work was supported by grant PICOSI09-209 from the Instituto de Ciencia y Tecnología del Distrito Federal, México, awarded to PI, and by grant IN213410 from UNAM, DGAPA (<https://dgapa.unam.mx/>) awarded to PI. LFP was supported by a scholarship from CONACYT-Mexico (<https://conacyt.mx/>).

Conflicts of interest

The authors have no conflict of interest to declare in regard to this publication.

Author contributions

Conceptualization: P Iša

Methodology: P Iša, LF Paulin

Supervision: S López, CF Arias

Formal analysis: P Iša, LF Paulin

Funding acquisition: P Iša

Resources: J Hernandez, M Montalvo-Corral

Writing – original draft: P Iša, LF Paulin, M Montalvo-Corral

Writing – review and editing: P Iša, LF Paulin, J Hernandez, M Montalvo-Corral, S López, CF Arias

References

1. Lipatov AS, Govorkova EA, Webby RJ, Ozaki H, Peiris M, Guan Y, Poon L, Webster RG. Influenza: emergence and control. *Journal of Virology*. 2004;78(17):8951-8959. doi: 10.1128/JVI.78.17.8951-8959.2004.
2. Olsen B, Munster VJ, Wallensten A, Waldenstrom J, Osterhaus AD, Fouchier RA. Global patterns of influenza A virus in wild birds. *Science*. 2006;312(5772):384-388. doi: 10.1126/science.1122438.
3. Webster RG, Bean WJ, Gorman OT, Chambers TM, Kawaoka Y. Evolution and ecology of influenza A viruses. *Microbiology Reviews*. 1992;56:152-179.
4. Dusek RJ, Hallgrímsson GT, Ip HS, Jónsson JE, Sreevatsan S, Nashold SW, TeSlaa JL, Enomoto S, Halpin RA, Lin X, Fedorova N, Stockwell TB, Dugan VG, Wentworth DE, Hall JS. North Atlantic migratory bird flyways provide routes for intercontinental movement of avian influenza viruses. *PLoS One*. 2014;9(3):e92075. doi: 10.1371/journal.pone.0092075.
5. Girard YA, Runstadler JA, Aldehoff F, Boyce W. Genetic structure of Pacific Flyway avian influenza viruses is shaped by geographic location, host species, and sampling period. *Virus Genes*. 2012;44(3):415-428. doi: 10.1007/s11262-011-0706-5.

6. Pearce JM, Ramey AM, Flint PL, Koehler AV, Fleskes JP, Franson JC, Hall JS, Derksen DV, Ip HS. Avian influenza at both ends of a migratory flyway: characterizing viral genomic diversity to optimize surveillance plans for North America. *Evolutionary Applications*. 2009;2(4):457-468. doi: 10.1111/j.1752-4571.2009.00071.x.
7. Lang AS, Kelly A, Runstadle, JA. Prevalence and diversity of avian influenza viruses in environmental reservoirs. *Journal of General Virology*. 2008;89(Pt 2):509-519. doi: 10.1099/vir.0.83369-0.
8. Fusaro A, Monne I, Salviato A, Valastro V, Schivo A, Amarin NM, Gonzalez C, Ismail MM, Al-Ankari AR, Al-Blawi MH, Khan OA, Maken Ali AS, Hedayati A, Garcia Garcia J, Ziy GM, Shoushtari A, Qahtani KN, Capua I, Holmes EC, Cattoli G. Phylogeography and evolutionary history of reassortant H9N2 viruses with potential human health implications. *Journal of Virology*. 2011;85(16):8413-8421. doi: 10.1128/JVI.00219-11.
9. Hurt AC, Su YC, Aban M, Peck H, Lau H, Baas C, Deng YM, Spirason N, Ellström P, Hernandez J, Olsen B, Barr IG, Vijaykrishna D, Gonzalez-Acuna D. Evidence for the introduction, reassortment, and persistence of diverse Influenza A viruses in Antarctica. *Journal of Virology*. 2016;90(21):9674-9682. doi: 10.1128/JVI.01404-16.
10. Kapczynski DR, Pantin-Jackwood M, Guzman SG, Ricardez Y, Spackman E, Bertran K, Suarez DL, Swayne DE. Characterization of the 2012 highly pathogenic avian influenza H7N3 virus isolated from poultry in an outbreak in Mexico: pathobiology and vaccine protection. *Journal of Virology*. 2013;87(16):9086-9096. doi: 10.1128/JVI.00666-13.
11. Poovorawan Y, Pyungporn S, Prachayangprecha S, Makkoch J. Global alert to avian influenza virus infection: from H5N1 to H7N9. *Pathogens and Global Health*. 2013;107(5):217-223. doi: 10.1179/2047773213Y.0000000103.
12. Gonzalez-Reiche AS, Muller ML, Ortiz L, Cordon-Rosales C, Perez DR. Prevalence and diversity of low pathogenicity avian influenza viruses in wild birds in Guatemala, 2010-2013. *Avian Diseases*. 2016;601 (Suppl):359-364. doi: 10.1637/11130-050715-Reg.
13. Jhung MA, Nelson DI. Centers for Disease Control and Prevention (CDC). Outbreaks of avian influenza A (H5N2), (H5N8), and (H5N1) among birds – United States, December 2014-January 2015. *MMWR Morbidity and Mortal Wkly Report*. 2015;64(4):111. PMID: PMC4584850.
14. Montalvo-Corral M, Hernandez J. Genetic analysis of avian influenza virus from migratory waterfowl in Mexico. *Archives of Virology*. 2010;155(1):97-101. doi: 10.1007/s00705-009-0554-y.
15. Montalvo-Corral M, Lopez-Robles G, Hernandez J. Avian influenza survey in migrating waterfowl in Sonora, Mexico. *Transboundary and Emerging Diseases*. 2011;58(1):63-8. doi: 10.1111/j.1865-1682.2010.01182.x.
16. Escalera-Zamudio M, Nelson MI, Cobián-Güemes AG, López-Martínez I, Cruz-Ortiz N, Iguala-Vidales M, García ER, Barrera-Badillo G, Díaz-Quinonez JA, López S, Arias CF, Isa P. Molecular epidemiology of influenza A/H3N2 viruses circulating in Mexico from 2003 to 2012. *PLoS One*. 2014;9(7):e102453. doi: 10.1371/journal.pone.0102453.
17. Paulin LF, M. de los D Soto-Del Río, Sánchez I, Hernández J, Gutiérrez-Ríos RM, López-Martínez I, Wong-Chew RM, Parissi-Crivelli A, Isa P, López S, Arias CF. PhyloFlu, a DNA microarray for determining the phylogenetic origin of influenza

- A virus gene segments and the genomic fingerprint of viral strains. *Journal of Clinical Microbiology*. 2014;52(3):803-813. doi: 10.1128/JCM.03134-13.
18. Ponstingl H, Ning Z. SMALT-A new mapper for DNA sequencing reads. Poster exhibido en: Sequencing Informatics Division, The Wellcome Trust Sanger Institute; 02 agosto 2010; Hinxton, Cambridge CB10 1SA, UK.
 19. Fu L, Niu B, Zhu Z, Wu S, Li W. CD-HIT: accelerated for clustering the next-generation sequencing data. *Bioinformatics*. 2012;28(23):3150-3152. doi: 10.1093/bioinformatics/bts565.
 20. Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Research*. 2004;32(5):1792-1797. doi: 10.1093/nar/gkh340.
 21. Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermini LS. ModelFinder: fast model selection for accurate phylogenetic estimates. *Nature Methods*. 2017;14(6):587-589. doi: 10.1038/nmeth.4285.
 22. Nguyen LT, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: a fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Molecular Biology and Evolution*. 2015;32(1):268-274. doi: 10.1093/molbev/msu300.
 23. Ali R, Blackburn RM, Kozlakidis Z. Next-generation sequencing and influenza virus: a short review of the published implementation attempts. *HAYATI Journal of Biosciences*. 2016;23:155-159. doi: 10.1016/j.hjb.2016.12.007.
 24. Bevins SN, Pedersen K, Lutman MW, Baroch JA, Schmit BS, Kohler D, Gidlewski T, Nolte DL, Swafford SR, DeLiberto TJ. Large-scale avian influenza surveillance in wild birds throughout the United States. *PLoS One*. 2014;9(8):e104360. doi: 10.1371/journal.pone.0104360.
 25. Bevins SN, Dusek RJ, White CL, Gidlewski T, Bodenstein B, Mansfield KG, DeBruyn P, Kraege D, Rowan E, Gillin C, Thomas B, Chandler S, Baroch J, Schmit B, Grady MJ, Miller RS, Drew ML, Stopak S, Zscheile B, Bennett J, Sengl J, Brady C, Ip HS, Spackman E, Killian ML, Torchetti MK, Sleeman JM, Deliberto TJ. Widespread detection of highly pathogenic H5 influenza viruses in wild birds from the Pacific flyway of the United States. *Sci Rep*. 2016;6:28980. doi: 10.1038/srep28980.
 26. Fries AC, Nolting JM, Bowman AS, Lin X, Halpin RA, Wester E, Fedorova N, Stockwell TB, Das SR, Dugan VG, Wentworth DE, Gibbs HL, Slemons RD. Spread and persistence of influenza A viruses in waterfowl hosts in the North American Mississippi migratory flyway. *J Virol*. 2015;89(10):5371-5381. doi: 10.1128/JVI.03249-14.
 27. Kydyrmanov A, Sayatov M, Karamendin K, Zhumatov K, Asanova S, Daulbayeva K, Starick E, Fereidouni S. Monitoring of influenza A viruses in wild bird populations in Kazakhstan in 2002-2009. *Arch Virol*. 2017;162(1):147-155. doi: 10.1007/s00705-016-3076-4.
 28. Afanador-Villamizar A, Gomez-Romero C, Diaz A, Ruiz-Saenz J. Avian influenza in Latin America: a systematic review of serological and molecular studies from 2000-2015. *PLoS ONE*. 2017;12(6): e0179573. doi: 10.1371/journal.pone.0179573.