



Revista mexicana de biodiversidad

ISSN: 1870-3453

ISSN: 2007-8706

Instituto de Biología

Iglesias, Ricardo; Palacios-Vargas, José G.; Castaño-Meneses, Gabriela
Comparison of oribatid mites from agricultural soils with
contrasting irrigation types in Hidalgo State, Mexico: a case study
Revista mexicana de biodiversidad, vol. 90, 2019
Instituto de Biología

DOI: 10.22201/ib.20078706e.2019.90.2780

Available in: <http://www.redalyc.org/articulo.oa?id=42562784020>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

UAEM  redalyc.org

Scientific Information System Redalyc

Network of Scientific Journals from Latin America and the Caribbean, Spain and Portugal

Project academic non-profit, developed under the open access initiative

Ecology

Comparison of oribatid mites from agricultural soils with contrasting irrigation types in Hidalgo State, Mexico: a case study

Comparación de ácaros oribátidos de suelos agrícolas con tipos de riego contrastante en el estado de Hidalgo, México: un estudio de caso

Ricardo Iglesias ^a, José G. Palacios-Vargas ^a, Gabriela Castaño-Meneses ^{a, b, *}

^a Laboratorio de Ecología y Sistemática de Microartrópodos, Departamento de Ecología y Recursos Naturales, Facultad de Ciencias, Universidad Nacional Autónoma de México, Circuito Exterior Universitario, Ciudad Universitaria, Coyoacán, 04510 Ciudad de México, Mexico

^b Ecología de Artrópodos en Ambientes Extremos, Unidad Multidisciplinaria de Docencia e Investigación, Facultad de Ciencias, Campus Juriquilla, Boulevard Juriquilla 3001, Col. Jurica La Mesa 76230, Juriquilla, Querétaro, Mexico

*Corresponding author: gabycast99@hotmail.com (G. Castaño-Meneses)

Received: 10 August 2018; accepted: 26 January 2019

Abstract

The objective of this work is to know the structure (composition and density) of the community of the oribatid mites and their relationship to edaphic parameters in 2 agricultural parcels with different type of irrigation. It constitutes the first work done with oribatid mites in agricultural plots irrigated with waste water in Mexico. Two agricultural parcels were selected: one is irrigated with waste water (San Salvador) and the other with well water (El Bondho) from the municipality of San Salvador. The density of oribatids in San Salvador was higher than that from El Bondho (2,360 and 1,935 ind/m², respectively). The maximum abundances in percentages in San Salvador were for 3 species: *Tectocepheus velatus elegans* (48%), *Scheloribates* sp. (15%) and *Oppiella nova* (13%); but for El Bondho, there were only 2 species: *Zygoribatula connexa* (74%) and *Ramusella* sp. (10%). Due to soil conditions by the irrigation type, results gave information, on one hand, about the tolerance of some species like *O. nova*, *R. ardua*, and *T. velatus elegans* to heavy metals, and on the other hand, about the low resistance to heavy metals of *Epilohmannia pallida* and *Z. connexa*, which were present in the less polluted parcel, but are tolerant to saline soils. Results can be useful to support the use of some species of oribatid mites as bioindicators in ecosystems with similar conditions.

Keywords: Bioindicators; Communities; Heavy metals; Seasonal variation

Resumen

El presente trabajo tiene como objetivo conocer la estructura (composición y densidad) de la comunidad de ácaros oribátidos y su relación con parámetros edáficos en 2 parcelas agrícolas con riego diferente, y constituye el primer trabajo sobre ácaros oribátidos en suelos regados con aguas residuales en México. Se seleccionaron 2 parcelas agrícolas,

una regada con aguas residuales (San Salvador) y otra con aguas de pozo (El Bondho) del municipio de San Salvador. La densidad de oribátidos en San Salvador fue mayor que en El Bondho (2,360 y 1,935 ind/m², respectivamente). Las abundancias máximas en porcentajes fueron para 3 especies en San Salvador: *Tectocephus velatus elegans* (48%), *Scheloribates* sp. (15%) y *Oppiella nova* (13%), pero para El Bondho, fueron sólo 2, *Zygoribatula connexa* (74%) y *Ramusella* sp. (10%). Debido a las condiciones del suelo por el tipo de irrigación, se obtuvieron resultados que dan información, por un lado, sobre la tolerancia de algunas especies como *O. nova*, *R. ardua* y *T. velatus elegans* a los metales pesados, y por el otro, la baja resistencia a tales metales pesados de *Epilohmannia pallida* y *Z. connexa*, las que están presentes en la parcela con menos contaminación, pero que son tolerantes a suelos salinos. Los resultados obtenidos pueden ser útiles para apoyar el uso de algunas especies de ácaros oribátidos como bioindicadores en ambientes con condiciones similares.

Palabras clave: Bioindicadores; Comunidades; Metales pesados; Variación estacional

Introduction

The Mezquital Valley, a semiarid zone in the state of Hidalgo, Mexico, is a very important agricultural area with about 90,000 ha irrigated with waste water from Mexico City. This area represents the largest scheme of waste water use in Latin America (Braatz & Kandiah, 1996; Prado et al., 2015). The municipality of San Salvador, Hidalgo, has an important agricultural surface where the main crops are alfalfa (*Medicago sativa* L.), maize (*Zea mays* L.) and beans (*Phaseolus vulgaris* L.). According to the “Main Development Plan” of the Municipality (Ayuntamiento de San Salvador, Hgo., 2012) some plots have been irrigated with waste water for more than 100 years (Prado et al., 2015).

The use of waste water in the Mezquital Valley is an undervalued resource for agriculture due to the high concentration of nutrients contained in the water (Romero, 1997; Sagasta et al., 2013; Siebe & Cifuentes, 1995). Nevertheless, the frequent irrigation with such waters for a long period has led to the accumulation of several metals and other elements in the agricultural soils in Mexico. Such accumulation can be toxic for the animals living there (Aktar et al., 2009; Prieto-García et al., 2007; Siebe & Fisher, 1996).

Oribatid mites (Cryptostigmata) are, by their number and ecological role, one of the most dominant groups in most of the organic layers of soil where their densities can reach several hundreds of thousand specimens per m² (Borah & Kakati, 2013; Norton, 1990). Those mites can usually be found where there is an accumulation of organic matter with certain humidity and temperature ranges. Many oribatid mite species live in natural and altered conditions. Other species have a limited distribution or have a large abundance in places with an anthropogenic impact (Aoki, 1979; Eeva & Penttinen, 2009; Hunter, 2007). These mites are usually characterized by having a low fecundity and a very low capacity to increase their populations in a short time. Very few species have modifications for dispersion,

so they cannot escape from the stress of their environments. This characteristic has made them very important as bioindicators due to the incorporation of information about the habitat and niche history and profiles (Behan-Pelletier, 1999; Gulvik, 2007). One example of oribatid mites as bioindicators is *Humerobates rostralamellatus* which has a highly restricted relation to heavy metal pollutants (Lebrun & van Straalen, 1995). On the other hand, Al-Assiuty et al. (2000) have shown that species such as *Rhysotritia ardua*, *Oppia bifurcata*, *Niloppia sticta*, *Striatoppia niliaca*, and *Microzetes alces*, are restricted to residual muds, pointing out that *Scheloribates laevigatus* and *Epilohmannia cylindrica cylindrica* are very tolerant to soils with residual mud with high levels of organic matter, and heavy metals like Fe, Zn, Cu, Mg and Cd mainly.

Studies carried out in the Mezquital Valley on the quality of the irrigation water have shown that after 80 years of use, the content of metals in the plots irrigated with wastewater is 3 to 6 times higher than in the plots irrigated with well water, and particularly Cd, Pb, Cr, and Zn, are incorporated in the soil from the irrigation accumulated in the arable layer (Herre et al., 2004; Siebe, 1994). Among soil invertebrates, oribatid mites have been recorded as the group that can accumulate the highest heavy metal concentrations in different soils (van Straalen et al., 2001). The high concentration of heavy metals in soils can have a negative effect on oribatid mite richness and their reproduction (El-Sharabasy & Ibraim, 2010). Those negative effects can be direct by the poisoning of the specimens or indirect through the disturbance of the trophic web (Parmelee et al., 1993; Vladislav et al., 2015).

Due to the poor knowledge of the ecological function of the oribatid mites in agricultural regions of Mexico, this study of the oribatid mites was done in 2 different parcels with very contrasting irrigation types: one with residual water and the other with well water. In this way, we would be able to detect which species could be used as indicators of the perturbation of the soils. According to the literature, there are some oribatid mite species more tolerant to heavy

metal concentrations in soils, thus, we would expect a different composition of oribatid mites in both parcels due to their irrigation type and heavy metal contents, with more species tolerant in the residual water than in the well water parcel, and also more diversity in residual water parcel, since oribatid mites can develop in soils with high contents of heavy metals (van Straalen et al., 2001).

The knowledge of the structure and composition of the population of the oribatid mites and their relationship with edaphic parameters will be a basis for future studies on the environmental impact and can be of use in the agricultural strategies of Mexico and the recognition of oribatid mite species as bioindicators.

Materials and methods

The 2 plots are in the “Distrito de Riego 063” (DDR 063, 1997), in San Salvador Municipality (19°40'–20°29' N; 99°57'–99°27' W), situated in the Mezquital Valley, Hidalgo, at an average altitude of 1,985 m (Fig. 1). The weather of the region is semiarid dry, with an annual average temperature of 17 °C and an annual average precipitation of 475 mm (DDR 063, 1997; Ayuntamiento de San Salvador, Hgo., 2012). The municipality is in the Trans-Mexican Volcanic Belt. The soils are derived from basaltic, sedimentary and metamorphic rocks; among the intrusive rocks there are andesitic basaltic, basalts and diabasas; among the sedimentary there are hydroclastic, slates, argillaceous, limestone, limestone slates, marls, conglomerates, breaches, sands and alluviums; among the metamorphic rocks there is only marble (CNA, 1995). The site receives residual water from the most important draining of the river Tula and Mexico Valley. In the natural channel of the Tula River, the Endhó dam was built with agricultural aims. The Tula River starts from the infiltrations of the Requena dam which also stores water draining from the Taxhimay dam of the Tepeji River, the Central emitter and the El Salto River, which contributes to an annual water volume average of 498,3 million of m³ (Conagua, 2015).

Two plots were selected in the area, each one with a 7,000 m² extension. The 2 studied plots are similar in their pedogenic characteristics and the distance between them is about 1 km (CNA, 1995). One plot, named San Salvador (SS; 20°17'40" N, 99°00'01" W) has been irrigated with waste water for 80 years, and the other plot, El Bondho (B; 20°17'09" N, 99°00'31" W), has been irrigated with well water for 100 years. In each plot, an area of 50 × 20 m was selected for each sampling data. In that area, 10 samples (95 cm²) were taken randomly bimonthly from December 1998 to October 1999 for a total of 120 samples for all the study. On the same day of the collection, samples were




Figure 1. Localization of the plots San Salvador and El Bondho, in Hidalgo, Mexico (modified from INEGI, 1998).

taken to the laboratory where the fauna was extracted by Berlese-Tullgren funnels for 6 days. For 3 days, the samples were kept at room temperature, for the other 3 days, they were kept within the light from a 60-watt lamp. The oribatid mites were isolated, counted and mounted for their identification.

The physical-chemical analysis of the soil-like organic matter, pH, heavy metals, interchanging cations, porosity and electric conductivity (EC) were done in the Laboratorio de Edafología “Nicolás Aguilera” of the Faculty of Sciences, UNAM.

In order to characterize the community structure, the species richness (S), the Shannon diversity index (H'), the Simpson dominance index (λ), the Pielou's evenness (J'), and the Sørensen similarity index (Ludwig & Reynolds, 1988) were calculated. The Shannon diversity indices between localities were compared by a modified *t*-student test (Magurran, 1988).

The mean values of the edaphic parameters and the heavy metal contents in the 2 plots were compared with a *t*-student test for independent samples.

In order to detect the effect of the plot and the collection date on the oribatid mite abundance and the soil

parameters (pH, porosity, EC, Mg, Ca, K, Na), a factorial Anova test was performed in order to reduce the effect of pseudoreplication, using the interaction term as an error term (Camus & Lima, 1995; Manly, 1991). A Tukey's post hoc test was used to compare the months of collecting. The relationship between the edaphic parameters and the density of oribatid mites was evaluated by multiple correlation tests. A principal component test was performed to identify the variables that explain most of the variability observed in the communities of the oribatid mites. The data were normalized by $\sqrt{(X + 0.5)}$, except percentage that was transformed by $\text{ARCOSEN } \sqrt{X}$ (Zar, 1984). The

analyses were performed using the software Statistica, ver 6.0 (Statsoft, 1999).

Results

A total of 39,487 mites were collected in the 2 plots, the oribatids represented only the 6% of the total (2,448 individuals) including immature stages. In the SS plot, 1,345 oribatid mites were collected, while in the B plot 1,103 oribatids were recorded. A total of 20 species of oribatid mites belonging to 19 genera and 15 families were found in the study (Table 1).

Table 1

Abundance (number of individuals), density (ind/m²) and percentage of oribatid mites density in San Salvador (SS) and El Bondho (B), agricultural plots in Hidalgo, Mexico. H' = Shannon's diversity index; J' = Pielou's evenness index; S = species richness; λ = Simpson's dominance index.

Taxa	Abundance		Density		Percentage	
	SS	B	SS	B	SS	B
Thyrisomidae						
<i>Gemmazetes cavatica</i> (Kunst, 1962)	53	4	558	42	4	0.40
Oppiidae						
<i>Oppiella nova</i> (Oudemans, 1902)	181	37	1905	389	13	3
<i>Ramusella</i> sp.	1	116	10	1221	0.01	10
<i>Brachioppia</i> sp.	1	-	10	-	0.01	-
<i>Microppia</i> sp.	-	9	-	95	-	0.80
Oribatulidae						
<i>Zygoribatula connexa</i> (Berlese, 1904)	96	814	1010	8568	7	74
<i>Z. ca. Bonairensis</i>	1	4	10	42	0.01	0.40
Epilohmanniidae						
<i>Epilohmannia pallida</i> Balogh et Mahunka, 1980	3	36	31	379	0.02	3
Carabodidae						
<i>Carabodes ecuadoriensis</i> Balogh, 1988	3	-	31	-	0.02	-
Euphthiracaridae						
<i>Rhysotritia ardua</i> (C. L. Koch, 1841)	23	12	242	126	2	1
Scheloribatidae						
<i>Scheloribates</i> sp.	198	10	2,084	105	15	1
<i>Setobates</i> sp.	6	-	63	-	0.40	-
Xylobatidae						
<i>Xylobates</i> sp.	107	10	1,126	105	8	0.9
Hypochthoniidae						
<i>Hypochthonius</i> sp.	1	-	10	-	0.01	-
Lohmanniidae						
<i>Lohmannia banksi</i> Norton et al., 1978	6	-	63	-	0.40	-
Trhypochthoniidae						
<i>Allonothrus</i> sp.	-	-	-	10	-	0.10
Haplozetidae						
<i>Rostrozetes</i> sp.	7	24	74	253	0.50	2

Table 1
Continued

Taxa	Abundance		Density		Percentage	
	SS	B	SS	B	SS	B
Tectocephidae						
<i>Tectocephus velatus elegans</i> Ohkubo, 1981	651	19	6852	200	48	1.70
Ceratozetidae						
<i>Ceratozetes</i> sp.	4	7	42	74	0.30	1
Galumnidae						
<i>Galumna</i> sp.	3	-	31	-	0.20	-
Total	1,345	1,103	14,157	11,610	100	100
H'			1.064	1.092		
S			18	14		
J'			0.57	0.41		
λ			0.29	0.56		

In the SS plot, 18 species were recorded and 14 in the B plot. The most abundant species in SS was *Tectocephus velatus elegans*, while in B, *Zygoribatula connexa* was the most abundant. The highest values of abundance, diversity, evenness, and species richness were found in SS, only the dominance was higher in B. The exclusive species for SS were *Galumna* sp., *Lohmannia banksi*, *Hypochothonius* sp., *Setobates* sp., *Carabodes ecuadoriensis*, and *Brachiopopia* sp., and for B, *Microppia* sp. and *Allonothrus* sp. were exclusive (Table 1).

There were variations in the composition and the abundance of the oribatid mites in both plots. The highest species richness values were recorded in December and the lowest values were found in April in both plots. The highest values of Shannon's diversity index were observed in December and August and the lowest in February and June in B and SS, respectively. Nevertheless, significant differences between the Shannon diversity indices in the 2 localities ($t = 1.493$, $p > 0.05$) were not found.

The species *O. nova*, *Z. connexa*, and *Xylobates* sp. were present in all the collecting dates in SS, while in B, *Epilohmannia pallida* was the most constant species recorded in all the dates except during April. *Gemmazetes cavatica* was recorded in both plots only during December.

The Sørensen similarity coefficient between both communities was 75%. The highest values registered were in December and August, when more species are shared in both plots and the lowest value was recorded in February. The Anova used to evaluate the effect of the plot on the general density of oribatid mites did not show a statistically significant value. Nevertheless, when the species densities are compared and analyzed, there

is a significant effect of the locality, the data and the interaction of both variables on the densities of some species as *Oppiella nova*, *Ramusella* sp., *Microppia* sp., *Zygoribatula connexa*, *Scheloribates* sp., and *Xylobates* sp. (Table 2). For *Epilohmannia pallida*, only a significant effect of plot on density ($F_{1,108} = 13.21$; $p < 0.05$) was detected. The significant effect of the collecting date was detected on the density of *Gemmazetes cavatica* ($F_{5,108} = 2.75$; $p < 0.05$), *Carabodes ecuadoriensis* ($F_{5,108} = 3.87$; $p < 0.05$), and *Rhysotritia ardua* ($F_{5,108} = 3.47$; $p < 0.05$). The density of *Tectocephus velatus elegans* is affected significantly by the type of plot ($F_{1,108} = 46.39$; $p < 0.05$) and the collecting date ($F_{5,108} = 19.40$; $p < 0.05$), however, the interaction between them is not significant ($F_{5,108} = 1.01$; $p > 0.05$). In most of the species, the differences, according to the post hoc Tukey's test, were present between December and October ($p < 0.05$).

The average of the edaphic parameters and the heavy metals of the 2 plots is shown in table 3. The *t*-student test results show that there are significant differences in most of the edaphic parameters, except in the percentage of the organic matter and the porosity, and in relation with the heavy metals, only in Cr and Mn differences between plots were not detected (Table 3).

According with the Anova test, we found that in almost all edaphic parameters the plot (irrigation type) and the collection date have a significant effect on values, except on the organic matter; also there is a significant effect of plot to porosity percentage and only for the date for Na (Table 4), and in those parameters we did not find a significant effect with the interaction between the 2 variables. The post hoc Tukey's test shows that for

Table 2

Factorial Anova test results evaluate the effect of plot (P) and collecting date (D) on oribatid mites species in San Salvador and El Bondho, Hidalgo, Mexico. Pt = Plot; D = collecting data; P × D = interaction; * = $p < 0.05$.

Effect	df	F value
<i>Oppiella nova</i>		
P	1,108	12.55*
D	5,108	10.81*
P × D	5,108	6.81*
<i>Ramusella</i> sp.		
P	1,108	25.26*
D	5,108	9.24*
P × D	5,108	8.34*
<i>Microppia</i> sp.		
P	1,108	12.01*
D	5,108	3.87*
P × D	5,108	3.87*
<i>Zygoribatula connexa</i>		
P	1,108	19.30*
D	5,108	25.09*
P × D	5,108	21.11*
<i>Scheloribates</i> sp.		
P	1,108	15.80*
D	5,108	7.45*
P × D	5,108	5.72*
<i>Xylobates</i> sp.		
P	1,108	12.61*
D	5,108	3.84*
P × D	5,108	3.79*

Table 3

Average ± SD and *t*-student test values for edaphic parameters and heavy metals in San Salvador (SS) and El Bondho (B) plots. Mg = Magnesium; Ca = calcium; K = potassium; Na = sodium; EC = electric conductivity; OM = organic matter percentage; Po = porosity; Fe = iron; Cu = copper; Zn = zinc; Mn = manganese; Cr = chromium; Cd = cadmium; Ni = nickel; Pb = lead. N = 60; df = 118; < 0.05; ns = no significant.

Parameter	SS	B	t
Mg meq/100 g soil	24.20±12.92	44.33±16.39	7.47 *
Ca meq/100 g soil	16.85±8.20	23.04±14.05	2.95*
K meq/100 g soil	3.15±1.20	5.56±2.69	6.35*
Na meq/100 g soil	2.69±0.91	5.12±5.53	3.34*
pH	7.58±0.47	8.67±0.26	15.93*
EC ds/m	2.12±0.76	2.97±1.04	5.06*
OM %	2.77±3.63	2.84±2.72	0.11 ns
Po %	49.33±3.54	50.13±3.71	1.21 ns
Heavy metal			
Fe mgKg ⁻¹	5.76±4.17	1.69±1.15	7.28*
Cu mgKg ⁻¹	4.57±4.09	0.93±1.04	6.68*
Zn mgKg ⁻¹	14.13±13.12	1.63±1.32	7.35*
Mn mgKg ⁻¹	14.27±17.19	13.41±20.47	0.25 ns
Cr mgKg ⁻¹	0.03±0.05	0.03±0.05	0.005 ns
Cd mgKg ⁻¹	0.23±0.17	0.05±0.06	7.86*
Ni mgKg ⁻¹	1.36±0.84	0.36±0.41	8.25*
Pb mgKg ⁻¹	2.74±2.32	0.97±0.76	5.61*

interchanged cations (Mg, Ca, K and Na), the differences are present in February ($p < 0.05$), while for the porosity percentage, the differences are detected in April for both plots. For the pH, the most different months are April and December, for the SS and the B plots, respectively; for EC, the differences are detected in December for the SS, and in April for the B plot. In the case of the heavy metals, significant effects of the plot and the collection date were found, and the interaction between the 2 variables on the contents of all elements, except for Mn in which only the collecting date had a significant effect, and for Cr in which no effect was found (Table 5).

The correlation analysis between the edaphic parameters, heavy metals, and oribatid mites shows that the most significant relationship in SS was presented by *T. velatus elegans* ($r = 0.80$), which was positively correlated

with Mn and Cr, and negatively with Zn. *Scheloribates* sp. had a positive and significant correlation with K ($r = 0.74$). *Rhysotritia ardua* also had a negative correlation with K and Mn, and positive with Cu and Ni ($r = 0.69$). On the other hand, in B, *Allonothrus* sp. had a positive correlation with the porosity and Cd ($r = 0.72$). *Ramusella* was negatively correlated with Ca and Mg, and had a positive correlation with porosity ($r = 0.69$).

The principal component analysis has shown that there are 2 main factors: the first one is related to the heavy metals (Fe, Cu, Zn, Cd, Ni, Pb) and the pH, and it explains 42.51% of the variation (eigenvalue = 8.80). The second factor is related to the content of K and Ca, and it explains 56.31% of variation (eigenvalue = 2.21), as shown in figure 2.

Table 4

Factorial Anova test results evaluate the effect of plot (P) and collecting date (D) and interaction between variables ($P \times D$) on edaphic parameters in San Salvador and El Bondho. Mg = Magnesium; Ca = calcium; K = potassium; Na = sodium; EC = electric conductivity; Po = porosity percentage; OM = organic matter percentage. * = $p < 0.05$, ns = no significant.

Effect	df	F value
Mg		
P	1,108	108.86*
D	5,108	20.17*
$P \times D$	5,108	4.23*
Ca		
P	1,108	17.84*
D	5,108	22.40*
$P \times D$	5,108	ns
K		
P	1,108	159.84*
D	5,108	61.04*
$P \times D$	5,108	10.99*
Na		
P	1,108	11.72*
D	5,108	ns
$P \times D$	5,108	ns
pH		
P	1,108	984.7*
D	5,108	61.5*
$P \times D$	5,108	8.5*
EC		
P	1,108	72.19*
D	5,108	37.62*
$P \times D$	5,108	7.24*
Po		
P	1,108	ns
D	5,108	5.61*
$P \times D$	5,108	ns
OM		
P	1,108	ns
D	5,108	ns
$P \times D$	5,108	ns

Table 5

Anova two- way test results, evaluate the effect of plot (P) and collecting date (D) on the heavy metals contents in San Salvador and El Bondho. Fe = Iron; Cu = copper; Zn = zinc; Mn = manganese; Cr = chromium; Cd = cadmium; Ni = nickel; Pb = lead. * = $p < 0.05$, ns = no significant.

Effect	df	F value
Fe		
P	1,108	121.33*
D	5,108	21.46*
P × D	5,108	10.90*
Cu		
P	1,108	137.89*
D	5,108	34.91*
P × D	5,108	16.35*
Zn		
P	1,108	162.61*
D	5,108	27.44*
P × D	5,108	22.05*
Mn		
P	1,108	ns
D	5,108	3.13*
P × D	5,108	ns
Cr		
P	1,108	ns
D	5,108	ns
P × D	5,108	ns
Cd		
P	1,108	123.90*
D	5,108	20.31*
P × D	5,108	5.47*
Ni		
P	1,108	82.85*
D	5,108	3.59*
P × D	5,108	3.51*
Pb		
P	1,108	102.41*
D	5,108	43.90*
P × D	5,108	11.22*




Figure 2. Principal components analysis for edaphic factors and heavy metals on the abundance of oribatid mites in Hidalgo, Mexico.

Discussion

The densities of oribatid mites recorded in this study were much lower than those found in forest soils and even in those in agricultural soils (Maribie et al., 2011; Yina et al., 2013) although it is known that this group of mites is usually the dominant species in the edaphic environments (Norton, 1990).

These results can be related to the use of pesticides and fertilizers in the agricultural practices which usually have a negative effect on the abundance, distribution and composition of the species (Desmond & Alex, 2013; Minor et al., 2004; Socarrás & Robaina, 2011). In the present study, despite the high concentrations of heavy metals in the parcel watered with residual water, it had the highest values of abundance and diversity. The amount of some oribatid mite species like *Tectocephus velatus elegans*, *Oppiella nova*, *Scheloribates* sp., *Gemmazetes cavatica*, and *Xylobates* sp., seems to increase with the presence of several elements like Cr and Mn. It was observed that the densities of such populations were lower with the diminution of the concentration of the heavy metals in the crops in August and October. These results confirm the resistance that oribatid mites show to concentrations of heavy metals. This resistance was found in other studies in which there are soil invertebrates that can accumulate most of those metals in their bodies (van Straalen et al., 2001). Also, the recorded species, along with *Zygoribatula connexa*, which was the most abundant in B, have been frequently recorded in agricultural soils (Hubert, 2000; Ivan & Călugăr, 2013); even *Scheloribates* has been cited from environments polluted with heavy metals and metallurgic accumulations (Al-Assiuty et al., 2000; Corral & Iturrondobeitia, 2012; El-Sharabasy & Ibraim, 2010; Skubala, 1995).

Some studies performed in forest soils in Germany established that *Tectocephus* sp. and some members of the family Oppiidae, mainly *Oppiella nova*, were not affected in their abundance after the soils were perturbed. This species has preference to soils with an acid pH which can indicate a high alteration. It has been suggested that such resistance can be related to its very high reproductive rate, because this species is parthenogenetic (El-Sharabasy & Ibraim 2010; Maraun et al., 2003; Vladislav et al., 2015). According to Aoki (1979), the oribatid mites which are more resistant to different kinds of pollution are members of the families Brachychthoniidae, Oppiidae, Oribatulidae, and Tectocephidae, which except for Brachychthoniidae, were well represented in this study.

The incorporation of low concentrations of Cu (42-418 mg) in the diet of some oribatid mites can increase their fertility, however, if they exceed certain limits (above 700

mg), their life cycle can be affected and cause their death (Seniczak et al., 1997, 1999). The concentration in both of our plots was lower than 15 mg, so it can be beneficial to those mites (Skubala & Zaleski, 2011).

Skubala and Kafel (2004) have studied the bioaccumulation of metals in oribatid mites in a gradient of forestry ecosystems. They have found that the concentration of Zn and Cu was higher in the mycophagous species (*Oppiella nova* and *Tectocephus velatus*) than in the panfitophagous species (*Oribatula tibialis* and *Pergalumna nervosa*) or in the macrofitophagous (*Atropacarus striculus*). Other studies have demonstrated that the communities of some oribatid mites are almost completely tolerant to the presence of Fe, Zn, Cu, Cd, and Pb in the soil (Vladislav et al., 2015; Zaitsev & van Straalen, 2001). It has been observed that Zn is the metal which accumulates the most in the microfitophagous oribatid mites (Zaitsev & van Straalen, 2001). This may explain the reason for a positive correlation of this metal and the densities of *Oppiella nova* and *Tectocephus elegans*, considered as microfitophagous, in the present work in SS.

As an answer to the effect of the pollutants on populations of oribatid mites, Rusek & Marshall (2000) proposed a classification system: a) sensitive species like *Adoristes ovatus*, *Eporibatula rauscheninsis* and *Oppiella minus*; b) susceptible species such as *Carabodes labyrinthicus* and *Oribatula tibialis*, and c) tolerant species like *Chamobates schueltzi*, *Liochthonius* sp., *Tectocephus velatus*, *Trichoribates trimaculatus*, and *Zygoribatula exilis*.

Following the system of Rusek & Marshall (2000), only *Tectocephus velatus elegans* and *Carabodes ecuadoriensis* could be included in our study. However, *Oppiella nova* and *Zygoribatula connexa* can also be considered as tolerant species because they are very abundant in SS and B.

After this study, we can say that many species of *Tectocephus*, *Oppiella*, *Zygoribatula*, and *Scheloribates* are abundant and constant in those environments where disturbance exists or that have been polluted with heavy metals (Al-Assiuty et al., 2000; Gan, 2013; Vladislav et al., 2015), or in those grounds that have been used for agricultural purposes (Norton & Sillman, 1985; Ruiz et al., 1986).

Studies on the accumulation and presence of heavy metals in soils in the area where the present study was carried out (until 1994) show that metals do not represent a risk for the productive potential and they have not been incorporated to trophic web (Flores-Magdaleno et al., 2011; Siebe, 1994). In view of our results, we believe that those levels of metals are still very similar.

The differences between the population abundance of both parcels may be due to the tolerance (or the intolerance) of some species to the presence of some heavy metals and to the microclimatic conditions of each parcel. Some species do not stand the high concentrations of some metals, however, some other species increase their populations where there are moderate concentrations of those metals (Skubala & Zaleski, 2011) and they most likely use and incorporate them in their metabolism (Corral & Iturrondobeitia, 2012; Skubala & Kafel, 2004).

In this study, *Oppeiella nova*, *Rhysotritia ardua*, *Scheloriabates* sp., and *T. velatus elegans* were the most abundant species in the heavy metal polluted parcel. On the other hand, they had a low density in the parcel watered with well water. *Epilohmannia pallida*, *Ramusella* sp., and *Zygoribatula connexa*, were the most abundant species in the less polluted parcel. The species of this last parcel have lower requirements of heavy metals or a low resistance to heavy metals, but they prefer soils with a higher pH.

To confirm the inferred relationships, it is necessary to make ecotoxicological studies at specific level in laboratory conditions (also in the field under natural conditions) to ascertain the accumulation percentage of heavy metals in the tissues of mites, in order to evaluate the impact of some pollutants in the residual waters used for agricultural purposes. We found information on the tolerance of some species such as *O. nova*, *R. ardua*, and *T. elegans* to heavy metals, and low resistance to heavy metals of *Epilohmannia pallida* and *Z. connexa*, that are present in the less polluted parcel, but are tolerant to saline soils. Results found in our study may be useful to support the use of some species of oribatid mites as bioindicators in environments with conditions similar to our study area.

Acknowledgements

This contribution was done with the support (project IN201598) of Dirección General de Asuntos del Personal Académico, UNAM, México, in charge of JGPV. The final version was reviewed by Luis Parra. Dr. Elizabeth Franklin (INPA-Brazil) and other anonymous reviewers gave valuable suggestions to improve the manuscript.

References

- Aktar, M. W., Sengupta, D., & Chowdhury, A. (2009). Impact of pesticides use in agriculture: their benefits and hazards. *Interdisciplinary Toxicology*, 1, 1–12.
- Al-Assiuty, A. I. M., Khalil, M. A., & Abdel-Lateif, H. M. (2000). Effects of dry sludge application on soil microarthropod communities in a reclaimed desert ecosystem. *Pedobiologia*, 44, 567–578.
- Aoki, J. (1979). Difference in sensitivities of oribatid families to environmental change by human impacts. *Revue d'Ecologie et Biologie du Sol*, 16, 415–422.
- Ayuntamiento de San Salvador, Hgo. (2012). *Plan de Desarrollo Municipal de San Salvador, Hgo. 2012-2016*. San Salvador, Hidalgo: Presidencia Municipal de San Salvador.
- Behan-Pelletier, V. M. (1999). Oribatid mite biodiversity in agroecosystems: role for bioindication. *Agriculture, Ecosystems and Environment*, 74, 411–423.
- Borah, M., & Kakati, L. N. (2013). Abundance and distribution of soil Acarina in natural and degraded forest ecosystems at Pathalipam, Lakhimpur, Assam. *International Journal of Scientific & Engineering Research*, 12, 1694–1709.
- Braatz, S., & Kandiah, A. (1996). The use of municipal waste water for forest and tree irrigation. *Unasylva*, 185, 45–51.
- Camus, P. A., & Lima, M. (1995). El uso de la experimentación en ecología: supuestos, limitaciones, fuentes de error, y su status como herramienta explicativa. *Revista Chilena de Historia Natural*, 68, 19–42.
- CNA (Comisión Nacional del Agua). (1995). *Impacto del riego con aguas residuales en las aguas subterráneas del Valle del Mezquital, Hidalgo*. Reporte Fase I. Departamento de Desarrollo Exterior. México D.F.: Comisión Nacional del Agua/ Brithis Geological Survey.
- Conagua (Comisión Nacional del Agua). (2015). *Actualización de la disponibilidad de agua en el acuífero Actopan-Santiago de Anaya (1313), Estado de Hidalgo*. México D.F.: Comisión Nacional del Agua.
- Corral, E. H., & Iturrondobeitia, J. C. (2012). Effects of cattle and industries on oribatid mites communities of grassland soil in the Basque Country (Spain). *International Journal of Acarology*, 3, 217–229.
- DDR 063 (Distrito de Desarrollo Rural). (1997). Características generales de los Distritos de Riego 003-Tula y 100 Alfajayucan. México D.F.: Comisión Nacional del Agua, Distrito de Riego 03 de Mixquiahuala.
- Desmond, A. O. D., & Alex, U. O. (2013). A comparative assessment of soil arthropod abundance and diversity in practical farmlands of University of Ibadan, Nigeria. *The International Journal of Environmental Resources Research*, 1, 17–29.
- Eeva, T., & Penttinen, R. (2009). Leg deformities of oribatid mites as an indicator of environmental pollution. *Science of the Total Environment*, 407, 4771–4776.
- El-Sharabasy, H. M., & Ibrahim, A. (2010). Communities of oribatid mites and heavy metal accumulation in oribatid species in agricultural soils in Egypt impacted by waste water. *Plant Protection Science*, 4, 159–170.
- Flores-Magdaleno, H., Mancilla-Villa, O. R., Mejía-Saenz, E., Olmedo-Bolaños, M. C., & Bautista-Olivas, A. L. (2011). Heavy metals in agricultural soils and irrigation wastewater of Mixquiahuala, Hidalgo, Mexico. *African Journal of Agricultural Research*, 24, 5505–5511.
- Gan, H. (2013). *Oribatid mite communities in soil: structure, function and response to global environmental change* (Ph.D.

- Thesis*). University of Michigan. Ann Arbor, Michigan, USA.
- Gulvik, M. E. (2007). Mites (Acari) as indicators of soil biodiversity and land use monitoring: a review. *Polish Journal of Ecology*, 3, 415–440.
- Herre, A., Siebe, C., & Kaupenjohann, M. (2004). Effect of irrigation water quality on organic matter, Cd and Cu mobility in soils of central Mexico. *Water Science and Technology*, 2, 277–284.
- Hubert, J. (2000). The oribatid community (Acari: Oribatida) on dry cow pasture. *Ekológia (Bratislava)*, 19, 354–364.
- Hunter, P. (2007). The human impact on biological diversity. How species adapt to urban challenges sheds light on evolution and provides clues about conservation. *Embo Reports*, 4, 316–318.
- INEGI (Instituto Nacional de Estadística Geografía e Informática). (1998). Cuaderno Estadístico Municipal. San Salvador, estado de Hidalgo, INEGI, Aguascalientes.
- Ivan, O., & Călugăr, A. (2013). A peculiarities of the edaphic mesofauna in some cultivated soils from the Central Moldavian Plateau. *Lucrări Științifice*, 2, 125–130.
- Lebrun, P., & van Straalen, N. M. (1995). Oribatid mites: prospects for their use in ecotoxicology. *Experimental and Applied Acarology*, 19, 361–380.
- Ludwig, J. A., & Reynolds, J. F. (1988). *Statistical ecology: a primer of methods and computing*. New York: Wiley Press.
- Magurran, A. (1988). *Ecological diversity and its measurement*. New Jersey: Princeton University Press.
- Manly, B. F. J. (1991). *Randomization and Monte Carlo methods in Biology*. New York: Chapman and Hall.
- Maraun, M., Salamon, J. A., Schneider, K., Schaefer, M., & Scheu, S. (2003). Oribatid mite and collembolan diversity, density and community structure in a moder beech forest (*Fagus sylvatica*): Effects of mechanical perturbations. *Soil Biology and Biochemistry*, 35, 1387–1394.
- Maribie, C. W., Nyamasyo, G. H. N., Ndegwa, P. N., Mung'atu, J. K., Lagerlöf, J., & Gikungu, M. (2011). Abundance and diversity of soils mites (Acari) a long gradient of land use types in Taita, Taveta, Kenya. *Tropical and Subtropical Agroecosystems*, 13, 11–26.
- Minor, M. A., Volk, T. A., & Norton, R. A. (2004). Effects of site preparation techniques on communities of soil mites (Acari: Oribatida, Acari: Gamasida) under short-rotation forestry plantings in New York, USA. *Applied Soil Ecology*, 25, 181–192.
- Norton, R. A. (1990). Acarina. Oribatida. In D. L. Dindal (Ed.), *Soil biology guide* (pp.779–803). New York: Wiley, Interscience Publication.
- Norton, R. A., & Sillman, D. Y. (1985). Impact of only waste application on the mite community of an arable soil. *Experimental Applied Acarology*, 1, 287–306.
- Parmelee, R. W., Wentsel, R. S., Phillips, C. T., Simini, M., & Checkai, R. T. (1993). Soil microcosm for testing the effects of chemical pollutants on soil fauna communities and trophic structure. *Environmental Toxicology and Chemistry*, 12, 1477–1486.
- Prado, B., Siebe, C., Bischoff, W. A., Hernández-Martínez, L., & Mora, L. (2015). El suelo: guardián de la calidad del agua subterránea. *Conabio. Biodiversitas*, 122, 6–9.
- Prieto-García, F., Méndez-Marzo, M. A., Martínez-Pezina, F. H., & Prieto-Méndez, J. (2007). Presencia de metales pesados en cultivos del Valle del Mezquital, México. *Revista Latinoamericana de Recursos Naturales*, 2, 100–110.
- Romero, A. H. (1997). El Valle del Mezquital. Estudio de caso. En R. Helmer e I. Espanhol (Eds.), *Water pollution control. A guide to the use of water quality management principles* (pp. 1–10). México D.F.: PNUMA/ CCAIS/ OMS.
- Ruiz, E., Mínguez, M. E., & Subías, L. S. (1986) Los Oribátidos (Acari, Oribatida) de los eriales de cultivo de una zona agrícola del sur de Madrid y efecto borde. *Actas VIII Jornadas de la Asociación Española de Entomología*. Sevilla, Spain.
- Rusek, J., & Marshall, V. G. (2000). Impacts of airborne pollutants on soil fauna. *Annual Review of Ecology and Systematics*, 31, 395–423.
- Sagasta, J., Medlicott, M. K., Qadir, M., Raschid-Sally, L., Dreschel, P., & Liebe, J. (2013). *Proceedings of the UN-Water project on the Safe Use of Wastewater in Agriculture*. UN-Water Decade Programme on Capacity Development (UNW-DPC). United Nations University.
- Seniczak, A., Seniczak, S., & Długosz, J. (1997). The influence of copper on the development, fertility and mortality of *Archeogozetes longisetosus* Aoki (Acari: Oribatida) in the laboratory conditions. *Arbeitsangun Mengen-spurenelemente*, 1997, 620–626.
- Seniczak, A., Seniczak, S., & Długosz, J. (1999). The effect of lead and copper on *Archeogozetes longisetosus* Aoki (Acari, Oribatida) in laboratory conditions. In K. Tajovský, K. Schlaghamerský, & V. Pízl (Eds.), *Contributions to Soil Zoology in Central Europe I* (pp. 289 –293). České Budejovice: ISB AS CR.
- Siebe, C. (1994). Acumulación y disponibilidad de metales pesados en suelos regados con aguas residuales, en el Distrito de Riego 03, Tula, Hgo., México. *Revista Internacional de Contaminación Ambiental*, 10, 15–21.
- Siebe, C. & Cifuentes, E. (1995). Environmental impact of wastewater irrigation in Central México: an overview. *International Journal of Environmental Health Research*, 5, 161–173.
- Siebe, C., & Fisher, W. R. (1996). Effect of long- term irrigation with untreated sewage effluents on soil properties and heavy metal adsorption of Leptosols and Vertisols in Central México. *Z. Pflanzenernähr. Bodenk.*, 159, 357–364.
- Skubala, P. (1995). Moss mites (Acarina: Oribatida) on industrial dumps of different age. *Pedobiologia*, 39, 170–184.
- Skubala, P., & Kafel, A. (2004). Oribatid mite communities and metal bioaccumulation in oribatid species (Acari, Oribatida) along the heavy metal gradient in forest ecosystems. *Environmental Pollution*, 132, 51–60.
- Skubala, P., & Zaleski, T. (2011). Heavy metal sensitivity and bioconcentration in oribatid mites (Acari, Oribatida) Gradient study in meadow ecosystems. *Science of the Total Environment*, 2012, 364–372.
- Socarrás, A. A., & Robaina, N. (2011). Caracterización de

- la mesofauna edáfica bajo diferentes usos de la tierra en suelo Ferralítico Rojo de Mayabeque y Artemisa. *Pastos y Forrajes*, 2, 185–198.
- Statsoft Inc. (1999). *Statistical user guide*. Complete Statistical System Statsoft. Oklahoma, USA.
- van Straalen, N. M., Butovsky, R. O., Pokarzhevski, A. D., Zaitsev, A. S., & Verhoef, S. C. (2001). Metal concentrations in soil and invertebrates in the vicinity of a metallurgical factory near Tula (Russia). *Pedobiologia*, 45, 451–466.
- Vladislav, S. A., Pavel, A., Barsukov, P. A., & Bashkin, V. N. (2015). Application of Soil Oribatid Mites as Bioindicators in Impact Areas of the Gas Industry in the West Siberian Tundra. *The Open Ecology Journal*, 2015 (Suppl 1-M4), 32–39.
- Yina, M. G. J., Castillo, J. A. F., & Bacca, T. (2013). Ácaros oribátidos presentes en seis sistemas de uso del suelo en Obonuco, Pasto (Nariño). *Boletín Científico Museo de Historia Natural*, 2, 60–68.
- Zaitsev, A. S., & van Straalen, N. M. (2001). Species diversity and metal accumulation in Oribatid mites (Acari: Oribatida) of forest affected by a metallurgical plant. *Pedobiologia*, 45, 467–479.
- Zar, J. H. (1984). *Biostatistical analysis*, 2nd. Edition. Nueva Jersey: Englewood Cliffs.