



Revista Boliviana de Química

ISSN: 0250-5460

ISSN: 2078-3949

revbolquim@outlook.com

Universidad Mayor de San Andrés

Bolivia

Bhushan Tewari, Brij

BASIC PHYTOCHEMICAL SCREENING AND ANTIBACTERIAL, ANTIFUNGAL AND
ANTIOXIDANT PROPERTIES OF SYZYGium CUMINI, A TREE FROM GUYANA

Revista Boliviana de Química, vol. 37, no. 3, 2020, July-, pp. 132-141

Universidad Mayor de San Andrés

Bolivia

Available in: <https://www.redalyc.org/articulo.oa?id=426365043001>

- How to cite
- Complete issue
- More information about this article
- Journal's webpage in redalyc.org

redalyc.org

Scientific Information System Redalyc

Network of Scientific Journals from Latin America and the Caribbean, Spain and
Portugal

Project academic non-profit, developed under the open access initiative



**BASIC PHYTOCHEMICAL
SCREENING AND ANTIBACTERIAL,
ANTIFUNGAL AND ANTIOXIDANT
PROPERTIES OF SYZYGIUM CUMINI,
A TREE FROM GUYANA**

**CRIBADO FITOQUÍMICO BÁSICO Y
PROPIEDADES ANTIBACTERIANAS,
ANTIFÚNGICAS Y ANTIOXIDANTES DE
SYZYGIUM CUMINI, UN ÁRBOL DE
GUYANA**

Received 03 24 2020
Accepted 08 26 2020
Published 08 26 2020

Vol. 37, No.3, pp. 132-141, Jul./Ago.2020
Revista Boliviana de Química

37(3), 132-141, Jul./Aug. 2020
Bolivian Journal of Chemistry
DOI: 10.34098/2078-3949.37.3.1



Full original article

Peer-reviewed

Brij Bhushan Tewari*

Department of Chemistry, Faculty of Natural Sciences, University of Guyana, Po Box 101110 Georgetown, Guyana. E-mail: brijtew@yahoo.com

Keywords: *Syzygium cumini*, Antioxidant power, Antifungal activity, Phytochemical screening, antibacterial property.

Palabras clave: *Syzygium cumini*, Poder antioxidante, Actividad antifúngica, Screening fitoquímico, Propiedades antibacterianas.

ABSTRACT

Syzygium cumini or Jamun is an evergreen tropical tree in ornamental plant family Myrtaceae. The plant material (leaves) of *Syzygium cumini* (Jamun) was collected at the Institute of Applied Science and Technology (IAST), University of Guyana, Turkeyen Campus. Leaves were dried in oven at 50-60 °C for 72 h. The moisture content was calculated. The dried leaves were grounded and extracted in ethanol, methanol, ethyl acetate, and chloroform solvents, succesively. Extracts were collected and evaporation of solvent was done on rotavapour. The respective solvent was added to viscous semi solid liquid extract to make up the desired volume of extract solution. The micro-organisms (*Escherichia coli*, *Staphylococcus aureus* and *Candida albicans*) were obtained from GPHC, Georgetown, Guyana. The antioxidant, antimicrobial and antifungal activity of the different extracts was assessed by methods reported in the literature. The maximum and the minimum antioxidant power was exhibited by the methanol and the chloroform extracts, respectively. The chloroform and the ethyl acetate extracts were found to have the maximum and the minimum antibacterial activity against *Escherichia coli*, *Staphylococcus aureus*, as well as the antifungal activity against *Candida albicans*, respectively by the disc diffusion method. Phytochemical analysis of the *Syzygium cumini* leave extracts revealed the presence of carbohydrates, terpenoids, proteins, amino acids and flavonoids.



*Corresponding author: brijtew@yahoo.com

RESUMEN

Syzygium cumini o Jamun es un árbol tropical de hoja perenne de la familia de plantas ornamentales Myrtaceae. El material vegetal (hojas) de *Syzygium cumini* (Jamun) se colectó en el Instituto de Ciencia y Tecnología Aplicadas (IAST) de la Universidad de Guyana, campus de Turkeyen. Las hojas se secaron en estufa a 50-60 °C durante 72 h. Se calculó el contenido de humedad. Las hojas secas se trituraron y se extrajeron en etanol, metanol, acetato de etilo y disolventes de cloroformo, sucesivamente. Se recogieron los extractos y se evaporó el disolvente en rotavapor. Se añadió el disolvente respectivo al extracto líquido semisólido viscoso para completar el volumen deseado de solución de extracto. Los microorganismos (*Escherichia coli*, *Staphylococcus aureus* y *Candida albicans*) se obtuvieron de GPHC, Georgetown, Guyana. La actividad antioxidante, antimicrobiana y antifúngica de los diferentes extractos se evaluó mediante métodos descritos en la literatura. El poder antioxidante máximo y mínimo fue exhibido por los extractos de metanol y cloroformo, respectivamente. Se encontró que los extractos de cloroformo y acetato de etilo tenían la actividad antibacteriana máxima y mínima contra *Escherichia coli*, *Staphylococcus aureus*, así como la actividad antifúngica contra *Candida albicans*, respectivamente, por el método de difusión por disco. El análisis fitoquímico de los extractos de hojas de *Syzygium cumini* reveló la presencia de carbohidratos, terpenoides, proteínas, aminoácidos y flavonoides.

INTRODUCTION

Syzygium cumini or Jamun is an evergreen tropical tree belonging to the flowering plant family myrtaceae. The scientific classification of *Syzygium cumini* is given as follows:

Kingdom:	<i>Plantae</i>
	<i>Angiosperms</i>
	<i>Endicots</i>
	<i>Rosids</i>
Order:	<i>Myrtales</i>
Family:	<i>Myrtaceae</i>
Genus:	<i>Syzygium</i>
Species:	<i>Syzygium cumini</i>

Syzygium cumini is also known as black plum, java plum, duhat plum and malabar plum. More than one thousand species were classified as belonging to the genus *Syzygium*. It is a slow growing species. It can reach up to 30 meters and live more than 100 years. *Syzygium cumini* is original from India, Nepal, Pakistan, Bangladesh, Sri Lanka, Philippines and Indonesia. It was is man-introduced to Florida, USA, Suriname, Brazil, Trinidad & Tobago and Guyana. *Syzygium cumini* leaves are used as food for live stocks. Wine (and vinegar) are also made from the fruits. It is rich in vitamins A and C. Leaves and bark of *S. cumini* are used for controlling blood pressure. *Syzygium cumini* seeds are used in various alternative healing system like Unani, Chinese and Ayurveda medicine. *Syzygium cumini* wood is water resistance, so it is used in railway sleepers and to installing motors in wells. Kaur et al. [1] investigated drying (50, 70, 90 and - 55° C) characteristics and antioxidant properties of java plum seeds and skin waste. A review on ethnobotanical uses, antimicrobial potential, pharmacological properties and phytoconstituents of *Syzygium cumini* was reported by Singh and Navneet [2]. Mitra et al. [3] examined effects of summer, winter, rainy and autumn seasons on UV absorbing properties of the acetone extract of *S. cumini* leaves pointing out the rainy season as responsible of maximum absorbing properties and it can be regarded as anti-solar agent for preparation of sun screen lotions. The UV absorption properties of the methanolic, ethanolic, benzenic, acetone and ethyl acetate leave extract of *S. cumini* were examined by Mitra et al. [4]. The acetone leave extract of *S. cumini* was found to have maximum UV absorption potential. Arunpandian et al. [5] presented a review focused on the information on traditional and medicinal use of *S. cumini*. Phytochemical contents and antioxidant capacity of *S. cumini* from Indonesia were investigated by Sukmasari et al. [6]. They have also compared the properties of the leave extract of *S. cumini* and *S. polyanthum*. Prema et al. [7] studied the phytochemical constituents and antidiabetic properties of *S. cumini* seeds extract. *S. cumini* seeds powder was found to possess antidiabetic properties in type - 2 diabetic model rats. The potential use of *S. cumini*



methanolic leave extract for dengue by increasing platelet and leukocyte levels were described by Bandiola and Corpuz [8]. A review on pharmacological properties and therapeutic potential of *S. cumini* was discussed by Kumawat et al. [9]. Antimicrobial activity of *S. cumini* juice extract on enteric pathogenic bacteria (viz. *Salmonella typhimurium*, *Shigella flexneri*, *Staphylococcus aureus* and *Escherichia coli*) was described by Haque et al. [11]. Khan et al. [12] studied the antifungal potential of the methanolic extract of bark and leaves of *S. cumini* against *Rhizoctonia solani*. Kuhn. Hasanuzzaman et al. [13] performed phytochemical screening of methanolic, acetone, chloroform and n-hexane extracts of *S. cumini* seeds, root, stem, bark, leaves. The antimicrobial potential of methanolic, aqueous, and petroleum ether extracts of *S. cumini* leaves was tested by Elfadil et al. [14] against bacteria (viz. *Staphylococcus aureus*, *Escherichia coli*, *Pseudomonas aeruginosa*, *Bacillus subtilis*) and fungi (*Aspergillus niger* and *Candida albicans*). Ramos and Bandiola [10] has examined phytochemicals in extracts from methanol, ethanol, water, ethyl acetate and hexane macerations out of the leaves of *S. cumini*. Praphakaran et al. [15] reported that ethanolic extract of leaves and aqueous extracts of seeds *S. cumini* were found to have very high antimicrobial property for wide range of gram positive and gram negative bacterial strains. The natural antimicrobial potential of extract of all parts of *Syzygium cumini* (L) over the chemical synthetic antibiotic was studied by Mohit et al. [16]. Jabeen and Javaid [17] evaluated the antifungal potential of the ethanolic, aqueous and n-hexane extracts from leaves, fruit, root bark and stem bark of *Syzygium cumini* (L) against *Ascochyta arabiei*, the cause of blight disease of chick pea (*Cicer arietinum*). The antioxidant activity of *S. cumini* leaf extracts was investigated using the 2, 2-diphenyl-1-picrylhydrazyl (DPPH) free radical-scavenging and ferric-reducing antioxidant power (FRAP) arrays was described by Ruan et al. [18]. A significant linear relationship between antioxidant potency, free radical-scavenging ability and the content of phenolic compounds of leaves extracts were observed. The plant of *Syzygium cumini* with fruits is shown in Figure 1.



Figure 1. The *Syzygium cumini* plant with fruits

RESULTS AND DISCUSSION

Moisture contents

The moisture contents were calculated and are exposed in Table1



Reducing power

The reducing power serves to measure the reductive ability of an antioxidant, and it is evaluated by the transformation of Fe(III) to Fe(II) in the presence of the sample extracts, Gulcin et al. [19]. The reducing power of *Syzygium cumini* leaves extracts are summarized in Table 2. From Table 2 it is clear that reducing power increased with an increase in the concentration of plant extracts. The ability to reduce Fe (III) may be attributed to hydrogen donation from phenolic compounds, according to Shimada et al. [20], which is also related to the presence of the reducing agent, Duh [21]. In addition, the number and position of the hydroxyl group of phenolic compounds also rule their antioxidant activity, Sawaddiwong et al. [22]. Some deviation from the increase in the reducing power of leave extract with an increase in concentration may be due to decrease in hydrogen donor ability of phenolic compounds.

Table 1. Percentage moisture content for *Syzygium cumini* leaves

No. of Leaves Packets	Wt. of green leaves grams	Wt. of leaves after 24h (gr)	Wt. of leaves after 48h (gr)	Wt. of leaves after 72h (gr)	Percentage moisture content
1	62.61	43.68	43.53	44.01	30.47
2	56.57	43.82	43.92	44.02	22.54
3	57.04	46.29	46.39	46.92	18.85
4	96.73	66.68	66.49	67.01	31.26
5	77.57	48.58	48.56	49.06	37.40
6	82.80	52.26	52.21	52.50	36.94
7	80.98	49.01	49.10	49.62	39.37
8	88.26	51.27	51.25	51.30	41.93
9	139.12	75.83	74.29	74.81	46.60
10	109.17	59.99	60.10	60.32	45.05
11	87.76	62.98	62.10	62.32	29.24

$$\text{Percentage moisture content} = \frac{\text{weight of green leaves} - \text{weight of dry leaves}}{\text{weight of green leaves}} \times 100$$

It is observed from Table 2 that the antioxidant power of *Syzygium cumini* leaves extract in various solvents follow the order:

Methanol > ethyl acetate > ethanol > chloroform

In the ethanol extract high absorbance was observed at 1.0 µL concentration. From 2.0 to 10.0 µL concentration no definite order of reducing antioxidant power was observed. The methanol extract showed high reducing anti-oxidant power or high difference in the absorbance of leaves extract and control. In ethyl acetate extract maximum difference in the absorbance of leaves extract and control was observed at 7.0 µL concentration. In chloroform extract minimum reducing or antioxidant power or minimum difference in absorbance of leaves extract and control was observed.



Table 2. Reducing antioxidant power of ethanolic, methanolic, ethyl acetate and chloroform leave extracts of *Syzygium cumini* (Jamun)

S. N0.	Leaf extract (µL)	<i>Syzygium cumini</i> (nm)*			
		Ethanol (control)	Methanol (control)	Ethyl acetate (control)	Chloroform (control)
1	Control	0.00	0.00	0.00	0.00
2	1	0.019	0.019	0.009	0.009
3	2	0.006	0.026	0.010	0.004
4	3	0.005	0.035	0.015	0.006
5	4	0.008	0.048	0.017	0.009
6	5	0.009	0.059	0.001	0.007
7	6	0.003	0.053	0.003	0.006
8	7	0.006	0.056	0.071	0.002
9	8	0.007	0.067	0.067	0.005
10	9	0.001	0.061	0.061	0.002
11	10	0.001	0.081	0.068	0.001

*Absorbance *Syzygium cumini* leave extracts

Antimicrobial activity

Antimicrobial activity of *Syzygium cumini* leaves extract against gram positive and negative bacteria and *Candida albicans* fungus are summarized in Table 3 and in the Table 4 by disc diffusion and poison plate methods, respectively.

The results in Table 3 in disc diffusion and Table 4 poison plate methods indicated that all plants extracts showed that antimicrobial activities toward the gram positive bacteria *Staphylococcus aureus* as well as gram negative bacteria *Escherichia coli* and *Candida albicans*.

It is observed from Table 3 that inhibitory zone follows the order:

Chloroform > ethanol > methanol > ethyl acetate

Chloroform extract showed more effective inhibitory zone against bacteria and fungi in comparison to other solvents extracts. In ethanol extract maximum (22.98 nm; 900 µL) and minimum (11.12 nm; 300 µL) inhibitory zone was observed for *Candida albicans* and *Staphylococcus aureus*, respectively. In methanol extract maximum (21.25 nm; 900 µL) and minimum (10.16 nm; 300 µL) inhibitory zone observed for *Candida albicans* and *Staphylococcus aureus*, respectively. In ethyl acetate extract maximum (18.99 nm, 900 µL) and minimum (13.78 nm; 300 µL) inhibitory zone was observed for *Escherichia coli* and *Candida albicans*, respectively. In chloroform extract maximum (24.03 nm, 900 µL) and minimum (17.00 nm, .00 µL) inhibitory zone observed for *Escherichia coli* and *Candida albicans*, respectively.

It is clear from Table 4 that inhibitory zone follows the order:

Ethanol > ethyl acetate > chloroform > methanol

All extracts showed nearly same inhibitory zone or antibacterial activity. In ethanol extract maximum (24.93 nm; 900 µL) and minimum (17.45 nm; 300 µL) inhibitory zone was observed for *Escherichia coli* and *Candida albicans*, respectively. In methanol extract maximum (24.43 nm; 900 µL) and minimum (17.16 nm; 300µL) inhibitory zone was observed for *Escherichia coli* and *Candida albicans*, respectively.



Table 3. Antimicrobial activity of four crude leaves extract of *Syzygium cumini* (Jamun) compared by disc diffusion method

Extract solvent (μL)	Diameter of inhibitory zone (nm)*		
	<i>E. coli</i>	<i>S. aureus</i>	<i>C. albicans</i>
Ethanol	0.00	0.00	0.00
300	16.96	11.12	18.67
600	17.98	13.24	20.87
900	19.90	15.45	22.98
Methanol	0.00	0.00	0.00
300	16.69	10.16	19.00
600	18.60	12.17	20.02
900	19.10	15.89	21.25
Ethyl acetate	0.00	0.00	0.00
300	16.90	14.50	13.78
600	18.90	15.60	14.34
900	18.99	16.00	14.78
Chloroform	0.00	0.00	0.00
300	21.00	19.00	17.00
600	22.34	20.56	17.98
900	24.03	23.57	18.20

* Replicate

In ethyl acetate extract maximum (24.03 nm; 900 μL) and minimum (17.00 nm; 300 μL) inhibitory zone was observed for *Escherichia coli* and *Candida albicans*, respectively. In chloroform extract maximum (24.33 nm; 900 μL) and minimum (17.30 nm; 300 μL) inhibitory zone was observed for *Escherichia coli* and *Candida albicans*, respectively.

Phytochemical analysis

Phytochemical analysis Table 5 of the *Syzygium cumini* (Jamun) leaves extracts revealed the presence of carbohydrates, flavonoids, terpenoids, saponins, tannins, proteins and amino acids etc.

It is observed from Table 5 that Phyto-constituent (carbohydrate) is present in the extract of each solvent. Alkaloids are not detected in all four (ethanol, methanol, ethyl acetate and chloroform) solvent extracts. Glycoside is not detected in the ethanol solvent extract. Glycosides tests for methanol, ethyl acetate and chloroform extracts could not be performed. Saponins are only tested in methanol solvent. Tannins and flavonoids are tested in ethanol extracts



only. Ethanol solvent is found to be more effective solvent for the phytochemical screening of *Syzygium cumini* leaves extract.

Table 4. Antimicrobial activity of leaves extract of *Syzygium cumini* (Jamun) compared with control by poison plate method

Plant	Extract solvent (μL)	Diameter of inhibitory zone (nm)*		
		E. coli	S. aureus	C. albicans
<i>Syzygium cumini</i> (Jamun)	Ethanol	0.00	0.00	0.00
	300	21.50	19.60	17.45
	600	22.64	20.76	17.88
	900	24.93	23.07	18.10
	Methanol	0.00	0.00	0.00
	300	21.12	19.14	17.16
	600	22.24	20.26	17.28
	900	24.43	23.47	18.40
	Ethyl acetate	0.00	0.00	0.00
	300	21.00	19.00	17.00
	600	22.34	23.57	17.98
	900	24.03	23.57	18.20
	Chloroform	0.00	0.00	0.00
	300	20.10	19.20	17.30
	600	22.44	20.66	17.99
	900	24.33	23.47	18.60

* Replicate

Table 5. Phytochemical analysis of *Syzygium cumini* (Jamun) leaves extract

S.No	Phyto constituents	Ethanol	Methanol	Ethyl acetate	Chloroform
1	Alkaloids	–	–	–	–
2	Carbohydrate	+	+	+	+
3	Saponins	–	–	–	–
4	Protein and amino acids	–	–	–	–
5	Tannins	+	–	–	–
6	Flavonoids	+	–	–	–
7	Glycosides	–	–	–	–
8	Terpenoids	+	–	–	–

– = Absence + = Presence

CONCLUDING REMARKS

- Syzygium cumini* leave extracts were found to have antimicrobial, antifungal and antioxidant properties.
- The antioxidant power of *Syzygium cumini*'s leaves in extracts in various solvents follows the order: methanol > ethyl acetate > ethanol > chloroform
- The antioxidant power of *Syzygium cumini*'s leaves in extracts in various solvents was found to increase with increase of its concentrations.



- (iv) The chloroform extract was found to have more antibacterial property in comparison to methanol, ethylacetate, and ethanol extracts.
- (v) In disc diffusion method inhibitory zone for antibacterial activity follows the order:
Chloroform > ethanol > methanol > ethyl acetate
- (vi) In poison plate method inhibitory zone for antibacterial activity follows the order:
Ethanol > ethyl acetate > chloroform > methanol
- (vii) In poison plate method all extracts have showed nearly same inhibitory zone or antibacterial activity.

EXPERIMENTAL

Collecting of plant material

The plant material (leaves) of *Syzygium cumini* was collected at the Institute of Applied Science and Technology (IAST), University of Guyana, Turkeyen Campus, Georgetown. Guyana.

Preparation of plant materials:

The collected leaves sample of *Syzygium cumini* was weighted on Citizen CTG 3000E electronic balance. The leaves dried in oven (Gallenkamp Incubator Model IH-150) at 50-60°C. The dried leaves were cooled at room temperature and weighted again on Citizen electronic balance. Weight of green leaves, dried leaves and value of percentage moisture content in various samples of *Syzygium cumini* is given in Table 1. The weight of ground leaves of *Syzygium cumini* was found to be 560 grams.

Collection of test organism

Three micro-organisms (*Escherichia coli*, *Staphylococcus aureus* and *Candidus albicans*) were used for the study. All the tested strains are reference strains and were collected from the microbiology laboratory of Georgetown Public Hospital Corporation, Georgetown (GHPC). All cultures are maintained in nutrient broth (Himedia, M002) at 37°C and maintained on nutrient agar (Himedia MM012) slants at 4°C.

Extraction and preparation of test solutions

The grounded leaves of *Syzygium cumini* were extracted in each solvent, ethanol, ethyl acetate, chloroform and methanol. 20 g of dried pulverized leaves were soaked with 200 mL of each solvent for 48 h separately. Solvent is decanted each time and residue again soaked with same solvent for 24 h. The total extract is combined and filtered. The evaporation of solvent was done on rotavapour (Buchi). The respective solvent was added to viscous semi solid liquid extract to make up the desired volume of extract solution.

Reducing antioxidant power

Ammonium thiocyanite, ferric chloride, linoleic acid (99.5 %), thiobarbituric acid, trichloroacetic acid, potassium ferrocyanide, butylated hydroxyl anisol, sodium dihydrophosphate, sodium monhydrophosphate, potassium dihydrophosphate obtained from Aldrich. The reducing antioxidant power of the plants methanolic, ethanolic, ethyl acetate and chloroform extract was determined by the method reported in the literature [23,24]. Different concentration of plant extracts (100 - 1000 µL) in 1 mL of distilled water were mixed with phosphate buffer (2.5 mL, 0.2 M ,pH 6.6) and potassium ferricynaide $K_3[Fe(CN)_6]$, (2.5 mL, 1 %) . The mixture was incubated at 50° C for 20 min. Then 2.5 mL of trichloroacetic acid 10 % was added to mixture, which was then centrifuged for 10 min at 3000 rpm. The upper layer (2.5 mL) was mixed with distilled water 92.5 mL and $FeCl_3$, 0.5 mL, 1 %. The absorbance was measured at 700 nm against a blank using UV-Vis spectrophotometer (Phillips X500). Increased absorbance of the reaction mixture indicates increase in the reducing power.

Anti-Microbial Assay



Materials

Mueller Hinton Agar, Agar plates and microbial discs were purchased from the International Pharmaceutical Association (IPA) in Guyana. Solvents like chloroform, ethanol, ethyl acetate and methanol obtained from Aldrich. Scitilation vials 20 mL were obtained from Meditron Scientific Sales, Georgetown, Guyana.

Aseptic chamber

Aseptic chamber consists of a wooden box of L = 1 meter, B = 1 meter and D = 0.5 meter area. Chamber is cleaned with 70 % ethanol twice and irradiated with short wave UV light for 1 h.

Potato dextrose agar (PDA) medium

Potato Dextrose Agar (PDA) medium prepared according to method reported by Talaro [25]. This is the medium on which cultured bacteria *Escherichia coli* and *Staphylococcus aureus* were grown. The 200 g potato was peeled finely chopped and boiled to a mash in distilled water. Each 12.5 g dextrose and 12.5 g agar was placed in a 1 L measuring cylinder. The potato mash was stirred and poured in to the same measuring cylinder. Distilled water was added to make the solution up to 500 mL. The content was stirred until the consistency of solution mixture. The stirred mixture poured into conical flasks, plugged with cotton wool and tightly wrapped by aluminum foil. The flasks were autoclaved at 121° C, 15 psi, for 15 minutes.

Mother plates

Mother plates were prepared by pouring PDA mixture into Petri dishes and to cool at room temperature, in the aseptic chamber.

Antimicrobial assay was done by disc diffusion and poisons plate method [26] using plants *Syzygium cumini* extracts (ethanol, methanol, ethyl acetate and chloroform) and commonly used antibiotics. The test quantities of specific extracts were dissolved in depending upon the solubility of the extracts. The dissolution of the organic extracts (chloroform, methanol, ethanol and ethyl acetate) was aided by water, which did not affect the growth of microorganism, in accordance with our control experiments. The surfaces of media were inoculated with bacterial from a broth culture. After 18 h of incubation at a specific temperature 28° C - 30° C for *Escherichia coli*, *Staphylococcus aureus* and *Candidus albicans* the plates were examined and the diameters of the inhibition zones were measured to the nearest millimeter.

Phytochemical analysis of the plant extracts

Materials

Glacial acetic acid, thionil chloride, dichloromethane, copper sulfate, lead acetate, diethyl ether, ferric chloride, acetic anhydride, antimony chloride, all obtained from Aldrich.

Method

Phytochemical analysis of all the aqueous plant extracts was carried out by suitable methodologies in search of active ingredient responsible for antimicrobial toxicity. The phytochemicals include under study were saponins, terpenoids, alkaloid, glycoside, carbohydrates, protein and amino acids, tannins, and flavonoids the analysis was carried out according to the methodologies of Edeoga et al. [27].

REFERENCES

1. Kaur, A., Singh, D. Kaur, H., Sogi, J. 2018, *Stored Products Postharvest Res.*, 9 (4), 36.



2. Navneet, S.A., **2018**, *IJIPSR*, 6 (1), 32.
3. Mitra, P., Mitra P.K., Ghosh, T. **2018**, *Acta Sci. Agri.*, 2 (11), 59.
4. Mitra, P., Mitra P.K., Ghosh, T. **2018**, *Acta Sci. Param Sci.*, 2 (9), 69.
5. Arunpaudiyani, J., Roselin, R.B., Chitra, S.R., Kaniga, P. **2018**, *WJPPS*, 7 (4), 499.
6. Sukmasari, S., Mohd, F.N., Doolaanea, A.A., Jabbar, O.A., Qader, A., Rahman, M.N.A., **2018**, *J. Pharm. Sci. & Res.* 10 (1), 31.
7. Proma, N.M., Naima, J., Islam, J.A., Papel, M.R., Rahman, M., Hossain, M.K. **2018**, *IJIPSR*, 9 (5), 1806.
8. Bandiola, T.M.B., Carpuz, M.J.T. **2018**, *Pharm. Anal. Acta.*, 9 (5), 1.
9. Kumawat, M., Damar, J., Kachchhwaha, J., Garg A. K., Singh C., **2018**, *WJPPS*, 7 (03), 312.
10. Ramos, I.L., Bandiola, T.M.B. **2017**, *Der Pharm. Lett.*, 9 (2), 74.
11. Haque, R., Sumiya, M.K., Sakib, N., Sarkar, O.S., Siddique, T.T.T., Hossain, S., Islam, A., Parvez, A.K., Talukder, A.A., Dey, S. K. **2017**, *Adv. Microbiol.* 7, 193.
12. Khan, A., Jabeen, K., Iqbal, S., **2016**, *Pure Appl. Biol.*, 5 (2), 193.
13. Hasanuzzaman, M., Islam, U., Islam, M.B. **2016**, *J. Biol. Sci.*, 24, 11.
14. Elfadil, A.G., Karamallah, A.A., Abualhassan, A.M., Hamid, A.A., Sabahelkhier, M.K. **2015**, *Nova J. Med. Biol. Sci.*, 4 (2), 1.
15. Prabhakaran, S., Gothandam, K.M., Sivashanmugam, K. **2011**, *Res. Pharm.*, 1 (11), 22.
16. Mohit, Y., Kadtan, K., Wakte, P. **2011**, *J. Pharm. Res.*, 4 (8), 278.
17. Jabeen K., Javaid, A. **2010**, *Nat. Prod. Res.*, 24 (12), 1158.
18. Ruan, Z.P., Zhang, L.L., Lin, Y.M. **2008**, *Molecules*, 13, 2545.
19. Gulcin, L., Oktay, M., Kirecci, E., Kufrevioglu, O.I. **2003**, *Food Chem.*, 83 (3), 371.
20. Shimada, K., Fujikawa, K., Yahara, K., Nakamura, T. **1992**, *J. Agri. Food Chem.*, 40, 945.
21. Duh, P. **1998**, *J. Am. Oil Chem. Soc.* 75, 455.
22. Sawaddiwong, R., Jongjareorak, A., Bazakul, S. **2008**, *KMITL Sci. J.*, 8 (2), 54.
23. Oyaizu, M. **1986**, *Jpn. J. Nutr.*, 44, 307.
24. Huda-faujan, H. Noriham, A., Narrakaih, A.S., Badji, A. **2009**, *S Afr. J. Biotechnol.*, 8 (3), 484.
25. Talaro, K.P., Wm. C. Brown Publishers, USA, **1993**.
26. Awoyinka, A.O., Balagun, A.O., Ogunnowo, A. A. **2007**, *J. Med. Plants Res.*, 1, 063.
27. Edeoga, H.O., Okwu, D.E., Mbalble, B.O. **2005**, *Afr. J. Biotechnol.*, 4 (7), 685.