



Revista Boliviana de Química

ISSN: 0250-5460

ISSN: 2078-3949

revbolquim@outlook.com

Universidad Mayor de San Andrés

Bolivia

Paredes de Gómez, Betty; Huanca Z., Paul K.; Salas-Arias, Erika; Pérez-Montaño, H. Saúl; del Carpio, Graciano
ASSESSMENT OF THE PRESENCE OF HEAVY METALS, AS FOR AN INVENTORY
ON GARLIC, ONION AND IN SEMIURBAN SOILS OF AREQUIPA, PERU
Revista Boliviana de Química, vol. 37, no. 4, 2020, September-October, pp. 202-208
Universidad Mayor de San Andrés
Bolivia

Available in: <https://www.redalyc.org/articulo.oa?id=426365221003>

- How to cite
- Complete issue
- More information about this article
- Journal's webpage in redalyc.org

UAEH  redalyc.org

Scientific Information System Redalyc
Network of Scientific Journals from Latin America and the Caribbean, Spain and
Portugal

Project academic non-profit, developed under the open access initiative



ASSESSMENT OF THE PRESENCE OF HEAVY METALS, AS FOR AN INVENTORY ON GARLIC, ONION AND IN SEMIURBAN SOILS OF AREQUIPA, PERU

EVALUACIÓN DE LA PRESENCIA DE METALES PESADOS, COMO PARA UN INVENTARIO EN AJO, CEBOLLA Y SUELOS SEMIURBANOS DE AREQUIPA, PERÚ

Received 06 02 2020
Accepted 08 04 2020
Published 10 30 2020

Vol. 37, No.4, pp. 202-208, Sep./Oct.2020
Revista Boliviana de Química

37(4), 202-208, Sep./Oct. 2020
Bolivian Journal of Chemistry
DOI: 10.34098/2078-3949.37.4.3



Full original article

Peer-reviewed

Betty Paredes de Gómez¹, Paul K. Huanca Z.^{2,*}, Erika Salas-Arias³, H. Saúl Pérez-Montaño³, Graciano del Carpio⁴

¹Escuela Profesional de Química, Facultad de Ciencias Naturales y Formales FCNF, Universidad Nacional San Agustín de Arequipa UNSA, Av. Venezuela s/n, phone +5154404197, Arequipa, Perú, quimica@unsa.edu.pe, <http://fip.unsa.edu.pe/quimica/>

²Escuela Profesional de Ingeniería de Materiales, Facultad de Ingeniería de Procesos FIP, Universidad Nacional San Agustín de Arequipa UNSA, Av. Independencia s/n-Pab. Ing. Materiales, phone +5154200037, Arequipa, Perú, materiales@unsa.edu.pe, <http://fip.unsa.edu.pe/ingmateriales/>

³Universidad Católica San Pablo, Urb. Campiña Paisalista s/n Quinta Vivanco – Barrio de San Lázaro, phone +5154605630 anexo 266, Arequipa, Perú, <http://ucsp.edu.pe>

⁴Escuela Profesional de Biología, Facultad de Ciencias Biológicas FCB, Universidad Nacional San Agustín de Arequipa UNSA, Av. Daniel Alcides Carreón s/n-Pab. Biología, phone +51959627875, Arequipa, Perú, biologia@unsa.edu.pe, <http://fip.unsa.edu.pe/biologia/>

Keywords: Heavy metals, Garlic, Onion, Soils accumulation, ICP-MS.

Palabras clave: Metales pesados, Ajo, Cebolla, Acumulación de suelos, ICP-MS.

ABSTRACT

The objective of this study is the evaluation of the accumulation of heavy elements, namely Pb, Cr, Cd, Tl, Cu, Co in garlic, onion and soil of semi-rural soils and its potential effect on the health of the population of the city of Arequipa. The results showed concentrations below the permissible limits for all of them in garlic, onion and soils in accordance with international standards, showing that the accumulation of heavy metals in the soil does not exceed the limits according to international standards.



*Correspondent author: phuanca@unsa.edu.pe

RESUMEN

El objetivo de este estudio es la evaluación de la acumulación de elementos pesados, Pb, Cr, Cd, Tl, Cu, Co en ajo, cebolla y suelo de suelos semirurales y su efecto potencial en la salud de la población de la ciudad de Arequipa. Los resultados mostraron concentraciones por debajo de los límites permisibles para todos ellos en ajo, cebolla y suelos de acuerdo con los estándares internacionales demostrando que la acumulación de metales pesados en el suelo no supera los límites según los estándares internacionales.

INTRODUCTION

Contamination of agriculture soils by the long-term accumulation of different transitions element is considerate of the great importance around the world [1]. The ecotoxicity and persistence of heavy metals and metalloids through natural and anthropogenic contributions on agriculture soils is very important and has been attracted the attention of the global science community [2]. Heavy metals are defined as an inorganic chemical hazard, the most common are: lead (Pb), chromium (Cr), arsenic (As), cadmium (Cd), copper (Cu), mercury (Hg) and nickel (Ni) [3,4]. Natural inventory of the heavy metals is inherited from parent rocks. Anthropogenic activities are sources that heavily influenced the nature inventory. These include coal combustion, traffic emissions, industrial activities, mining operations, and agriculture activities [1,5]. It is true that the anthropogenic contribution exceeds sometimes the natural contribution due to the crescent urban and industrial rising in the last decades.

Understanding the spatial distribution and accumulation of heavy elements is critical. It will help to evaluate the complex and different mechanism of transition of elements from soils to vegetals, their mobility and bioavailability, leaching losses, uptake for plants and their influence on human health by ingesting grazing livestock meat [2,3]. Quantify the number of heavy metals is essential. Their input and output on agriculture soils would provide information of the source apportionment and create an inventory about the source of heavy elements to agricultural soils, atmospheric depositions, fertilizer application, organic and inorganic waste disposal and the contribution of the industry and human activity, helping to diagnose the current situation of heavy metals on agriculture soils [6]. Worldwide concern about the crescent inventory of heavy metals have shown the importance to keep an update record it started in 1974 in Netherlands [7]. Since then, European nations have been performed periodical research, for example, France and Germany [8], in United States of America and lately China [9,10].

An inventory of the presence of the heavy elements is significant because it indicates the accumulation or depletion of the different elements in specific geographical areas, ecosystems, and provide a valuable knowledge base to improve the management and recommendation about norms and policies to regulate the anthropogenic sources [10]. Leclerc and Laurel published new global scale inventory for the period 2000-2014, indicating the accumulation of heavy metals on agricultural soils in 2017 [11]. However, the total release of the harmful heavy chemicals and their quantification result from the specific studies and for specific cases, and the data cannot be extrapolated for specific geographic locations because of several parameters that will affect the results of the analysis.

In this study, our team will aim to (1) Develop a framework to quantify the presence of heavy metals on livestock and agriculture soil in the urban region of Arequipa- Peru. (2) Contribute to creating a national inventory regionally and for the world. (3) Quantify the impact of heavy metals on livestock and their potential harm to human health.

EXPERIMENTAL

Data collection and materials

Our study evaluated the accumulation of heavy metals in livestock and soils in the urban region of Arequipa. The study was performed in the summer of 2016, the soil (onions and garlic) and livestock were collected from three places Tiabaya 71.631669 West; 16.426431 South Sachaca 71.562181 West; 16.424828 South and Uchumayo 71.631669 West; 16.426431 South. Soil samples were collected randomly and chemically analyzed by triplicate.



Livestock samples were collected, carefully cleaned, cut, and dried using a lyophilization; the chemical analyzed was performed by triplicated as well.

Heavy metals concentration on the soil and livestock were measured digesting the samples in nitric acid and hydrogen peroxide at 150°C using digester into a microwave oven. The samples with this treatment were diluted and measured with ICP-MS. For quality control, an internal control (Spinach NBS 1570 and Sea Lettuce BCR 279) were run in parallel to the livestock and soil samples.

The following seven heavy metals considerate as a priority pollutant element by the USEPA (United State Environmental Protection Agency) have been identified in this study as: chromium, copper, nickel, cobalt, lead, cadmium, and thallium.

RESULTS

Soils are an essential reservoir of chemical elements and a living matrix for compounds in their different states of the matter. Chemical analysis of the agriculture areas, plants, and soils show essential information about the inventory data for heavy metal in the Arequipa region.

Heavy metal concentration was determined using the weight of fresh vegetables and after drying. We used the following equation to calculate the concentration.

$$[C_{p.f.}] = \frac{(100 - X - Y)}{100} [C_{p.d.}] \quad \text{equation 1}$$

Where:

X= Percentage loss of water after dry

Y= Percentage of residual water

$C_{p.d.}$ = Concentration mg/kg of the sample dry weight.

$C_{p.f.}$ = Concentration mg/kg of the sample fresh weight

Table 1: Summary of concentration of heavy metals concentrate on mg/kg for garlic, onion, and soils from the semi-urban area of Arequipa

		Cr mg/kg	SD	Pb mg/kg	SD	Cd mg/kg	SD	Tl mg/kg	SD	Ni mg/kg	SD	Cu mg/kg	SD	Co mg/kg	SD
Sachaca	Onion	0.095	10.14	0.037	1.87	0.01	5.85	0.0001	0.06	0.034	1.58	0.302	15.56	0.008	0.06
Tiabaya	Onion	0.176	2.22	0.156	1.37	0.017	1.01	0.0008	0.02	0.013	0.52	0.432	19.82	0	0.00
Tiabaya	Garlic	0.484	12.93	0.189	7.86	0.014	1.07	0.00047	0.07	0.0028	0.50	1.259	35.47	0.0028	0.33
Uchumayo	Garlic	0.58	29.00	0.15	11.69	0.1	3.29	0.002	0.55	0.007	0.82	1.03	53.40	0.003	0.26
	Soil	39.41	6.85	10.99	1.26	<1	1.05	<1	0.00	11.38	1.52	34.5945	3.32	9.615	0.82
NBS 1570	Standard														
Measure				0,176±0.023		2,93±0.12				2.02±0.11		12.0±0.4		0.41±0.03	
Certified				[0,20]		2,84±0.07				2.14±0.10		12.2±0.6		0.39±0.05	
Sea lettuce BCR 279	Standard														
Measure				2.06±0.08		1.66±0.07				1.68±0.11				0.59±0.09	
Certified				1.99±0.06		1.52±0.04				1,59±0.02				0.57±0.02	

Statistical Analysis

The statistical analysis was performed evaluating the distribution t- Student, calculating the confidence intervals of the concentration average of the heavy metals with 98% of confidence. Results are shown in table 1 and 2 for the distribution of heavy metals on onion and garlic respectively on Tiabaya, Sachaca and Uchumayo semirural places of Arequipa Peru.



Table 2. Statistical results of concentration of heavy metals on onion.

Pb in Onion (Sachaca)			Cr in Onion (Sachaca)		Cd in Onion (Sachaca)		Ni in Onion (Sachaca)	
Low limit		High limit	Low limit	High limit	Low limit	High limit	Low limit	High limit
IC(u;0.98)	0.02949358	0.04455268	0.05405020	0.13559697	0.01343154	0.03358234	0.02751816	0.04023940
t(0.99;2)	6.96456		6.96456		6.96456		6.96456	
Av dev	0.00108112		0.005854409		0.00337522		9.1328E-04	
Est error	0.00752955		0.040773385		0.02350694		0.00636062	

Tl in Onion (Sachaca)			Cu in Onion (Sachaca)		Co in Onion (Sachaca)	
Low limit		High limit	Low Limit	High limit	Low Limit	High limit
IC(u;0.98)	-0.00009317	0.00038338	0.23973107	0.36486933	0.00053110	0.00100813
t(0.99;2)	6.96456		6.96456		6.96456	
Av dev	3.4213E-05		8.9839E-03		3.4247E-05	
Est error	0.00023828		0.06256913		0.00023851	

Pb in Onion (Tiabaya)			Cr in Onion (Tiabaya)		Cd in Onion (Tiabaya)		Ni in Onion (Tiabaya)	
Low limit		High limit	Low limit	High limit	Low limit	High limit	Low limit	High limit
IC(u;0.98)	0.15075141	0.16176005	0.16700422	0.18488458	0.01258763	0.02069264	0.01078956	0.01496291
t(0.99;2)	6.96456		6.96456		6.96456		6.96456	
Av dev	0.00079033		0.001283668		0.00058187		2.9961E-04	
Est error	0.00550432		0.00894018		0.0040525		0.002086673	

Tl in Onion (Tiabaya)			Cu in Onion (Tiabaya)		Co in Onion (Tiabaya)	
Low limit		High limit	Low limit	High limit	Low limit	High limit
IC(u;0.98)	0.00000867	0.00014418	0.35233739	0.51171031	0.00000000	0.00000000
t(0.99;2)	6.96456		6.96456		6.96456	
Av dev	9.7285E-06		1.1442E-02		0.0000E+00	
Est error	6.77548E-05		0.079686456		0	

DISCUSSION

Trace elements in livestock and soil

Chromium

The mean content of chromium in onion for Sachaca and Tiabaya was 0.095 mg/kg and 0.176 mg/kg respectively, in the case of garlic for Tiabaya was 0.484 mg/kg and Uchumayo 0.58 mg/kg. The content of chromium in soil was 39.41 mg/kg on average; this result is between the average value for the background concentration of chromium on Europe's soils report that signals limits of 5 to 65 mg/kg [12,13]. The presence of Cr(VI) reported in Plants (*Alyssum obovatum*, *Cochlearia arctica* (Brassicaceae), *Armeria scabra* (Limoniaceae), *Salix arctica* (Salicaceae) showed values from 0.09–0.59 mg/kg [14]. Our research did not target the presence of different ionic species of chromium, the samples assayed indicated that onion from Sachaca had values between the limits reported by Panichev et al [14]. However, samples from Tiabaya onion and garlic and Uchumayo garlic showed the highest value compared with onion from Sachaca. The highest value found in the garlic sample in Tiabaya had close values found in the same species reported by Sakizadeh et al. 2018 [15]. The daily amount of chromium intake per day in our diet according to the European Food Safety Authority (EFSA) is 57.3–83.8 µg/kg and for the World Health Organization (WHO) is 47 µg/kg [1,16].

Lead

The concentration of lead in onion samples shown values of 0.037 mg/kg and 0.156 mg/kg for Sachaca and Tiabaya, whilst garlic showed 0.189 mg/kg for Tiabaya and 0.15 for Uchumayo. The determined content of lead on the soil



was 10.99 mg/kg. The limit permissible for lead in soils according to the World Health Organization (WHO) is 50 mg/kg. The amount on the lead registered on garlic was 0.21 mg/kg reported for Sakizadeh et al. 2018 [15]. Our data show that the concentration of lead is low, besides onion sample from Sachaca that show the lowest content on lead, the other samples show very close values. Cai Li et al [17], report that vegetable on China depending on its kind allows between 0.3 to 0.1 mg/kg. According to the FAO/WHO the maximum intake for lead is 25 ug/kg body weight per week [18].

Table 3. Statistical results of concentration of heavy metals on garlic.

Pb in Garlic (Tiabaya)			Cr in Garlic (Tiabaya)		Cd in Garlic (Tiabaya)		Ni in Garlic (Sachaca)	
Low limit		High limit	Low limit	High limit	Low limit	High limit	Low limit	High limit
IC(u;0.98)	0.15776423	0.22098166	0.43209324	0.53611145	0.00958078	0.01816291	0.00077297	0.00478184
t(0.99;2)	6.96456		6.96456		6.96456		6.96456	
Av dev	0.00453851		0.00746768		0.00061613		2.8780E-04	
Est error	0.03160871		0.052009108		0.00429107		0.002004432	

Tl in Garlic (Tiabaya)		Cu in Garlic (Tiabaya)		Co in Garlic (Tiabaya)	
Low limit	High limit	Low limit	High limit	Low limit	High limit
IC(u;0.98)	0.00021031	0.00074039	1.11619461	1.40141010	0.00145688
t(0.99;2)	6.96456		6.96456		6.96456
Av dev	3.8056E-05		2.0476E-02		1.8856E-04
Est error	0.00026504		0.142607744		0.0013132

Pb in Garlic (Uchumayo)		Cr in Garlic (Uchumayo)		Cd in Garlic (Uchumayo)		Ni in Garlic (Uchumayo)	
Low limit	High limit	Low limit	High limit	Low limit	High limit	Low limit	High limit
IC(u;0.98)	0.10303576	0.19706734	0.46402363	0.69724771	0.08714731	0.11358056	0.00331994
t(0.99;2)	6.96456		6.96456		6.96456		6.96456
Av dev	0.00675072		0.016743634		0.0018977		4.7296E-04
Est error	0.047015789		0.116612041		0.01321663		0.003293933

Tl in Garlic (Uchumayo)		Cu in Garlic (Uchumayo)		Co in Garlic (Uchumayo)	
Low limit	High limit	Low limit	High limit	Low limit	High limit
IC(u;0.98)	0.00030555	0.00468898	0.81268662	1.24210686	0.00159131
t(0.99;2)	6.96456		6.96456		6.96456
Av dev	3.1470E-04		3.0829E-02		1.4955E-04
Est error	0.00219171		0.214710122		0.00104157

Cadmium

Similarly, the concentration of cadmium ranged from 0.014 to 0.1 mg/kg in garlic and onion, in comparison, Sakizadet et al., 2018 [15] reported cadmium in fresh vegetables with an average value of 0.01mg/kg and Xinyu Li et al 2018 [19] reported 0.21 mg/kg on the edible part of vegetables cultivated on contaminated areas but 0.05 mg/kg on vegetable cultivated on controlled soils. The values of Sachaca, Tiabaya for onion showed very close values, 0.01 and 0.014 mg/kg respectively. The value of the amount of cadmium of garlic from Tiabaya was similar to the content in onions. However, the average value on Uchumayo had shown one magnitude higher on Garlic 0.1 mg/kg. The mean concentration of cadmium on soil was determined as less than 1 mg/kg and it's below to the value recommended for the WHO/ FAO that is 3mg/kg. The daily amount of cadmium consumed recommended by the USEPA 2009 is 1ug/kg/day [18].

Thallium

The average content of thallium in onions from Tiabaya and Sachaca is 0.0008 and 0.0001 mg/k, respectively. The content in Garlic showed differences between samples from Tiabaya 0.00047 mg /kg and Uchumayo with 0.002 mg/kg. Huang Yun Yu et al. 2018 [20], reported the average content of thallium as 0.051 mg/kg and for cabbage from 0.01 to 0.06 mg/kg. These values corresponded to samples coming from farmlands close to mining and



industrial activities. Our data indicated that there is no contamination in onion and garlic with thallium, in accordance to the German and Chinese guide food for thallium that indicated the permissible limit as 0.05mg/kg. The content of thallium on soils according to the Canadian and Swiss regulations [21] is 1mg/kg; our data showed that the concentration of thallium is less than 1 mg/kg. The daily amount recommended for consumption is 0.00001 mg/kg/per day [18].

Nickel

The content of nickel in onions in Tiabaya and Sachaca showed values of 0.034 and 0.011 mg/kg respectively. Values found on garlic were 0.0028 and 0.007 mg/kg for Tiabaya and Uchumayo, respectively. The concentration of nickel in plant cells was reported to be in the rank 0.5 to 5 mg/kg [22], the concentration in onions and garlic showed that the concentration is below the limits for nickel with values from 0.007 and 0.034 mg/kg. The permissible limit for Nickel on soils according to WHO is 50mg/kg. The average content of nickel in the soil in the present study was 11.38 mg/kg. The recommended daily amount of consumption of nickel is 0.02mg/kg/per day [18].

Copper

The average concentration of copper in onions from Tiabaya en Sachaca were 0.302 and 0.432 mg/kg, the concentration of copper in garlic was higher, samples from Tiabaya en Uchumayo showed 1.259 and 1.03 mg/kg. Copper concentration on livestock has been reported with ranges between 1.75 to 9.26 mg/kg [23,24], the maximum permissible amount of copper in vegetable is 50 mg/kg [25]. The content of copper in onion and garlic are below the permitted limits. In the case of the soil, the maximum amount of copper is 100mg/kg according to the WHO [18]. Our sample had a value of 34.5945 mg/kg, indicating that there is no contamination with copper. The amount of copper for human ingestion is 0.04 mg/kg/per day [18].

Cobalt

The concentration of cobalt in onions ranged from 0.008 mg/k to no detected for samples of Sachaca and Tiabaya. In the case of garlic, the values varied from 0.0028 and 0.003 mg/kg for Tiabaya and Uchumayo. Amin et al. 2013 [26], reported that the concentration of cobalt in onion and garlic in contaminated areas were of 0.03 to 0.06 mg/kg and 0.01 to 1mg/kg. Khan et al. 2008 [27] and Chekri et al. 2019 [28] reported the presence of cobalt in toddler food with values between 0.7 to 8.6 ug/kg similar to the values of Melo et al., 2008 [29]. The concentration of the analyzed soil sample fall between the results reported by the literature. The content of cobalt in the examined soil was 9.3 mg/kg which is below the maximum amount according to USEPA or 80mg/kg. The daily intake of cobalt is 0.0003 mg/kg per day [18].

CONCLUSIONS

The presence of seven heavy inorganic elements Pb, Cr, Cd, Tl, Ni, Cu, and Co was assessed on garlic, onion and soils from semiurban areas of Arequipa city. The concentration of these elements were all between the limits for vegetables and soils according to the international standards. The concentration of the heavy elements in the vegetables does not show potential harm to the health of the population, and the same condition can be expressed for the evaluated soils samples. The results indicated that the presence of heavy metals in vegetables and soils are low. However, the accumulation, physical and chemical behavior of the soils, and the identification of the sources still remains a pending task.

ACKNOWLEDGMENTS

The authors want to thank Universidad Nacional de San Agustín de Arequipa for their financial support of the project contract N° 05-2017-UNSA.

REFERENCES



1. Lv, J., Liu, Y. **2019**, An integrated approach to identify quantitative sources and hazardous areas of heavy metals in soils, *Science of the Total Environment*, **646**, 19-28.
2. Nicholson, F.A., Smith, S.R., Alloway, B., Carlton-Smith, C., Chambers, B. **2003**, An inventory of heavy metals inputs to agricultural soils in England and Wales, *Science of the Total Environment*, **311**, 205-219.
3. Wuna, R.A., Okieimen, F.E., **2011**, Heavy metals in contaminated soils: a review of sources, chemistry, risks and best available strategies for remediation. *International Scholarly Research Notices*, Article ID 402647. DOI: <https://doi.org/10.5402/2011/402647>.
4. Evanko, C.R., Dzombak, D.A. Remediation of metals-contaminated soils and groundwater, *Technology evaluation report, TE-97-01, GWRTAC E-Series, Ground-water remediation technologies analysis center*, <http://www.gwrtac.org>, **1997**, Pittsburg, U.S.A. pp. 61. <https://clu-in.org/download/toolkit/metals.pdf>. Access date: May, 2020.
5. Alloway, B.J. *Heavy Metals and Metalloids as Micronutrients for Plants and Animals*, In: *Heavy metals in soils: trace metals and metalloids in soils and their bioavailability*, *Environmental pollution Series*, Vol, 22, ed by Alloway, B.J. **2012**, Springer, Netherlands, 195-209. DOI: 10.1007/978-94-007-4470-7
6. Shi, T., Ma, J., Wu, F., Ju, T., Gong, Y., Zhang, Y., Wu, X., Hou, H., Zhao, L., Shi, H. **2019**, Mass balance-based inventory of heavy metals inputs to and outputs from agricultural soils in Zhejiang Province, China, *Science of the Total Environment*, **649**, 1269-1280.
7. Misseyer, M.P. & Van der Most, P.F.J. (1993, October) *Decision Support Systems for Environmental Impact Assessment; Addressing the Industrial Need*. Paper presented at the CSEIA '93: Proceedings of the IFIP TC5/WG5.11 Working Conference on Computer Support for Environmental Impact Assessment, ed by Guariso, G., Page, B., North-Holland Publishing Co., Elsevier Science Publishers, Amsterdam, Netherlands, pp. 301-310. Retrieved from: <https://dl.acm.org/doi/10.5555/647036.715555>
8. Belon, E., Boisson, M., De Portes, I., Eglin, T., Feix, I., Bispo, A., Galsomies, L., Leblond, S., Guellier, C. **2012**, An inventory of trace elements inputs to French agricultural soils, *Science of the Total Environment*, **439**, 87-95.
9. Hou, J., Shen, Y., Cao, S., Cheng, X., Wang, W. **2013**, Heavy metal flux research in Tianjin suburb vegetable farmland, *Journal of Anhui Agricultural Sciences*, **41**, 5764-5773.
10. Shi, T., Ma, J., Wu, X., Ju, T., Lin, X., Zhang, Y., Li, X., Gong, Y., Hou, H., Zhao, L. **2018**, Inventories of heavy metal inputs and outputs to and from agricultural soils: A review, *Ecotoxicology and environmental safety*, **164**, 118-124.
11. Leclerc, A., Laurent, A. **2017**, Framework for estimating toxic releases from the application of manure on agricultural soil: National release inventories for heavy metals in 2000–2014, *Science of the Total Environment*, **590**, 452-460.
12. Utermann, J., Düwel, O., Nagel, I., **2006**, Background values in European soils and sewage sludges. Results of a JRC-coordinated study on background values (Part II). European Commission DG-JRC., EUR, 22265
13. Hamilton, E.M.; Young, S.D., Bailey, E.H., Watts, M J. **2018**, Chromium speciation in foodstuffs: a review, *Food Chemistry*, **250**, 105-112
14. Panichev, N., Mandiwana, K., Kataeva, M., Siebert, S. **2005**, Determination of Cr (VI) in plants by electrothermal atomic absorption spectrometry after leaching with sodium carbonate, *Spectrochimica Acta Part B: Atomic Spectroscopy*, **60**, 699-703.
15. Sakizadeh, M., Martin, J.A.R., Zhang, C., Sharafabadi, F.M., Ghorbani, H. **2018**, Trace elements concentrations in soil, desert-adapted and non-desert plants in central Iran: Spatial patterns and uncertainty analysis, *Environmental pollution*, **243**, 270-281.
16. Filippini, T., Cilloni, S., Malavolti, M., Violi, F., Malagoli, C., Tesaro, M., Bottecchi, I., Ferrari, A., Vescovi, L., Vinceti, M. **2018**, Dietary intake of cadmium, chromium, copper, manganese, selenium and zinc in a Northern Italy community, *Journal of Trace Elements in Medicine and Biology*, **50**, 508-517.
17. Cai, L.M., Wang, Q.S., Wen, H.H., Luo, J., Wang, S. **2019**, Heavy metals in agricultural soils from a typical township in Guangdong Province, China: Occurrences and spatial distribution, *Ecotoxicology and environmental safety*, **168**, 184-191.
18. Fosu-Mensah, B.Y., Addae, E., Yirenya-Tawiah, D., Nyame, F. **2017**, Heavy metals concentration and distribution in soils and vegetation at Korle Lagoon area in Accra, Ghana, *Cogent Environmental Science*, **3**, Article ID 1405887. DOI: <https://doi.org/10.1080/23311843.2017.1405887>
19. Li, X., Li, Z., Lin, C.-J., Bi, X., Liu, J., Feng, X., Zhang, H., Chen, J., Wu, T. **2018**, Health risks of heavy metal exposure through vegetable consumption near a large-scale Pb/Zn smelter in central China, *Ecotoxicology and environmental safety*, **161**, 99-110.
20. Yu, H.-Y., Chang, C., Li, F., Wang, Q., Chen, M., Zhang, J. **2018**, Thallium in flowering cabbage and lettuce: Potential health risks for local residents of the Pearl River Delta, South China, *Environmental pollution*, **241**, 626-635.
21. Liu, J., Li, N., Zhang, W., Wei, X., Tsang, D.C., Sun, Y., Luo, X., Bao, Z.A., Zheng, W., Wang, J. **2019**, Thallium contamination in farmlands and common vegetables in a pyrite mining city and potential health risks, *Environmental pollution*, **248**, 906-915.
22. Pozzatti, M., Nakadi, F.V., Vale, M.G.R., Welz, B. **2017**, Simultaneous determination of nickel and iron in vegetables of Solanaceae family using high-resolution continuum source graphite furnace atomic absorption spectrometry and direct solid sample analysis, *Microchemical Journal*, **133**, 162-167.
23. Srinivas, N., Rao, S. R., Kumar, K.S. **2009**, Trace metal accumulation in vegetables grown in industrial and semi-urban areas-a case study, *Applied Ecology and Environmental Research*, **7**, 131-139.
24. Zhang, H., Li, H., Wang, Z., Zhou, L. **2011**, Accumulation characteristics of copper and cadmium in greenhouse vegetable soils in Tongzhou District of Beijing, *Procedia Environmental Sciences*, **10**, 289-294.
25. Samara, C., Misaelides, P., Tsalev, D., Anousis, I., Kouimtzi, T. **1992**, Trace element distribution in vegetables grown in the industrial area of Thessaloniki/Greece. *Fresenius Environmental Bulletin*, **1**, 577-582.
26. Hussain, A., Alamzeb, S., Begum, S. **2013**, Accumulation of heavy metals in edible parts of vegetables irrigated with waste water and their daily intake to adults and children, District Mardan, Pakistan, *Food Chemistry*, **136**, 1515-1523.
27. Khan, S., Cao, Q., Zheng, Y., Huang, Y., Zhu, Y. **2008**, Health risks of heavy metals in contaminated soils and food crops irrigated with wastewater in Beijing, China, *Environmental pollution*, **152**, 686-692.
28. Chekri, R., Le Calvez, E., Zinck, J., Leblanc, J.-C., Sirot, V., Hulin, M., Noël, L., Guérin, T. **2019**, Trace element contents in foods from the first French total diet study on infants and toddlers, *Journal of Food Composition and Analysis*, **78**, 108-120.
29. Melø, R., Gellein, K., Evje, L., Syversen, T. **2008**, Minerals and trace elements in commercial infant food, *Food and chemical toxicology*, **46**, 3339-3342.