



Eclética Química

ISSN: 0100-4670

ISSN: 1678-4618

ecletica@journal.iq.unesp.br

Universidade Estadual Paulista Júlio de Mesquita Filho
Brasil

Soares Emídio, Elissandro; Pereira da Silva, Claudia; Rodrigues de Marchi, Mary Rosa

Estrogenic mycotoxins in surface waters of the Rico Stream micro-basin, São Paulo, Brazil: occurrence and potential estrogenic contribution

Eclética Química, vol. 45, no. 2, 2020, pp. 23-32

Universidade Estadual Paulista Júlio de Mesquita Filho
Brasil

DOI: <https://doi.org/10.26850/1678-4618eqj.v45.2.2020.p23-32>

Available in: <https://www.redalyc.org/articulo.oa?id=42963115003>

- How to cite
- Complete issue
- More information about this article
- Journal's webpage in [redalyc.org](https://www.redalyc.org)




 [redalyc.org](https://www.redalyc.org)

Scientific Information System Redalyc

Network of Scientific Journals from Latin America and the Caribbean, Spain and Portugal

Project academic non-profit, developed under the open access initiative

Estrogenic mycotoxins in surface waters of the Rico Stream microbasin, São Paulo, Brazil: occurrence and potential estrogenic contribution

Elissandro Soares Emídio⁺, Claudia Pereira da Silva⁺, Mary Rosa Rodrigues de Marchi⁺

São Paulo State University (Unesp), Institute of Chemistry, 55 Prof. Francisco Degni St, Araraquara, São Paulo, Brazil

⁺Corresponding author: Elissandro Soares Emídio, Phone: +55 16 3301-9735 email address: elissandro_se@yahoo.com.br

ARTICLE INFO

Article history:

Received: July 21, 2019

Accepted: December 11, 2019


Published: April 1, 2020

Keywords:

1. estrogenic mycotoxins
2. occurrence
3. surface waters
4. estrogenicity
5. southeast Brazil

ABSTRACT: The aim of this study was to assess the occurrence of zearalenone (ZEN) and its metabolites zearalanone (ZAN), α -zearalenol (α -ZEL), β -zearalenol (β -ZEL), α -zearalanol (α -ZAL), and β -zearalanol (β -ZAL) in the surface waters of northern São Paulo state (Brazil) and to evaluate the associated potential estrogenic contribution to the aquatic environment. The determination of the estrogenic mycotoxins in water samples from the Rico Stream microbasin yielded levels of up to 59 ng L⁻¹ and their corresponding calculated

estrogenic equivalent (cEEQ) values were between < 0.03 and 1.4 ng L⁻¹, which are associated with negative effects on the reproduction and growth of some fish species. The physicochemical and microbiological parameters were evaluated to determine the water quality in the Rico stream region. This study revealed the first data about the presence of estrogenic mycotoxins in the Brazilian river waters and emphasizes the need to include these substances in future public policies concerning water quality, since these compounds are not yet legally regulated. From an environmental aspect, it is necessary to take into account the continuous introduction into surface water of microcontaminants associated with wastewater effluent, such as estrogenic mycotoxins.



1. Introduction

Among the hundreds of mycotoxin-producing fungi, the genus *Fusarium* is considered one of the most prevalent in terms of animal health implications and economic damage¹. Resorcylic acid lactones (RALs) are compounds that exhibit endocrine disruptive behavior and are produced by fungi of the genus *Fusarium*². The representative mycotoxin of this class of RALs is zearalenone (ZEN), cited among the most common mycotoxins worldwide³.

In Brazil, there have been few studies on the incidence of fusariotoxins. For example, the

occurrence of ZEN was reported in maize⁴, rice⁵ and wheat⁶ in the south and southeast Brazilian regions. Fusariotoxins are mostly produced under high humidity and at temperatures of approximately 20 to 26 °C. The Brazilian climate offers good conditions for these toxins to be present in grains used in the diets of animals and humans⁷.

The incidence of ZEN and its metabolites has been reviewed extensively in food for humans and animals⁸. However, little is known about its impact or environmental distribution, with few studies having been carried out concerning these attributes⁹. Some publications have reported the occurrence of ZEN in surface water, groundwater

and effluent from sewage treatment plants and industries, with concentrations of ZEN or its metabolites ranging from undetected levels to 4120 ng L⁻¹ ¹⁰. Zearalanone (ZAN) and β -zearealenol (β -ZEL) have rarely been detected or analyzed in aqueous environmental samples. ZEN and its metabolites can contribute to the overall estrogenic activity in the environment and therefore could pose a risk to wild fish in their natural habitat, as indicated by a zebrafish study in which there was a change in the reproductive capacity of animals exposed to low concentrations of estrogenic compounds (100 to 3200 ng L⁻¹) for 21 days³.

Some studies conducted in Brazil have reported the occurrence of endocrine disruptors in surface waters, with monitoring results showing the periodic presence of these substances in natural waters, suggesting that different sources in addition to untreated sewage contribute to the input into water sources¹¹. Possible aqueous environmental contamination routes of estrogenic mycotoxins include: drainage water from fields cultivated with infected plants, runoff from livestock facilities or manure applications to field crops and human excretions in urban wastewaters¹².

The purpose of this study was to examine for the first time the occurrence of estrogenic mycotoxins in Brazilian water samples, using dispersive liquid-liquid microextraction (DLLME) followed by liquid chromatography–tandem mass spectrometry and to identify possible sources of contamination, seasonal variations and potential contribution to total estrogenic activity in the southeast region of Brazil (São Paulo state).

2. Experimental

2.1 Standards and reagents

ZEN, α -ZEL, β -ZEL, α -ZAL, β -ZAL and ZAN standards were all obtained from Sigma Aldrich (St. Louis, MO, USA) with purities of at least 98%. The zearalenone-d6 (surrogate internal standard, ZEN-d6) was supplied from Toronto Research Chemicals (North York, ON, Canada). Acetonitrile (ACN) and methanol (MeOH), both HPLC grade were provided from J. T. Baker (Phillipsburg, NJ, USA). Bromocyclohexane was

acquired from Alfa Aesar (MA, USA). Hydrochloric acid (37% m/v) and sodium hydroxide were purchased from J.T. Baker (Phillipsburg, NJ, USA) to adjust the pH of water samples. Deionized water was obtained using a Milli-Q purification system (Bedford, MA, USA).

2.2 Sampling area

The Rico Stream microbasin is localized in the northeast region of São Paulo state and consists of the municipalities Jaboticabal, Taquaritinga, Monte Alto, Guariba and Santa Ernestina (Fig. 1), occupying an area of approximately 563 km² at an altitude between 498 and 754 m.

The Rico Stream is the main source of water in the micro-basin, with a total of 59.2 km of watercourse length discharging into the Mogi Guaçu River. Because of its high representativeness in the region, Rico Stream stands out as the main supply of water for the city of Jaboticabal (population ca. 75,000). Anthropogenic impacts on the environment may occur both directly or indirectly, due to effluent discharges from domestic sewage treatment plants of the Monte Alto municipality and animal husbandry (poultry, swine and cattle)¹³.

In accordance with the classification proposed by Köeppen, the region possesses a climate defined as Aw, characterized as tropical with summer rainfall and dry winters. The average temperature is 23 °C, and the total average annual rainfall is 1405 mm occurring mainly during the period from October to March, with the dry season extending from April to September¹⁴.




Figure 1. Map of the studied area showing the location of four sampling sites (Adapted from Pissarra *et al.*¹⁵ and Takahashi *et al.*¹⁶).

2.3 Sample collection and pretreatment

The collection of water samples was performed from October 2014 to July 2015 (ten sampling campaigns) at four different sampling sites located in the Rico Stream, except one that was located in Tijuco Stream (stream tributary) between the municipalities of Monte Alto and Jaboticabal. Reference sites: P1 situated near the Rico Stream source with minor contamination impacts and upstream of wastewater treatment plant (WWTP) effluent discharge (21°18'37.26"S; 48°27'39.49"W); P2 situated at the downstream discharge point of the WWTP, which also represents the influence of input from agricultural industry and runoff from livestock facilities or fields receiving livestock manure applications (21°18'46.08"S; 48°26'56.33"W); P3 located in the Tijuco Stream, upstream of the public water supply and near the confluence with the Rico Stream (21°17'56.08"S; 48°22'31.69"W) and P4 located freshwater abstraction site for the public water

supply to Jaboticabal city (21°18'36.07"S; 48°19'26.11"W).

The water samples were collected at 30 cm depth in 1000 mL amber glass bottles, previously rinsed thoroughly with ultrapure water. After collection, samples were transported to the laboratory under refrigeration at 4 °C (ice packs) and processed within 48 h. In the laboratory prior to extraction, the samples were filtered through a glass fiber filter with a pore size of 0.6 µm (Macherey-Nagel GF-3), acidified to pH 4 and spiked to 200 ng L⁻¹ with an internal surrogate (ZEN-d₆). Samples were then immediately extracted and analyzed using the previously developed and validated dispersive liquid-liquid microextraction (DLLME) method and liquid chromatography coupled to tandem mass spectrometry (LC-MS/MS)¹⁷.

2.4 Determination of physicochemical and microbiological parameters

A total of twelve physicochemical and microbiological parameters were used to assess water quality control and pollution. Parameters such as fecal coliforms, total organic carbon, chemical oxygen demand (COD), biochemical oxygen demand (BOD), total phosphorous, total nitrogen, nitrate, nitrite, ammoniacal nitrogen, potassium, turbidity and total dissolved solids were determined using standard procedures described by the American Public Health Association¹⁸.

2.5 Analytical Methodology

An aliquot of 10 mL of the water sample was added into a 12 mL conical glass centrifuge tube with screw cap. Thereafter, 100 µL of bromocyclohexane (extraction solvent) was added to the sample solution. The mixture was then vigorously shaken using a vortex mixer at 2500 rpm for 120 s. After centrifugation for 10 min at 3500 rpm, the organic sedimented phase was collected in a glass insert (300 µL) using a 100 µL Hamilton microsyringe and was evaporated in a N₂ flow. The residue was reconstituted in 80 µL of MeOH/0.1% formic acid in H₂O (50:50, v/v) and injected into the LC-ESI-MS/MS for analysis.

2.6 Instrumentation

The HPLC analyses were carried out on an Agilent 1200 series LC system coupled with a 3200 QTRAP mass spectrometer (Applied Biosystems/MDS Sciex Instruments) with an electrospray ionization source (ESI). The mass spectrometric parameters selected for experiments of full scan and ion fragmentation were as follow: ion spray voltage of -4 kV, curtain gas (nitrogen) at 15 psi, auxiliary gas at 40 psi, nebulizer gas at 50 psi, source temperature at 600 °C, interface

heater and entrance potential of -10 V. Chromatographic separation of the estrogenic mycotoxins was achieved at 40 °C using a Gemini C₁₈ column (150 × 4.6 mm, 5 µm; Phenomenex, Torrance, CA, USA) preceded by a Phenomenex C₁₈ column guard (4 × 3 mm). The mobile phase consisted of 0.1% formic acid in water (48%, v/v), acetonitrile (25%, v/v) and methanol (27%, v/v), with a flow rate of 1.0 mL min⁻¹ for 17 min and injection volume was set to 20 µL. Applied Biosystems Analyst 1.5 software was used for data acquisition.

2.7 Quality assurance/quality control

The calibration was performed by internal standardization method with matrix-matched blank samples. The calibration curve consists of eight different concentration values, in the range 8 to 680 ng L⁻¹. The analytes were extracted by DLLME prior to chromatographic analysis for each point of the calibration curve. The calibration curve was obtained from the ratio between the peak areas of analytes and the internal standard versus the analyte concentrations (ng L⁻¹). Every experimental point was analyzed in triplicate. The extraction efficiency (recovery percentage) of the analytes was evaluated by spiked with 200 ng L⁻¹ of the internal surrogate (ZEN-d6) to all river water samples.

2.8 Estrogenicity

The estrogenic potential of the aqueous samples was calculated in equivalent activity compared to the reference substance (natural hormone 17-β-estradiol)¹⁹. The potential contribution of each estrogenic mycotoxin to the total estrogenic activity of the surface waters was calculated according to the Eq. 1:

$$cEEQ = \text{concentration of estrogenic mycotoxin} \times RP \quad (1)$$

where cEEQ is the calculated estrogenic equivalent (17-β-estradiol) of a specific estrogenic mycotoxin and RP is the relative potency of the mycotoxin in an individual bioassay.

3. Results and Discussion

3.1 Occurrence of estrogenic mycotoxins in environmental samples

The results obtained from the collected samples (during the period of October 2014 to July 2015) are shown in Tab. 1. Surrogate recovery values (ZEN-d6) ranged from 71 to 124% from the extraction method. These recovery values are

suitable when dealing with complex environmental samples such as the river water analyzed in this study.

Table 1. Estrogenic mycotoxin concentrations (ng L⁻¹) at four sampling sites in the Rico Stream micro-basin.

Compounds	October/2014				November/2014			
	P1	P2	P3	P4	P1	P2	P3	P4
β-ZAL	n.d. ^c	n.d.	n.d.	n.d.	48	n.d.	n.d.	n.d.
β-ZEL	n.d.	<40 ^b	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
α-ZAL	n.d.	n.d.	n.d.	n.d.	<20 ^b	<20 ^b	27	<20 ^b
α-ZEL	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
ZAN	n.d.	n.d.	n.d.	n.d.	<8 ^b	n.d.	n.d.	n.d.
ZEN-d ₆ ^a	112	108	107	106	107	96	76	91
ZEN	<8 ^b	<8 ^b	<8 ^b	<8 ^b	<8 ^b	<8 ^b	<8 ^b	<8 ^b
Compounds	December/2014				January/2015			
	P1	P2	P3	P4	P1	P2	P3	P4
β-ZAL	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
β-ZEL	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
α-ZAL	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
α-ZEL	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
ZAN	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
ZEN-d ₆ ^a	89	80	91	93	112	81	74	77
ZEN	n.d.	n.d.	n.d.	n.d.	22	29	25	27
Compounds	February/2015				March/2015			
	P1	P2	P3	P4	P1	P2	P3	P4
β-ZAL	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
β-ZEL	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
α-ZAL	n.d.	n.d.	n.d.	n.d.	<20 ^b	n.d.	n.d.	n.d.
α-ZEL	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
ZAN	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
ZEN-d ₆ ^a	118	105	97	97	73	71	115	100
ZEN	n.d.	n.d.	n.d.	n.d.	59	46	51	42
Compounds	April/2015				May/2015			
	P1	P2	P3	P4	P1	P2	P3	P4
β-ZAL	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
β-ZEL	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
α-ZAL	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
α-ZEL	n.d.	n.d.	n.d.	n.d.	<20 ^b	n.d.	n.d.	n.d.
ZAN	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
ZEN-d ₆ ^a	91	81	73	78	114	124	101	127
ZEN	n.d.	n.d.	n.d.	n.d.	18.8	n.d.	n.d.	n.d.
Compounds	June/2015				July/2015			
	P1	P2	P3	P4	P1	P2	P3	P4
β-ZAL	n.d.	n.d.	<40 ^b	n.d.	n.d.	n.d.	n.d.	n.d.
β-ZEL	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
α-ZAL	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
α-ZEL	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
ZAN	n.d.	n.d.	19	n.d.	n.d.	n.d.	n.d.	n.d.
ZEN-d ₆ ^a	117	119	119	88	102	103	98	88
ZEN	14	12	11	<8 ^b	n.d.	n.d.	n.d.	n.d.

^aZEN-d₆ - Recovery (%), spiked at 200 ng L⁻¹ in triplicate

^b limit of quantification value

^c n.d. – not detected

All analytes were detected at least once at each point over the sampling period. ZEN compound had a higher frequency of detection (52.5%), followed by α -ZAL (12.5%), β -ZAL (5.0%), ZAN (5.0%), α -ZEL (5.0%) and β -ZEL (2.5%). A total of 47.5% of the samples had concentrations above the limit of quantification (LOQ), ranging from 11 to 59 ng L⁻¹. Among the six analytes, ZEN was determined at all sampling points in five of the sampling campaigns performed. The highest levels were determined during the period of greatest rainfall. In addition, most ZEN metabolites were detected during the rainy season. The compounds β -ZAL, α -ZAL and ZAN were quantified only once, whereas α -ZEL and β -ZEL did not present in concentrations above the LOQ (Tab. 1). The limits of detection were for β -ZAL and β -ZEL of 20 ng L⁻¹, for α -ZAL and α -ZEL of 8 ng L⁻¹ and for ZAN and ZEN of 4 ng L⁻¹.

The results obtained in this work are comparable to those obtained by other researchers. In previous studies, the detection frequencies for ZEN ranged from 0 to 30% and differed according to the seasons during sampling^{9,20-22}. The highest concentrations of ZEN (up to 96 ng L⁻¹) were detected in small streams or channels near agricultural fields with crops contaminated by *Fusarium* sp.^{9,20,23}. Low levels (up to 35 ng L⁻¹) were found in river and lake waters^{18,22-25}. The metabolites β -ZAL and α -ZAL were investigated in two rivers contaminated by wastewater effluents²⁴. The maximum concentration of both compounds was 3 ng L⁻¹. The concentrations of β -ZEL and α -ZEL metabolites were investigated in agricultural streams upstream and downstream of the WWTP in New York (USA). In agricultural fields, the concentrations of these compounds were generally below the limit of detection, but the highest detected concentrations reached tens or even hundreds of ng L⁻¹. In summary, the levels of estrogenic mycotoxins detected in rivers, streams and drainage channels occurred generally at levels of dozens of ng L⁻¹. However, much higher concentrations have occasionally been reported (single study), such as the detection of hundreds of ng L⁻¹ for α -ZEL and β -ZEL in streams near agricultural areas or wastewater effluents in New York, USA⁹.

The results obtained for physical, chemical and microbiological parameters at the sampling sites (Tab. 2) showed that variables such as fecal coliforms (December–July), BOD₅ (October and January), total phosphorus (October), nitrite

concentration (December), and total ammoniacal nitrogen concentration (October) exceeded the values established by CONAMA for class 2 freshwater (Brazilian Environmental Council)²⁶.

According to the results of these parameters (Tab. 2), there was increase in the total phosphorus, potassium concentration, total dissolved solids and turbidity during the rainy season (November to March), which was characteristic of rainfall events and the corresponding transport of fine soil particles into the body of water. There was an increase in the concentration total nitrogen during the extensive dry season in October 2014. It is assumed that the high concentration of ammoniacal nitrogen (42 mg L⁻¹ for P2 site) probably originated from runoff waters that carried residues from livestock animals or by the drainage of agricultural soils where animal excrement was used as fertilizer.

The fecal contamination indicator (fecal coliforms) exceeded the maximum limit of 1000 CFU 100 mL⁻¹²⁴ in samples from all months, with the exception of October and November 2014. These results indicated that Rico Stream is subject to a great anthropogenic influence, probably related to the use and soil occupation of this microbasin. The Rico Stream microbasin presents serious problems of environmental degradation, characterized by residential and rural areas with large deforestation events and lack of adequate management of animal husbandry waste, resulting in contamination of the whole drainage network.

The P2 site had the highest concentrations of eight of the hydrological variables (fecal coliforms, BOD, total phosphorous, nitrate, nitrite, potassium, total dissolved solids, ammoniacal nitrogen concentrations), including 14 values above CONAMA legislation. This sampling site is characterized by tributaries that drain from the urban area of the Monte Alto municipality and is located downstream of the effluent discharge of the wastewater treatment plant (WWTP) (site that receives the highest contribution among others).

Soil occupation and land use in the area of the Rico Stream consists of small farms with diversified production. According to field observations during the sampling period, horticulture, swine and several agricultural crops (especially sugarcane and peanuts) were present in the area. The presence of mechanized irrigation of agricultural crops during both the dry and rainy periods was also observed during the sampling period. It may be inferred that the presence of

estrogenic mycotoxins in the Rico Stream was probably due to effluent discharge from the WWTP of Monte Alto municipality (detection in downstream of the WWTP – P2 site), runoff from livestock facilities or fields receiving livestock

manure applications (detection in upstream of the WWTP – P1 site). Rico Stream contamination from fungi in agricultural crops is unlikely since maize (temporary tillage) is insignificant in the region.

Table 2. Data on physicochemical and microbiological parameters determined in Rico Stream micro-basin over the period of sampling.

Month	Sampling sites	Fecal coliforms (CFU 100 mL ⁻¹)	Total organic carbon (mg L ⁻¹)	Chemical oxygen demand (mg L ⁻¹)	Biological oxygen demand (mg L ⁻¹)	Total phosphorus (mg L ⁻¹)	Nitrate (mg L ⁻¹)	Nitrite (mg L ⁻¹)	Total nitrogen (mg L ⁻¹)	Total ammoniacal nitrogen (mg L ⁻¹)	Total dissolved solids (mg L ⁻¹)	Potassium (mg L ⁻¹)	Turbidity (NTU)
October 2014	P1	850	10	24	5	0.098	1.0	<0.011	46	0.30	112	<1.109	11
	P2	65	72	180	81	1.820	3.4	0.621	46	42	343	<1.109	16
	P3	22	16	40	15	0.230	<1.0	<0.011	0.5	0.51	132	<1.109	7
	P4	40	21	52	30	0.176	5.5	0.040	6.0	0.50	159	<1.109	10
November 2014	P1	<1.0	9	23	5	0.123	6.3	<0.011	6.5	0.21	210	6.162	13
	P2	500	16	40	<5.0	0.420	7.2	<0.011	10	2.90	184	5.217	12
	P3	<1.0	22	55	<5.0	0.120	5.7	<0.011	5.9	0.20	168	6.801	12
	P4	70	8	20	<5.0	0.181	6.2	<0.011	6.4	0.25	198	5.209	13
December 2014	P1	2300	5	12	<5.0	0.055	0.33	0.024	0.5	0.16	48	2.482	14
	P2	2200	5	12	<5.0	0.707	0.78	1.301	2.4	0.26	83	4.091	16
	P3	3300	6	17	<5.0	0.059	0.75	0.032	0.9	0.13	48	2.719	15
	P4	3900	2	4	<5.0	0.072	2.6	0.026	2.7	0.11	47	3.073	15
January 2015	P1	600	14	37	15	<0.050	<1.0	0.020	0.4	0.12	94	1.593	4
	P2	20000	72	187	94	1.157	1.1	0.355	6.4	<0.05	111	3.892	7
	P3	5000	16	41	21	<0.050	<1.0	0.036	0.3	<0.05	138	1.946	46
	P4	6000	11	29	13	<0.050	1.6	0.032	2.3	0.25	132	2.034	58
February 2015	P1	95	<0.04	18	<5.0	<0.050	<1.0	0.013	0.2	<0.05	53	7.53	12
	P2	48	<0.04	18	<5.0	<0.050	<1.0	0.118	0.5	0.06	92	11.07	19
	P3	165	<0.04	22	<5.0	<0.050	<1.0	0.029	0.2	<0.05	54	7.699	14
	P4	1100	<0.04	11	<5.0	<0.050	3.8	0.071	4.0	0.14	54	9.994	15
March 2015	P1	61	8	19	<5.0	<0.050	<1.0	0.068	0.4	0.15	44	3.237	10
	P2	620	22	54	<5.0	0.196	<1.0	0.143	0.6	0.24	62	4.396	11
	P3	380	5	12	<5.0	<0.050	<1.0	0.069	0.6	0.36	40	3.886	56
	P4	3100	5	12	<5.0	0.037	1.3	0.100	2.0	0.34	40	3.611	37
April 2015	P1	140	<0.04	<4	<5.0	<0.050	1.0	<0.011	1.9	0.54	50	2.165	10
	P2	510	22	54	<5.0	0.237	12	0.108	14	1.82	90	4.569	10
	P3	1100	19	46	<5.0	<0.050	2.3	0.036	2.8	0.29	50	2.286	92
	P4	900	35	84	<5.0	<0.050	4.0	0.016	4.5	0.15	46	2.796	74
May 2015	P1	410	69	186	<5.0	<0.050	0.31	0.015	0.7	0.33	90	2.420	9
	P2	2000	136	367	<5.0	<0.050	9.0	0.019	10	0.44	152	3.230	11
	P3	4300	2	4	<5.0	<0.050	0.70	0.014	1.3	0.40	84	2.119	21
	P4	1200	40	108	<5.0	<0.050	2.3	0.011	2.5	0.09	84	2.276	19
June 2015	P1	1600	16	40	<5.0	<0.050	1.0	0.015	1.2	0.16	47	1.79	8
	P2	420	35	88	<5.0	<0.050	1.5	0.232	2.4	0.64	96	3.991	14
	P3	5000	22	56	<5.0	<0.050	0.8	0.020	0.9	0.07	42	1.892	14
	P4	960	21	52	<5.0	<0.050	3.4	0.047	3.4	<0.05	42	2.232	13
July 2015	P1	200	42	86	<5.0	<LQ	0.74	<LQ	0.7	<LQ	65	2.196	5.3
	P2	3600	57	93	<5.0	1.649	2.0	0.374	2.4	<LQ	106	3.671	9.7
	P3	2700	9	33	<5.0	<LQ	0.87	<LQ	0.9	<LQ	89	2.536	12.2
	P4	5200	12	29	<5.0	0.033	4.6	0.044	4.6	<LQ	76	2.474	9.26

3.2 Estrogenicity

The calculated cEEQ values ranged from < 0.03 to 1.4 ng L^{-1} (mean value) (Tab. 3). The highest values of cEEQ were for α -ZAL, even though the maximum concentration calculated in the samples was lower than those obtained for ZEN, as α -ZAL showed the highest relative estrogenic potency among the quantified mycotoxins. The cEEQ values for the estrogenic mycotoxins in this study (Tab. 3) were higher than those reported in natural waters in Europe and USA (< 0.01 to 0.63 ng L^{-1})²⁷, except for the study performed by Kolpin *et al.*⁹, in

which the highest value of cEEQ was calculated (New York, USA) for α -ZEL (< 8.8 to 388 ng L^{-1}).

In the specific case of ZEN, some adverse effects have been detected *in vivo* at concentrations comparable in this study. ZEN concentrations between 2 and 50 ng L^{-1} (lowest observed effect concentration - LOEC), activation of genes that regulate the growth and reproduction of fish (*Pimephales promelas*) has been reported²⁸. The concentrations of ZEN reported in this work (Tab. 1) were higher or comparable to the LOEC from *in vivo* studies (fish)²⁸.

Table 3. Concentration range, cEEQ and relative estrogenic potency of the analytes in the Rico Stream microbasin.

Compounds	Concentration range (ng L^{-1})	cEEQ ^a (mean value) (ng L^{-1})	cEEQ range ^b (ng L^{-1})	RP ^c
ZEN	$< 4.0^d$ –59	< 0.03 –0.47	< 0.01 –5.0	7.87×10^{-3}
ZAN	$< 4.0^d$ –19	< 0.07 –0.33	< 0.02 –3.1	1.76×10^{-2}
α -ZAL	$< 8.0^d$ –27	< 0.42 –1.4	< 0.15 –4.7	5.20×10^{-2}
β -ZAL	$< 20^d$ –48	< 0.43 –1.0	< 0.09 –2.3	2.14×10^{-2}

^a Mean cEEQs were calculated from individual mycotoxin concentrations multiplied by the geometric mean of relative estrogenic potencies (RPs) of analytes from different *in vitro* systems

^b cEEQs ranges were calculated as the concentrations of mycotoxins multiplied by the minimum and maximum RPs of the analytes from different *in vitro* systems

^c Geometric mean of the relative estrogenic potencies²⁷

^d LOD value

4. Conclusions

In this study, it was concluded that estrogenic mycotoxins are present along the Rico Stream microbasin and that the contribution routes of these compounds are the run-off from livestock farming practices, drainage of agricultural soils from animal manure or wastewater treatment plant effluent from Monte Alto city. Additionally, this is the first report of estrogenic mycotoxins in environmental waters in Brazil. All estrogenic mycotoxins were detected at some point during the study, and quantifiable concentrations (11 to 59 ng L^{-1}) were similar to those found in studies conducted in other countries that are influenced by agricultural areas, livestock around river waters, as well as waterbodies that receive WWTP effluents. Thus, our results can contribute to discussion of the improvement of the water and sewage treatment in Brazil, as well as to the development of global environmental legislation.

5. Acknowledgments

E. S. Emídio is grateful for the doctoral student fellowship from CNPq (Brazilian National Counsel of Technological and Scientific Development). Research support was provided by the FATE - Foundation for Science, Technology and Education Support. The authors are grateful to Dr. Laudicéia Giacometti Lopes and SAAJ for providing the sampling logistics.

6. References

- [1] Escrivá, L., Font, G., Manyes, L., *In vivo* toxicity studies of *Fusarium* mycotoxins in the last decade: a review, *Food and Chemical Toxicology* 78 (2015) 85–206. <https://doi.org/10.1016/j.fct.2015.02.005>.
- [2] Pamplona-Silva, M. T., Mazzeo, D. E. C., Bianchi, J., Marin-Morales, M. A., Estrogenic compounds: chemical characteristics, detection methods, biological and environmental effects, *Water, Air, & Soil Pollution* 229 (144) (2018) 1–27. <https://doi.org/10.1007/s11270-018-3796-z>.

- [3] Schwartz, P., Thorpe, K. L., Bucheli, T. D., Wettstein, F. E., Burkhardt-Holm, P., Short-term exposure to the environmentally relevant estrogenic mycotoxin zearalenone impairs reproduction in fish, *Science of The Total Environment* 409 (2) (2010) 326-333. <https://doi.org/10.1016/j.scitotenv.2010.10.017>.
- [4] Oliveira, M. S., Rocha, A., Sulyok, M., Krska, R., Mallmann, C. A., Natural mycotoxin contamination of maize (*Zea mays* L.) in the South region of Brazil, *Food Control* 73 (Part B) (2017) 127-132. <https://doi.org/10.1016/j.foodcont.2016.07.033>.
- [5] Dors, G. C., Bierhals, V. da S., Badiale-Furlong, E., Parboiled rice: chemical composition and the occurrence of mycotoxins, *Food Science and Technology* 31 (1) (2011) 172-177. <https://doi.org/10.1590/S0101-20612011000100025>.
- [6] Tibola, C. S., Fernandes, J. M. C., Guarienti, E. M., Effect of cleaning, sorting and milling processes in wheat mycotoxin content, *Food Control* 60 (2016) 174-179. <https://doi.org/10.1016/j.foodcont.2015.07.031>.
- [7] Santin, E., Maiorka, A., Zanella, I., Magon, L., Micotoxinas do *Fusarium* spp na avicultura comercial, *Ciência Rural* 31 (1) (2001) 185-190. <https://doi.org/10.1590/S0103-84782001000100030>.
- [8] Shore, L. S., Shemesh, M., Estrogen as an environmental pollutant, *Bulletin of Environmental Contamination and Toxicology* 97 (2006) 447-448. <https://doi.org/10.1007/s00128-016-1873-9>.
- [9] Kolpin, D. W., Schenzel, J., Meyer, M. T., Phillips, P. J., Hubbard, L. E., Scott, T.-M., Bucheli, T. D., Mycotoxins: diffuse and point source contributions of natural contaminants of emerging concern to streams, *Science of The Total Environment* 470-471 (2014) 669-676. <https://doi.org/10.1016/j.scitotenv.2013.09.062>.
- [10] Lundgren, M. S., Novak, P. J., Quantification of phytoestrogens in industrial waste streams, *Environmental Toxicology and Chemistry* 28 (11) (2009) 2318-2323. <https://doi.org/10.1897/09-029.1>.
- [11] Pessoa, G. P., Santos, A. B. dos, Souza, N. C. de, Alves, J. A. C., Nascimento, R. F. do, Development of methodology to determine estrogens in wastewater treatment plants, *Química Nova* 35 (5) (2012) 968-973. <https://doi.org/10.1590/S0100-40422012000500019>.
- [12] Schenzel, J., Hungerbühler, K., Bucheli, T. D., Mycotoxins in the environment: II. Occurrence and origin in Swiss river waters, *Environmental Science & Technology* 46 (24) (2012) 13076-13084. <https://doi.org/10.1021/es301558v>.
- [13] Zanini, H. L. H. T., Amaral, L. A. do, Zanini, J. R., Tavares, L. H. S., Caracterização da água da microbacia do córrego rico avaliada pelo índice de qualidade de água e de estado trófico, *Engenharia Agrícola* 30 (4) (2010) 732-741. <https://doi.org/10.1590/S0100-69162010000400017>.
- [14] Centro de Pesquisas Meteorológicas e Climáticas Aplicadas à Agricultura (CEPAGRI), Clima dos Municípios Paulistas. 2018.
- [15] Pissarra, T. C. T., Rodrigues, F. M., Politano, W., Galbiatti, J. A., Morfometria de microbacias hidrográficas do córrego Rico, afluente do Rio Mogi-Guaçu, Estado de São Paulo, Brasil, *Revista Árvore* 34 (4) (2010) 669-676. <https://doi.org/10.1590/S0100-67622010000400011>.
- [16] Takahashi, E. L. H., Rosa, F. R. T., Langeani, F., Nakaghi, L. S. O., Spatial and seasonal patterns in fish assemblage in Córrego Rico, upper Paraná River basin, *Neotropical Ichthyology* 11 (1) (2013) 143-152. <https://doi.org/10.1590/S1679-62252013000100017>.
- [17] Emídio, E. S., Silva, C. P. da, Marchi, M. R. R. de, Determination of estrogenic mycotoxins in environmental water samples by low-toxicity dispersive liquid-liquid microextraction and liquid chromatography-tandem mass spectrometry, *Journal of Chromatography A* 1391 (2015) 1-8. <https://doi.org/10.1016/j.chroma.2015.02.067>.
- [18] American Public Health Association (APHA), American Water Works Association (AWWA), Water Environment Federation (WEF), Standard Methods for the Examination of Water and Wastewater, American Public Health Association, Washington, 21th ed., 2005.
- [19] Jarošová, B., Bláha, L., Giesy, J. P., Hilscherová, K., What level of estrogenic activity determined by *in vitro* assays in municipal waste waters can be considered as safe?, *Environment International* 64 (2014) 98-109. <https://doi.org/10.1016/j.envint.2013.12.009>.
- [20] Gromadzka, K., Waśkiewicz, A., Goliński, P., Świetlik, J., Occurrence of estrogenic mycotoxin - zearalenone in aqueous environmental samples with various NOM content, *Water Research* 43 (4) (2009) 1051-1059. <https://doi.org/10.1016/j.watres.2008.11.042>.
- [21] Hartmann, N., Erbs, M., Forrer, H.-R., Vogelgsang, S., Wettstein, F. E., Schwarzenbach, R. P., Bucheli, T. D., Occurrence of zearalenone on *Fusarium graminearum* infected wheat and maize fields in crop organs, soil, and drainage water, *Environmental Science & Technology* 42 (15) (2008) 5455-5460. <https://doi.org/10.1021/es8007326>.

- [22] Maragos, C. M., Zearalenone occurrence in surface waters in central Illinois, USA, *Journal Food Additives & Contaminants: Part B Surveillance* 5 (1) (2012) 55-64. <https://doi.org/10.1080/19393210.2012.659764>.
- [23] Bucheli, T. D., Wettstein, F. E., Hartmann, N., Erbs, M., Vogelgsang, S., Forrer, H.-R., Schwarzenbach, R. P., *Fusarium* mycotoxins: overlooked aquatic micropollutants?, *Journal of Agricultural and Food Chemistry* 56 (3) (2008) 1029-1034. <https://doi.org/10.1021/jf073082k>.
- [24] Laganà, A., Bacaloni, A., De Leva, I., Faberi, A., Fago, G., Marino, A., Analytical methodologies for determining the occurrence of endocrine disrupting chemicals in sewage treatment plants and natural waters, *Analytica Chimica Acta* 501 (1) (2004) 79-88. <https://doi.org/10.1016/j.aca.2003.09.020>.
- [25] Kolpin, D. W., Hoerger, C. C., Meyer, M. T., Wettstein, F. E., Hubbard, L. E., Bucheli, T. D., Phytoestrogens and mycotoxins in Iowa Streams: an examination of under investigated compounds in agricultural basins, *Journal of Environmental Quality* 39 (6) (2010) 2089-2099. <https://doi.org/10.2134/jeq2010.0121>.
- [26] Conselho Nacional do Meio Ambiente (CONAMA), Resolução nº 357 de 17 de março de 2005, Ministério do Meio Ambiente (2005) 1-27. <http://www.mma.gov.br/port/conama/res/res05/res35705.pdf>.
- [27] Jarošová, B., Javůrek, J., Adamovský, O., Hilscherová, K., Phytoestrogens and mycoestrogens in surface waters-their sources, occurrence, and potential contribution to estrogenic activity, *Environment International* 81 (2015) 26-44. <https://doi.org/10.1016/j.envint.2015.03.019>.
- [28] Johns, S. M., Denslow, N. D., Kane, M. D., Watanabe, K. H., Orlando, E. F., Sepúlveda, M. S., Effects of estrogens and antiestrogens on gene expression of fathead minnow (*Pimephales promelas*) early life stages, *Environmental Toxicology* 26 (2) (2011) 195-206. <https://doi.org/10.1002/tox.20545>.