



Revista de Biología Tropical

ISSN: 0034-7744

ISSN: 0034-7744

Universidad de Costa Rica

Antunes, Altamir Rocha; Elias, Guilherme Alves; Pezente, Gisele; Dos Santos, Robson
Scientific literature on *Inga* (Fabaceae) from Santa Catarina state, Southern Brazil (1983-2017)

Revista de Biología Tropical, vol. 67, no. 6, 2019, pp. 1247-1256

Universidad de Costa Rica

DOI: DOI 10.15517/RBT.V67I6.34731

Available in: <http://www.redalyc.org/articulo.oa?id=44965908012>

- ▶ How to cite
- ▶ Complete issue
- ▶ More information about this article
- ▶ Journal's webpage in redalyc.org

UDEM 

Scientific Information System Redalyc

Network of Scientific Journals from Latin America and the Caribbean, Spain and Portugal

Project academic non-profit, developed under the open access initiative

Scientific literature on *Inga* (Fabaceae) from Santa Catarina state, Southern Brazil (1983-2017)

Altamir Rocha Antunes*, Guilherme Alves Elias, Gisele Pezente & Robson dos Santos

Herbário Pe. Dr. Raulino Reitz (CRI), Programa de Pós-Graduação em Ciências Ambientais, Universidade do Extremo Sul Catarinense – UNESC, Criciúma, SC, Brasil; altamirmad@hotmail.com, guilherme@unesc.net, gipezente@hotmail.com, rsa@unesc.net

* Correspondence

Received 08-X-2018. Corrected 19-VII-2019. Accepted 18-IX-2019.

ABSTRACT. The genus *Inga* Mill. belongs to the mimosoid clade (Fabaceae, Caesalpinioideae) that includes 131 species in Brazil. It is the most important genus of Fabaceae. In this sense, this study aimed to perform a bibliometric analysis on *Inga* from Santa Catarina state. A survey of the published literature was conducted using the electronic databases of the Web of Science, Scopus and SciELO with the accepted names of *Inga* species and its synonyms. Papers were distributed in four subject categories: C1 (Ecological), C2 (morphology, anatomy, taxonomy, histology, physiology and genetics), C3 (production and use) and C4 (biochemical and nutritional properties). We registered 232 papers for 13 species of *Inga*. C1 was the most studied subject category, mainly in topics such as nutrient supply, shade and nitrogen fixing capacity. We also noticed that the subjects diversified over the years, with registered papers in all categories. *Inga edulis*, *I. vera* and *I. marginata* were the most registered species in our survey. Our results showed an increase in the number of articles on *Inga* over time, especially in the last 13 years. However, some important gaps need to be addressed, such as the relatively small number and/or lack of studies conducted for some species.

Key words: bibliometrics, *Inga edulis*, legumes, Ingeae.

Rocha Antunes, A., Alves Elias, G., Pezente, G., & dos Santos, R. (2019). Scientific literature on *Inga* (Fabaceae) from Santa Catarina state, Southern Brazil (1983-2017). *Revista de Biología Tropical*, 67(6), 1247-1256.

The Fabaceae Lindl. (legumes) family is the richest in Brazil with 751 genus and 19 000 species; it is one of the most representative in the world (Bruneau et al., 2013). Among them, *Inga* is the most expressive and larger genus of Ingeae (Possette & Rodrigues, 2010). It can be distinguished from other genera for their arboreal habit, paripinnate leaves with nectary located between each pair of leaflets, fleshy and indehiscent fruits with seeds surrounded by a sweet and fleshy sarcotesta (Vasconcelos, 2014). *Inga* is exclusively neotropical and Brazil is one of the main centers of genetic diversity (Pennington, 1997). It has economic

potential for reforestation, phytotherapy, energy production and food (Fernandes, Dondoni Da Costa, Araújo, & Lopes, 2016). Besides that, legumes are essential for fertilization, fodder, wood, tannins, oils, resins in the manufacture of varnishes, paint and dyes and in horticultural trade (Gepts et al., 2005; LPWG, 2017).

As most information about *Inga* is being published and made available in electronic databases, studies have been focused on nitrogen fixation capacity (Yatazawa, Uchino, & Hambali, 1983), shading plantations (Bishop, 1983; Staver, 1989; Alegre & Rao, 1996; Kettler, 1996), medicinal potential (Tauchen et al.,

2016) and nutritional and biochemical properties (Lima, Santos, & Porta, 2018). Bibliometric studies are very important and are being increasingly used to measure and analyze the scientific development in a specific research field (Hood & Wilson, 2001; Elias, Corrêa, Citadini-Zanette, & Santos, 2015).

The state of Santa Catarina has been a pioneer in the study of its forests and flora, exemplified by the Illustrated Flora of Santa Catarina - one of the most comprehensive works on Brazilian plants - developed by Reitz (1965), and the Barbosa Rodrigues Herbarium (HBR). Currently, the Floristic and Forest Inventory of Santa Catarina (IFFSC) (Vibrans, Sevegnani, Gasper, & Lingner, 2012a; Vibrans, Sevegnani, Gasper, & Lingner, 2012b; Vibrans, Sevegnani, Gasper, Müller, & Reis et al. 2013a; Vibrans, Sevegnani, Gasper, & Lingner, 2013b; Vibrans, Sevegnani, Gasper, & Lingner, 2013c) is responsible of disseminating current comprehensive data on Santa Catarina forest flora. In this context, this paper aimed to carry out a bibliometric analysis of the genus *Inga* naturally

occurring in Santa Catarina, southern Brazil. We expect that our results may provide support for future researches on *Inga* species in Brazil.

Literature reviewed: The survey of the published literature on *Inga* was conducted using the databases Web of Science, Scopus and SciELO. Accepted names and the synonyms of naturally occurring species in Santa Catarina (Tropicos, 2013; Flora do Brasil 2020 Under Construction, 2019) were used as keywords (Table 1). The search was performed in March 2018 and all papers published until December 31st, 2017 were compiled.

As mentioned before, the study area is located in the state of Santa Catarina, southern Brazil because *Inga* has great representation in the tree component in this area (Burkart, 1979). However, few articles deal specifically with the genus in the state. The Köppen climate classification for Santa Catarina is humid subtropical with no defined dry season and hot summers (Cfa) or balmy summers (Cfb), corresponding to 40 and 60 % of the study

TABLE 1
Inga species distributed in Santa Catarina with its synonyms and vernacular names

TABLA 1
Especies de *Inga* distribuidas en Santa Catarina con sus sinónimos y nombres vernaculares

Accepted names	Synonyms	Vernacular names
<i>Inga edulis</i> Mart.	<i>Inga scabriuscula</i> Benth.	ingá-cipó
<i>Inga edwallii</i> (Harms) T.D.Penn.	none	ingá
<i>Inga lentiscifolia</i> Benth.	none	ingá
<i>Inga marginata</i> Willd.	none	ingá-feijão
<i>Inga sellowiana</i> Benth.	none	ingá-ferro
<i>Inga sessilis</i> (Vell.) Mart.	none	ingá-macaco
<i>Inga striata</i> Benth.	<i>Inga nuda</i> Salzm. <i>Inga salzmanniana</i> Benth.	ingá-quadrado
<i>Inga subnuda</i> Salzm. ex Benth.		ingá
<i>Inga subnuda</i> subsp. <i>luschnathiana</i> (Benth.) T.D.Penn.	<i>Inga luschnathiana</i> Salzm. ex Benth.	ingá
<i>Inga vera</i> Willd.	<i>Inga spuria</i> Humb. & Bonpl. ex Willd.	ingá-do-brejo
<i>Inga vera</i> subsp. <i>affinis</i> (DC.) T.D.Penn.	<i>Inga arinensis</i> Hoehne <i>Inga meissneriana</i> Miq. <i>Inga uraguensis</i> Hook. & Arn. <i>Inga uruguensis</i> Hook. & Arn. <i>Inga affinis</i> DC.	ingá
<i>Inga virescens</i> Benth.	none	ingá-torcido
<i>Inga vulpina</i> Mart. ex Benth.	<i>Inga guilleminiana</i> Benth.	ingá-estrela

TABLE 2
Study categories for *Inga* species distributed in Santa Catarina

TABLA 2
Categorías de estudio para las especies de *Inga* distribuidas en Santa Catarina

Category	Study categories	Description
C1	Ecological	Floristics, phytosociology, phenology, conservation, population dynamics and interactions.
C2	Biological properties	Morphology, anatomy, histology, taxonomy, plant physiology and genetics.
C3	Production and use	Production and transformation of forest products, use by communities or commercialization.
C4	Biochemical and nutritional properties	Plant services used as feedstock for the isolation of one or more biochemical substances.




Fig. 1. Scientific production overview on *Inga* species distributed in Santa Catarina, southern Brazil.

Fig. 1. Producción científica sobre especies de *Inga* distribuidas en Santa Catarina, sur de Brasil.

area respectively (Alvares, Stape, Sentelhas, Gonçalves, & Sparovek, 2013). Rainfall is well distributed throughout the state with an annual average of 1 250 - 2 000 mm /year with no areas of regular drought (Nimer, 1990). Each paper was identified by: (i) the year of publication, (ii) the scientific journal of publication, (iii) authors and (iv) study categories (Table 2) (Elias et al., 2015).

Trends in the literature: We found 232 papers for 13 species of *Inga* between 1983 and 2017. (Fig. 1). All papers were distributed in 133 scientific journals from different

locations. All articles resulted in 934 collaborations among the 232 articles.

From 1983 to 1989 few papers were published and, for many years, no papers appeared on the subject. During this period, papers were mainly focused in the genus consortium with other species, highlighting the supply of nutrients and shading, which increases the yield of productions of coffee, cocoa and banana (Bishop, 1983; Staver, 1989). The following decade, publications maintained the same trend. Exceptionally, in 1996, the amount of articles published increased. At this time, papers called attention on the consortium between *Inga* and

monocultures, such as beans and rice (Alegre & Rao, 1996; Kettler, 1996).

From the year 2000 until 2010, the number of papers grew: in 2006, 14 papers were published, mainly on biochemical and nutritional properties of *Inga* (Arévalo-Pinedo, Dos Santos, Salles Arévalo, Zuniga, & Pinedo, 2006; Faria et al., 2006). These works dealt with *Inga*'s increase in nutritional compounds as well as its conservation of germplasm. Despite the rise in publications, 2009 showed a decrease. The articles focused on *Inga*'s ecological relation with other organisms, such as the preference of birds for *Inga* species (Fink, Lindell, Morrison, Zahawi, & Holl, 2009).

During the years 2010-2017, we registered the highest scientific production on *Inga*. Most papers were focused on properties of secondary metabolism and applications in plant biological activities (De Sousa Dias, De Souza, & Rogez, 2010; Pinto, Souza, & Oliveira, 2010). The scientific production decreased exclusively in 2012 and 2015. However, they followed the same trend, where they focused on exploring the biochemical properties of *Inga* and their applications (dos Santos Barros, do Nascimento, & de Azevedo, 2012; Guillermo-Ferreira, Cardoso-Leite, & Gandolfo, 2012; De Freitas et al., 2015; Nygren & Leblanc, 2015).

Over the years, the growth in scientific production has been intensified in several areas of knowledge, such as ecology, biological activities, production and phytochemistry (De Sousa Dias et al., 2010; da Silva et al., 2014; Lamarca & Barbedo, 2015; Avila Jr., Pinheiro, & Sazima, 2015; Machado, de Oliveira, Zério, Parra, & Macedo, 2017). All papers were distributed in 133 scientific journals in several areas of knowledge. Agroforestry Systems was the main journal with 17 published papers, followed by Forest Ecology and Management (8), Revista Árvore (7), Rodriguésia (7) and Biotropica (6). These journals published articles on Forest Sciences, particularly the environment and nature conservation, forestry, use of forest products, taxonomy and forest management. Our results suggest a relatively dispersed paper distribution, where 19.6 %

represents the five most influential journals. Other papers were distributed between one or two publications per journal.

The most productive author on *Inga* was Barbedo (on plant physiology) with 12 papers, followed by Leblanc (plant breeding) and Rogez (interactions of phenolic compounds) with nine papers each, Nygren (ecophysiology) with eight papers and Holl (restoration ecology) with six papers. On the other hand, the five most representative authors (those who have published more than six papers) mainly were focused on C1, the most representative in this study with 178 papers (Fig. 2). In this class, we found several research lines in ecological approaches, such as the floristic composition and phytosociological structure of a riparian forest in Southern Brazil, where *Inga marginata* was one of the most important species in the sample (Nakajima, Soares-Silva, Medri, Goldenberg, & Correa, 1996). Another study evaluated *Inga edulis* in processes of increasing soil fertility, highlighting its positive




Fig. 2. Distribution of published papers per study categories on *Inga* species distributed in Santa Catarina, southern Brazil. C1 - Ecological; C2 - Morphology, anatomy, histology, taxonomy physiology and genetics; C3 - Production and use; C4 - Biochemical and nutritional properties.

Fig. 2. Distribución de artículos publicados por categoría de estudio sobre especies de *Inga* distribuidas en Santa Catarina, sur de Brasil. C1 - Ecológico; C2 - Morfología, anatomía, histología, taxonomía, fisiología y genética; C3 - Producción y uso; C4 - Propiedades bioquímicas y nutricionales.

performance (Kanmegne, Bayomock, Duguma, & Ladipo, 2000). Also, some studies called out *Inga*'s ability to resist disturbances such as herbivory (Thomaz Heerd & Ferreira de Melo Junior, 2016) and herbicides (Cabral et al., 2017), besides facilitating regeneration of forests due to mycorrhizal associations (Iglesias, Salas, Leblanc, & Nygren, 2011; Nygren, Leblanc, Lu, & Luciano, 2013).

Conversely, C4 was the second place most researched field with 41 species, such as *Inga edulis*, *Inga vera* and *Inga marginata* (Fig. 2). The studies pointed out that Fabaceae is well known for the compounds of the secondary metabolism and their use in biological assays (Molares & Ladio, 2012). The majority of studies were published from 2010 onwards and they highlighted the active principles of *Inga* species (Lima et al., 2018). Research with methanolic extracts of the barks, leaves, and flowers of *Inga marginata* evidenced reduction of the severity of anthracnose in bean plantations to values below 35 % observed in the control group (Andrade Pinto, Souza, & Oliveira, 2010). Chemical studies showed phenolic compounds on the leaves of genus *Inga* used in traditional medicine as laxatives. On the other hand, another study shows that the leaves of *Inga edulis*, *I. marginata* and *I. laurina* presented high antioxidant capacity, probably related to the presence of phenolic substances (Lima et al., 2018).

These types of studies are important precursors for investigations on biological activities (Antunes, 2018). For instance, indigenous communities use *Inga* species for inflammation and rheumatism as reported in an ethnobotanical study (Lima et al., 2018). Besides that, it is a much appreciated genus in Northern Brazil as food, whereby it is important to know their nutritional values. Those studies have shown that some of its secondary compounds have a high content of phenolic compounds, elevating its pharmacological potential, especially its antioxidant capacity (Pompeu, Rogez, Monteiro, Tinti, & Carvalho, 2012). *Inga edulis*, particularly, exhibited a high content of phenolic compounds and high antioxidant

capacity, capable of having antiulcerogenic activity (Pompeu et al., 2012). *I. vera*, on the other hand, provides a defense strategy – with peptidase inhibitors - that interferes in the physiology of insects. This mechanism has been considered as a biotechnological alternative for the control of pests (Bezerra, Oliveira, & Macedo, 2017).

We also registered eight papers on the C2 category (Fig. 2). In general, it is an underrepresented class; however, it is very relevant for the research line, particularly taxonomic studies (Rapini, 2004). Other papers investigated germination capacities, its break of dormancy and seed viability (Parisi, Biagi, Medina, & Barbedo, 2016; Delgado, Da Silva, & da Silva, 2017). On the other hand, C3 was the least representative class of our sample, with only five published papers (Fig. 2). All of them focused on timber and food uses (Clement, Cristo-Araújo, Eeckenbrugge, Pereira, & Picanço-Rodrigues, 2010; Lopes, Crepaldi, & Lobão, 2017; de Freitas et al., 2018).

The species *Inga edulis*, *I. vera* and *I. marginata* were the most cited species in our study, mainly in C1 class (Table 3). *Inga edulis* is a tree naturally distributed in tropical America, largely known by its wood quality and used, in Brazil, for food and shading of coffee plantations (Burkart, 1979). The strong ecological interactions, highlighted in our results, were related to animals, especially birds (Lindell, Reid, & Cole, 2013). Besides that, some studies emphasize the dispersers preference for *I. edulis* than other tree species in the same environment (Lindell et al., 2013). In addition, it shows antioxidant activities with moderate biological activities (Souza et al., 2007; De Freitas et al., 2018).

Studies on *I. vera* followed the same pattern of *I. edulis*: it stands out as a preferential diet for birds (Ragusa-Netto, 2004), and other ecological interactions. Patterns between morphological parameters of birds and flowers of *Inga* suggests a close mutualistic and complex relationship of pollination and environment (Githiru, Lens, Bennur, & Ogol, 2002; Cruz-Neto, Machado, Duarte Jr, & Lopes,

TABLE 3
Number of papers registered per species and study categories

TABLA 3
Número de artículos registrados por especie y categorías de estudio

Species	C1	C2	C3	C4
<i>Inga edulis</i> Mart.	86	2	3	31
<i>Inga edwallii</i> (Harms) T.D.Penn.	0	0	0	0
<i>Inga lentiscifolia</i> Benth.	1	0	0	0
<i>Inga marginata</i> Willd.	22	0	0	4
<i>Inga sellowiana</i> Benth.	0	0	0	0
<i>Inga sessilis</i> (Vell.) Mart.	3	0	0	0
<i>Inga striata</i> Benth.	5	0	0	0
<i>Inga subnuda</i> Salzm. ex Benth.	8	0	1	1
<i>Inga subnuda</i> subsp. <i>luschnathiana</i> (Benth.) T.D.Penn.	2	0	0	0
<i>Inga vera</i> Willd.	45	4	1	5
<i>Inga vera</i> subsp. <i>affinis</i> (DC.) T.D.Penn.	5	2	0	0
<i>Inga virescens</i> Benth.	1	0	0	0
<i>Inga vulpina</i> Mart. ex Benth.	0	0	0	0
Total	178	8	5	41

2011). *I. marginata* also stood out in ecological approaches, exhibiting potential as a facilitator of natural regeneration in the Atlantic Forest, with the highest number of species growing under its individuals (Spadeto, Wilson Fernandes, Negreiros, & Kunz, 2017) as well as *I. edulis* and *I. vera*. Besides that, *I. marginata* has potential for phytoremediation, being used for ecological restoration, especially in environments contaminated by herbicides (Cabral et al., 2017).

Finally, few investigations have been conducted on *I. lentiscifolia*, *I. sessilis*, *I. striata*, *I. subnuda*, *I. subnuda* subsp. *luschnathiana*, *I. vera* subsp. *affinis*, *I. virescens*, while no published paper were found for *I. edwallii*, *I. sellowiana* and *I. vulpina*. These species are more restrict, occurring in a few places in Santa Catarina. In this context, the bibliometric analysis showed an increase in the number of papers on *Inga* over time, especially in the last 13 years. However, some important gaps need to be addressed, such as the relatively small number and/or lack of studies conducted

for some species. In addition, we expect that more researches will be performed with *Inga*, emphasizing the less studied categories and species to reveal the real potential of this genus in Santa Catarina.

ACKNOWLEDGMENTS

We acknowledge financial support from the Coordination for the Improvement of Higher Education Personnel (FAPESC/CAPEs) for the first author's scholarship funding and the Universidade do Extremo Sul Catarinense (UNESC) for infrastructure.

RESUMEN

Literatura científica sobre *Inga* (Fabaceae) en el estado de Santa Catarina al sur de Brasil (1983-2017). El género *Inga* Mill. pertenece al clado mimosoide (Fabaceae, Caesalpinioideae) con 131 especies en Brasil. Es el género más importante de las Fabáceas. En este sentido, el objetivo de este estudio fue realizar un análisis bibliométrico de *Inga* en el Estado de Santa Catarina. Se condujo un estudio de la literatura publicada utilizando las bases de

datos electrónicas de la Web of Science, Scopus y SciELO con los nombres aceptados de las especies *Inga* y sus sinónimos. Los trabajos se distribuyeron en cuatro categorías temáticas: C1 (ecológico), C2 (morfoloía, anatomía, taxonomía, histología, fisiología y genética), C3 (producción y uso) y C4 (propiedades bioquímicas y nutricionales). Se registraron 232 trabajos para 13 especies de *Inga*, donde se exhibió un notable incremento de publicaciones. C1 fue la categoría temática más estudiada, principalmente en temas tales como: suministro de nutrientes, sombra y capacidad de fijación de nitrógeno. *Inga edulis*, *I. vera* e *I. marginata* fueron las especies más registradas en nuestro estudio. Nuestros resultados mostraron un aumento en el número de artículos sobre *Inga* con el tiempo, especialmente en los últimos 13 años. Sin embargo, es necesario abordar algunos vacíos importantes como el número relativamente pequeño y/o la falta de estudios realizados para algunas especies.

Palabras clave: bibliometría, *Inga edulis*, legumbres, Ingeae.

REFERENCES

- Alegre, J. C., & Rao, M. R. (1996). Soil and water conservation by contour hedging in the humid tropics of Peru. *Agriculture, Ecosystems & Environment*, 57(1), 17-25. DOI: [https://doi.org/10.1016/0167-8809\(95\)01012-2](https://doi.org/10.1016/0167-8809(95)01012-2)
- Andrade Pinto, J. M., Souza, E. A., & Oliveira, D. F. (2010). Use of plant extracts in the control of common bean anthracnose. *Crop Protection*, 29(8), 838-842. DOI: <https://doi.org/10.1016/j.cropro.2010.03.006>
- Alvares, C. A., Stape, J. L., Sentelhas, P. C., Gonçalves, J. L. M., & Sparovek, G. (2013). Köppen's climate classification map for Brazil. *Meteorologische Zeitschrift*, 22(6), 711-728. DOI: <https://doi.org/10.1127/0941-2948/2013/0507>
- Antunes, A. R., & Citadini-Zanette, V. (2018). *Erythrina falcata* Benth. (Fabaceae): Estudo Etnobotânico, Fitoquímico e Biológico. *Revista Brasileira de Agroecologia*, 13(14), 192-193.
- Arévalo-Pinedo, A., Dos Santos, F. L., Salles Arévalo, Z. D., Zuniga, A. D. G., & Pinedo, R. A. (2006). Description isotherms for murici (*Byrsonima sericea*) and *Inga* (*Inga edulis*) pulps. *Journal of Food Engineering*, 76(4), 611-615. DOI: <https://doi.org/10.1016/j.jfoodeng.2005.06.011>
- Ávila Jr., R., Pinheiro, M., & Sazima, M. (2015). The generalist *Inga subnuda* subsp. *luschnathiana* (Fabaceae): negative effect of floral visitors on reproductive success? *Plant Biology*, 17(3), 728-733. DOI: <https://doi.org/10.1111/plb.12291>
- Bezerra, C. D. S., Oliveira, C. T., & Macedo, M. L. R. (2017). *Inga vera* trypsin inhibitor interferes in the proteolytic activity and nutritional physiology of *Ephestia kuehniella* larvae. *Entomologia Experimentalis et Applicata*, 165(2-3), 109-119. DOI: <https://doi.org/10.1111/eea.12605>
- Bishop, J. P. (1983). Tropical forest sheep on legume forage/fuelwood fallows. *Agroforestry Systems*, 1(2), 79-84. DOI: <https://doi.org/10.1007/BF00596350>
- Bruneau, A., Doyle, J. J., Herendeen, P., Hughes, C., Kenicer, G., Lewis, G., ... Pennington, T. (2013). Legume phylogeny and classification in the 21st century: Progress, prospects and lessons for other species-rich clades. *Tiina Särkinen TAXON*, 6281921(622), 217-248.
- Burkart, A. (1979). *Leguminosas Mimosoideas*. Itajaí, Brasil: Flora Ilustrada Catarinense.
- Cabral, C. M., dos Santos, J. B., Ferreira, E. A., Machado, V. M., Pereira, I. M., Silva, D. V., & Souza, M. F. (2017). Tolerance to the herbicide clomazone and potential for changes of forest species. *Bioscience Journal*, 33(4), 897-904. DOI: <https://doi.org/10.14393/BJ-v33n4a2017-36750>
- Clement, C. R., de Cristo-Araújo, M., d'Eeckenbrugge, G. C., Pereira, A. A., & Picanço-Rodrigues, D. (2010). Origin and domestication of native Amazonian crops. *Diversity*, 2(1), 72-106. DOI: <https://doi.org/10.3390/d201007210.1371/journal.pone.0003311>
- Cruz-Neto, O., Machado, I. C., Duarte Jr., J. A., & Lopes, A. V. (2011). Synchronous phenology of hawkmoths (Sphingidae) and *Inga* species (Fabaceae-Mimosoideae): Implications for the restoration of the Atlantic forest of northeastern Brazil. *Biodiversity and Conservation*, 20(4), 751-765. DOI: <https://doi.org/10.1007/s10531-010-9975-x>
- Da Silva, K., De Meyer, S. E., Rouws, L. F., Farias, E. N., dos Santos, M. A., O'Hara, G., ... Zilli, J. E. (2014). *Bradyrhizobium ingae* sp. nov., isolated from effective nodules of *Inga laurina* grown in Cerrado soil. *International Journal of Systematic and Evolutionary Microbiology*, 64, 3395-3401. DOI: <https://doi.org/10.1099/ijs.0.063727-0>
- Da Silva Possette, R. F., & Rodrigues, W. A. (2010). O gênero *Inga* Mill. (Leguminosae - Mimosoideae) no estado do Paraná, Brasil. *Acta Botanica Brasileira*, 24(2), 354-368. DOI: <https://doi.org/10.1590/S0102-33062010000200006>
- De Freitas, F. A., Araujo, R. C., Soares, E. R., Nunomura, R. C. S., da Silva, F. M. A., da Silva, S. R. S., ... Koolen, H. H. F. (2018). Biological evaluation and quantitative analysis of antioxidant compounds in pulps of the Amazonian fruits bacuri (*Platonia insignis* Mart.), inga (*Inga edulis* Mart.), and uchi (*Sacoglottis uchi* Huber) by UHPLC-ESI-MS/MS.

- Journal of Food Biochemistry*, 42(1). DOI: <https://doi.org/10.1111/jfbc.12455>
- De Freitas, T. A., Franca, M. G. C., de Almeida, A. A. F., de Oliveira, S. J. R., de Jesus, R. M., Souza, V. L., ... Mangabeira, P. A. (2015). Morphology, ultrastructure and mineral uptake is affected by copper toxicity in young plants of *Inga subnuda* subs. *luschnathiana* (Benth.) TD Penn. *Environmental Science and Pollution Research*, 22(20), 15479-15494. DOI: <https://doi.org/10.1007/s11356-015-4610-8>
- De Sousa Dias, A. L., De Souza, J. N. S., & Rogez, H. (2010). Purification of phenolic compounds from *Inga edulis* leaves using solid-phase extraction: major compounds quantification and antioxidant capacity evaluation. *Quimica Nova*, 33(1), 38-42.
- De Vasconcelos, G. C. L. (2014). *A tribo Ingeae Benth. (Mimosoideae, Leguminosae) no Estado da Paraíba* (Master's thesis). Universidade Federal de Viçosa, Minas Gerais, Brazil.
- Delgado, L. G. M., Da Silva, R. B. G., & da Silva, M. R. (2017). Morphological quality of *Inga vera* seedlings under different water managements. *IRRIGA*, 22(3), 420-429. DOI: <https://doi.org/10.15809/irriga.2017v22n3p420-429>
- Dos Santos Barros, S. V. do Nascimento, C. C., & de Azevedo, C. P. (2012). Energetic characterization of native and exotic forest species cultivated at Amazonas. *Floresta*, 42(4), 725-732.
- Elias, G. A., Corrêa, P., Citadini-Zanette, V., & Santos, R. (2015). Arecaceae: Análise bibliométrica das espécies nativas do estado de Santa Catarina. *Ciência e Natura*, 37(1), 85-92. DOI: <https://doi.org/10.5902/2179460X15399>
- Faria, J. M. R., Davide, L. C., Da Silva, E. A. A., Davide, A. C., Pereira, R. C., van Lammeren, A. A. M., & Hilhorst, H. W. M. (2006). Physiological and cytological aspects of *Inga vera* subsp. *affinis* embryos during storage. *Brazilian Journal of Plant Physiology*, 18(4), 503-513. DOI: <https://doi.org/10.1590/S1677-04202006000400008>
- Fernandes, J. M., Dondoni Da Costa, R., Araújo, C. R., & Lopes, S. (2016). Taxonomia de *Inga macrophylla* Humb. & Bonpl. ex Willd. (Leguminosae, Mimosoideae): Uma Nova Ocorrência Para Mato Grosso, Brasil. *Enciclopédia Biosfera*, 13(24), 1329-1335. DOI: https://doi.org/10.18677/EnciBio_2016B_123
- Fink, R. D., Lindell, C. A., Morrison, E. B., Zahawi, R. A., & Holl, K. D. (2009). Patch size and tree species influence the number and duration of bird visits in forest restoration plots in southern Costa Rica. *Restoration Ecology*, 17(4), 479-486. DOI: <https://doi.org/10.1111/j.1526-100X.2008.00383.x>
- Flora do Brasil 2020. (2019). Jardim Botânico do Rio de Janeiro. Recuperado de <http://floradobrasil.jbrj.gov.br>
- Gepts, P., Beavis, W. D., Brummer, E. C., Shoemaker, R. C., Stalker, H. T., Weeden, N. F., & Young, N. D. (2005). Legumes as a model plant family. Genomics for food and feed report of the Cross-Legume Advances Through Genomics Conference. *Plant Physiology*, 137(4), 1228-1235. DOI: <https://doi.org/10.1104/pp.105.060871>
- Githiru, M., Lens, L., Bennur, L. A., & Ogot, C. P. K. O. (2002). Effects of site and fruit size on the composition of avian frugivore assemblages in a fragmented afro-tropical forest. *Oikos*, 96(2), 320-330. DOI: <https://doi.org/10.1034/j.1600-0706.2002.960214.x>
- Guillermo-Ferreira, R., Cardoso-Leite, R., & Gandolfo, R. (2012). First observation of alternative food usage (extrafloral nectar) by the assassin bug *Atopozelus opsimus* (Hemiptera, Reduviidae). *Revista Brasileira de Entomologia*, 56(4), 489-491. DOI: <https://doi.org/10.1590/S0085-56262012000400001410.1111/j.1463-6395.2011.00522.x>
- Hood, W. W., & Wilson, C. S. (2001). The literature of bibliometrics, scientometrics, and informetrics. *Scientometrics*, 52(2), 291-314. DOI: <https://doi.org/10.1023/A:1017919924342>
- Iglesias, L., Salas, E., Leblanc, H. A., & Nygren, P. (2011). Response of *Theobroma cacao* and *Inga edulis* seedlings to cross-inoculated populations of arbuscular mycorrhizal fungi. *Agroforestry Systems*, 83(1), 63-73. DOI: <https://doi.org/10.1007/s10457-011-9400-9>
- Kanmegne, J., Bayomock, L. A., Duguma, B., & Ladipo, D. O. (2000). Screening of 18 agroforestry species for highly acid and aluminum toxic soils of the humid tropics. *Agroforestry Systems*, 49(1), 31-39. DOI: <https://doi.org/10.1023/A:1006334931018>
- Kettler, J. S. (1996). Fallow enrichment of a traditional slash/mulch system in southern Costa Rica: Comparisons of biomass production and crop yield. *Agroforestry Systems*, 35(2), 165-176. DOI: <https://doi.org/10.1007/BF00122777>
- Lamarca, E. V., & Barbedo, C. J. (2015). Desiccation sensitivity of embryos of *Inga vera* Willd. obtained from different environmental conditions. *Revista Arvore*, 39(6), 1083-1092. DOI: <https://doi.org/10.1590/0100-67622015000600011>
- Lima, N. M., Santos, V. N. C., & Porta, L. (2018). Quimiodiversidade, Bioatividade e Quimiosistemática do Gênero *Inga* (FABACEAE): Uma Breve Revisão. *Revista Virtual Química*, 10(3), 459-473.
- Lindell, C. A., Reid, J. L., & Cole, R. J. (2013). Planting design effects on avian seed dispersers in a tropical forest restoration experiment.

- Restoration Ecology*, 21(4), 515-522. DOI: <https://doi.org/10.1111/j.1526-100X.2012.00905.x>
- Lopes, L. C. M., Crepaldi, M. O. S., & Lobão, A. Q. (2017). Useful woody species and its environmental availability: the case of artisanal fishermen in Itaúnas, Brazil. *Acta Scientiarum - Biological Sciences*, 39(2), 227-234. DOI: <https://doi.org/10.4025/actasciobiolsci.v39i2.33187>
- LPWG. (2017). A new subfamily classification of the Leguminosae based on a taxonomically comprehensive phylogeny – The Legume Phylogeny Working Group (LPWG). *Taxon*, 66(1), 44-77. DOI: <https://doi.org/10.12705/661.3>
- Machado, S. W., de Oliveira, C. F. R., Zério, N. G., Parra, J. R. P., & Macedo, M. L. R. (2017). *Inga laurina* trypsin inhibitor (ILTI) obstructs *Spodoptera frugiperda* trypsins expressed during adaptive mechanisms against plant protease inhibitors. *Archives of Insect Biochemistry and Physiology*, 95(4). DOI: <https://doi.org/10.1002/arch.21393>
- Molares, S., & Ladio, A. (2012). The usefulness of edible and medicinal fabaceae in argentine and chilean patagonia: environmental availability and other sources of supply. *Evidence-Based Complementary and Alternative Medicine*, 2012, 1-12. DOI: <https://doi.org/10.1155/2012/901918>
- Nakajima, J. N., Soares-Silva, L. H., Medri, M. E., Goldenberg, R., & Correa, G. T. (1996). Floristic composition and phytosociological structure in a riparian forest of Tibagi River, Telêmaco Borba, Paraná, Brazil. *Arquivos de Biologia E Tecnologia*, 39(4), 933-948.
- Nimer, E. (1990). Climatologia da região sul. In E. Nimer (Ed.), *Climatologia do Brasil* (pp. 3-65). Rio de Janeiro, Brasil: IBGE/Departamento de Recursos Naturais e Estudos Ambientais.
- Nygren, P., & Leblanc, H. A. (2015). Dinitrogen fixation by legume shade trees and direct transfer of fixed N to associated cacao in a tropical agroforestry system. *Tree Physiology*, 35(2), 134-147. DOI: <https://doi.org/10.1093/treephys/tpu116>
- Nygren, P., Leblanc, H. A., Lu, M. E., & Luciano, C. A. G. (2013). Distribution of coarse and fine roots of *Theobroma cacao* and shade tree *Inga edulis* in a cocoa plantation. *Annals of Forest Science*, 70(3), 229-239. DOI: <https://doi.org/10.1007/s13595-012-0250-z>
- Parisi, J. J. D., Biagi, J. D., Medina, P. F., & Barbedo, C. J. (2016). Fungicide and drying effects on the viability of recalcitrant seeds of *Inga vera* subsp. *affinis*. *Tropical Plant Pathology*, 41(3), 177-182. DOI: <https://doi.org/10.1007/s40858-016-0084-6>
- Pennington, T. (1997). *The Genus Inga*. Richmond, UK: Royal Botanic Gardens Kew.
- Pinto, J. M. A., Souza, E. A., & Oliveira, D. F. (2010). Use of plant extracts in the control of common bean anthracnose. *Crop Protection*, 29(8), 838-842. DOI: <https://doi.org/10.1016/j.cropro.2010.03.006>
- Pompeu, D. R., Rogez, H., Monteiro, K. M., Tinti, S. V., & Carvalho, J. E. (2012). Antioxidant capacity and pharmacologic screening of crude extracts of *Byrsonima crassifolia* and *Inga edulis* leaves. *Acta Amazonica*, 42(1), 165-172.
- Ragusa-Netto, J. (2004). Flowers, fruits, and the abundance of the yellow-chevroned parakeet (*Brotogeris chiriri*) at a gallery forest in the South Pantanal (Brazil). *Revista Brasileira de Biologia*, 64(4), 867-877.
- Rapini, A. (2004). Modernizando a taxonomia. *Biota Neotropica*, 4(1), 1-4. DOI: <https://doi.org/10.1590/S1676-06032004000100002>
- Reitz, R. (1965). Plano de coleção. In R. Reitz (Ed.), *Flora Ilustrada Catarinense* (pp. 1-71). Itajaí, Brasil: Herbario Barbosa Rodrigues.
- Souza, J. N. S., Silva, E. M., da Silva, M. N., Arruda, M. S. R., Larondelle, Y., & Rogez, H. (2007). Identification and antioxidant activity of several flavonoids of *Inga edulis* leaves. *Journal of the Brazilian Chemical Society*, 18(6), 1276-1280. DOI: <https://doi.org/10.1590/s0103-50532007000600025>
- Spadeto, C., Wilson Fernandes, G., Negreiros, D., & Kunz, S. H. (2017). Facilitative effects of tree species on natural regeneration in an endangered biodiversity hotspot. *Revista Brasileira de Botânica*, 40(4), 943-950. DOI: <https://doi.org/10.1007/s40415-017-0408-x>
- Staver, C. (1989). Shortened bush fallow rotations with relay-cropped *Inga edulis* and *Desmodium ovalifolium* in wet central Amazonian Peru. *Agroforestry Systems*, 8(2), 173-196. DOI: <https://doi.org/10.1007/BF00123120>
- Tauchen, J., Bortl, L., Huml, L., Miksatkova, P., Doskocil, I., Marsik, P., ... Kokoska, L. (2016). Phenolic composition, antioxidant and anti-proliferative activities of edible and medicinal plants from the Peruvian Amazon. *Revista Brasileira de Farmacognosia*, 26(6), 728-737. DOI: <https://doi.org/10.1016/j.bjp.2016.03.016>
- Thomaz Heerd, S., & Ferreira de Melo Junior, J. C. (2016). Strategies of defense and level of herbivory in strata of the crown of *Inga edulis* Mart. (Fabaceae) in a remnant urban forest. *Acta Botanica Venezuelica*, 39(1), 101-117.
- Tropicos. (2013). Missouri Botanical Garden. Recuperado de www.tropicos.org.
- Vibrans, A. C., Sevegnani, L., Gasper, A. L., & Lingner, D. V. (2012a). *Inventário florístico florestal de Santa*

- Catarina: diversidade e conservação dos remanescentes florestais*. Blumenau, Brasil: Edifurb.
- Vibrans, A. C., Sevegnani, L., Gasper, A. L., & Lingner, D. V. (2012b). *Inventário florístico florestal de Santa Catarina: Floresta Estacional Decidual*. Blumenau, Brasil: Edifurb.
- Vibrans, A. C., Sevegnani, L., Gasper, A. L., Müller, J. J. V., & Reis, M. S. (2013a). *Inventário florístico florestal de Santa Catarina: resultados resumidos*. Blumenau, Brasil: Edifurb.
- Vibrans, A. C., Sevegnani, L., Gasper, A. L., & Lingner, D. V. (2013b). *Inventário florístico florestal de Santa Catarina: Floresta Ombrófila Mista*. Blumenau, Brasil: Edifurb.
- Vibrans, A. C., Sevegnani, L., Gasper, A. L., & Lingner, D. V. (2013c). *Inventário florístico florestal de Santa Catarina: Floresta Ombrófila Densa*. Blumenau, Brasil: Edifurb.
- Yatazawa, M., Uchino, F., & Hambali, G. G. (1983). Nitrogen fixing activity in warty lenticellate tree barks. *Soil Science and Plant Nutrition*, 29(3), 285-294.