



Revista de Biología Tropical

ISSN: 0034-7744

ISSN: 2215-2075

Universidad de Costa Rica

Malaviya, Devendra Ram; Baig, Mirza Jaynul; Kumar, Bijendra; Kaushal, Pankaj  
Effects of shade on guinea grass genotypes *Megathyrsus maximus* (Poales: Poaceae)  
Revista de Biología Tropical, vol. 68, no. 2, 2020, April-June, pp. 563-572  
Universidad de Costa Rica

DOI: <https://doi.org/10.15517/RBT.V68I2.38362>

Available in: <https://www.redalyc.org/articulo.oa?id=44965969016>

- How to cite
- Complete issue
- More information about this article
- Journal's webpage in redalyc.org

UDEM  redalyc.org

Scientific Information System Redalyc  
Network of Scientific Journals from Latin America and the Caribbean, Spain and  
Portugal

Project academic non-profit, developed under the open access initiative

## Effects of shade on guinea grass genotypes *Megathyrus maximus* (Poales: Poaceae)

Devendra Ram Malaviya<sup>1\*</sup>, Mirza Jaynul Baig<sup>2</sup>, Bijendra Kumar<sup>3</sup> & Pankaj Kaushal<sup>4</sup>

1. ICAR-Indian Grassland and Fodder Research Institute, Jhansi-284003, India, \*\*ICAR-Indian Institute of Sugarcane Research, Lucknow-226002, India; drmalaviya47@rediffmail.com
  2. National Rice Research Institute (ICAR), Cuttack, 753006, India; mjbaigerri@gmail.com
  3. Indian Grassland and Fodder Research Institute (ICAR), Jhansi, 284003, India; ujlayanbijendrakumar71@gmail.com
  4. National Institute of Biotic Stress Management (ICAR), Raipur, 493225, India; pkaushal70@gmail.com
- \* Correspondence. \*\*present address.

Received 16-VII-2019.

Corrected 13-I-2020.

Accepted 17-III-2020.

**ABSTRACT. Introduction:** Light stress is an important factor limiting the biomass yield while combining forage production with crops or forestry. Guinea grass is a widely adapted perennial fodder grass. The species exhibits high degree of variation for morphology, adaptation and biomass yield. **Objective:** Since there is a need in identifying shade adapted forage grasses for the expanding area under agroforestry/silvipastures, the present investigation took the task of understanding how the morphologically distinct genotypes of guinea grass respond under different shaded intensities. **Methods:** In the present study, forty-four genotypes related with the shade response were studied in varying shading conditions (pure sunlight, 25, 50 and 75 % shade) created artificially. **Results:** Based on green and dry matter yields ranking, the genotype IG 01-98 performed the best followed by genotypes IG 01-92, IG 97-5, IG 97-6 and IG 01-89 in decreasing order. Particularly, IG 01-93 was identified as the best performing under 50 % shading conditions. On the other hand, most of the top ranking genotypes performed well both under open and up to 50 % of shade. Morphologically, these genotypes were taller and possessed longer and broader leaves. Under shaded conditions (over 50 %), leaf length and width showed an increasing trend compared to open conditions. Also, chlorophyll content increased with shading intensity. **Conclusions:** Most of the genotypes collected from the southern Indian humid tropical environment with early flowering nature were tolerant to shade. Differential genotypic response was observed for biomass yield and yield attributes under shade. The study established appreciable variability for shade tolerance among genotypes.

**Key words:** biomass, genetic diversity, leaf length, leaf width, plant height, shade tolerance.

Ram Malaviya, D., Jaynul Baig, M., Kumar, B., & Kaushal, P. (2020). Effects of shade on guinea grass genotypes *Megathyrus maximus* (Poales: Poaceae). *Revista de Biología Tropical*, 68(2), 563-572.

Among various abiotic stresses, light stress is emerging as one of the important factors that limits the biomass yield. This is important in the context of shrinking land area under cultivation and increasing demand for cropping systems to enhance productivity per unit area. Combining pastures with plantation crops or with forest trees is likely to increase the availability of forage and use the land more efficiently. The area under agro-forestry and silvipasture systems is

fast increasing in India and around the world. The extent of agroforestry in India was estimated to be 14.46 million ha (excluding fallow land), whereas the potential area under agroforestry was estimated to be about 17.45 million ha (Newaj et al., 2017). *Panicum maximum* cultivars, particularly Petrie (Green Panic) and Embu (Creeping Guinea), have been used for grazing under coconuts (Stfir & Shelton, 1991; Plucknett, 1979). Shading intensity

varies significantly in various agro-forestry, silvipasture, and social forestry systems. Shade intensity also varies according to the age of plantation. Hence, suitable understory crop for varying degree of shade tolerance is required.

Among many perennial grasses, *Megathyrsus maximus* (guinea grass), formerly known as *Panicum maximum* Jacq., has proven to be a high biomass producing grass in agroforestry and silvipasture systems. Singh and Chaturvedi (2011) identified the *Alnus nepalensis*, *Stylosanthesis guyensis* and *P. maximum* system as the best for high rainfall hilly regions with the potential of 13.5 t<sup>ha</sup><sup>-1</sup>(DM) fodder per year. The grass is known to be tolerant to shading conditions (Wong & Wilson, 1980; Kaushal, Malaviya, & Singh, 2000) and is recommended for cultivation in agroforestry and forestry systems.

This grass is a widely adapted perennial grass for utilization as fodder because of its ease of propagation, fast re-growth, high biomass and nutritional quality. Under rain-fed condition (400-1 500 mm rainfall per year), it yields 40-60 t/ha dry matter with over 14 % crude protein content and 41-72 % dry matter digestibility (Bogdan, 1977; Sukhchain & Sidhu, 1992). *In vitro* dry matter digestibility of organic matter from first harvest to third harvest is reported to vary from 56.9 to 87.7 % among different varieties (Aganga & Tshwenyane, 2004). It is also responsive to higher input conditions and also suitable for cultivated/irrigated conditions. The species is highly responsive to fertilizers, which results in an increased number of cuts of the grass (Pereira et al., 2012). Despite the latter information and being considered to be shade tolerant, some studies had reported different results. For instance, in some of the studies, shading decreased the yield in guinea grass with a significant increase in the N content under shade (Norton, Wilson, Shelton, & Hiu, 1991; Deinum et al., 1996). On the other hand, Lowry, Lowry and Jones (1988) reported an increase up to 250 %. Hence, there is particular need of studies that take into account the available

genetic diversity to identify what genotype is more for shaded condition.

The crops' primary gene pool exhibits high degree of variation for several traits including morphology, adaptation and biomass yield. Therefore, the morphological, biochemical, reproductive and physiological parameters were characterized and the distinct morphotypes were identified in the sets of germplasm used at the Indian Grassland and Fodder Research Institute in Jhansi, India (Jain, Roy, Kaushal, Malaviya, & Zadoo, 2003, 2006; Kaushal, Malaviya, & Singh, 1999; Kaushal et al., 2008; Malaviya, 1996, 1998). This gene pool was considered ideal for studies that evaluate this factor under abiotic stress because such diverse lines may respond differently to the stress condition. Hence, in view of the expanding area under agroforestry/silvipastures and the necessity of identifying the shade adapted forage grasses, the present investigation tried to understand how the morphologically distinct genotypes of guinea grass respond to various shaded environments. Additionally, the correlation between morphology and the biomass yield under abiotic stress was also envisaged to be established.

## MATERIALS AND METHODS

**Site of study:** The analysis was carried out at the experimental farm of ICAR - Indian Grassland and Fodder Research Institute (IGFRI) in Jhansi, India (25.27° N & 78.35° E), 271 m.a.s.l. under rain-fed conditions. The annual rain-fall during the first and the second year of the experiment was 477 mm and 1 147 mm respectively. The rain-fall distribution was mainly during July to September. Minimum and maximum temperatures during the productive months (July to September) in the first and second year also ranged between 21-28 °C and 29-38 °C, respectively. The bright sunlight hours during the growth period (July, August and September) in the first year were 5.3, 4.0 and 8.0 hours/day, whereas during the second year they were 5.2, 5.1 and 2.7 hours/day, respectively. The sunlight intensity in open

conditions at 12 pm in September was observed to be 582.3  $\mu\text{mol}/\text{m}^2/\text{s}$ .

**Experimental design:** The experiment was conducted by planting the genotypes in open as and shaded conditions. Three shaded conditions were created by using a shade net which intercepted 25 %, 50 % and 75 % of the sunlight. Shade nets were put at a height of 2 m and the crop was also covered from the sides. A total of 44 genotypes representing the collections from Ethiopia and India were studied. This material included exotic genotypes from Ethiopia (IG 97-6, IG 97-5, IG 97-14, IG 96-240, IG 97-35), North India (IG 01-83, IG 01-134, IG 01-180, IG 01-183, IG 01-181, IG 01-187, IG 01-141), Central India (IG 01-207-1, IG 01-207-2, IG 01-207-3, IG 01-207-4, IG 01-204-1, IG 01-204-2, IG 01-200, IG 01-87) and South India (IG 01-155, IG 01-212, IG 01-171, IG 01-221, IG 01-205, IG 01-191, IG 01-152, IG 01-98, IG 01-96, IG 01-93, IG 01-92, IG 01-89, IG 01-206, IG 01-201, IG 01-185, IG 01-176, IG 01-220, IG 01-198, IG 01-197, IG 01-190, IG 01-160, IG 01-122, IG 01-116, IG 01-85). The seeds containing those genotypes were procured from the gene bank of IGFRI, Jhansi, India. Nursery of the genotypes was raised and six week old seedlings were transplanted.

**Raising of plants and observations:** Genotypes were transplanted in 3 x 1 m plots that had 3 m long paired rows, spaced 50 cm apart and accommodating six tussocks spaced at 50 cm in the row. The plants were harvested above ground at 15 cm of height once during the first year of establishment and thrice during the second year in the rainy season. For identification of shade tolerant genotypes, cumulative yield over four cuttings was considered. Dry matter percent, at each harvest, was estimated by drying 500 g green sample of each accession at 60 °C in hot air oven until obtaining a constant sample weight. Data on morphological observations such as plant height, leaf blade length and leaf blade width were recorded on three plants in each plot during the second year, just

before the second harvest. Plant height was recorded as the total length of the longest tiller from the ground to the top (the tip of the last emerged leaf). The leaf width and length were recorded on the third leaf from the top of the three randomly selected plants. The first and the last plants in the row were not considered for this data recording. Chlorophyll content among genotypes in ambient light condition and that at three shade levels was recorded on three plants in each plot using CCM-200 by placing the third leaf from the top between the sensor points provided in the instrument. This data was recorded between 9 to 11 am.

**Statistical analysis:** Paired t-test, quadratic regression and correlation analysis were performed following standard statistical methods using MS Excel program. Ranking of genotypes was done based on their score for green and dry biomass. The analysis of variance and the interaction effect was calculated following Kharrazi-Kopaei and Sadooghi-Alvandi (2007). The effect of ante-dependence was judged following Kenwood (1987). The correlation matrix and variance was generated on the residual basis. The t-test was performed on yield data within treatments over different cuttings and between the cumulative yield over the cuts with the yield in individual cuts. It was also performed between the two groups of the genotypes: the genotypes with an increase in yield up to 50 % shade and the remaining genotypes showing yield increase even at 75 % of shade.

## RESULTS

**Biomass production under open vis-à-vis shaded condition:** The total green forage yield (GFY), in four harvests for two years, under open conditions ranged from 14.9 to 77.5 MT/ha across the genotypes with a mean of 43.5 MT/ha. The mean green biomass increased to 73.8 MT/ha and 100.7 MT/ha under 25 % and 50 % shade, respectively, and declined to 91.8 MT/ha under 75 % shading (Digital Appendix 1, Fig. 1). The paired t-test revealed significant difference ( $< 0.01$  % level) only at 25 % and 50

% shading when comparing with the open and shading at 25 % respectively (Digital Appendix 1). Since the fresh and the dry biomass declined at 75 % shading, further correlations and the identification of genotypes was done considering yield in open vis-à-vis at 50 % shading conditions. There was a positive significant correlation between green biomass yield under open vis-à-vis at 50 % shading conditions. The quadratic trend shown in Fig. 1 and the equation in the Digital Appendix 1 also show the trend of decline at 75 % shade.


The dry matter yield (DMY), in the four harvests during two years, ranged from 3.9 to 19.46 MT/ha under ambient sunlight condition with a mean of 11.31 MT/ha across the genotypes. The mean dry matter production was 16.09 MT/ha, 20.42 MT/ha and 18.78 MT/ha, respectively, under the three shaded conditions (25 %, 50 % and 75 %) (Digital Appendix 1). However, there was no significant difference between the yields at 50 % shade and 75 % shade. The positive correlation was established between dry matter yields under open conditions with that under 50 % shaded condition. The quadratic trend shown in Fig. 1 and the equation in Digital Appendix 1 also show the trend of nominal decline at 75 % shade.

The interaction effect calculated ( $F = 2.78$ , d.f. = 78, 48) was highly significant ( $P = 0.000114$ ) (Digital Appendix 2). Since the

p-value was too low, the highly significant interaction effects for green forage yield among treatments and the genotypes were confirmed, as suggested by Kharrati-Kopaei and Sadooghi-Alvandi (2007). Similarly, for the dry forage yield, it was highly significant ( $F = 2.097$ ,  $P = 0.003$ , d.f. = 78, 48).

In the repeated observation, the relations were likely to be serially correlated, hence, the residual correlation matrix was generated from the yield data observed at different cuttings. In the present study, the correlation pattern was not the typical for repeated measurements over different cuttings in open and shaded conditions (Digital Appendix 3). Since they did not follow a trend, it suggests that there were other factors influencing the regeneration such as seasonal variation, its regeneration ability and the genotypic differences. Similar trends were also observed for the correlation of dry matter productivity.

The t-test between yield at different cuttings with their preceding cutting yield were found to be significant in open as well as shaded conditions. The trend was similar for the GFY and dry matter yield (DMY) (Digital Appendix 4). On the other hand, the t-test between the first and second cutting for green fodder yield (GFY) and that between the third and fourth cutting for DMY were the only values with non-significant differences.



**Fig. 1.** Yield and yield attributes of guinea grass grown under open and shaded conditions. (Lines connecting the data point are quadratic trend lines).

**Fig. 1.** Componentes de rendimiento y rendimiento del zacate guinea cultivados en condiciones abiertas y sombreadas (Las líneas que conectan el punto de datos son líneas de tendencia cuadráticas).

For the ante-dependence test, the genotypes were grouped on the basis of biomass yield: genotypes that showed decline in yield under 50 % shade and those that showed yield increase even at 75 % shade. The t-test showed non-significant differences between the groups in open, 25 % shade and 50 % shaded condition. However, at 75 % shade, significant differences between the groups were observed (Digital Appendix 5). Nevertheless, the increase of variance over time was typical of cumulative yield data over the cuttings, which indicated that the covariance structure was not stationary.

**Chlorophyll content and morphological adaptation in shaded condition:** The average plant height observed among genotypes in open condition was 122.3 cm, which increased to 154.3 cm and 172.0 cm respectively at 25 % and 50 % shading (Digital Appendix 1). Marginal numerical and non-significant increase of plant height to 174.0 cm was noticed at 75 % shading. The plant height showed positive correlation when the shading was increased. The average leaf length when the shading was increased to more than 50 %. There was no significant difference between leaf length at 50 % shade and 75 %. Additionally, the leaf length had a positive correlation and regression with increasing shade. The leaf width also had similar trends as the leaf length (Digital Appendix 1).

The genotypes grown under open and shaded conditions showed that the chlorophyll content, as indicated by CCI, increased with the shading intensity. Average CCI in the open environments was 20.9. It increased to 32.1, 38.7 and 45.4 respectively under 25 %, 50 % and 75 % shading conditions (Digital Appendix 1). On the other hand, there was no correlation between chlorophyll content in open vis-à-vis shaded conditions; however, there was linear increase in chlorophyll content with increasing shade.

## DISCUSSION

The mean green biomass increased under 25 and 50 % shading conditions, but declined under 75 % shading. However, there was no significant difference between yield at 50 and 75 % shade. Guinea grass under dense shade generally gives 27 % of light, according to previous studies, and have 11 % higher biomass than under full sunlight conditions (Eriksen & Whitney, 1981). In this investigation, the increase in yield under shade was significantly higher, which follows the results of Wong and Wilson (1980) who found a 27 % increase in shoot yield and a 76 % increase in nitrogen yield of 8-week cut green panic (*P. maximum* var. *trichoglume*) grown under 40 % of light transmission, compared to full sunlight conditions. On the other hand, there was an increase in yield up to 250 % in guinea grass under the shade of the canopy trees *Albizia lebbek*, compared to those in full sun outside the canopy, according to Lowry and collaborators (1988).

In regards to experiments conducted under artificially created shaded condition with a 50 % cloth, Wilson, Catchpole and Weier (1986) showed an increase in dry weight and nitrogen yield on green panic, similar to the one obtained in this study. Such an increase in biomass yield under shade is also reported for other tropical grasses. Wilson, Hill, Cameron and Shelton (1990) also reported a 35 % increase in growth for bahia grass (*Paspalum notatum*) under approximately 55 % light transmission within a plantation of *Eucalyptus grandis* trees, in addition to a 67 % increase in nitrogen yield of the herbage.

Most importantly, the genotypes that give high yield under open conditions were also more abundant under shaded condition, although it was not a consistent pattern. For instance, the genotypes IG 97-5 and IG 01-93, which ranked 8<sup>th</sup> and 15<sup>th</sup> respectively under open conditions, ranked first and second respectively



TABLE 1  
Top ranking genotypes based on their rank in green and dry matter yield in 50 % shade and open condition

TABLA 1  
Genotipos de alto rango basados en su rendimiento de materia verde y seca en 50 % de sombra y la condición abierta

Genotype	Rank in Green fodder yield		Rank in Dry matter yield		Total	Overall Rank
	open	50 % shade	open	50 % shade		
IG 01-98	3	3	2	3	11	1
IG 01-92	2	6	3	6	17	2
IG 97-5	8	1	8	1	18	3
IG 97-6	5	4	5	8	22	4
IG 01-89	4	13	4	10	31	5
IG01-220	1	17	1	17	36	6
IG 01-93	15	2	17	2	36	6
IG 01-155	6	15	10	11	42	7
IG 01-212	9	19	6	18	52	8
IG 01-187	16	12	11	13	52	8
IG 01-171	13	11	20	16	60	9
IG 01-116	7	24	7	26	64	10
IG 01-207-2	12	28	12	29	81	11
IG 01-96	11	30	16	25	82	12
IG 01-191	10	34	9	32	85	13
IG 01-221	14	38	21	37	110	14
IG 96-240	17	36	28	40	121	15

under 50 % shade (Table 1). The high value of the regression coefficient observed for green biomass yields under open and the three shaded conditions suggests the involvement of additional factors affecting its yield under different light intensities. The top three genotypes (IG 01-92, IG 01-98 and IG 97-5), on the basis of overall score, also performed well under 50 % shaded condition. Dry matter accumulation also increased significantly to more than 50 % shade and showed a positive correlation between dry matter yields under open conditions and those under shade.

The higher herbage yields obtained by Odedire and Babayemi (2008) for *P. maximum*, when cropped with *Leucaena leucocephala*, was attributed to the ability of the grass to withstand the shading effect, since the dry matter (DM) production of tropical forage grasses is influenced by photoperiods by stimulating floral induction (Taiz, Zeiger, Møller, & Murphy, 2015). This might be the reason why there is a lower yield in open conditions compared

to those in shaded conditions. The top three genotypes for dry matter yield under open conditions were identified as IG 01-220, IG 01-98 and IG 01-92, whereas, under 50 % shade condition, the top three genotypes were IG 97-5, IG 01-93 and IG 01-98. Thus, IG 01-98 performed well under open and shaded conditions.

In regards to the genotypic response to shaded conditions, the genotypes IG 97-5, IG 01-93 and IG 01-98 ranked first, second and third respectively for the green and dry biomass yield under 50 % shading conditions (Table 1). However, considering the overall ranking - based on the green and dry matter yield under open and shaded condition - IG 01-98 was the best followed by IG 01-92, IG 97-5, IG 97-6 and IG 01-89 in decreasing order. Of these genotypes, excepting IG 97-5 (which ranked 8<sup>th</sup> under open conditions), the rest of the genotypes were good both under open and 50 % shade conditions. This shows that the genotypes, irrespective of the percentage point reduction under shade, were a high yielder

under shade intensity over 50 %. In contrast, the genotype IG 01-93, which ranked 15th and 17th respectively for green and dry matter yield in open conditions, ranked second for both traits under shaded conditions. This shows that IG 01-93 recorded a minimum percentage point reduction under shaded conditions. Such genotypes can be treated as lines to be specifically grown under shades, in addition to their inclusion in breeding programmes for development of shade tolerant lines.

The microclimate created by the shade has a direct impact on the wind speed, temperature regime, humidity, evapo-transpiration and soil moisture levels. Hence, these conditions are likely to impact the biomass production of any crop growing under shade. If the light intensity is not reduced below the threshold level, other factors will increase biomass production. In the present study, green and dry biomass of many genotypes increased under shading. It appears that the less light intensity under shade was compensated with higher moisture availability compared to open conditions. Araujo, Santos, Rodriguez, and Pezzopane (2018) have also identified the water availability index and the day length as key seasonal factor for dry matter production in guinea grass. Increase in yield under moderate light reduction has also been reported by Samarakoon, Wilson and Shelton (1990). Genotypic variation in guinea grass, involving a few genotypes, for shade tolerance was observed in earlier reports also (Kaushal et al., 2000).

Most of the genotypes were responsible for the increase in the plant height under shade over 50 % compared to open conditions. It was observed that most of the genotypes, which ranked in the top five for yield parameters under open conditions and shade, were also characterized by higher values of plant height, leaf length and leaf width than the general mean for their respective traits (Table 1). Increase in plant height of guinea grass under shade was also reported by Anita and Lakshmi (2014). However, the genotype IG 01-207-2, which showed maximum plant height in open conditions, was associated with highly

reduced height under shade. On the other hand, the genotype IG 01-93, which ranked second for green forage yield (GFY) and dry matter yield (DMY) under shade, was not tall and exhibited further reduction in height under shade. However, as per visual observation, this genotype was marked for increased number of tillers under shade. Additionally, under shading conditions over 50 %, leaf length and leaf width showed an increasing trend. Leaves of the three genotypes among the top five ranking were longer and wider than most of the other genotypes, thus indicating that long and broad leaved genotypes may perform better in shaded environment.

Chlorophyll content is a crucial factor in plant growth. Non-destructive, *in situ*, optical techniques are widely used to provide a relative indication of leaf chlorophyll concentration (Parry, Blonquist, & Bugbee, 2014). We used CCM-200 for comparing chlorophyll content. It gives the ratio of transmission of radiation at 931 nm to that at 653 nm, which is defined as the chlorophyll content index (CCI) and requires chlorophyll prediction equations for getting chlorophyll content (Cericovic, Masdoumier, Ghazlen, & Latouche, 2012). However, as per requirement of the present study, CCI were directly used for comparison among genotypes as well as the change in CCI affected by shading conditions. According to our results, the average chlorophyll content increased with shading intensity, but there was no correlation between chlorophyll content in open vis-à-vis shaded conditions. The top ranking genotypes for yield parameters were not among the top ranking genotypes for CCI either. Also, no clear trend for increase or decrease of CCI in open vs shaded condition could be established as the genotypes behaved differently for this trait. Therefore, we were unable to make generalizations.

Among the top five efficient (better performing) genotypes, IG 01-98, IG 01-92, IG 01-89, IG 97-5 and IG 97-6, the first three belonged to collections from south Indian humid tropical environments, whereas the latter two were from an exotic collection from



Ethiopia. For instance, the genotype IG 01-93, which ranked second for green and dry matter yield under shaded conditions, also was in the collection from South India. This genotype possessed high shade tolerance because, in open conditions, it ranked 15th and 17th respectively for green and dry matter production. All these south Indian genotypes possessed early flowering nature. Thus, it appears that, with early flowering nature, they are more adapted to shading tolerance. Genotype IG01-220 was considered highly susceptible to shade because it ranked first in open conditions for dry and green matter production; however, they ranked 17th under 50 % shading conditions. Similarly, many other genotypes, which ranked moderately high in open conditions, showed a high reduction in yield under shade (Table 1).

In general, *P. maximum* is reported to have high degree of shade adaptability (Wong, Sharudin, & Rahim, 1985). The species shows linear increase in leaf area with increasing shade (Durr & Rangel, 2000). Our results also confirm that, with increasing shade intensity, leaf-length and leaf-width augmented, but with differential genotypic response. *P. maximum*'s genotypes represent C3-C4 intermediate types but some of them are close to C3, while others are close to C4 anatomy (our unpublished anatomical study on these genotypes). Hence, a differential response to shade is expected.

In conclusion, the study clearly established the trend of some accession of genotypes being more suitable under shade than other. Thus, genotypes IG 01-98, IG 01-92, IG 97-5, IG 97-6 and IG 01-89 were successfully identified suitable both under shaded and open conditions, whereas IG 01-93 was identified as suitable only in 50 % shade conditions. Morphological features such as tall nature, long and broad leaves were associated with shade tolerance. The study also established appreciable variability for shade tolerance among genotypes under study which can be utilized further in breeding programmes. The genotypes that were identified can be valuable genetic resources to be utilized in India, Africa and any other place with iso-climatic conditions.

**Ethical statement:** authors declare that they all agree with this publication and made significant contributions; that there is no conflict of interest of any kind; and that we followed all pertinent ethical and legal procedures and requirements. All financial sources are fully and clearly stated in the acknowledgements section. A signed document has been filed in the journal archives.

## ACKNOWLEDGMENTS

The authors are thankful with the Indian Council of Agricultural Research, New Delhi, India for financial support of the work under the National Agriculture Technology Project. Special thanks to Mrs. Madhavi Chaubey for helping in the statistical analysis.

## RESUMEN

**Efectos de la sombra sobre genotipos del zacate guinea *Megathyrsus maximus* (Poales: Poaceae).** **Introducción:** El estrés leve es un factor importante que limita el rendimiento de la biomasa al tiempo que combina la producción forrajera con los cultivos o la silvicultura. El zacate guinea es una planta de forraje perenne ampliamente adaptada. La especie presenta un alto grado de variación en su morfología, adaptación y rendimiento de biomasa. **Objetivo:** Como existe la necesidad de la identificación de pastos forrajeros adaptados a la sombra para el área de expansión bajo agroforestería /silvicultura, la presente investigación se realizó para entender cómo los genotipos morfológicamente distintos al zacate guinea responden a diferentes intensidades de sombra. **Métodos:** Se estudiaron 44 genotipos y su respuesta a los niveles de sombra: luz solar pura y 25, 50 y 75 % de sombra (creados artificialmente). **Resultados:** Basado en la clasificación de rendimientos de materia verde y seca, el genotipo IG 01-98 fue el mejor, seguido de los genotipos IG 01-92, IG 97-5, IG 97-6 e IG 01-89 en orden decreciente. Por su parte, el IG 01-93 fue identificado como el de mayor rendimiento bajo el 50 % de condición de sombra. La mayoría de los genotipos de primer nivel tuvieron un buen desempeño, tanto bajo sombra abierta como al 50 %. Morfológicamente, estos genotipos fueron más altos y poseían hojas más largas y más anchas. En condiciones de sombra, de más del 50 %, la longitud y ancho de la hoja mostraron una tendencia creciente en comparación con la condición abierta. El contenido de clorofila aumentó con la intensidad de la sombra. **Conclusiones:** La mayoría de los genotipos recolectados al sur de la India en un ambiente tropical húmedo y tienen una floración temprana, fueron tolerantes a la sombra. Se observó una respuesta genotípica diferencial para los atributos de rendimiento y rendimiento

de biomasa bajo sombra. El estudio mostró la existencia de una variabilidad apreciable para la tolerancia a la sombra entre los genotipos.

**Palabras clave:** biomasa, diversidad genética, altura de planta, longitud de la hoja, anchura de la hoja, tolerancia a la sombra.

## REFERENCES

- Aganga, A.A., & Tshwenyane, S. (2004). Potentials of Guinea Grass (*Panicum maximum*) as Forage Crop in Livestock Production. *Pakistan Journal of Nutrition*, 3, 1-4.
- Anita, M.R., & Lakshmi, S. (2014). Growth characters of guinea grass (*Panicum maximum* Jacq.) under varying light intensity and potassium. In A.B. Khan (Ed.), *e-Proceedings of International Conference on "Green India: Strategic Knowledge for Combating Climate Change: Prospects & Challenges"* (pp. 338-342). Pondicherry, India: Excel India Publications.
- Araujo, L.C., Santos, P.M., Rodriguez, D., & Pezzopane, J.R.M. (2018). Key factors that influence for seasonal production of Guinea grass. *Scientia Agricola*, 75, 191-196. DOI: <http://dx.doi.org/10.1590/1678-992X-2016-0413>
- Bogdan, A.B. (1977). *Tropical Pastures and Fodder Plants*. London, United Kingdom: Longman.
- Cerovic, Z.G., Masdoumier, G., Ghazlen, N.B., & Latouche, G. (2012). A new optical leaf-clip meter for simultaneous non-destructive assessment of leaf chlorophyll and epidermal flavonoids. *Physiologia Plantarum*, 146(3), 251-260. DOI: [doi: 10.1111/j.1399-3054.2012.01639.x](https://doi.org/10.1111/j.1399-3054.2012.01639.x)
- Deinum, B., Sulastri, R.D., Zeinab, M.H.J., & Maassen, A. (1996). Effect of light intensity on growth, anatomy and forage quality of two tropical grasses (*Brachiaria brizantha* and *Panicum maximum* var *trichoglume*). *Netherlands Journal of Agricultural Sciences*, 44, 111-124.
- Durr, P., & Rangel, J. (2000). The response of *Panicum maximum* to a simulated subcanopy environment. 1. Soil x shade interaction. *Tropical Grasslands*, 34, 110-117.
- Eriksen, F.L., & Whitney, A.S. (1981). Effects of light intensity on the growth of some tropical forage species. 1. Interaction of light intensity and nitrogen fertilization on six forage grasses. *Agronomy Journal*, 73, 423-433.
- Jain, A., Roy, A.K., Kaushal, P., Malaviya, D.R., & Zadoo, S.N. (2003). Principal component analysis in guinea grass (*Panicum maximum* Jacq.) germplasm. *Indian Journal of Plant Genetic Resources*, 16, 32-35.
- Jain, A., Roy, A.K., Kaushal, P., Malaviya, D.R., & Zadoo, S.N. (2006). Isozyme banding pattern and estimation of genetic diversity among guinea grass germplasm. *Genetic Resources Crop Evolution*, 53, 339-347.
- Kaushal, P., Malaviya, D.R., & Singh, K.K. (1999). Evaluation of exotic accessions of guinea grass (*Panicum maximum* Jacq.). *Indian Journal of Plant Genetic Resources*, 12, 215-218.
- Kaushal, P., Malaviya, D.R., & Singh, K.K. (2000). Identification of shade tolerant genotypes and alterations in nutrient contents under shade in guinea grass. *Range Management & Agroforestry*, 21, 74-78.
- Kaushal, P., Malaviya, D.R., Roy, A.K., Pathak, S., Agrawal, A., Khare, A., & Siddiqui, S.A. (2008). Reproductive pathways of seed development in apomictic guinea grass (*Panicum maximum* Jacq.) reveal uncoupling of apomixis components. *Euphytica*, 164, 81-92.
- Kenwood, M.G. (1987). A method for comparing profiles of repeated measurements. *Applied Statistics*, 36, 296-308.
- Kharrati-Kopaei, M., & Sadooghi-Alvandi, S.M. (2007). A new method for testing interaction in unreplicated two-way analysis of variance. *Communications in Statistics – Theory and Methods*, 36, 2787-2803.
- Lowry, J.B., Lowry, J.B.C., & Jones, R.J. (1988). Enhanced grass growth below canopy of *Albizia lebbek*. *Nitrogen Fixing Research Reports*, 6, 45-46.
- Malaviya, D.R. (1996). Distribution of morphological diversity among germplasm lines of *Panicum maximum*. *Indian Journal of Plant Genetic Resources*, 9, 193-196.
- Malaviya, D.R. (1998). Evaluation of *Panicum maximum* lines for sustained productivity. *Range Management & Agroforestry*, 19, 126-132.
- Newaj, R., Rizvi, R.H., Chaturvedi, O.P., Alam, B., Prasad, R., Kumar, D., & Handa, A.K. (2017). *A country level assessment of area under agroforestry and its carbon sequestration potential*. Jhansi, India: ICAR - Central Agroforestry Research Institute.
- Norton, B.W., Wilson, J.R., Shelton, H.M., & Hiu, K.D. (1991). The Effect of Shade on Forage Quality. In H.M. Shelton & W.W. Stiller (Eds.), *Forages for Plantation Crops. Proceedings of workshop* (pp. 83-88). Bali, Indonesia, ACIAR Proceedings No. 32.
- Odedire, J.A., & Babayemi, O.J. (2008). Comparative studies on the yield and chemical composition of *Panicum maximum* and *Andropogon gayanus* as influenced by *Tephrosia candida* and *Leucaena leucocephala*. *Livestock Research for Rural Development*, 20(2), 27.

- Parry, C., Mark Blonquist Jr., J., & Bugbee, B. (2014). *In situ* measurement of leaf chlorophyll concentration: analysis of the optical/absolute relationship. *Plant, Cell and Environment*, 37, 2508-2520. DOI:10.1111/pce.12324
- Pereira, V.V., da Fonseca, D.M., Martuscello, J.A., Cecon, P.R., Santos, M.V., & Braz, T.G.S. (2012). Biomass accumulation in mombasa guinea grass plants under different levels of nitrogen supply and plant densities. *Revista Brasileira de Zootecnia*, 41, 1118-1126.
- Plucknett, D.L. (1979). *Managing Pastures and Cattle Under Coconuts*. Westview Tropical Agriculture Series 2. Boulder, United States: Westview Press.
- Samarakoon, S.P., Wilson, J.R., & Shelton, H.M. (1990). Growth, morphology and nutritive value of shaded *Stenotaphrum secundatum*, *Axonopus compressus* and *Pennisetum clandestinum*. *The Journal of Agricultural Science*, 114, 161-169. DOI: 10.1017/S0021859600072154
- Singh, C., & Chaturvedi, O.P. (2011). Rehabilitation of degraded lands through silvipastoral systems – a balanced approach for alternative land use in the north-western Himalayas. In P. Panwar, A.K. Tiwari, & K.S. Dadhwal (Eds.). *Agroforestry Systems for Resources Conservation and Livelihood Security in Lower Himalayas* (pp. 273-296). New Delhi, India: New India Publishing Agency.
- Stfir, W.W., & Shelton, H.M. (1991). Review of Forage Resources in Plantation Crops of Southeast Asia and the Pacific. In H.M. Shelton & W.W. Stfir (Eds.), *Forages for Plantation Crops. Proceedings of Workshop* (pp. 22-31). Bali, Indonesia, ACIAR Proceedings.
- Sukhchain, & Sidhu, B.S. (1992). Correlation and path coefficients analysis for reproductive traits in Guinea grass. *Euphytica*, 60, 57-60.
- Taiz, L., Zeiger, E., Møller, I.M., & Murphy, A. (2015). *Plant Physiology and Development*. Sunderland, United States: Sinauer.
- Wilson, J.R., Catchpole, V.R., & Weier, K.L. (1986). Stimulation of growth and nitrogen uptake by shading a rundown green panic pasture on brigalow clay soil. *Tropical Grasslands*, 20, 134-143.
- Wilson, J.R., Hill, K., Cameron, D.M., & Shelton, H.M. (1990). The growth of *Paspalum notatum* under the shade of a *Eucalyptus grandis* plantation canopy or in full sun. *Tropical Grasslands*, 24, 24-28.
- Wong, C.C., & Wilson, J.R. (1980). Effects of shading on the growth and nitrogen content of green panic and siratro in pure and mixed swards defoliated at two frequencies. *Australian Journal of Agricultural Research*, 31, 269-285.
- Wong, C.C., Sharudin, M.A., & Rahim, H. (1985). Shade tolerance potential of some tropical forages for integration with plantations. *MARDI Research Bulletin*, 13, 225-247.