



Revista de Biología Tropical

ISSN: 0034-7744

ISSN: 2215-2075

Universidad de Costa Rica

Velázquez, María-S.; Fabisik, Juan-C.; Barrera, Marcelo; Allegrucci, Natalia; Valdés, Fabricio-E.; Abarca, Camila-L.; Cabello, Marta
Diversity and abundance of arbuscular mycorrhizal fungi (Glomeromycota) associated with *Ilex paraguarensis* in Northeastern Argentina
Revista de Biología Tropical, vol. 68, no. 4, 2020, October-, pp. 1231-1240
Universidad de Costa Rica

DOI: <https://doi.org/10.15517/RBT.V68I4.41543>

Available in: <https://www.redalyc.org/articulo.oa?id=44966323018>

- How to cite
- Complete issue
- More information about this article
- Journal's webpage in redalyc.org

UAEH
redalyc.org

Scientific Information System Redalyc
Network of Scientific Journals from Latin America and the Caribbean, Spain and Portugal

Project academic non-profit, developed under the open access initiative

Diversity and abundance of arbuscular mycorrhizal fungi (Glomeromycota) associated with *Ilex paraguarensis* in Northeastern Argentina

María S. Velázquez^{1*}, Juan C. Fabisik^{2†}, Marcelo Barrera³, Natalia Allegrucci¹,
Fabricio E. Valdés^{1,4}, Camila L. Abarca¹ & Marta Cabello^{1,4}

1. Instituto de Botánica Carlos Spegazzini, Facultad de Ciencias Naturales y Museo, Universidad Nacional de La Plata, Anexo 1, Avenida 60 y 122 S/N, La Plata, Argentina; mariasvelazquez@yahoo.com.ar, nataliaallegrucci@yahoo.com, camila.abrc@gmail.com
2. Instituto Nacional de Tecnología Agropecuaria, Agencia de Extensión Rural Virasoro, Calle Gobernador Ferre N° 2842, Gobernador Virasoro, Corrientes, Argentina
3. Laboratorio de Investigación en Sistemas Ecológicos y Ambientales, Facultad de Ciencias Agrarias y Forestales, Facultad de Ciencias Naturales y Museo, Universidad Nacional de La Plata. Calle 113 N°469, La Plata, Argentina; mbarrera@fcnym.unlp.edu.ar
4. Comisión de Investigaciones Científicas, Calle 526 entre 10 y 11, La Plata, Argentina; iamrondii@gmail.com, mcabello@gmail.com

* Correspondence

Received 25-IV-2020.

Corrected 18-VIII-2020.

Accepted 26-VIII-2020.

ABSTRACT. Introduction: The expansion and intensification of agriculture causes profound changes at a global scale, which generates a strong impact on crop productivity and consequently, a decrease in the biodiversity associated. *Ilex paraguariensis*, known as yerba mate, is a native species from South America and it has been introduced in the world market (e.g. Middle East, Europe, and United States) due to its multiple nutritional benefits and antioxidant properties. The association of plants of *I. paraguarensis* with arbuscular mycorrhizal fungi (AMF), especially under field conditions, has been scarcely reported so far. **Objective:** The aim of this work was to assess the species composition, richness, spore density and diversity of Glomeromycota communities of yerba mate under different crop and natural conditions, seasonally over a 2-year period. **Methods:** Soil samples were extracted in winter and summer from five productive sites (situations 1-5), with contrasting crop conditions with regard to historical management, as a reference situation (situation 6). Spores were identified following morphological criteria. **Results:** Acaulosporaceae and Glomeraceae presented the highest spore densities in all sampling sites/ seasons/years. The lowest spore density and diversity index was recorded in 2014, when rainfall was higher than in 2013. **Discussion:** We reported *Acaulospora capsicula* detected by morphological analysis, for the first time in South America. Differences between years could be attributed to rainfall. This study contributes to the knowledge of the dynamics and factors that influence the structure of AMF communities over time. This information would be valuable to generate conservation strategies for this group of microorganisms, which are key to the sustainable development of yerba mate cultivation systems.

Key words: arbuscular fungi; Glomeromycota communities; yerba mate; spore density; species richness.

Velázquez, M.S., Fabisik, J.C., Barrera, M. Allegrucci, N., Valdés, F.E., Abarca, C.L., & Cabello, M. (2020). Diversity and abundance of arbuscular mycorrhizal fungi (Glomeromycota) associated with *Ilex paraguarensis* in Northeastern Argentina. *Revista de Biología Tropical*, 68(4), 1231-1240.

The expansion and intensification of agriculture causes profound changes at a global scale, including the degradation of agricultural soils, which generates a strong impact on crop

productivity and consequently, a decrease in the biodiversity associated (De la Fuente & Suárez, 2008). The productivity of *Ilex paraguariensis* St Hil. (Aquifoliaceae), popularly

known as “yerba mate”, is severely affected by soil degradation (Bergotini et al., 2017). This species is native to Argentina, Brazil, Paraguay and Uruguay in South America. It has been introduced in the world market (e.g. Middle East, Europe, and United States) due to its multiple nutritional benefits and antioxidant properties (Heck & De Mejia, 2007).

It is essential to preserve the physical, chemical and biological soil properties in an optimum state for their agricultural sustainability. Plants rely on their interaction with a wide range of soil microorganisms for growth and survival. The microbial interactions and the relationship with plant growth responses and nutrition in the rhizosphere have been widely studied (Larsen, Jaramillo-López, Nájera-Rincon, & González-Esquivel, 2015). The arbuscular mycorrhizal fungi (AMF) (Phylum Glomeromycota) (Schüßler, Schwarzott, & Walker, 2001) are one of the rhizosphere microorganisms symbiotically associated with plants, which promote key ecosystem services related to soil fertility, plant health, and nutrition (Gianinazzi et al., 2010).

The association of plants of *I. paraguayensis* with AMF, especially under field conditions, has been scarcely reported so far. Andrade, Queiroz, Hermes, and Oliveira (2000) registered the presence of mycorrhizal colonization in roots of this species in Santa Catarina (Brazil) and Bergottini et al., (2017) detected the presence of AMF with high throughput sequencing of its root-associated microbiome. In addition, Velázquez, Fabisik, Abarca, Allegrucci, and Cabello (2018) studied the arbuscular mycorrhizal colonization in *I. paraguayensis* plants growing under different crop conditions and in natural environments. However, so far there are few data available on the species composition of AMF communities under field conditions.

Numerous edaphic factors, such as changes in soil pH, humidity, and availability of nutrients, anthropogenic disturbances and seasonality affect the structure and dynamics of Glomeromycota communities (Smith & Read, 2008). Jefwa et al. (2012) demonstrated that

changes in land use can affect the spore density of AMF, and the species richness and diversity of their communities. Besides soil disturbance, intrinsic factors related to AMF life strategies, host physiology and environmental conditions can also modify their sporulation patterns. Gai, Feng, Cai, Christie, and Li (2006) reported that the AMF sporulation is reduced under drought conditions, whereas Shi et al. (2006) consider that their sporulation does not follow a predictable pattern.

In Argentina, more than 60 % of the surface cultivated with yerba mate has serious problems of soil degradation due to inadequate agricultural practices maintained over time. Their recoveries demand not only the application of already developed technologies, but also the advance in research and innovations of existing tools. In this sense, AMF would play an important role. For this reason, it is necessary to know the composition of Glomeromycota communities and identify the factors that can affect their structure and dynamics. The aim of this work was to assess the species composition, richness, spore density and diversity of Glomeromycota communities of *I. paraguayensis* under different crop and natural conditions, seasonally over a 2-year period.

MATERIALS AND METHODS

Study area and Sampling sites: This study was carried out in the provinces of Corrientes and Misiones, Northeastern Argentina. In this region, the soils are typical Kandihumultes of the Ultisol Order, and the climate is classified as humid subtropical, according to the Köppen classification. Samples were collected from five productive sites of *I. paraguayensis* in Corrientes, with contrasting crop situations with regard to historical management. As a reference situation (situation 6), we used an area located in Misiones, with native trees of this species that grow spontaneously in the undergrowth forest. The chemical characterization of the sampling sites is found in Velázquez et al. (2018). The sampling sites corresponded to the following situations: *i*) Situation 1 (Y28) was

implanted in 1928, with a density of 450 plants/ha, there is no fertilizer application and there is continuous weed control by harrow disc. An intensive harvest is carried out at this site and the soil showed extreme deterioration; (ii) Situation 2 (Y24) is the oldest crop, implanted in 1920, with a density of 600 plants/ha, no fertilizers applied. In the last 15 years, it has been managed sustainably, and the harvest is moderate; (iii) Situation 3 (CU) was implanted in 1958, with a density of 1 000 plants/ha. It has fewer extractive harvests than the previous situations and the disk harrow was replaced by chemical weed control 10 years ago. Fertilizers are applied alternatively depending on resource availability; (iv) Situation 4 (CN) was implanted in 1984, with a density of 2 600 plants/ha, fertilization with 300 kg/ha/year N:P:K, and mechanical harvest; (v) Situation 5 (L49) was implanted in 1992, with a density of 4 000 plants/ha, fertilization with 400 kg/ha/year of N:P:K, and mechanical harvest; and (vi) Situation 6 (NAT), correspond to native trees growing spontaneously in the rainforest and not exploited commercially.

Sampling: Samples of soil from the rhizosphere of *I. paraguayensis* plants were collected in winter (July-August) and summer (November-December), which are respectively the stages of lower and higher physiological activity of yerba mate, during the years 2013 and 2014. At situations 1-5, we defined three transects along the plantation lines, which

constituted the repetitions. Each transect was separated from the other by 10 plantation lines. At each transect, we collected a composite sample that consisted of 5-6 subsamples of rhizosphere soil. In the case of situation 6 (NAT), the distance between repetitions was 100 meters. We collected a total of 72 samples (6 sampling situations x 3 repetitions x 2 seasons x 2 years).

In the productive sites (situations 1-5), the mean rainfall of the last 15 years was 1 923 mm, being the mean annual rainfall of 1 697 mm and 2 462.9 mm in 2013 and 2014, respectively. The mean temperatures at these sites were 20.7 °C and 20.8 °C for 2013 and 2014, respectively. In the case of situation 6 (NAT), the mean rainfall of the last 15 years was 1 995 mm, being the mean annual rainfall of 1 995.5 mm and 2 925 mm in 2013 and 2014, respectively. The mean annual temperatures at NAT for 2013 and 2014 were 20.9 °C and 20.7 °C, respectively. Fig. 1 shows the rainfall distribution during 2013-2014 in the sampling sites.

Isolation and identification of AMF spores: Arbuscular mycorrhizal fungi spores were obtained from 100 g (dry weight) of each soil sample by wet sieving using sieves of different mesh sizes (450-105-75-30 µm), and decanting (Gerdemann & Nicolson, 1963). The supernatant was then centrifuged in a sucrose gradient (Walker, Mize, & Nabb, 1982). Arbuscular mycorrhizal fungi species were identified following original species descriptions,




Fig. 1. Distribution of monthly rainfall (2013-2014). **A.** Situation 1-5, productive sites. **B.** Situation 6, native trees. The arrows indicate sampling dates.

taxonomically accepted criteria (Blaszkowski, 2012) of INVAM (International Culture Collection of Arbuscular and Vesicular-Arbuscular Mycorrhizal Fungi: <http://invam.caf.wvu.edu>), and http://www.arbuscular-mycorrhiza.net/amphylo_home.html

The structure of AMF community was analysed by the species richness (S, total number of different AMF taxa present in a sample), abundance (number of spores of each taxon in the sample), frequency observed (percentage of samples in which a taxon is recorded), total density of spores (total number of spores recorded), and Shannon's diversity index (H' ; $H' = -\sum p_i \ln p_i$, where p_i is the relative abundance of the its species compared with the total of the species identified in the sample) (Magurran, 1988). All parameters were referred to 100 g of dry soil.

Statistical analyses: To evaluate the effect of the different sampling situations, seasons and years on the total spore density and diversity index, a multi-factor analysis of variance (ANOVA model I) followed by Tukey's test ($P \leq 0.05$) was used. All analyses were performed with InfoStat version 2012 (Di Renzo et al., 2012). A principal component analysis (PCA) was employed to explore the effect of years on the structure of AMF communities (MVSP, version 3.1).

RESULTS

Twenty-three taxa of Glomeromycota belonging to 11 genera in the families Acaulosporaceae, Claroideoglomeraceae, Diversisporaceae, Entrophosporaceae, Gigasporaceae, Glomeraceae, and Pacisporaceae were recovered (Table 1).

All families were recovered throughout the 2-year period in at least one of the sampling sites (Table 2). Acaulosporaceae and Glomeraceae presented the highest spore densities in all sampling sites/ seasons/years. The rest of the families were scarcely represented, and some of them were present at only one sampling site per year. Six of the seven families were

recorded during the first year (2013), in which Entrophosporaceae and Pacisporaceae were absent only in winter and summer, respectively. The lowest spore density was recorded in winter 2014, which is consistent with the lowest diversity of families found at this sampling date, due to the absence of Claroideoglomeraceae, Entrophosporaceae, and Gigasporaceae. Entrophosporaceae was the only family not reported in summer 2014.

Three species, *Acaulospora capsicula* (96%), *Glomus* sp. (92 %), and *Diversispora spurca* (79%), showed frequencies higher than 50 % and were registered in all sampling situations. *Acaulospora capsicula* is recorded for the first time in Argentina. *Acaulospora denticulata*, *A. excavata*, *A. rhemii*, *A. spinosa*, *Acaulospora* sp.1, *Funnelformis geosporum*, and *Rhizophagus clarum* were recorded with low frequency only during the first year of sampling, whereas *Claroideoglomus claroideum* and *Scutellospora gilmorei* were only found during the second year.

The total number of spores was significantly higher in 2013 than in 2014 ($F = 24.7$, $df = 1$, $P = < 0.0001$). No significant differences in the spore number were found between seasons or sampling sites ($F = 3.51$, $df = 1$, $P = 0.067$ and $F = 1.3$, $df = 5$, $P = 0.2785$, respectively). Significant differences were recorded only in the interaction Year x Season ($F = 8.84$, $df = 1$, $P = 0.0046$), with the highest numbers in winter 2013.

Fig. 2 shows the diversity index (H') and the species richness (S) analysed by sampling site/seasons/years. Both indices showed higher values in 2013 ($x_{H'} = 0.95$ and $x_s = 4.5$) than in 2014 ($x_{H'} = 0.77$ and $x_s = 3.11$), but only the species richness (S) was significantly different between years ($F = 19.08$, $df = 1$, $P = 0.0001$).

Not significant differences were detected in H' and S index at sites ($F = 2.77$, $df = 5$, $P = 0.02$ and $F = 2.15$, $df = 5$, $P = 0.07$ respectively). The H' and S index was great in Y28 ($x_{H'} = 1.08$ and $x_s = 4$ respectively) and the lowest in Y24 ($x_{H'} = 0.5$) and L49 ($x_s = 2$).

The PCA biplot (Fig. 3) showed that 52 % of the total variation was explained by the first

TABLE 1
Mean spore abundance (number of spores/100g of dry soil) and frequency (F %) of each taxon of Glomeromycota obtained over the study period (2013-2014) and in the two seasons (winter-summer)

Phylum	2013												2014												F%												
	Y28			L49			Y24			CN			NAT			CU			Y28			L49				Y24			CN			NAT			CU		
	WIN	SUM		WIN	SUM		WIN	SUM		WIN	SUM		WIN	SUM		WIN	SUM		WIN	SUM		WIN	SUM			WIN	SUM		WIN	SUM		WIN	SUM				
Acaulosporaceae	101	43	89	9	83	127	68	26	69	0	84	95	1	<1	5	19	23	11	15	44	26	61	13	6	30	53	8										
<i>Acaulospora bireticulata</i>			<1		<1																						4										
<i>A. capsicula</i>			<1	5																							4										
<i>A. denticulate</i>			<1	2	2	4										2								2	2	2	29										
<i>A. laevis</i>	5	8	16	4	15	9	<1	2	34	5	10	1			3								3	2	2	71											
<i>A. mellea</i>	<1																									4											
<i>A. rehmtii</i>	1						1		3		2		1	1												29											
<i>A. scrobiculata</i>										1																4											
<i>A. spinose</i>										1																4											
<i>Acaulospora</i> sp	<1										1															4											
Claroideglomeraceae																																					
<i>Clareidoglomus clarioideum</i>			<1			2									2												4										
<i>C. etunicatum</i>																											84										
Diversisporaceae																																					
<i>Diversispora spurca</i>	6	6		2	2	4	<1	2	3	4	4	4	4	2	4	7	9	3	6	5	2	5	5	5	5	79											
Entrophosporaceae																																					
<i>Entrophospora infrequens</i>						1																		<1			4										
Gigasporaceae																																					
<i>Gigaspora</i> sp.	2	5																									13										
<i>Racocetra fulgida</i>		3	2											1					1								21										
<i>Scutellospora Gilmore</i>														1						2							8										
Glomeraceae																																					
<i>Funneliformis geosporum</i>	18		6	5																							25										
<i>F. mosseae</i>	7	1												2								1	1				33										
<i>Glomus microaggregatum</i>		1	<1																	2		2					33										
<i>Glomus</i> sp.		21	7	8										11	3	7	19	1	7	28	<1	3	4	12	92	13											
<i>Rhizophagus clarus</i>			3																																		
Pacisporaceae																																					
<i>Pacispora</i> sp.			2																					<1			4										
Total abundance	147	88	130	32	126	164	97	47	146	26	154	107	25	32	46	38	16	56	57	82	12	10	41	72													

Site 1 (Y28); Site 2 (Y24); Site 3 (CU); Site 4 (CN); Site 5 (L49) and Site 6 (NAT). The data correspond to the mean of three replicate.




Fig. 2. Shannon's diversity index (H') and species richness (S) of Glomeromycota over the study period (2013- 2014) and in the two seasons (winter-summer). Site 1 (Y28); Site 2 (Y24); Site 3 (CU); Site 4 (CN); Site 5 (L49) and Site 6 (NAT). (Species richness; Shannon's diversity index). The data correspond to the mean of three replicates.

two axes, which separated the samples into two groups on opposite sides of the graph according to the years (2013 on the right and 2014 on the left). The species *A. capsicula*, *A. mellea*,

F. geosporum and *Glomus* sp. correlated positively with the first axis (0.48, 0.60, 0.43 and 0.34 respectively) and were associated to 2013 samples, while *G. microaggregatum* showed a




Fig. 3. Principal component analysis (PCA) of the relative abundances of AMF species over the study period (2013-2014) and in the two seasons (INV: winter-VE: summer). Site 1 (Y28); Site 2 (Y24); Site 3 (CU); Site 4 (CN); Site 5 (L49) and Site 6 (NAT). 1: 2013; 2: 2014.

TABLE 2

Site 1 (Y28); Site 2 (Y24); Site 3 (CU); Site 4 (CN); Site 5 (L49) and Site 6 (NAT).

DISCUSSION

Entrophosporaceae, Gigasporaceae, and Pacisporaceae showed a low frequency, and their presence was occasional in the different crop situations/seasons/years, without showing any tendency. According to Lovelock and Ewel (2005), the distribution patterns of the relative abundances of fungal species are difficult to interpret, because there is not enough data

on the ecology and physiology of most species. Among the species identified, the frequency of *A. capsicula*, *Glomus* sp. and *D. spurca* was higher than 50 % and were recorded in all crop situations/seasons/years analysed. *Acaulospora capsicula* is a new record for Argentina and South America. This species was originally described from the rhizosphere of *Ficaria verna* growing in a forest in northern Poland (Błaszowski, 1990) and posteriorly, Schultz, Bever, and Morton (1999) found *A. capsicula* together with *D. spurca* associated with *Juncus conglomeratus*, *Salix triandra*, *Thuja occidentalis*, and *Triticum aestivum*, in the campus of Duke University, North Carolina. *Diversispora spurca* has been commonly reported worldwide, with already 20 records for South America. These two species together with *Glomus* sp., are included within the families Acaulosporaceae (*A. capsicula*), Diversisporaceae (*D. spurca*), and Glomeraceae (*Glomus* sp.), which are characterized by producing a high number of spores of small size and rapid germination. In general, they are considered to be opportunistic, because they can rapidly take advantage of favourable conditions under varied and fluctuating environments (Velázquez & Cabello, 2011).

In this study, the total abundance of spores was analysed seasonally in sites with different agricultural practices, over a 2-year period. Of the variables analysed, only the effect of the years influenced the spore density. A 77 % of the total spore number was recorded in 2013. The mean density of spores was greater in 2013 (105 spores/100 g of dry soil) than in 2014 (40 spores/100 g of dry soil). Rivera and Fernández (2003) classified the spore density as low (< 200 spores/50 g of soil), average (between 300-500 spores/50 g of soil) and high (> 600 spores/50 g of soil). However, these numbers might vary depending on whether they are natural ecosystems or agroecosystems. The differences in spore density that were found between the first and second year could be attributed to rainfall. Velázquez et al. (2018) highlighted that when analysing the effect of rainfall on the AMF colonization, it should

be considered not only the annual rainfall but also its distribution, particularly during the three or four months prior to sampling. These authors observed that the vegetative state of yerba mate crop at a given time better reflects the water status of the soil during the previous three to four months. In this study, the lowest number of AMF spores was recorded in 2014, when rainfall was 89 % higher than in 2013. García, Mendoza, and Pomar (2008) observed that excessive water availability reduces the spore number in soil and that fluctuations in spore density are more related to this factor than to seasonal changes. The low number of spores reported herein for *I. paraguayensis* is comparable with our previous work (Velázquez et al., 2018), in which a significantly less spore number and higher values of colonization were recorded in 2014 than in 2013. Ragupathy and Mahvedan (1993) observed that the presence of internal and external mycelium increases during the rainy season when germination is favoured, with the consequent increase in colonization and decrease in spore abundance and diversity in the soil during these periods.

Similarly, the species richness (S) and diversity (H') showed the same trend mentioned above for the spore density. That is, the highest values of both indices were recorded during the first study year. The greater spore density was due to the greater number of families of Glomeromycota that led to higher species diversity.

In summary, the present study represents a diversity survey of Glomeromycota associated with *I. paraguayensis*, under different crop situations and natural conditions. *Acaulospora capsicula* is reported for the first time in South America. This study also contributes to the knowledge of the dynamics and factors that influence the structure of AMF communities over time. We hope that this information would be valuable to generate conservation strategies for this group of microorganisms, which are key players in the sustainable development of yerba mate cultivation systems, as well as in other tropical crops with similar characteristics.

Ethical statement: authors declare that they all agree with this publication and made significant contributions; that there is no conflict of interest of any kind; and that we followed all pertinent ethical and legal procedures and requirements. All financial sources are fully and clearly stated in the acknowledgements section. A signed document has been filed in the journal archives.

ACKNOWLEDGMENTS

This study was partially supported by Comisión de Investigaciones Científicas de la Provincia de Buenos Aires and Universidad Nacional de La Plata (11/N 773).

RESUMEN

Diversidad y abundancia de hongos micorrízicos arbusculares (Glomeromycota) asociados con *Ilex paraguariensis* en el noreste de Argentina. **Introducción:** la expansión e intensificación de la agricultura provoca profundos cambios a escala mundial, lo que genera un fuerte impacto en la productividad de los cultivos y, en consecuencia, una disminución de la biodiversidad asociada. *Ilex paraguariensis*, conocida como yerba mate, es una especie nativa de América del Sur y se ha introducido en el mercado mundial (por ejemplo, Medio Oriente, Europa y Estados Unidos) debido a sus múltiples beneficios nutricionales y propiedades antioxidantes. La asociación de plantas de *I. paraguariensis* con hongos arbusculares (AMF), especialmente en condiciones de campo, ha sido escasamente documentada. **Objetivo:** El objetivo de este trabajo fue evaluar estacionalmente durante un período de 2 años: la composición de especies, la riqueza, la densidad de esporas y la diversidad de las comunidades Glomeromycota de yerba mate en diferentes cultivos y condiciones naturales. **Métodos:** Se extrajeron muestras de suelo en invierno y verano de cinco sitios productivos con condiciones de cultivo (situaciones 1-5), en contraste con manejo histórico como situación de referencia (situación 6). Las esporas se identificaron siguiendo criterios morfológicos. **Resultados:** Acaulosporaceae y Glomeraceae presentaron las mayores densidades de esporas en todos los sitios de muestreo / estaciones / años. El índice más bajo de densidad y diversidad de esporas se registró en 2014, cuando las precipitaciones fueron más altas que en 2013. **Discusión:** Se registra por primera vez para América del Sur a *Acaulospora capsicula* identificada por caracteres morfológicos. Las diferencias entre años podrían atribuirse a las precipitaciones. Este estudio contribuye al conocimiento de la dinámica y los factores que influyen en la

estructura de las comunidades AMF a lo largo del tiempo. Esta información sería valiosa para generar estrategias de conservación para este grupo de microorganismos, que son clave para el desarrollo sostenible de los sistemas de cultivo de yerba mate.

Palabras clave: hongos arbusculares; comunidades de Glomeromycota; yerba mate; densidad de esporas; riqueza de especies.

REFERENCES

- Aidar, M.P.M., Carrenho, R., & Joly, C.A. (2004). Aspects of arbuscular mycorrhizal fungi in an atlantic forest chronosequence parque estadual turístico do Alto Ribeira (Petar), SP. *Biota Neotropica*, 4(2), 1-15.
- Andrade, A.C.S., Queiroz, M.H., Hermes, R.A.L., & Oliveira, V.L. (2000). Mycorrhizal status of some plants of the Araucaria forest and the Atlantic rainforest in Santa Catarina, Brazil. *Mycorrhiza* 10(3), 131-36.
- Bergottini, V.M., Hervé, V., Sosa, D.A., Otegui, M.B., Zapata, P.D., & Junier, P. (2017). Exploring the diversity of the root-associated microbiome of *Ilex paraguariensis* St. Hil. (Yerba Mate). *Applied Soil Ecology*, 109, 23-31.
- Błaszowski, J. (1990). Polish Endogonaceae. VII. *Acaulospora capsicula* sp. nov. *Mycologia*, 82(6), 794-798.
- Błaszowski, J. (2012). *Glomeromycota*. Institute of Botany, Kraków, Poland: Polish Academy of Sciences.
- De La Fuente, E.B., & Suárez, S.A. (2008). Problemas ambientales asociados a la actividad humana: la agricultura. *Ecología Austral*, 18(2), 239-252.
- Di Rienzo, J.A., Casanoves, F., Balzarini, M.G., Gonzalez, L., Tablada, M., & Robledo, C.W. (2012). *InfoStat versión 2012*. Grupo InfoStat, FCA, Universidad Nacional de Córdoba, Argentina.
- Gai, J.P., Feng, G., Cai, X.B., Christie, P., & Li, X.L. (2006). A preliminary survey of the arbuscular mycorrhizal status of grassland plants in southern Tibet. *Mycorrhiza*, 16(3), 191-196.
- García, I., Mendoza, R., & Pomar, M.C. (2008). Deficit and excess of soil water impact on plant growth of *Lotus tenuis* by affecting nutrient uptake and arbuscular mycorrhizal symbiosis. *Plant Soil*, 304(1-2), 117-131.
- Gerdeman, J.W., & Nicolson, T.H. (1963). Spores of mycorrhizal *Endogone* species extracted from soil by wet sieving and decanting. *Transaction of the British Mycological Society*, 46, 235-239.
- Gianinazzi, S., Gollotte, A., Binet, M., van Tuinen, D., Redecker, D., & Wipf, D. (2010). Agroecology:

- the key role of arbuscular mycorrhizas in ecosystem services. *Mycorrhiza*, 20(8), 519-530.
- Heck, C.I., & De Mejia, E.G. (2007). Yerba Mate Tea (*Ilex paraguariensis*): a comprehensive review on chemistry, health implications, and technological considerations. *International Journal of Food Science*, 72(9), 138-151.
- Jansa, J., Mozafar, A., Anken, T., Ruh, R., Sanders, I.R., & Frossard, E. (2002). Diversity and structure of AMF communities as affected by tillage in a temperate soil. *Mycorrhiza*, 12, 225-234.
- Jefwa, J.M., Okoth, S., Wachira, P., Karanja, N., Kahindi, J., Njuguni, S., ... Huising, J. (2012). Impact of land use types and farming practices on occurrence of arbuscular mycorrhizal fungi (AMF) Taita-Taveta district in Kenya. *Agriculture Ecosystem & Environment*, 157, 32-39.
- Larsen, J., Jaramillo-López, P., Nájera-Rincon, M., & González-Esquivel, C. (2015). Biotic interactions in the rhizosphere in relation to plant and soil nutrient dynamics. *Journal of Soil Science and Plant Nutrition*, 15(2), 449-463.
- Lovelock, C.E., & Ewel, J.J. (2005). Links between tree species, symbiotic fungal diversity and ecosystem functioning in simplified tropical ecosystems. *New Phytologist*, 167(1), 219-228.
- Magurran, A.E. (1988). *Ecological diversity and its measurement*. New Jersey, United States: Princeton University Press.
- Oehl, F., Sieverding, E., Ineichen, K., Mäder, P., Boller, T., & Wiemken, A. (2003). Impact of land use intensity on the species diversity of arbuscular mycorrhizal fungi in agroecosystems of Central Europe. *Applied and Environmental Microbiology*, 69(5), 2816-2824.
- Palenzuela, J., Ferrol, N., Boller, T., Azcón-Aguilar, C., & Oehl, F. (2008). *Otospora barei*, a new fungal species in the Glomeromycetes from a dolomitic shrub land in a Sierra de Baza National Park (Granada, Spain). *Mycologia*, 100(2), 296-305.
- Raguphathy, S., & Mahvedan, A. (1993). Distribution of vesicular-arbuscular mycorrhizae in the plants and rhizosphere soils of the tropical plains, Tamil Nadu, India. *Mycorrhiza*, 3(3), 123-136.
- Rivera, R., & Fernández, K. (2003). *El manejo efectivo de la simbiosis micorrízica, una vía hacia la agricultura sostenible, estudio de caso*. Ciudad de La Habana, Cuba: INCA Ediciones.
- Schultz, P.A., Bever, J.D., & Morton, J.B. (1999). *Acaulospora colossica* sp. nov. from an old field in North Carolina and morphological comparisons with similar species, *A. laevis* and *A. koskei*. *Mycologia*, 91, 676-683.
- Schüßler, A., Schwarzott, D., & Walker, C. (2001). A new fungal phylum, the Glomeromycota: phylogeny and evolution. *Mycological Research*, 105(12), 1413-1421.
- Schüßler, A., & Walker C. (2010). *The Glomeromycota. A Species List with New Families and New Genera*. Munich, and Oregon State University: The Royal Botanic Garden Kew, Botanische Staatssammlung.
- Shi, Z.Y., Chen, Y.L., Feng, G., Liu, R.J., Christie, P., & Li, X.L. (2006). Arbuscular mycorrhizal fungi associated with the Meliaceae on Hainan Island, China. *Mycorrhiza*, 16(2), 81-87.
- Smith, S.E., & Read, D.J. (2008). *Mycorrhizal symbiosis*. London, England: Academic Press.
- Stürmer, S.L., & Siqueira, J.O. (2011). Species richness and spore abundance of arbuscular mycorrhizal fungi across distinct land uses in Western Brazilian Amazon. *Mycorrhiza*, 21(4), 255-267.
- Turrini, A., & Giovannetti, M. (2012). Arbuscular mycorrhizal fungi in national parks, nature reserves and protected areas worldwide: a strategic perspective for their in situ conservation. *Mycorrhiza*, 22(2), 81-97.
- Velázquez, M.S. (2010). *Comunidades de hongos formadores de micorrizas arbusculares en el Parque Nacional El Palmar* (Doctoral Dissertation). National University of La Plata, Argentina.
- Velázquez, M.S., & Cabello, M.N. (2011). Occurrence and diversity of arbuscular mycorrhizal fungi in trap cultures from El Palmar National Park soils. *European Journal of Soil Biology*, 47(4), 230-235.
- Velázquez, M.S., Cabello, M., Irrazabal, G.B., & Godeas, A. (2008). *Acaulosporaceae* from el Palmar National Park, Entre Ríos, Argentina. *Mycotaxon*, 103(1), 171-188.
- Velázquez, M.S., Fabisik, J.C., Abarca, C.L., Allegrucci, N., & Cabello, M. (2018). Colonization dynamics of arbuscular mycorrhizal fungi (AMF) in *Ilex paraguariensis* crops: Seasonality and influence of management practices. *Journal of King Saud University – Science*, 32(1), 183-188. DOI: 10.1016/j.jksus.2018.03.017
- Walker, C., Mize, W., & Mc Nabb, H.S. (1982). Populations of endogonaceous fungi at two populations in central Iowa. *Canadian Journal of Botany*, 60, 2518-2529.