



Revista Geológica de América Central

ISSN: 0256-7024

Universidad de Costa Rica

Ascanio D, Rincón; Ana L, Valerio; César A, Laurito
First fossil record of a Megatheriidae-Megatheriinae in the Early Hemphillian
(Late Miocene) from San Gerardo de Limoncito, Curré Formation, Costa Rica
Revista Geológica de América Central, no. 62, 2020, pp. 1-24
Universidad de Costa Rica

DOI: <https://doi.org/10.15517/rgac.v62i0.41278>

Available in: <https://www.redalyc.org/articulo.oa?id=45466248001>

- How to cite
- Complete issue
- More information about this article
- Journal's webpage in redalyc.org

UAEH
redalyc.org

Scientific Information System Redalyc
Network of Scientific Journals from Latin America and the Caribbean, Spain and
Portugal

Project academic non-profit, developed under the open access initiative

First fossil record of a Megatheriidae-Megatheriinae in the Early Hemphillian (Late Miocene) from San Gerardo de Limoncito, Curré Formation, Costa Rica

Primer registro fósil de un Megatheriidae-Megatheriinae para el Hemphilliano (Mioceno Tardío) de San Gerardo de Limoncito, Formación Curré, Costa Rica

Ascanio D. Rincón^{1*}, Ana L. Valerio² & César A. Laurito^{3, 4}

¹Instituto Venezolano de Investigaciones Científicas (IVIC), Laboratorio de Paleontología, Centro de Ecología, Ado. postal 20632, Caracas 1020-A - Venezuela

²Museo Nacional de Costa Rica, Sección de Geología, Departamento de Historia Natural, Apdo. postal 749-1000, San José, Costa Rica

³Instituto Nacional de Aprendizaje (INA), Centro Nacional Especializado en Turismo (CENETUR), San José, Costa Rica

⁴Museo Nacional de Costa Rica, Departamento de Historia Natural (investigador asociado), San José, Costa Rica

*Autor de contacto: paleosur1974@gmail.com

(Recibido: 4/03/2020; aceptado: 18/03/2020)

ABSTRACT: We report the first record of Megatheriidae-Megatheriinae fossil record in Central and North America, which also represents the earliest record of a Megatheriidae-Megatheriinae outside of South America at 5.8 Ma, and before the closure of the Isthmus of Panama. The fossil record of the Megatheriidae record comes from the Curré Formation, Late Hemphillian (Hh3) (Late Miocene), San Gerardo de Limoncito locality, Canton of Coto Brus, Puntarenas Province, Costa Rica, and shows a combination of plesiomorphic and advance characters whose combination is unique among the Megatheriinae. These Megatheriidae remains are scarce and fragmentary, however, they have enough morphological information and key morphological characters to allow us to propose a new genus and species of Megatheriinae, *Sibotherium ka* gen. et sp. nov. Based on the morphology preserved in these remains, *Sibotherium ka* is more closely related to *Megathericulus*, *Pyramiodontherium* or *Anisodontherium* than to the more advanced Megatheriinae genera *Urumaquia*, *Megatherium* or *Eremotherium*. We explain the early record of this xenarthran in Costa Rica by the presence of a land corridor established for short periods of time as a result of tectonic uplift or by a drop in sea level or a mixture of both at approximately 5.8 Ma in Southern Central America.

Keywords: Mammalia; Xenarthra; Megatheriinae; Curré Formation; Costa Rica.



Resumen: Reportamos el primer registro fósil de Megatheriidae-Megatheriinae en América Central y del Norte, que además representa el registro más temprano de un Megatheriidae-Megatheriinae fuera de América del Sur con 5,8 Ma, y antes del cierre del Istmo de Panamá. El registro fósil de este Megatheriidae proviene de la Formación Curré, Hemphilliano tardío (Hh3) (Mioceno Tardío), de la localidad de San Gerardo de Limoncito, Cantón de Coto Brus, Provincia de Puntarenas de Costa Rica, y muestra una combinación de caracteres plesiomórficos y avanzados cuya combinación es única entre los Megatheriinae. Estos restos de Megatheriidae son escasos y fragmentarios, sin embargo, tienen suficiente información morfológica y caracteres morfológicos clave que nos permiten proponer un nuevo género y especie de Megatheriinae, que llamaremos *Sibotherium ka* gen. et sp. nov. En cuanto a la morfología preservada en estos restos, *Sibotherium ka* estaría más estrechamente relacionado con *Megathericulus*, *Pyramiodontherium* o *Anisodontherium* que con los Megatheriinae más avanzados *Urumaquia*, *Megatherium* o *Eremotherium*. Explicamos el registro temprano de este Xenarthra en Costa Rica, mediante la presencia de un corredor terrestre, establecido por breves períodos de tiempo producto del levantamiento tectónico o una caída del nivel del mar o una mezcla de ambos, hace aproximadamente 5.8 Ma en América Central meridional.

Palabras clave: Mammalia; Xenarthra; Megatheriinae; Curré Formation; Costa Rica.

INTRODUCTION

The Xenarthra are characterized by a particular suite of skeletal modifications, e. g. shape and relationship of the vertebral metapophysis on the last thoracic and lumbar vertebra, presence of synsacrum, ischio-sacral union, sacroischial fenestra, reduction or absence of enamel, monophiodonty and hypsodonty (McDonald, 2003). The Xenarthrans constitute one of the most peculiar and characteristic clades of South American mammals (Pujos, et al., 2013). Xenarthra, including fossil and living species, consist of two major clades: Pilosa (Sloths, and Anteaters), without dermal armor, but covering with a dense hair, which comprise Vermilingua and Eufolivora, and Cingulata (Armadillos, pampatheres and glyptodonts), characterized by the development of bony dermal armor (armoured Xenarthra) (Carlini, Brandoni & Molin, 2013; Pujos & De Iuliis, 2007; Pujos, et al., 2013; Varela, Tambusso, McDonald & Fariña, 2019).

The last hypotheses of the Eufolivora or Sloths phylogenetic relationship, based on bayesian morphological clock analysis, are grouping them in five major clades: Mylodontidae, Nothotheriidae, Megatheriidae, Megalonychidae and Bradypodidae (Varela et al., 2019).

The Megatheriidae are small to giant ground sloths, and they are diagnosed by eight unequivocal synapomorphies. These include an elongate mandibular symphysis and spout [62(3) and 68(2), Fig. 8 (Gaudin, 2004)], the lack of a clear demarcation between the posterior end of the symphysis and the ventral edge of the horizontal ramus of the mandible in lateral view [69(1), Gaudin, 2004].

Dental characters have been of particular importance in dividing the diverse assemblage of extinct ground sloths into its traditional constituent families: the Megatheriidae, typified by large square teeth with parallel transverse crests and a thickened external layer of cementum; the Mylodontidae, diagnosed by their peculiar lobate dentition; and the Megalonychidae, identified by their quadrate or ovate molariform teeth with subparallel crests, and by their enlarged caniniform or incisiform first upper and lower teeth (Gaudin, 2004).

In addition, the three auditory region synapomorphies of the Megatheriidae identified by Gaudin (1995) also serve to diagnose this node. The sister-group relationship between *Eremotherium* and *Megatherium* (subfamily Megatheriinae, Node 25) is one of the most robust nodes on the cladogram. It is diagnosed by 48 unambiguous synapomorphies, among them nine unique to the megatheriines. Of these nine, seven are dental characters: the third lower molariform is the smallest molariform; the upper caniniform is square in cross-section; the lower caniniform is rectangular in cross-section; the lower first molariform is square in cross-section; the second and third upper molariforms are square in cross-section; the second lower molariform is square in cross-section; and the third lower molariform is trapezoidal in cross-section.

The other two include a posterior external opening of the mandibular canal that faces anteromedial, lying on the internal surface of the ascending ramus, and a lacrimal foramen that opens into a ventrally directed canal on the surface of the lacrimal bone (Gaudin, 2004).

The megatheriidae include two Subfamilies: Planopsinae and Megatheriinae (Varela, et al., 2019). Overall, the list of valid megatheriine genera from the Middle Miocene–Pliocene of South America currently comprises *Megathericulus*

Ameghino, (1904), *Eomegatherium* Kraglievich (1926), *Promegatherium* Ameghino (1883), *Megatheridium* Cabrera (1928), *Pliomegatherium* Kraglievich (1930), *Megatheriops* Ameghino and Kraglievich (1921), *Plesiomegatherium* Roth (1911), *Megatheridium* Cabrera (1928), *Pyramiodontherium* Rovereto (1914), *Anisodontherium* Brandoni & De Iuliis (2007) and Brandoni, Powell & González (2012), *Urumaquia* Carlini, Brandoni & Sánchez (2006) and *Proeremotherium* Carlini et al. (2006).

There are also megatherioid fossils from the Middle Miocene of Cerdas, Bolivia, reported as Megatheriidae indeterminate (Croft et al., 2016). Hirschfeld (1985) reported some material of indeterminate Megatheriinae from the middle Miocene of La Venta in Colombia.

Some features in the skull of the Pliocene (Codore Formation, Urumaco desert Falcon state Venezuela) taxa *Proeremotherium eljebe* could indicate an ancestral or close relationship with the geographically-widespread megatheriine *Eremotherium laurillardii* (Carlini et al., 2006).

Eremotherium comprises three species: *E. eomigrans* De Iuliis and Cartelle, 1999, from the late Pliocene (Blancan) of the USA; *E. sefvei* De Iuliis and St-André, 1997, from the Pleistocene of Bolivia, and *E. laurillardii* from several late Pleistocene localities of North America (e.g., southern Mexico and west coast of USA up to South Carolina), Central America and low latitudes of South America (De Iuliis, 1996).

The only one report of a Megatheriidae in the late Miocene of Central America was made by Laurito and Valerio (2012a), who reported some teeth of Megatheriidae genus and species indeterminate, from San Gerardo de Limoncito, Curré Formation, (Early Hemphillian Hh2 - Late Miocene) from Costa Rica. Also, a juvenile specimen of *Pliometanastes* has been reported from that locality by Valerio and Laurito (2014), very similar to? *Pliometanastes galushai*, which is why they considered the specimen as a juvenile of *Pliometanastes* sp.

AGE, GEOLOGY AND LOCALITY

The rock unit that contains the fossiliferous remains corresponds to the Curré Formation. Henningsen (1965) defines this formation as a series of tuffaceous sandstones with intercalations of conglomerates, siltstones and occasionally shales, dark green and brownish green when altered. Mora (1979), describes two subunits according to their sedimentological characteristics: The subunit of volcanoclastic conglomerates of gray-green color, overlain by shales that settled in a transitional environment between delta and beach. The subunit of black shales, interspersed with strata of sandstones and fine conglomerates, was deposited in a paralic environment of estuary or swamp type. They are very chloritized and basically consist of subvolcanic products.

The paleontological material recovered is housed at the Geology Section of the Natural History Department at The National Museum of Costa Rica, and comes from a level of fine conglomerate that is located towards the roof of a sequence of blue clays, which in turn is overlain by a sequence of medium to fine sandstones, which confirms its association with the superior subunit or subunit of shales (*sensu* Mora, 1979).

Granados and Aguilar (1983), based on the presence of molluscs, obtained a Middle-Late Miocene age for the Curré Formation.

With regard to sedimentology, in the San Gerardo de Limoncito locality the following sedimentary facies are established (Figs. 1, 2).

Neritic or Proximal Platform

The lithofacies of glauconitic blue clay characterizes this depositional environment and suggests a slow sedimentation rate. This lithofacies is constituted by a bank of 2 meters of visible thickness, of decimetric strata of clays and silts, finely laminated with abundant plant detritus and wood reworked by the waves, and floating wood perforated by *Teredo* sp.

No other fossil traces are observed into the sediments, which, together with the abundant presence of glauconite and carbonaceous material, is interpreted as a predominantly reducing environment (Chafetz & Reid, 2000).

Subaquatic fan delta

The lithofacies of conglomerates is what characterizes this depositional environment and is composed of blue clay intraclasts, coming from the lower level which it overlies at an erosional discordance. The clasts have diameters that vary between 5 and 13 cm; they are well rounded, little or not spherical; some have point contact, but 90% of them float in a matrix of glauconitic clay. The clasts are arranged with their long axis parallel to the layer of stratification, and it is possible to observe some degree of imbrication. In this sequence are frequent bone and dental remains of various vertebrates, both terrestrial and marine. The *Trypanites* ichnofacies, in this case, is inferred indirectly from the blue clay intraclasts that are usually perforated by bivalves of the family Pholadidae, from which it is still possible to find remains of the valves within the fossil traces of *Gastrochaenolites* sp.




Fig. 1: Geographic location of studied area, Geological map of the San Gerardo de Limoncito area, modified from Denyer & Alvarado (2007).




Fig. 2: Stratigraphic column and sedimentological interpretation of the locality of San Gerardo de Limoncito, Costa Rica (Laurito & Valerio, 2010) where *Sibotherium ka* gen. et sp. nov. was found.

This suggests the existence of lithified bottoms in marine environments at depths not greater than 35 m, suggested by the Pholadidae traces and associated with rocky coasts, hardgrounds, unconformities and other surfaces of omission; this is also confirmed by the presence of oysters and parautochthonous Cirripedia, as well as the shell of a male crab identified as *Callinectes* sp. However, it is clear that these underwater, shallow, hardgrounds were located in areas adjacent to or directly below where the sedimentary lobes of the subtidal fan delta, associated with a fluvial system, were deposited. The energy of the deposit was associated with one or several periods of rainfall, which reached the traction necessary to erode the hardgrounds and produce the “rip up clasts” or intraclasts, and even to detach fragments of the blue clays and integrate them within the thick sediments of the fan delta.

Infralittoral

The lithofacies of fine-to medium-grained sandstones, well selected and clean, is the predominant lithology and forms a bank of 2.5 m thickness, constituted by metric and homogeneous strata in which a diffuse parallel lamination and low angle foresets associated with small tidal channels is observed. Other levels present scarce carbonaceous material and infrequent ichnofossils. This sequence overlaps the conglomerate facies by a sedimentary discordance and presents two centimeter cycles of coarse gravels at its base that quickly grade to medium and fine sands. Also, it was verified in the field that this sequence interdigitates laterally with the facies of the internal platform, since it was observed that these fine sands partially

reach the blue clay facies, directly overlying it at some points. The *Skolithos* ichnofacies, in this case, is deduced from small and scarce traces of *Thalassinoides* sp., suggesting a high energy environment, associated with clean and well selected sediments with parallel lamination and low angle foresets. This is in accordance with the definition given by Seilacher (1967) for this ichnofacies.

SYSTEMATIC PALEONTOLOGY

Xenarthra Cope (1889)
Phyllophaga Owen (1842)
Megatherioidea Gray (1821)
Megatheriidae Gray (1821)
Megatheriinae Gray (1821)

Sibotherium Gen. Nov.

DIAGNOSIS. — *Sibotherium* differs from other Megatheriinae in the following character combination: Large size Megatheriidae-Megatheriinae; the posterior end of the symphysis lies well anterior to the first molariform; the m4 is posterior to the anterior margin of the ascending ramus, and not visible in lateral view; molariforms 6th to 22 % wider than long; the valleys between the lophs of the occlusal surface of the molariforms are V-shaped and shallow; astragalus 12,5% wider than long than any other medium size Megatheriinae; the navicular facet lies dorsally, with plane of discoid facet intersecting it approximately at the middle of the navicular depression; navicular facet lies posterior to the most anterior border of discoid facet, in dorsal view; odontoid process and odontoid facet lateral to the navicular facet; Navicular facet projects medially more than in any other Megatheriinae; sulcus talis most mesial, and short; the odontoid process is 110° from the discoid facet.

ETYMOLOGY. — *Sibö*, Bribri word from Talamancan Mythology means good creator, and the Greek *Therios* for beast, meaning the beast of *Sibö*, in allusion of the first mammals appearing before the connection of the Americas. Cf. Jara Murillo, C. V. 2003. *Diccionario de Mitología Bribri*. Edition: 2.^a reimp.(2014), 336 pp.

TYPE SPECIES. — *Sibotherium ka* gen. et sp. nov.

Sibotherium ka gen. et sp. nov.

2012. Megatheriidae indeterminate. Laurito & Valerio (2012a)

DIAGNOSIS. — As for the genus

ETYMOLOGY. — *Ká* from Bribri mythology, which mean site, referred to the Coto Brus Valley, the site of *Sibö*, *Sibotherium ka*, means the Beast of the site of *Sibö*. In the Bribri Myth, *Sibö* created in the beginning the earth and main land, and all the animals and plants. The new Megatheriinae from Coto Brus Valley, in Costa Rica, Central America, symbolizes one of the first South American Heralds that appeared previous to the closure of the Panama Isthmus.

HOLOTYPE. — CFM-994, left astragalus.

Referred material— CFM-2607, posterior fragment of a right mandible, just with the posterior part of the alveolus for m4, and the base of the coronoid process; CFM-3731, most anterior part of a mandible with the most anterior part of alveolus for m1; fragments of molariforms with the occlusal surface, CFM-3634, 3000, 1532, 1530, 1447, 1858; fragments of molariforms, CFM-1531, 2604, 2606, 1696; CFM-1654, left astragalus, lacking the odontoid process, and the most dorsal part of the navicular facet; CFM-993, right astragalus lacking the odontoid process, navicular process, and sustentacular facet; CFM-3813, fragment of navicular process of right astragalus; CFM-995, right calcaneum without the sustentacular facet and the posterior half of tuber calcis; CFM-998, left tuber calcis of calcaneum; CFM-1243, fragment of calcaneus just with the octal facet; CFM-1239, proximal part of right fourth metatarsal; CFM-1301, distal part of fourth right metatarsal; CFM-2498, proximal part of fifth metatarsal; CFM-1645, left navicular; CFM-3058, right navicular of young individual; CFM-3795, clavicle; CFM-1840, proximal part of right tibia; CFM-1302, fragment of lateral trochlea of right tibia; CFM-2625, left magnum; CFM-1785, left lunar; CFM-3713, left lunar.

TYPE LOCALITY. — Curré Formation, San Gerardo de Limoncito, Costa Rica.

AGE. — Late Hemphillian (Hh3) (Late Miocene), Curré Formation, San Gerardo de Limoncito, Costa Rica. $^{87}\text{Sr}/^{86}\text{Sr}$ 5.80 Ma $\pm$ 0.60/-0.80 (Table 1)

Table 1

Age of Curré Formation, San Gerardo de Limoncito, Costa Rica.

Sample	Lab code	Type of material	$^{87}\text{Sr}/^{86}\text{Sr}$	1 sd*	2SE(M)*	n	Weight grams	Ma	$\pm$ Ma
CR-1	6058 AR CR	Oyster	0.709004	28	7	59	0.05857	5.80	0.60/-0.80
CR-2	6059 AR CR	Mollusc	0.708854	25	7	57	0.06235	11.15	1.55/-1.10

Sr isotope analyses were performed with a THERMO SCIENTIFIC TRITON PLUS Thermal Ionization Mass Spectrometer (TIMS).

Laboratory NBS987 standard: $0.710253 \pm 12^*$ n = 81

1 sd = 1 standard deviation; *) in the last two figures

$2 \text{ SE (M)} = 2\text{sd} / \text{root n}$

n = number of relationships measured per run

Ages calculated with: LOWESS 5 Fit 26 03 13

McArthur, J.M.; Howarth, R.J.; Shields, G.A. 2012. Strontium Isotope Stratigraphy. In: The Geologic Time Scale Vols 1 & 2 Pages: 127-144; Gradstein, FM; Ogg, JG; Schmitz, MD; et al. (Eds).

DESCRIPTION

Molariforms and mandible. — The molariforms (Fig. 3) in *Sibotherium* are slightly larger than in *Anisodontherium*; the valleys between the transverse crests in the molariforms of *Sibotherium* are shallow but V-shaped, as in *Pyramiodontherium*, *Megatheriops*, *Eremotherium* or *Megatherium*, but the *Anisodontherium* molariforms have a markedly deep V-shaped valley between the transverse crests, particularly in m1 (Brandoni & De Iuliis, 2007; Brandoni et al., 2012; Saint-André & De Iuliis, 2001).

The *Sibotherium* molariforms are mesiodistally compressed, and rectangular in section rather than squared, so they are 6 to 22 % wider than long, but they are less compressed mesiodistally than *Anisodontherium* (Brandoni et al., 2012), where are extremely compressed (13 to 37% wider than long), *Megathericulus patagonicus* (38% wider than long) (Brandoni et al., 2017), *Eomegatherium andinum*, or La Ventan Megatheriidae. This is in marked contrast to the nearly squared molariforms present in all other known megatheriines, such as *Pyramiodontherium*, *Megatheriops*, *Megatherium*, *Eremotherium*, *Plesiomegatherium hansmeyeri*, *Pliomegatherium lelongi* and *Proeremotherium* (Brandoni & De Iuliis, 2007; Carlini, Brandoni, Sánchez & Sánchez-Villagra, 2018). The teeth of *Sibotherium* have a V-shaped occlusal surface, but not so deep or highly rectangular as in *Anisodontherium* (Brandoni et al., 2012). In *Proeremotherium eljebe* M1–4 are essentially as wide as long, and M5 is longer than wide (Carlini et al., 2006), however, the measurements reported by Carlini et al. of the teeth reveals a tendency to be 3 to 38 % longer than wide, and differs from *Sibotherium* in which the teeth are wider than long and thus rectangular in shape.

The first upper molariform (CFM-1530) is trapezoidal in occlusal view, with the lingual and labial wall of the tooth oblique with respect to the midline of the tooth. The mesial loph is narrower and higher than the distal loph, which is longer and very low.

The M1-M2 (CFM-3000; CFM-3634), are slightly rectangular, or nearly quadrangular, 6.5 and 6.9 % wider than long, with the lophs equal in size and at the same level.

The first right lower molariform CFM-1858 of *Sibotherium* is trapezoidal, but rectangular (17% wider than long) in occlusal view, with the mesial or anterior loph narrow, highly oblique with respect to the lingual wall and slightly higher than the posterior loph, which is wider than the mesial, perpendicular to the lingual wall of the tooth, low than mesial, and both lophs are less sharp than in *Eremotherium laurillardi*. The lingual wall of the first molariform is parallel with respect to the midline of the tooth, but the labial wall is very oblique, and resembles the first lower tooth of *Eremotherium* and *Megatherium* (De Iuliis, 1996).

The second lower molariform of *Sibotherium* (CFM-1532 right) resembles the first lower molariform in its morphology, but the mesial loph and the labial wall of the tooth is less oblique than the first one. However, it is one of the most compressed teeth of *Sibotherium* being 22% wider than long, but is less compressed than *Anisodontherium* in which m2 is 32.14% wider than long (Brandoni et al., 2012).

The third lower molariform (CFM-1447 right) is rectangular in shape in occlusal view, 13.8 % wider than long, and less compressed than *Anisodontherium* in which is 34.6% wider than long.

In the anterior fragment of the mandible of *Sibotherium* (CFM-3731) the posterior margin of the mandibular symphysis lies well anterior (antero-posteriorly) to the first molariform, and approximately in the same position as in *Megathericulus patagonicus*, *Anisodontherium halmyronomum* and *Eomegatherium andinum* (according to the characters indicated by Brandoni et al., 2012). In other megatheriines for which this character is known, the symphysis ends more posteriorly than the first molariform, reaching at least to the level of the mesial margin of m1 and extending, as in *Eremotherium*, or *Megatherium americanum*, as far back as the middle of m2 (Pujos & Salas, 2004; Saint-André & De Iuliis, 2001). In *Pliomegatherium lelongi* and *Eremotherium laurillardi* the symphysis reaches at least at middle to the m1.

In *Plesiomegatherium hansmeyeri* the posterior border of the mandibular symphysis lies at the level of the alveolar septum between m1 and m2 (Brandoni, Carlini, Anaya, Gans & Croft, 2017), and in *Sibotherium* the symphysis is well anterior to m1. However, Pujos & Salas (2004) suggested that posterior end of mandibular symphysis in *Plesiomegatherium hansmeyeri* is anterior to m1.

In *Sibotherium* the end of the mandibular symphysis starts above the most ventral end of the root of m1, differing from *Eremotherium laurillardi* where the end of mandibular symphysis starts at the same level of the most ventral end of the m1.

In *Sibotherium* the mandibular symphysis is very vertically positioned with respect to the anterior border wall of m1 about 30°, as in *Pliomegatherium lelongi* (Brandoni, 2006, 2013), differing from *Eremotherium laurillardi* in which the




Fig. 3: Molariforms of *Sibotherium ka* gen. et sp. nov. in occlusal view A. CFM-1530, first upper molariform, B. CFM-1447, third lower molariform, C. CFM-1532 second lower molariform, and D. CFM-3000 first or second upper molariform, E. CFM-3634, first or second upper molariform; F. CFM-1858 first lower molariform), and labial views (A', B', C' and D', E', F').

mandibular symphysis is about 60° with respect to the anterior border wall of m1. Besides, in *Sibotherium* the mandibular symphyseal spout is shorter than in any Megatheriidae in which the mandible is preserved.

In *Megathericulus patagonicus*, *Pliomegatherium lelongi*, *Eremotherium laurillardi*, *Megatherium americanum*, *Megatherium altiplanicum*, *Anisodontherium* sp. the symphyseal spout is prominent and long (Brandoni, 2006, 2013; Brandoni et al., 2012; De Iuliis et al., 2008; Saint-André & De Iuliis, 2001).

In *Sibotherium* the m4 is posterior to the anterior margin of the ascending ramus, and not visible in lateral view. In *Anisodontherium* the m4 is anterior to anterior margin of ascending ramus, so that it is visible in lateral view (Brandoni et al., 2012).




Fig. 4: Fragments of the mandible of *Sibotherium ka* gen. et sp. nov. Anterior part in mesial (A), lateral (A'), and dorsal (A'') views. Posterior part in mesial (B), lateral (B') and dorsal (B'') views. **1Mf**, first molariform; **Sy**, mandibular symphysis; **PLO** posterolateral opening of mandible; **4Mf**, fourth molariform.

In *Sibotherium* the distal part of m4, in lateral view, is covered by the anterior margin of the ascending ramus (see Fig. 4.B), as also occurs in *Pl. hansmeyeri*, *Pyramiodontherium*, *Megatheriops*, and several Quaternary Megatheriinae (Brandoni & De Iuliis, 2007; Pujos & Salas, 2004).




Fig. 5: *Sibotherium ka* gen. et sp. nov. Holotype CFM-994. Left astragalus in A, anteromedial; B, fibular; C, dorsomedial; D, posteriomedial; E, ventral; and F, posterior, views. AEF, astragalar ectal facet; ASF, astragalar sustentacular facet; Cub, cuboid facet; DF, discoid facet; FF, fibular facet; Nav, navicular facet; OF, odontoid facet; OP, odontoid process; OT, odontoid tuberosity; ST, sulcus tali.

Astragalus. — The left astragalus of *Sibotherium* (CFM-994, Fig. 5) has the typical shape of Megatheriinae, with a well-developed and relatively central odontoid process in dorsomedial view, and a navicular facet in anterior view (*sensu* Brandoni, Carlini, Pujos & Scillato-Yané, 2004).

However, in the *Sibotherium* astragalus (CFM-994) the odontoid process is not completely centered, and the posterior part of the discoid facet, posterior to the odontoid process, is 50% smaller than the anterior part of that facet anterior to the odontoid process, a condition that is similar to the Megatheriidae from La Venta (Colombia) (UCMP 39593) reported and figured by Hirschfeld (1985), and reported in *Megathericulus patagonicus* (De Iuliis et al., 2008), but differs from *Urumaquia*, *Megatherium*, *Eremotherium*, *Pliomegatherium*, *Eomegatherium*, *Pyramiodontherium*, where the odontoid process is centered, and the discoid facet posterior and anterior parts to the odontoid process is more or less equivalent in longitude.

Regarding Hirschfeld (1985) the astragalus of *Megathericulus patagonicus* is smaller and has a less well-developed odontoid process than in the La Venta astragalus (UCMP 39593). The La Venta Megatheriidae astragalus (UCMP 39593) is similar in size to *Eomegatherium nanum* which is at least 120 mm long (Hirschfeld, 1985). *Sibotherium* has a well-developed odontoid process larger than that of the La Venta megatheres.

In *Sibotherium* the astragalus is wider than long (CFM-994; length = 153.2mm; wide = 185.6mm; CFM-1654, length = 146.7mm; wide = 159mm), differing from *Urumaquia* where the astragalus is longer than wide (length = 175; wide = 144mm), as in *Eremotherium laurillardi* (CFM-850 left, length = 265mm; wide = 257mm; CFM-849 right, length = 270; wide = 253; CFM-1871 right, length = 227mm; wide = 224mm) (Figs. 6, 7, 8).

In the *Sibotherium* astragalus (CFM-994), the fibular facet is divided in two main areas. As in other Megatheriinae, an anteroposteriorly elongated dorsal portion joined with the discoid facet dorsally, and a ventral portion extended ventrally and is not in contact with the ectal facet as in many Megatheriinae, except in *Megathericulus patagonicus* in which the ectal facet and the ventral portion of the fibular facet are in contact (*sensu* Brandoni, González & Bucher, 2019).

In *Sibotherium* (CFM-994) the angle between the odontoid and discoid facets in the astragalus is 110°, but in *Eremotherium* (CFM-1871 collected in Palmares, Costa Rica) it is 115°, and in *Urumaquia* it is approximately 100°, 90° in *Pyramiodontherium bergi* and *P. scilatoyanei*, and between 100° and 120° in *Megatherium* and *Eremotherium laurillardi*.

In the left astragalus of *Sibotherium* (CFM-994) the sulcus tali is short and deep, and in *Eremotherium* it is as wide as long, and the sulcus tali is longer and shallow.

In *Megathericulus patagonicus* the navicular facet lies dorsally, with the plane of the discoid facet intersecting it approximately at its middle or two third, as in *Megathericulus primaevus* Cabrera, 1939; in contrast, it is relatively more ventral in other megatheriines (Brandoni and Carlini, 2009; Brandoni et al., 2019; De Iuliis et al., 2008). The navicular facet is well dorsal to the plane of the discoid facet, that plesiomorphic feature distinguish *M. patagonicus*, *Urumaquia*, and *Sibotherium* from more derived megatheriine where the navicular facet is most ventrally located (De Iuliis et al., 2008; Brandoni, et al., 2019).

The astragalar depression is deep and conical with a very sharp apex. In *Sibotherium* (CFM-994), *Urumaquia* and *Megathericulus* nearly the middle of the navicular facet is dorsal with respect to the plane of the discoid facet, whereas in *Pyramiodontherium* spp., *Eremotherium laurillardi* and *Megatherium urbinai*, only one third is dorsal to this plane (Brandoni and Carlini, 2009). Carlini, Brandoni & Sánchez (2008a) suggest that the navicular facet in *Urumaquia* is one third dorsally with respect to the discoid facet, but in their figure 2B it is clear that in *Urumaquia* the navicular facet is 50% dorsally, as in *Megathericulus* or *Sibotherium*.

Sibotherium differs from *Urumaquia* in having the navicular process and the navicular facet completely to the posterior border of the discoid facet for the tibia, differing from *Urumaquia* in which both the navicular process and navicular facet are completely anterior to the anterior border of the discoid facet.

The odontoid tuberosity in *Sibotherium* is not raised as in *Megathericulus*, or *Pyramiodontherium*, but is most robust and developed than in *Eremotherium* or *Megatherium*, but smaller than in *Pyramiodontherium*, where it is very robust and highly developed and strongly projected medially raised (*sensu* Brandoni and Carlini, 2009; De Iuliis et al., 2008).

In *Sibotherium* the discoid facet and navicular facet are separated by a wide and deep channel more than in *Eremotherium eomigrans* or *E. laurillardi*, and the navicular process projects more medially than in any other Megatheriinae, giving the astragalus the appearance of having a neck. The odontoid process and the odontoid facet in *Sibotherium* are lateral to the navicular facet, differing from many other Megatheriinae (*Urumaquia*, *Megathericulus*, *Pyramiodontherium*, *Anisodontherium*, *Eomegatherium*, *Eremotherium*, and *Megatherium*) where the odontoid process is posterior to the medial plane of the navicular facet.




Fig. 6: *Sibotherium ka* gen. et sp. nov. CFM-1654. Left astragalus in A, anteromedial; B, fibular; C, dorsomedial; D, posteromedial; E, ventral; and F, posterior, views. **AEF**, astragalar ectal facet; **ASF**, astragalar sustentacular facet; **Cub**, cuboid facet; **DF**, discoide facet; **FF**, fibular facet; **Nav**, navicular facet; **ST**, sulcus tali.




Fig. 7: *Sibotherium ka* gen. et sp. nov. astragalus length by wide graph compare with different Megatheriidae.

In *Sibotherium* the ventral fibular facet of the astragalus is not connected with the ectal facet as in *Anisodontherium*, which contrasts with *Megathericulus patagonicus* where they are connected. (Brandoni & De Iuliis, 2007).

In *Sibotherium* the sulcus tali in the astragalus is short, narrow and deep and the most anterior part of the sulcus tali is more anteromesial than in *Eremotherium* where is more anterolateral, longer and shallow.

The sustentacular facet in *Sibotherium* is short, oval in shape, and its long axis is obliquely positioned with respect to the long axis of the ectal facet, and the *sulcus tali* is short and also obliquely positioned with respect to the lateral border of the ectal facet. In *Eremotherium* the sustentacular facet is short and triangular in shape, and its long axis is parallel with respect to the long axis of the ectal facet, and the sulcus tali is parallel with respect to the lateral border of the ectal facet.

CFM-995 corresponds to the right calcaneum without the sustentacular facet and the posterior half of the tuber calcis; CFM-998, left tuber calcis of calcaneum; CFM-1243, fragment of calcaneus just with the ectal facet. The calcaneum is short, robust and widened posteriorly as in *Pyramiodontherium scillatoyanei*, differing from the other megatheriines where this bone is longer and less robust (De Iuliis, Ré, & Vizcaíno, 2004).

DISCUSION

Phylogenetic approach

A new Xenarthra taxa is questionable or doubtful when that taxa is erected on single or very few isolated remains, partial or poorly preserve remains, and without consideration that the diagnostic characters fall within the range of known variation in other sloth species (De Iuliis, 2017).



Fig. 8: Hypothetical paleogeographical reconstruction of the Chortis Block, Isthmus of Panama and North West of South America at the Upper Miocene. SGL: San Gerardo de Limoncito Locality and site of the found of *Sibotherium ka* gen. et sp. nov. The Northern portion of Southern Central America probably looked more like a peninsula that allowed the arrival of North American fauna, while the Panamanian territory probably looked like an archipelago that when the sea level dropped or a tectonic rising was happening, or both situations coincided, allowed the xenarthrans to overcome the narrow oceanic passages, swimming or rafting. The map was made by one of the authors, Laurito based on the geological maps and other data, published by Albert, Val & Hoom, 2018, Alvarado & Cárdenes, 2016; Alcarado & Cárdenes, 2016; Alvarado et al., 2007; Coates, Aubry, Berggren, Collins & Kunk, 2003; Coates, Collins, Aubry & Berggren, 2004; Denyer & Alvarado (2007); Laurito & Valerio (2012a).

Even though the fossil material is fragmentary, the characters used to diagnose *Sibotherium* have been used by many authors, including Brandoni & De Iuliis (2007), Carlini et al. (2006, 2018), De Iuliis et al. (2008) and Pujos et al. (2017), among others. Who diagnosed other taxa under similar conditions, from characters that have been interpreted as discrete morphological characters.

For example, Brandoni et al. (2019) recognize the importance of the navicular facet depending on the astragalar discoid facet plane; the shape and proportion of the molariforms; the position of the posterior part of the mandibular symphysis respect to the first molariform. All of them, very important discrete morphological characters, used to diagnose basal or advance Megatheriidae, and have been used, depending on their degree of development, their absence or presence, to diagnose taxa. Likewise, De Iuliis et al., 2008 propose that the compression or not of the molariforms in Megatheriidae, as well as their relative position respect to the mandibular symphysis and the relative position of the fifth molariform respect to the anterior part of the ascending ramus of the mandible, or even the shape of the lophs on the occlusal surfaces are valid discrete characters to be used to diagnose taxa.

The new Megatheriidae from San Gerardo de Limoncito, *Sibotherium ka*, is erected on the base of few isolated and partially preserved remain, but we can recognize in those remains many diagnostic features where its unique combination resulting on the distinguishing of a new taxon.

For example, the astragalus of megatheriines is conservative, and few consistent morphologically diagnostic features at the specific level are recognized in this element in Plio- Pleistocene megatheriines (De Iuliis, 1996). The isolated astragali can rarely be confidently identified to species, because of general morphological similarities (De Iuliis, 1996). But, apparently for pre Plio-Pleistocene megatheriines (Middle and Late Miocene megatheriines) the astragalus has diagnostic value, and can be used to distinguish between taxa. For example, De Iuliis et al. (2008) used some astragalus features, combined with some other cranial and postcranial features in the revised diagnosis proposed for *Megathericulus patagonicus*, considered the earliest certain megatheriine, dated at 11.8 Ma (Mayoan – Laventan SALMA, Middle Miocene).

However, in Curré Formation 3 different astragali have been found belonging to the same taxon, and associated with cranial and other postcranial elements which allow us to name the new Megatheriinae taxon. It should be clarified that all specimens were recovered within the same outcrop, in an area of not more than 200 square meters, and they keep the same morphology and proportions.

Sibotherium ka shows two plesiomorphic characters of Megatheriinae, the molars mesiodistally compressed (wider than long), and the navicular facet of the astragalus dorsally located with respect to the discoid facet (Brandoni et al., 2019; De Iuliis 1996; De Iuliis et al., 2008; Pujos et al., 2013), which suggests that *Sibotherium* is a basal member of Megatheriinae as are *Megathericulus*, *Pyramiodontherium*, or *Anisodontherium*.

Probably another plesiomorphic character in Megatheriinae is the developed odontoid tuberosity, which is strongly developed in Miocene Pliocene Megatheriinae (*M. patagonicus* and *Pyramiodontherium bergi*), and poorly developed in advanced Megatheriinae (*Eremotherium* and *Megatherium*) (Brandoni et al., 2019; De Iuliis et al., 2008). The odontoid tuberosity in *Sibotherium* is well developed, but not as strongly as in *M. patagonicus* or *P. bergi*.

In *Sibotherium* the posterior end of the mandibular symphysis lies far anterior to the m1, as in *Megathericulus patagonicus*, *Anisodontherium halmyronomum*, and *Eomegatherium andinum*. This is considered by De Iuliis et al. (2008) as another plesiomorphic character opposed to the more posterior position of the symphysis with respect to the m1 in more derived megatheriines (e.g. *Eremotherium*, *Megatherium*) (De Iuliis, et al., 2008), but the position of the posterior end of the mandibular symphysis in *Pyramiodontherium brevirostrum* reaches the plane of the alveolar septum between m1 and m2. That character has been treated as plesiomorphic by De Iuliis (1996).

Pujos et al., (2013) suggested that *Megathericulus* is a small-sized Megatheriinae (Megatheriinae half the size of *Megatheriops* or *Pyramiodontherium*), and *Eomegatherium nanum* is a middle-sized Megatheriinae. However, De Iuliis et al. (2004) suggest that whereas *Pyramiodontherium scillatoyanei* is a medium-sized Megatheriinae, with a body mass of approximately 2,500 kg, they considered *Pyramiodontherium bergi*, *Megatherium americanum*, *Eremotherium laurillardi*, and *Eremotherium eomigrans* as large-sized Megatheriinae. Farina et al. (1998) reported a body mass of *Megatherium americanum* in a range of 6,073 kg and 2,745 kg. Carlini et al. (2006) suggested that *Urumaquia robusta* is a large-sized Megatheriinae. The astragalus of *Sibotherium ka* is similar in size of *Pyramiodontherium bergi*, *Urumaquia robusta*, or *Megatherium lundii*, which allow us to suggest that *Sibotherium* is a large size Megatheriinae.

Biogeographic and chronological implications

Until now the earliest record of a Megatheriinae in North America, after the formation of the Panamanian Land Bridge, is *Eremotherium eomigrans* from Haile 7C, latest Blancan (probably between 2.2 and 1.7 Ma), located approximately 6 km NE of Newberry, Alachua County, Florida, USA, and apparently was among the earliest mammalian immigrants to North America (De Iuliis & Cartelle, 1999).

Then San Gerardo de Limoncito ground sloth *Sibotherium ka* from Costa Rica, represents the first Megatheriidae-Megatheriinae fossil record in Central and North America with a radiometric age ($^{87}\text{Sr}/^{86}\text{Sr}$) of 5.80 Ma $\pm$ 0.60/-0.80, and apparently before the closure of the Isthmus of Panama around 2.8 Ma according to the O'Dea et al. (2016) hypothesis.

However, the earliest certain late Tertiary mammalian immigrant in North America from South America is the Megalonychidae *Pliometanastes* from California and Florida at 8 Ma (Hirschfeld & Webb, 1968; Morgan, 2005), and the Mylodontidae *Thinobadistes*, known from Florida and Texas, probably with the same age (Webb, 1989). A southern record of *Pliometanastes* has been reported from Juchipila, El Resbalón, Mexico with 5.59 Ma (Carranza-Castañeda, Aranda-Gómez, Wang & Iriando, 2013), and from San Gerardo Limoncito (Late Miocene), Costa Rica (Laurito & Valerio, 2012b), with 5.8 Ma old.

The closure of the Panama seaway around 2.8 Ma ago implied that the first dispersal event of *Pliometanastes* and *Thinobadistes*, with 8 Ma in North America, must have immigrated by a water route, as island hoppers (Hirschfeld, 1985) or rafting immigrants (Laurito and Valerio, 2012a), with the assumption that ground sloths were durable swimmers and adapted to live in a wide variety of environments (Webb, 1997).

After 2.8 Ma ago the Isthmus of Panama originated as a landmass, so that fully walking mammals and birds (35 families of mammals and one of bird were involved in the GABI) began dispersal across the Isthmus about 2.4 Ma (Webb, 1976,

1997). Marshall (1988) stated that the first great contingent of South American immigrants appeared in North America near the end of the Blancan at about 2.6 to 2.8 Ma. The “walking” immigrants included two ground sloths, *Eremotherium* (De Iuliis & Cartelle, 1999) and *Glossotherium* (Robertson, 1976), armadillos, pampatheres, glyptodonts, capybaras, porcupines and a phorusrhacid bird. The peak of the interchange probably occurred sometime in the Irvingtonian NALMA, but continued into the Rancholabrean.

If we follow the hypothesis of 2.8 Ma for the closure of the Panama seaway, then the ancestor of the *Sibotherium ka*, from Costa Rica, was dispersed swimming or rafting through the Panama seaway from South America to the ancient Costa Rica terrestrial land mass.

Another hypothesis is that the Panama seaway was closed during middle Miocene times (Montes et al., 2015). A variation of that hypothesis implies that over the past 10 Ma there has been substantial dispersal and vicariance, maybe associated with landscape formation, volcanism, climate change, and/or sea level fluctuations, across the Isthmus of Panama and that pulses of dispersal of terrestrial organisms occurred over at least three periods during the last 30 Ma (Bacon et al., 2015).

If we follow this hypothesis of an early closure of the Panama seaway and origin of Isthmus of Panama, it implies that the Xenarthra dispersed to North and Central America walking by a terrestrial route, and not swimming or rafting.

The sharks and crocodyles as predators reported in the Curré Formation could represent an ecological barrier for the swimming dispersal route for terrestrial mammals (e. g. Giant Ground Sloths) in the case that Costa Rica and Panama were separate at 5.8 million years old.

There is evidence that some turtle shells and mammalian long bones in the Curré Formation from San Gerardo de Limoncito bone bed deposit, Costa Rica, are broken, some with fractured edges that are slightly abraded, indicating breakage prior to fluvial transport. Numerous bones of turtle, horse, and camel exhibit conical marks, which appear to be punctures and are consistent with those produced by isodont teeth. This suggests that many of the turtles, horses and camel could be the result of prey breakage by crocodilians, which have been reported in that formation (*sensu* Mead et al., 2006). This implies that those terrestrial mammals were living near water inhabited by crocodiles.

Also in the conglomeratic facies of the subaquatic Fan Delta of the Curré Formation in San Gerardo de Limoncito Costa Rica has been reported 7 Chondrichthyes taxa: *Isurus desori*, *Hemipristis serra*, *Carcharhinus priscus*, *Carcharhinus longimanus*, *Isogomphodon acuarius*, *Isogomphodon caunellensis*, *Sphyrna arambourgi* (Laurito & Valerio, 2008). However, there is no evidence of bite marks on terrestrial bones of mammals, but the sharks reported here probably represent another potential predator of terrestrial mammals in the Curré Formation about 5.8 million years ago.

On the other hand, stable isotopes of carbon and oxygen on the enamel fossils of the extinct equids *Calippus hondurensis*, *Dinohippus mexicanus*, and *Protohippus gidleyi*, the gomphothere *Gomphotherium hondurensis*, and the llama *Hemiauchenia vera* of the Early–Late Hemphillian (Hh3) from San Gerardo de Limoncito, Puntarenas province, Costa Rica, suggest that these mammals fed mainly on C3 plants and lived in clearings rainforests (Pérez-Crespo et al., 2018).

A probable paleogeographical scenario for migration

One hypothesis considered the possibility that a species morphologically close to *Proeremotherium eljebe* migrated into North America and gave rise to the early species of *Eremotherium* (i.e., *E. eomigrans* from the Blancan, latest Pliocene middle Pleistocene (Carlini et al., 2006, 2008a). That hypothesis has been supported by the presence of *Proeremotherium* sp. AMU- CURS 184 from the San Gregorio Formation, Urumaco Venezuela, which seems to be morphologically intermediate between *Proeremotherium eljebe* and *Eremotherium eomigrans*, and it was argued that after the cladogenetic process of *Eremotherium* giving rise to *E. laurillardi* in the Rancholabrean (Late Pleistocene) of North America, *E. laurillardi* migrated to South America and colonized most of the north and central lowlands of South America (Carlini et al., 2008a).

That biogeographic scenario of northern megatheriines, also has proposed for pampatheres (Scillato-Yané et al., 2005), glyptodonts (Carlini et al., 2008b; Carlini and Zurita, 2010; Zurita, Oliveira, Toriño & Krapovickas, 2011) and dasypodids (Castro, Carlini, Sánchez & Sánchez-Villagra, 2014), and assuming a migration from South America during mid/late Pliocene, dispersion and diversification into North America and re-ingression into South America during the latest Pleistocene.

However, *Proeremotherium* and *Eremotherium* are evidently close morphologically, and may be phylogenetically, but there is no evidence that *Proeremotherium* migrated to North America and diversified to *Eremotherium* and re-entered South America.

Also, the earliest records of *Eremotherium* in South America come from El Breal de Orocuál, which according to the faunal assemblage probably represent a late Pliocene-early Pleistocene timing (Rincón, Parra, Prevosti, Alberdi & Bell, 2009; Rincón, Prevosti, & Parra, 2011), that means that *Eremotherium* had been present in Northern South American environment in the late Pliocene/early Pleistocene, in opposition with the *Eremotherium* late Pleistocene re-entered hypothesis of Carlini et al. (2006, 2008a). The record of *Scirrotherium antelucanus* (Pampatheriidae) (Laurito & Valerio, 2013) and *Sibotherium ka* (Megatheriinae) from San Gerardo de Limoncito, Costa Rica, implies a much earliest presence of the ancestors of these taxa before the closure of Isthmus of Panama 5.8 Ma ago, in the late Miocene, and not in the middle-late Pliocene as Carlini et al. (2018) proposed.

The unknowns posed by the fossil record of mammals at the Upper Miocene of southern Costa Rica, basically are: the dispersion route that these faunas followed; the apparent absence of North American mammals in sediments of similar age in South America (beyond doubt); the apparent northward advanced migration of xenarthrans; and the geological viability of a continuous corridor between the Americas during the late Miocene.

The San Gerardo de Limoncito locality is just over 800 km from the northwestern corner of South America. Its fossil fauna, especially the mammal assemblage, constituted by large beasts and small ones, are indisputable evidence of an exchange between the Americas as early as the Late Miocene (5.8 Ma) and the existence of a terrestrial corridor, if not permanent at least temporary.

The fauna of North American ancestry is represented by the horses *Calippus hondurensis*, *Dinohippus mexicanus* and *Protohippus gidleyi*, the peccaries *Prosthennops serus* and *Protherohyus brachydontus* (Laurito & Valerio, 2010; Valerio & Laurito, 2020), the mastodon *Gomphotherium hondurensis* and the lamini camel *Hemiauchenia vera* (Valerio & Laurito, 2008; Laurito & Valerio, 2016). But there must be included within this migratory group *Pliometanastes*, a Xenarthra derived from a South American ancestor (Morgan, 2008; Laurito & Valerio, 2012b), which arrived in North America as early as 8.54 Ma (Hirschfeld, 1981), during the Early Hemphillian (Hh-1) and its fossil record in the late Miocene of Southern Central America could be interpreted as an example of reverse migration.

The South American fauna is represented by the Pampatheriidae *Scirrotherium antelucanus* Laurito & Valerio, 2013 and the Megatheriidae *Sibotherium ka*. These two species pose two possible migration scenarios: ones in which these animals were able to swim short distances, saving the sea gaps, and a second one, where indeed the land corridor was established for a short period of time either as a product of the tectonic uplift or a drop in sea level or a mixture of both.

Of the two scenarios, the first is the most difficult to demonstrate due to the lack of evidence, apart from the fact that their remains were found in coastal sediments. But the second one is much more likely.

The “widespread of neritic sediments throughout the Darien region during the late Miocene, and the presence of a regional unconformity suggests the completion of the Central American Arc collision with South America by 7.1 Ma”. After that, “no Pliocene deposits are recorded from either the Darien or the Panama Canal Basin, and no sediments younger than 4.8 Ma have been identified in the Atrato Basin of Colombia, suggesting rapid uplift and extensive emergence of the Central American isthmus in the latest Miocene” (Coates et al., 2004). This evidences a probable partial land corridor between the Northwest of Colombia, the Atrato Basin and the complexly folded and faulted synclinorium that forms the central Chucunaque-Tuira Basin of the Darien (cf. Coates et al., 2004).

On the other hand, a more or less continuous volcanic arc was developed since the Middle Miocene until the Pleistocene (De Boer, Defant, Stewart, & Bellon, 1991; MacMillan, Gans & Alvarado, 2004), which extended from the southern half of Costa Rica to immediately north of the Panama Canal basin.

This scenario allows hypothesizing the existence of long and narrow emerged territories linked to the subcontinents and interrupted by narrow and relatively deep and short interoceanic marine passages, bordered by small islands.

In other words, the presence of North American mammals in southern Costa Rica is sufficient evidence to assume the existence of a continuous land corridor between Southern Central America and the Chortis Block and therefore with North America.

The last element to complete the recipe are the oscillations of the sea level and Zachos, Pagani, Sloan, Thomas & Billups (2001) explain that “a gradual cooling and reestablishment of a major ice-sheet on Antarctica became by 10 Ma and the mean $\delta^{18}O$ values then continued to rise gently through the late Miocene until the early Pliocene (6 Ma), indicating additional cooling and small-scale ice-sheet expansion on West-Antarctica and in the Arctic”.

Extrapolating, we can assume that the continuous, but not so obvious drop of sea level, along the late Miocene, along with the continued tectonic uprising, driven by the subduction, and the tectonic coupling of the island arc, produced the temporary closure and narrowing of some marine passages at the south of the Panama Canal Basin and northwestern Colombian territories,

which allowed the migration of some xenarthrans and possibly additional biota, and this scenery could also explain the later biota exchanges and their episodic pattern, by way of migratory pulses prior to the definitive establishment of the land corridor.

Finally, it is possible that the climatic continuity between the south of Costa Rica, Panama and the Northwest of Colombia, facilitated the migration of South American elements and discouraged the migration of North American mammals adapted to more open forest and dry conditions. This is evidenced from the analysis of the $\delta^{13}\text{C}$ in the dental enamel from the horses, camels and mastodons associated with *Sibotherium ka* in SGL, where all of them fed on C3 plants, and it suggests that during the Late-early Hemphillian the site was more humid than today, thus favoring the presence of C3 plants. Together with the $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ values of the dental enamel in these fossils, the paleovegetation reconstructions made for this geological period suggest that these herbivorous mammals lived in a tropical rainforest, but in clearings of the forest where they fed on plants that developed there (*sensu* Pérez-Crespo et al., 2018).

ACKNOWLEDGEMENTS

Thanks to the Dr. Peter Schaaf and the Laboratorio Universitario de Geoquímica Isotópica (LUGIS) of the Universidad Nacional Autónoma de México (UNAM), to the staff who intervened in obtaining data: To Teodoro Hernández Treviño for the mechanical preparation of the samples. To Gabriela Solís Pichardo for performing isotopic analytical work in the ultrapure chemistry laboratory, data reduction and age calculation. To Gerardo Arrieta García for the isotopic measurements. To LUGIS for the partial funding of the analysis.

We thank to Dr. Spencer Lucas, New Mexico Museum of Natural History and Science for his critical reading of the text and his valuable comments, which helped improve this paper.

MATERIAL EXAMINATE TO THIS STUDY

Eremotherium eomigrans astragalus UF 46611, Alachua Co., Haile XVI, Florida, USA.

Eremotherium larillardi Astragalus AMNH 95753, San Miguel Lara state, Venezuela, Astragalus CFM-850, San Ramón, Costa Rica.

Eremotherium sp. Astragalus OR-1330, El Breal de Orocuál, ORS-16, and Astragalus OR-1331, El Breal de Orocuál, ORS-16, Monagas state, Venezuela.

Urumaquia robusta astragalus MCNC-74-72V (Holotype), Urumaco, Falcon state, Venezuela.

REFERENCES

- Albert, J. S., Val, P. & Hoorn, C. (2018). The changing course of the Amazon River in the Neogene: center stage for Neotropical diversification. *Neotropical Ichthyology*, 16(3): e180033. doi: 10.1590/1982-0224-20180033
- Alvarado, G. E. & Cárdenes, G. (2016). Geology, tectonics, and geomorphology of Costa Rica: A natural history approach. In M. Kapelle (Ed.), *Costa Rican Ecosystems* (pp.30-63). Chicago & London: The University of Chicago Press.
- Alvarado, G. E., Dengo, C., Martens, U., Bundschuh, J., Aguilar, T. & Bonis, S. B. (2007). Chapter 13: Stratigraphy and geologic history. In J. Bundschuh and G. E. Alvarado, (Eds.), *Central America: Geology, Resources and Hazards* (pp. 345-394). London: Taylor and Francis.
- Ameghino, F. (1904). Nuevas especies de mamíferos cretáceos y terciarios de la República Argentina. *Anales de la Sociedad Científica Argentina*, 58, 225-291.

- Bacon, C. D., Silvestro, D., Jaramillo, C., Tilston Smith, B., Chakrabarty, P. & Antonelli, A. (2015). Biological evidence supports an early and complex emergence of the Isthmus of Panama. *Proceeding of the National Academy of Sciences*, 112(24), 6110-6115.
- Brandoni, D. (2006). A review of *Pliomegatherium* Kraglievich, 1930 (Xenarthra: Phyllophaga: Megatheriidae). *Neues Jahrbuch Geologie Paläontologie*, 2006(4), 212-224.
- Brandoni, D. (2013). Los Tardigrada (Mammalia, Xenarthra) del Mioceno Tardío de Entre Ríos, Argentina. In D. Brandoni & J. I. Noriega. (Eds.), *El Neógeno de la Mesopotamia argentina* (pp. 135-144). Argentina: Asociación Paleontológica Argentina, Publicación Especial 14.
- Brandoni, D. & De Iuliis, G. (2007). A new genus for the Megatheriinae (Xenarthra, Tardigrada, Megatheriidae) from the Arroyo Chasicó Formation (Upper Miocene) of Buenos Aires Province, Argentina. *Neues Jahrbuch für Geologie und Paläontologie Abhandlungen*, 244, 53-64.
- Brandoni, D. & Carlini, A. A. (2009). On the presence of *Pyramiodontherium* (Mammalia, Tardigrada, Megatheriidae) in the "Conglomerado osífero" (late Miocene) of Argentina and its biogeographical implications. *Rivista Italiana di Paleontologia e Stratigrafia*, 115, 111-123.
- Brandoni, D., Powell, J. E. & González, O. E. (2012). *Anisodontherium* from the Late Miocene of north-western Argentina. *Acta Palaeontologica Polonica*, 57(2), 241-249.
- Brandoni, D., González Ruiz, L. & Bucher, J. (2019). Evolutive Implications of *Megathericulus patagonicus* (Xenarthra, Megatheriinae) from the Miocene of Patagonia Argentina. *Journal of Mammalian Evolution*. doi: 10.1007/s10914-019-09469-6
- Brandoni D., Carlini, A. A., Pujos, F. & Scillato-Yané, G. J. (2004). The pes of *Pyramiodontherium bergi* (Moreno & Mercerat, 1891) (Mammalia, Xenarthra, Phyllophaga): the most complete pes of a Tertiary Megatheriinae. *Geodiversitas*, 26(4), 643-659.
- Brandoni, D., Carlini, A. A., Anaya, F., Gans, P. & Croft, D. A. (2017). New remains of *Megathericulus patagonicus* Ameghino, 1904 (Xenarthra, Tardigrada) from the Serravallian (Middle Miocene) of Bolivia: chronological and biogeographical implications. *Journal of Mammalian Evolution*. doi: 10.1007/s10914-017-9384-y
- Carlini, A. A. & Zurita, A. E. (2010). An introduction to cingulate evolutionary history during the Great American Biotic Interchange: biogeographical clues from Venezuela. In M. R. Sánchez-Villagra, O. A. Aguilera & A. A. Carlini. (Eds.), *Urumaco and Venezuelan Paleontology - the Fossil Record of the Northern Neotropics* (pp. 233-255). Bloomington and Indianapolis, USA: Indiana University Press.
- Carlini, A. A., Brandoni, D. & Sánchez, R. (2006). First megatheriines (Xenarthra, Phyllophaga, Megatheriidae) from the Urumaco (late Miocene) and Codore (Pliocene) formations, Estado Falcón, Venezuela. *Journal of Systematic Palaeontology*, 4, 269-278.
- Carlini, A. A., Brandoni, D. & Sánchez, R. (2008a). Additions to the knowledge of *Urumaquia robusta* (Xenarthra, Phyllophaga, Megatheriidae) from the Urumaco Formation (Late Miocene), Estado Falcón, Venezuela. *Palaeontologische Zeitschrift*, 82, 153-162. doi: 10.1007/BF02988406
- Carlini, A. A., Brandoni, D. & Molin, C. N. D. (2013). A new genus and species of Planopinae (Xenarthra: Tardigrada) from the Miocene of Santa Cruz Province, Argentina. *Zootaxa*, 3694 (6), 565-578.

- Carlini, A. A., Brandoni, D., Sánchez, R. & Sánchez-Villagra, M. R. (2018). A new Megatheriinae skull (Xenarthra, Tardigrada) from the Pliocene of Northern Venezuela — implications for a giant sloth dispersal to Central and North America. *Palaeontologia Electronica*, 21.2(16A), 1-12. doi: 10.26879/771
- Carlini, A. A., Zurita, A. E., Scillato-Yané, G. J., Sánchez, R. & Aguilera, O. A. (2008b). New glyptodont from the Codore Formation (Pliocene), Falcón State, Venezuela, its relationship with the *Asterostemma* problem, and the paleobiogeography of the Glyptodontinae. *Palaeontologische Zeitschrift*, 82, 139-152. doi: 10.1007/BF02988405
- Carranza-Castañeda, O., Aranda-Gómez, J. J., Wang, X. & Iriondo, A. (2013). The Early-Late Hemphillian (Hh2) faunal assemblage from Juchipila Basin, State of Zacatecas, Mexico, and its biochronologic correlation with other Hemphillian faunas in Central Mexico. *Contributions in Science*, 521, 13-49.
- Castro, M. C., Carlini, A. A., Sánchez, R. & Sánchez-Villagra, M. R. (2014). A new Dasypodini armadillo (Xenarthra: Cingulata) from San Gregorio Formation, Pliocene of Venezuela: affinities and biogeographic interpretations. *Naturwissenschaften*, 101, 77-86. doi: 10.1007/s00114-013-1131-5
- Chafetz, H. S. & Reid, A. (2000). Syndepositional shallow-water precipitation of glauconite minerals. *Sedimentary Geology*, 136, 29-42.
- Coates, A. G., Collins, L. S., Aubry, M-P. & Berggren, W. A. (2004). The geology of the Darien, Panama, and the late Miocene-Pliocene collision of the Panama arc with northwestern South America. *Geological Society of America Bulletin*, 116, 1327-1344.
- Coates, A. G., Aubry, M-P, Berggren, W. A., Collins, L. S. & Kunk, M. (2003). Early Neogene history of the Central American arc from Bocas del Toro, western Panama. *Geological Society of America Bulletin*, 115, 271-287. doi: 10.1130/0016-7606(2003)115<0271:ENHOTC>2.0.CO;2
- Croft, D. A., Carlini, A. A., Ciancio, M. R., Brandoni, D., Drew, N. E., Engelman R. K. & Anaya, F. (2016). New mammal faunal data from Cerdas, Bolivia, a middle-latitude Neotropical site that chronicles the end of the Middle Miocene Climatic Optimum in South America, *Journal of Vertebrate Paleontology*. doi: 10.1080/02724634.2016.1163574
- De Boer, J. Z., Defant, M. J., Stewart, R. H. & Bellon, H. (1991). Evidence for active subduction below western Panama. *Geology*, 19, 649-652.
- De Iuliis, G. (1996). *A Systematic Review of the Megatheriinae (Mammalia: Xenarthra: Megatheriidae)* (Unpublished Ph.D. Dissertation). University of Toronto, Toronto, Canada.
- De Iuliis, G. (2017). Recent Progress and Future Prospects in Fossil Xenarthran Studies, with Emphasis on Current Methodology in Sloth Taxonomy. *Journal of Mammalian Evolution*. doi 10.1007/s10914-017-9407-8
- De Iuliis, G. & Saint-André, P. A. (1997). *Eremotherium sefvei* nov. sp. (Mammalia, Xenarthra, Megatheriidae) from the Pleistocene of Ulloma, Bolivia. *Geobios*, 30(3), 453-461.
- De Iuliis, G. & Cartelle, C. (1999). A new giant megatheriine ground sloth (Mammalia: Xenarthra: Megatheriidae) from the late Blancan to early Irvingtonian of Florida. *Zoological Journal of the Linnean Society*, 127, 495-151.
- De Iuliis, G., Ré, G. H. & Vizcaíno, S. F. (2004). The Toro Negro megatheriine (Mammalia: Xenarthra): a new species of *Pyramiodontherium* and a review of *Plesiomegatherium*. *Journal of Vertebrate Paleontology*, 24(1), 214-227.

- De Iuliis, G., Brandoni, D. & Scillato-Yané, G. J. (2008). New remains of *Megathericulus patagonicus* Ameghino 1904 (Xenarthra, Megatheriidae): information on primitive features of megatheriines. *Journal of Vertebrate Paleontology*, 28, 181-196.
- Denyer, P. & Alvarado, G. E. (2007). *Mapa Geológico de Costa Rica (1: 400 000)*. San José, Costa Rica: Librería Francesa S.A.
- Fariña, R. A., Vizcaíno, S. F. & Bargo, M. S. (1998). Body mass estimations in Lujanian (late Pleistocene-early Holocene of South America) mammal megafauna. *Mastozoología Neotropical*, 5, 87-108.
- Gaudin, T. J. (1995). The ear region of edentates and the phylogeny of the Tardigrada (Mammalia, Xenarthra). *Journal of Vertebrate Paleontology*, 15, 672-705. doi: 10.1080/02724634.1995.10011255
- Gaudin, T. J. (2004). Phylogenetic relationships among sloths (Mammalia, Xenarthra, Tardigrada): the craniodental evidence. *Zoological Journal of the Linnean Society*, 140, 255-305. doi: 10.1111/j.1096-3642.2003.00100.x
- Granados, R. & Aguilar, T. (1983). Geopaleontología de la cuenca baja del río Savegre Costa Rica. *Brenesia*, 21, 229-260.
- Henningsen, D. (1965). *La Fila Costeña del Pacífico y su posición dentro del sistema montañoso centroamericano meridional*. San José: Ministerio de Industria y Comercio, y Dirección de Geología Minas y Petróleo.
- Hirschfeld, S. E. (1981). *Pliometanastes protistus* (Edentata, Megalonychidae) from Knight's Ferry, California, with discussions of early Hemphillian megalonychids. *PaleoBios*, 36, 1-16.
- Hirschfeld, S. E. (1985). Ground sloths from the Friasian La Venta fauna, with additions to the Pre-Friasian Coyaima fauna of Colombia, South America. *University of California Publications, Geological Sciences*, 128, 1-91.
- Hirschfeld, S. E. & Webb, S. D. (1968). Plio-Pleistocene megalonychid sloths of North America. *Bulletin of the Florida State Museum, Biological Sciences*, 12, 213-296.
- Jara, C. (2003). *Diccionario de mitología bribri*. San José, Costa Rica: Editorial Universidad de Costa Rica.
- Laurito, C. A. & Valerio, A. L. (2008). Ictiofauna de la localidad de San Gerardo de Limoncito, Formación Curré, Mioceno Superior, cantón de Coto Brus, provincia de Puntarenas, Costa Rica. *Revista Geológica de América Central*, 39, 65-85.
- Laurito, C. A. & Valerio, A. L. (2010). *Los Caballos Fósiles de la Formación Curré, Cantón de Coto Brus, Costa Rica*. San José, Costa Rica: Museo Nacional de Costa Rica.
- Laurito, C. A. & Valerio, A. L. (2012a). Paleobiogeografía del arribo de mamíferos suramericanos al sur de América Central de previo al gran intercambio biótico americano: Un vistazo al GABI en América Central. *Revista Geológica de América Central*, 46, 123-144.
- Laurito, C. A. & Valerio, A. L. (2012b). Primer registro fósil de *Pliometanastes* sp. (Mammalia, Xenarthra, Megalonychidae) para el Mioceno Superior de Costa Rica, América Central. Una Nueva Pista en la Comprensión del Pre-GABI. *Revista Geológica de América Central*, 47, 95-108.
- Laurito, C. A. & Valerio, A. L. (2013). *Scirrotherium antelucanus*, una nueva especie de Pampatheriidae (Mammalia, Xenarthra, Cingulata) del Mioceno Superior de Costa Rica, América Central. *Revista Geológica de América Central*, 49, 45-62.

- Laurito, C. A. & Valerio, A. L. (2016). Camellos Laminos del Mioceno Tardío (Hemifiliano Temprano) de la Formación Curré, San Gerardo de Limoncito, cantón de Coto Brus, provincia de Puntarenas, Costa Rica. *Revista Geológica de América Central*, 54, 7-55. doi: 10.15517/rgac.v54i0.21148
- MacMillan, I., Gans, P. B. & Alvarado, G. (2004). Middle Miocene to present plate tectonic history of the southern Central American Volcanic Arc. *Tectonophysics*, 392 (1-4), 325-348. doi: 10.1016/j.tecto.2004.04.014
- Marshall, L. G. (1988). Land Mammals and the Great American Interchange. *American Scientist*, 76(4), 380-388.
- McDonald, H.G. (2003). Xenarthran skeletal anatomy: primitive or derived? (Mammalia, Xenarthra). In: Fariña R.A., Vizcaino S.F., & Storch G. (Eds), Morphological studies in fossil and extant Xenarthra (Mammalia). *Senckenbergiana Biologica*, 83, 5-17.
- Mead, J., Cubero, R., Valerio, A. L., Swift, S., Laurito, C. A. & Gómez, L. (2006). Plio-Pleistocene *Crocodylus* (Crocodylia) from southwestern Costa Rica. *Studies on Neotropical Fauna and Environment*, 41(1), 1-7.
- Montes, C., Cardona, A., Jaramillo, C., Pardo, A., Silva, J. C., Valencia, V., ...Niño, H. (2015). Middle Miocene closure of the Central American Seaway. *Science*, 348(6231), 226-229.
- Mora, S. (1979). *Estudio geológico de una parte de la región sureste del Valle del General, provincia de Puntarenas, Costa Rica* (Unpublished Lic. Thesis). Universidad de Costa Rica, San José, Costa Rica.
- Morgan, G. S. (2005). The Great American Biotic Interchange in Florida. *Bulletin of Florida Museum of Natural History*, 45(4), 271-311.
- Morgan, G. S. (2008). Vertebrate fauna and geochronology of the Great American Biotic Interchange in North America. In S. Lucas, G. Morgan, J. Spielmann & D. Prothero. (Eds.), *Bulletin New Mexico Museum of Natural History & Science* (44, 93-140). Albuquerque: New Mexico Museum of Natural History.
- O'Dea, A., Lessios, H. A., Coates, A. G., Eytan, R. I., Restrepo-Moreno, S. A., Cione, A. L. & Jackson, J. B. C. (2016). Formation of the Isthmus of Panama. *Science Advances*, 2(8), 1-11, e1600883. doi: 10.1126/sciadv.1600883
- Pérez-Crespo, V. A., Laurito, C. A., Arroyo-Cabrales, J., Valerio, A. L., Morales-Puente, P., Cienfuegos-Alvarado, E. & Otero, F. J. (2018). Feeding habits and habitat of herbivorous mammals from the Early-Late Hemphillian (Miocene) of Costa Rica. *Acta Palaeontologica Polonica*, 63(4), 645-652.
- Pujos, F. & Salas, R. (2004). A new species of *Megatherium* (Mammalia: Xenarthra: Megatheriidae) from the Pleistocene of Sacaco and Tres Ventanas, Peru. *Palaeontology*, 47, 579-604.
- Pujos, F. & De Iuliis, G. (2007). Late Oligocene Megatherioidea fauna (Mammalia: Xenarthra) from Salla-Luribay (Bolivia): new data on basal sloth radiation and Cingulata-Phyllophaga split. *Journal of Vertebrate Paleontology*, 27, 132-144.
- Pujos, F., Salas-Gismondi, R., Baby, G., Baby, P., Goillot, C., Tejada, J. & Antoine, P.O. (2013). Implication of the presence of *Megathericulus* (Xenarthra: Tardigrada: Megatheriidae) in the Laventan of Peruvian Amazonia. *Journal of Systematic Palaeontology*, 11, 973-991. doi: 10.1080/14772019.2012.743488
- Rincón, A. D., Parra, G. E., Prevosti, F. J., Alberdi, M. T. & Bell, C. J. (2009). A preliminary assessment of the mammalian fauna from the Pliocene-Pleistocene El Breal de Orocuál locality, Monagas State, Venezuela. *Museum of Northern Arizona Bulletin*, 64, 593-620.

- Rincón, A. D., Prevosti, F. J. & Parra, G. E. (2011). New saber-toothed cat records (Felidae: Machairodontinae) for the Pleistocene of Venezuela, and the Great American Biotic Interchange. *Journal of Vertebrate Paleontology*, 31, 468-478. doi: 10.1080/0272463 4.2011.550366
- Robertson, J. S. (1976). Latest Pliocene Mammals from Haile XVA, Alachua County, Florida. *Bulletin of Florida Museum, Biological Science*, 20, 111-186.
- Saint-André, P. A. & De Iuliis, G. (2001). The smallest and most ancient representative of the genus *Megatherium* Cuvier, 1796 (Xenarthra, Tardigrada, Megatheriidae), from the Pliocene of the Bolivian Altiplano. *Geodiversitas*, 23(4), 625-645.
- Scillato-Yané, G. J., Carlini, A. A., Tonni, E. P. & Noriega, J. I. (2005). Palaeobiogeography of the late Pleistocene pampatheres of South America. *Journal of South American Earth Sciences*, 20, 131-138. doi: 10.1016/j.jsames.2005.06.012
- Seilacher, A. (1967). Bathymetry of trace fossil. *Marine Geology*, 5, 413-429.
- Valerio, A. L. (2010). *Paleontología, bioestratigrafía y paleoecología de los caballos fósiles de la Formación Curré en el cantón de Coto Brus, Costa Rica (análisis basado en material dental)* (Unpublished Lic. Thesis). Universidad de Costa Rica, San José, Costa Rica.
- Valerio, A. L. & Laurito, C. A. (2008). Dental remains of immature individuals of *Rhynchotherium blicki* (Frick 1933) of San Gerardo de Limoncito, Coto Brus, Costa Rica. *Revista Geológica de América Central*, 39, 87-91.
- Valerio, A. L. & Laurito, C. A. (2014). Nueva evidencia de un estadio juvenil de *Pliometanastes protistus* y su relación con *?Pliomenastes galushai* Hirschfeld & Webb, 1968 (Xenarthra, Megalonychidae) para el Mioceno Superior de Costa Rica. *Revista Geológica de América Central*, 51, 159-163.
- Valerio, A. L. & Laurito, C. A. (2020). Primer registro de pecaríes fósiles (Artiodactyla, Tayassuidae) para el Mioceno Superior (Hh3: Hemphilliano tardío) de Costa Rica, América Central. *Revista Geológica de América Central*, 62, 25-47. doi: 10.15517/rgac.v62i0.40638
- Varela, L., Tambusso, P. S., McDonald, H. G., & Fariña, R. A. (2019). Phylogeny, macroevolutionary trends and historical biogeography of sloths: Insights from a Bayesian morphological clock analysis. *Systematic Biology*, 68, 204-218. doi: 10.1093/sysbio/syy058
- Webb, S. D. (1976). Mammalian faunal dynamics of the Great American Interchange. *Paleobiology*, 2, 216-234.
- Webb, S. D. (1989). Osteology and relationships of *Thinobadistes segnis*, the first mylodont sloth in North America. In K. H. Redford & J. F. Eisenberg. (Eds.), *Advances in Neotropical Mammalogy* (pp. 469-532). Gainesville, USA: The Sandhill Crane Press.
- Webb, S. D. (1997). The great American faunal interchange. In A. G. Coates. (Ed.), *Central America: A natural and cultural history* (pp. 97-122). New Haven, Connecticut, USA: Yale University Press.
- Zachos, J., Pagani, M., Sloan, L., Thomas, E. & Billups, K. (2001). Trends, rhythms, and aberrations in global climate 65 Ma to present. *Science*, 292, 686-693.
- Zurita, A. E., Oliveira, E. V., Toriño, P. & Krapovickas, J. (2011). On the taxonomic status of some Glyptodontidae (Mammalia, Xenarthra, Cingulata) from the Pleistocene of South America. *Annals of Paleontology*, 97, 63-83.