



SHILAP Revista de lepidopterología

ISSN: 0300-5267

ISSN: 2340-4078

avives@orange.es

Sociedad Hispano-Luso-Americana de Lepidopterología
España

Melia, T.; Sinulingga, N. G. H. B.; Maretha, M. V.; Wijaya, R.; Efendi, R. R.;
Oliveira, L. S. S.; Kkadan, S. K.; Tavares, W. de S.; Tarigan, M.; Duran, A.

First report of *Tetracona (Agrotera) amathealis* (Walker, 1859) on a *Eucalyptus grandis* W. Hill × *Eucalyptus pellita* F. Muell. (Myrtaceae) hybrid in Indonesia (Lepidoptera: Crambidae, Spilomelinae)

SHILAP Revista de lepidopterología, vol. 49, núm. 195, 2021, Julio-Septiembre, pp. 407-412

Sociedad Hispano-Luso-Americana de Lepidopterología
Madrid, España

Disponible en: <https://www.redalyc.org/articulo.oa?id=45569077004>

- Cómo citar el artículo
- Número completo
- Más información del artículo
- Página de la revista en redalyc.org

redalyc.org

Sistema de Información Científica Redalyc

Red de Revistas Científicas de América Latina y el Caribe, España y Portugal
Proyecto académico sin fines de lucro, desarrollado bajo la iniciativa de acceso
abierto

First report of *Tetracona (Agrotera) amathealis* (Walker, 1859) on a *Eucalyptus grandis* W. Hill × *Eucalyptus pellita* F. Muell. (Myrtaceae) hybrid in Indonesia (Lepidoptera: Crambidae, Spilomelinae)

T. Melia, N. G. H. B. Sinulingga, M. V. Maretha, R. Wijaya, R. R. Efendi, L. S. S. Oliveira, S. K. Kkadan, W. de S. Tavares, M. Tarigan & A. Duran

Abstract

Tetracona amathealis (Walker, 1859) (Lepidoptera: Crambidae) is previously known from Australia and New Guinea, where it feeds on the native *Eucalyptus tereticornis* Sm. (Myrtaceae). The objective of this study was the report, for the first time, *T. amathealis* in Indonesia as well as to record new host plants for this insect. Were collected manually from a *Eucalyptus grandis* W. Hill. × *Eucalyptus pellita* F. Muell. hybrid commercial plantation in Riau, Sumatra, Indonesia and a sample sent for identification via molecular analysis. Our findings revealed that the molecular approach used (through sequencing of its mt-COI gene) successfully allowed the identification of the species as *T. amathealis*. This insect is recorded for the first time in Indonesia on a new host, a *E. grandis* × *E. pellita* hybrid.

KEY WORDS: Lepidoptera, Crambidae, Spilomelinae, *Tetracona amathealis*, first record, Indonesia.

**Primer registro de *Tetracona (Agrotera) amathealis* (Walker, 1859) sobre un híbrido de *Eucalyptus grandis* W. Hill × *Eucalyptus pellita* F. Muell. (Myrtaceae) en Indonesia
(Lepidoptera: Crambidae: Spilomelinae)**

Resumen

Tetracona amathealis (Walker, 1859) (Lepidoptera: Crambidae) previamente es conocida de Australia y New Guinea, donde se alimenta del oriundo *Eucalyptus tereticornis* Sm. (Myrtaceae). El objetivo de este estudio fue registrar, por primera vez, *T. amathealis* en Indonesia así como registrar una nueva planta nutricia para el insecto. Manualmente, fueron recogidos de un híbrido de *Eucalyptus grandis* W. Hill. × *Eucalyptus pellita* F. Muell. Plantación comercial en Riau, Sumatra, Indonesia y una simple muestra enviada para identificación vía análisis molecular. Nuestras conclusiones revelaron que el enfoque molecular usado (a través del ordenación en serie de su gen mitocondrial mt-COI) permitió la identificación de la especie con éxito como *T. amathealis*. Este insecto se registra, por primera vez, en Indonesia sobre una nueva planta nutricia un híbrido de *E. grandis* × *E. pellita*.

PALABRAS CLAVE: Lepidoptera, Crambidae, Spilomelinae, *Tetracona amathealis*, primer registro, Indonesia.

Introduction

Eucalyptus L'Hér. (Myrtales: Myrtaceae) is a plant genus with most species native to Australia (PRASETYO *et al.*, 2017; INAIL *et al.*, 2019). It is largely planted in Riau, Sumatra, Indonesia, where is utilized to produce paper, pulp and viscose materials (TAVARES *et al.*, 2020). Several lepidopteran species infest *Eucalyptus* in Riau, including *Ophiusa disjungens* (Walker, 1858) (Erebidae)

(LAKSAMANA *et al.*, 2020), *Polyphagozerra coffeae* (Cossidae) (TAVARES *et al.*, 2020; TACHI *et al.*, 2020) and *Strepsicrates* sp. (Tortricidae) (KKADAN *et al.*, 2020a).

Edward Meyrick described the genus *Tetracona* in 1884 with *Aediodes amathealis* Walker, 1859 as type species (MEYRICK, 1884). The genus was for a long period treated as a synonym of *Agrotera* Schrank, 1802, but a recent taxonomic revision of *Agrotera* resulted in the re-instatement of *Tetracona* (CHEN *et al.*, 2017). The genus is placed in the tribe Agroterini based on the structure of the uncus, featuring a broad base and simple chaetae on the uncus head, a well-developed medial process of the gnathos, the rectangular, elongate valvae with parallel costal and ventral margins, and the notably elongate saccus (at least in *Tetracona amathealis* (Walker, 1859)) (MALLY *et al.*, 2019). The genus with its three species (*T. amathealis*, *Tetracona multispina* Jie & Li, 2020 and *Tetracona pictalis* Warren, 1896) is known from Australia, China and New Guinea (HERBISON-EVANS *et al.*, 2013; JIE *et al.*, 2020).

Francis Walker described *T. amathealis* in 1859 (synonyms *Pyralis ornatalis* Walker, 1866) based on material collected at Moreton Bay in Queensland, Australia (WALKER, 1859). It is found in Australia (Queensland, northern New South Wales and Western region) and New Guinea (CHEN *et al.*, 2017). The wingspan is about 20 mm. The basal half of the forewings is golden and the marginal half grey. An orange submarginal band divides these two areas. The hindwings are grey with a gold inner margin (HERBISON-EVANS *et al.*, 2013). The *T. amathealis* leafrollers are sometimes considered a *Eucalyptus tereticornis* Sm. pest (HERBISON-EVANS *et al.*, 2013), an economically important plant native to eastern Australia and southern New Guinea (BOOTH, 2019).

Molecular method is an important way to confirm identification of lepidopteran species such as Crambidae (BARRERA *et al.*, 2017) and Noctuidae (SULISTYONO *et al.*, 2020). Insects recovered from traps such as light and sticky traps often have their body parts damaged leading to a difficult species determination through morphology analysis (MORINIÈRE *et al.*, 2016; FUJIWARA *et al.*, 2017). The collection of insect immatures without a defined rearing protocol is also a challenge for species identification via morphology analysis (ANTONINI *et al.*, 2009; SHIN *et al.*, 2015), since most insects are identified based on their adult body morphology examination (TAHIR *et al.*, 2018).

The objective of this study was to report the findings of *T. amathealis* in Indonesia, confirmed by molecular technique, as well as its attack on a *Eucalyptus grandis* W. Hill. × *Eucalyptus pellita* F. Muell. hybrid. We provided also a key for *Tetracona* species identification in Indonesia.

Material and methods

COLLECTION OF *T. AMATHEALIS*

Tetracona amathealis leafrollers were collected manually from trees of an *E. grandis* × *E. pellita* hybrid in a commercial stand in the first semester of 2020 in Teso East area (0° 30' N × 101° 26' E, 33 m altitude) in Riau. Leafrollers were placed in one-liter plastic containers and taken to the Entomology Laboratory of the Asia Pacific Resources International Holdings Ltd. (APRIL) of the PT. Riau Andalan Pulp and Paper (RAPP), where they were kept in a room at 26 ± 2°C, 75 ± 15% RH and 14:10 h (L:D) photoperiod. Leafrollers received daily fresh *E. grandis* × *E. pellita* shoots as a food.

MOLECULAR IDENTIFICATION OF *T. AMATHEALIS*

One leafroller individual was sent to the Molecular Biology and Genomics Laboratory of APRIL in Pangkalan Kerinci, Riau for species name confirmation. DNA extraction and PCR of the mitochondrial cytochrome c oxidase subunit I (mt-COI) gene were carried out as described by SULISTYONO *et al.* (2020). The PCR product was then sent for Sanger Sequencing. Trimming and consensus sequence building were carried out using the sangeranalyseR package (CHAO *et al.*, 2020) with default settings. We used the consensus sequence to find the most similar sequence in genbank using blastn (ALTSCHUL *et al.*, 1990). The top hit, sorted by the highest score, is kept. The consensus

sequence, top hits from the blast search as well as several sequences from the *Strepsicrates* Meyrick, 1888 (Lepidoptera: Tortricidae, Eucosmini) were aligned using Clustal Omega (SIEVERS *et al.*, 2011), and the output was used to build a phylogenetic tree. The maximum likelihood approach (FELSENSTEIN *et al.*, 1981) was used to build a dendrogram, with a bootstrap value of 1,000 and GTR (LE *et al.*, 2008) as the DNA substitution model, both of which were implemented in the phangorn R package (SCHLIEP *et al.*, 2011).

Results

REPORT OF *T. AMATHEALIS* IN INDONESIA

This is the first report of *T. amathealis* (Fig. 1) as well as of a member in the genus *Tetracona* in Indonesia. We used molecular method to confirm the identity of this insect species, through sequencing of its mt-COI gene.

Report of *T. amathealis* on *E. grandis* × *E. pellita*

The *T. amathealis* were collected while feeding on a *E. grandis* × *E. pellita* hybrid, which represents two new host plants for this insect.

Discussion

REPORT OF *T. AMATHEALIS* IN INDONESIA

Tetracona amathealis, reported in the current study in Indonesia, has been previously reported in Australia (i.e., Queensland, northern New South Wales and Western region) and New Guinea. Other reported *Tetracona* species in Asia includes *T. multispina* in China, Jiangxi Province (i.e., Huangzihao, Fuliang; Tongboshan; Wuyuan, Shangbao; Dabali, Xunwu; Doushui, Shangyou; Shangyou Arboretum) (JIE *et al.*, 2020) and *T. pictalis* in Australia (i.e., Queensland) (NUSS *et al.*, 2003).

The mt-COI gene utilized to identify *T. amathealis* is often used in taxonomic studies due to its lower divergence within species and larger divergence across species. The most similar sequence to our sample is of that *T. amathealis* (Table 1), which was deposited at the Australian National Insect Collection. Phylogenetic analysis confirms that our sample cluster together with *T. amathealis* in all 1,000 bootstraps, indicating the similarity to *T. amathealis* is stable (Fig. 2).




Fig. 1.— Adult *Tetracona amathealis* (Walker, 1859).

Table 1.— Origin, score, E value, identity, and accession of *Tetracona* (*Agrotera*) *amathealis*.

Origin	Score	E value	Identity	Accession	
Teso East area, Riau, Sumatra, Indonesia	623	0.0	95%	HQ952617.1;	<i>Agrotera amathealis</i>

REPORT OF *T. AMATHEALIS* ON *E. GRANDIS* × *E. PELLITA*

Tetracona amathealis, collected in the present study while feeding on a *E. grandis* × *E. pellita* hybrid, has been previously recorded to feed on *E. tereticornis* (HERBISON-EVANS *et al.*, 2013). This insect was seen along with the tea mosquito bug, *Helopeltis theivora* Waterhouse, 1886 (Hemiptera: Miridae), a sapsucker and other mirid species on the commercial stands of *E. grandis* × *E. pellita* hybrid in Riau (KKADAN *et al.*, 2020b).



Acknowledgments

Thanks to Dr. Buck Richardson (LeapFrogOz, Kuranda Kreations, Queensland, Australia) for providing the figure 1. The following Indonesian companies provided financial support to the study: PT. Riau Andalan Pulp and Paper (RAPP) and Asia Pacific Resources International Holdings Ltd. (RAPP).

BIBLIOGRAPHY

- ALTSCHUL, S. F., GISH, W., MILLER, W., MYERS, E. W. & LIPMAN, D. J., 1990.– Basic local alignment search tool.– *Journal of Molecular Biology*, **215**(3): 403-410.
- ANTONINI, G., COLETTI, G., SERRANI, L., TRONCI, C., CRISTOFARO, M. & SMITH, L., 2009.– Using molecular genetics to identify immature specimens of the weevil *Ceratapion basicorne* (Coleoptera: Apionidae).– *Biological Control*, **51**(1): 152-157.
- BARRERA, G. P., VILLAMIZAR, L. F., ESPINEL, C., QUINTERO, E. M., BELAICH, M. N., TOLOZA, D. L., GHIRINGHELLI, P. D. & VARGAS, G., 2017.– Identification of *Diatraea* spp. (Lepidoptera: Crambidae) based on cytochrome oxidase II.– *PLoS ONE*, **12**(9): e0184053.
- BOOTH, T. H., 2019.– Assessing the thermal adaptability of tree provenances: An example using *Eucalyptus tereticornis*.– *Australian Forestry*, **82**(4): 176-180.
- CHAO, K. H., BARTON, B., PALMER, S. & LANFEAR, R., 2020.– SangeranalyseR: Simple and interactive analysis of Sanger sequencing data in R.– *Bioinformatics*: doi: <https://doi.org/10.1101/2020.05.18.102459> (preprint).

- CHEN, K., HORAK, M., DU, X. & ZHANG, D., 2017.– Revision of the Australian species of *Agrotera* Schrank (Lepidoptera: Pyraloidea: Crambidae: Spilomelinae).– *Zootaxa*, **4362**(2): 213-224.
- FELSENSTEIN, J., 1981.– Evolutionary trees from DNA sequences: A maximum likelihood approach.– *Journal of Molecular Evolution*, **17**(1): 368-376.
- FUJIWARA, K., UECHI, N., SHIMIZU, Y., TODA, S., INOUE, H., YAMAGUCHI, T., IWANAMI, T. & FUJIKAWA, T., 2017.– Effective molecular detection of *Diaphorina citri* Kuwayama (Hemiptera: Liviidae) in bulk insect samples from sticky traps.– *Journal of Applied Entomology*, **141**(1-2): 61-66.
- HERBISON-EVANS, D., HACOBIAN, B. & CROSSLEY, S., 2013.– *Agrotera amathealis* (Walker, 1859). Australian caterpillars and their butterflies and moths.– Available from <http://lepidoptera.butterflyhouse.com.au/spil/amathealis.html> (accessed 18 August 2020).
- INAIL, M. A., HARDIYANTO, E. B. & MENDHAM, D. S., 2019.– Growth responses of *Eucalyptus pellita* F. Muell plantations in South Sumatra to macronutrient fertilisers following several rotations of *Acacia mangium* Willd.– *Forests*, **10**(12): 1054.
- JIE, L. L., YANG, J. B. & LI, W. C., 2020.– First record of the genus *Tetracona* Meyrick (Lepidoptera, Crambidae) from China, with description of a new species.– *ZooKeys*, **941**: 101-105.
- KKADAN, S. K., SIRAIT, B. A., ASFA, R., TAVARES, W. DE S., TARIGAN, M., DURAN, A., WONG, C. Y. & SHARMA, M., 2020a.– Evaluation of a spinetoram-based insecticide against lepidopteran and thrips infesting acacia and *Eucalyptus* in Sumatra, Indonesia.– *Journal of Entomology and Zoology Studies*, **8**(2): 1345-1351.
- KKADAN, S. K., YESHWANTH, H. M., TAVARES, W. DE S., PASARIBU, I., ABAD, J. I. M., TARIGAN, M., DURAN, A., WONG, C. Y. & SHARMA, M., 2020b.– Mirid pests of *Eucalyptus* in Indonesia: Notes on damage symptoms, alternate hosts and parasitoid.– *Journal of the Kansas Entomological Society*, **92**(4): 577-588.
- LAKSAMANA, R. A., SINULINGGA, N. G. H. B., FERLIANDA, I. A., SAMOSIR, M. N., PANE, S. B., KKADAN, S. K., TAVARES, W. DE S., TARIGAN, M. & DURAN, A., 2020.– First report of *Ophiusa disjunctus* (Walker, 1858) on *Acacia mangium* (Fabaceae), and damage and notes of its biology on *Eucalyptus* (Myrtaceae) commercial plantations in Sumatra, Indonesia (Lepidoptera: Erebididae).– *SHILAP Revista de lepidopterologia*, **48**(191): 439-447.
- LE, S. Q. & GASCUEL, O., 2008.– LG: An improved, general amino-acid replacement matrix molecular biology and evolution.– *Molecular Biology and Evolution*, **25**(7): 1307-1320.
- MALLY, R., HAYDEN, J. E., NEINHUIS, C., JORDAL, B. H. & NUSS, M., 2019.– The phylogenetic systematics of Spilomelinae and Pyraustinae (Lepidoptera: Pyraloidea: Crambidae) inferred from DNA and morphology.– *Arthropod Systematics & Phylogeny*, **77**(1): 141-204.
- MEYRICK, E., 1884.– On the classification of the Australian Pyralidina.– *Transactions of the Entomological Society of London*, **1884**(61-80): 277-350.
- MORINIÈRE, J., CANCIAN DE ARAUJO, B., LAM, A. W., HAUSMANN, A., BALKE, M., SCHMIDT, S., HENDRICH, L., DOCKAL, D., FARTMANN, B., ARVIDSSON, S. & HASZPRUNAR, G., 2016.– Species identification in Malaise trap samples by DNA barcoding based on NGS technologies and a scoring matrix.– *PLoS ONE*, **11**(5): e0155497.
- NUSS, M., LANDRY, B., MALLY, R., VEGLIANTE, F., TRÄNKNER, A., BAUER, F., HAYDEN, J., SEGERER, A., SCHOUTEN, R., LI, H., TROFIMOVA, T., SOLIS, M. A., DE PRINS, J. & SPEIDEL, W., 2003-onwards.– *Global Information System on Pyraloidea* (*GlobIZ*).– Available from <http://www.pyraloidea.org/> (accessed 18 August 2020).
- PRASETYO, A., AISO, H., ISHIGURI, F., WAHYUDI, I., WIJAYA, I. P. G., OHSHIMA, J. & YOKOTA, S., 2017.– Variations on growth characteristics and wood properties of three *Eucalyptus* species planted for pulpwood in Indonesia.– *Tropics*, **26**(2): 59-69.
- SCHLIEP, K. P., 2011.– Phangorn: Phylogenetic analysis in R.– *Bioinformatics*, **27**(4): 592-593.
- SHIN, S., JUNG, S., HELLER, K., MENZEL, F., HONG, T. K., SHIN, J. S., LEE, S. H., LEE, H. & LEE, S., 2015.– DNA barcoding of *Bradysia* (Diptera: Sciaridae) for detection of the immature stages on agricultural crops.– *Journal of Applied Entomology*, **139**(8): 638-645.
- SIEVERS, F., WILM, A., DINEEN, D., GIBSON, T. J., KARPLUS, K., LI, W., LOPEZ, R., MCWILLIAM, H., REMMERT, M., SODING, J., THOMPSON, J. D. & HIGGINS, D. G., 2011.– Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega.– *Molecular Systems Biology*, **7**: 539.
- SULISTYONO, E., KKADAN, S. K., MARETHA, M. V., TAVARES, W. DE S., SIRAIT, B. A., SINULINGGA, N. G. H. B., TARIGAN, M. & DURAN, A., 2020.– *Spodoptera exigua* and *Spodoptera litura* (Lepidoptera, Noctuidae) on an *Acacia crassicaarpa* (Fabaceae) commercial nursery in Sumatra, Indonesia.– *Journal of the Lepidopterists' Society*, **74**(3): 176-182.
- TACHI, T., SHIMA, H., TAVARES, W. DE S. & TARIGAN, M., 2020.– A new species of *Cossidophaga* Baranov from Indonesia (Diptera: Tachinidae), a parasitoid of the carpenter moth, *Polyphagocarpa coffeae* (Nietner) (Lepidoptera: Cossidae).– *Oriental Insects*, **54**(2): 545-555.
- TAHIR, H. M., NOOR, A., MEHMOOD, S., SHERAWAT, S. M. & QAZI, M. A., 2018.– Evaluating the accuracy

- of morphological identification of insect pests of rice crops using DNA barcoding.– *Mitochondrial DNA Part B*, **3**(2): 1220-1224.
- TAVARES, W. DE S., KKADAN, S. K., HENDRIK, A. M., TARIGAN, M., ASFA, R., YAKOVLEV, R. V., TACHI, T., DURAN, A., WONG, C. Y. & SHARMA, M., 2020.– Notes on the biology and natural enemies of *Polyphagozerra coffeae* (Nietner, 1861) infesting *Eucalyptus pellita* F. Muell. (Myrtaceae) trees in Riau, Indonesia (Lepidoptera: Cossidae, Zeuzerinae).– *SHILAP Revista de lepidopterología*, **48**(190): 333-349.
- WALKER, F., 1859.– *Pyralites. List of the Specimens of Lepidopterous Insects in the Collection of the British Museum, London*, **17**(3): 255-508.

T. M., N. G. H. B. S., M. V. M., R. W., R. R. E., L. S. S. O., S. K. K., *W. S. T., M. T., A. D.
Asia Pacific Resources International Holdings Ltd. (APRIL)
PT. Riau Andalan Pulp and Paper (RAPP)
28300, Pangkalan Kerinci, Riau, Sumatra
INDONESIA / INDONESIA

E-mail: Tisha_Melia@aprilasia.com
<https://orcid.org/0000-0003-1926-9109>

E-mail: nike.grace.h@mail.ugm.ac.id
<https://orcid.org/0000-0002-0879-1930>

E-mail: Maggie_VM@aprilasia.com
<https://orcid.org/0000-0002-9135-5629>

E-mail: wijayariza12@gmail.com
<https://orcid.org/0000-0002-7485-9104>

E-mail: rudirio24@gmail.com
<https://orcid.org/0000-0001-8766-4800>

E-mail: Leonardo_Oliveira@aprilasia.com
<https://orcid.org/0000-0002-4056-6987>

E-mail: Srikumar@aprilasia.com
<https://orcid.org/0000-0002-1875-6899>

*E-mail: Wagner_Tavares@aprilasia.com
<https://orcid.org/0000-0002-8394-6808>

E-mail: Marthin_Tarigan@aprilasia.com
<https://orcid.org/0000-0002-9128-2650>

E-mail: Alvaro_Duran@aprilasia.com
<https://orcid.org/0000-0002-3035-9087>

*Autor para la correspondencia / *Corresponding author*

(Recibido para publicación / *Received for publication* 14-X-2020)
(Revisado y aceptado / *Revised and accepted* 18-XII-2020)
(Publicado / *Published* 30-IX-2021)