



SHILAP Revista de lepidopterología

ISSN: 0300-5267

ISSN: 2340-4078

avives@orange.es

Sociedad Hispano-Luso-Americana de Lepidopterología  
España

Efetov, K. A.; Knyazev, S. A.; Kucherenko, E. E.  
The first record of *Jordanita (Solaniterna) subsolana* (Staudinger,  
1862) from Kazakhstan (Lepidoptera: Zygaenidae, Procridinae)  
SHILAP Revista de lepidopterología, vol. 50, núm. 198, 2022, -Junio, pp. 233-239  
Sociedad Hispano-Luso-Americana de Lepidopterología  
Madrid, España

DOI: <https://doi.org/10.57065/shilap.127>

Disponible en: <https://www.redalyc.org/articulo.oa?id=45571929005>

- Cómo citar el artículo
- Número completo
- Más información del artículo
- Página de la revista en redalyc.org

redalyc.org

Sistema de Información Científica Redalyc

Red de Revistas Científicas de América Latina y el Caribe, España y Portugal  
Proyecto académico sin fines de lucro, desarrollado bajo la iniciativa de acceso  
abierto

# The first record of *Jordanita (Solaniterna) subsolana* (Staudinger, 1862) from Kazakhstan (Lepidoptera: Zygaenidae, Procridinae)

K. A. Efetov, S. A. Knyazev & E. E. Kucherenko

## Abstract

*Jordanita (Solaniterna) subsolana* (Staudinger, 1862) was discovered in eastern Kazakhstan using the sex attractant EFETOV-2 (2-butyl 2-dodecenoate) synthesised in the Crimean Federal University. This is the first record of *J. (S.) subsolana* from Kazakhstan. As this species is now found on the border of Kazakhstan and China, most probably that *J. (S.) subsolana* is also present in China. An overview of the Zygaenidae fauna of Kazakhstan is presented.

KEY WORDS: Lepidoptera, Zygaenidae, Procridinae, *Jordanita (Solaniterna) subsolana*, fauna, sex attractant, EFETOV-2, Kazakhstan, China.

## El primer registro de *Jordanita (Solaniterna) subsolana* (Staudinger, 1862) de Kazajstán (Lepidoptera: Zygaenidae, Procridinae)

## Resumen

*Jordanita (Solaniterna) subsolana* (Staudinger, 1862) fue descubierta en el este de Kazajstán usando el atrayente sexual EFETOV-2 (2-butil 2-dodecenoato) sintetizado en la Universidad Federal de Crimea. Este es el primer registro de *J. (S.) subsolana* de Kazajstán. Como esta especie se ahora se ha encontrado en la frontera de Kazajstán y China, es más que probable que *J. (S.) subsolana* esté también presente en China. Se presenta una visión general de la fauna de Zygaenidae de Kazajstán.

PALABRAS CLAVE: Lepidoptera, Zygaenidae, Procridinae, *Jordanita (Solaniterna) subsolana*, fauna, atracción sexual, EFETOV-2, Kazajstán, China.

## Introduction

New data on the biology, morphology, karyology and DNA barcoding of Zygaenidae species, which have been obtained over the past 20 years, have significantly increased our knowledge on this family and stimulated numerous systematic changes in it (EFETOV, 1999; EFETOV *et al.*, 2006, 2014a, 2014b, 2015a; EFETOV & SAVCHUK, 2009; EFETOV & TARMANN, 2014, 2016, 2017b; EFETOV & KNYAZEVA, 2014; KNYAZEVA *et al.*, 2015a, 2015b). As a consequence, the Zygaenidae family now includes five subfamilies: Inouelinae Efetov & Tarmann, 2017; Procridinae Boisduval, 1828; Chalcosiinae Walker, 1865; Callizygaeninae Alberti, 1954; and Zygaeninae Latreille, 1809 (EFETOV *et al.*, 2014b; EFETOV & TARMANN, 2017a). According to previous publications the Zygaenidae fauna of Kazakhstan is represented by two subfamilies: Procridinae (12 species) and Zygaeninae (13 species) (EFETOV, 1995, 1997, 2001, 2004; EFETOV & TARMANN, 1999; HOFMANN & TREMEWAN, 1996; MOLLET, 2008). New data presented in this paper on the distribution of zygaenids in Kazakhstan became possible due to the field experiments with attractant baits. It is well known that sex pheromones and sex attractants play an increasingly important role in

faunal investigations of Lepidoptera, because they allow researchers to discover new species, detect populations of pest or rare species, monitor and control the seasonal flight of moths (SUBCHEV *et al.*, 2012, 2013, 2016; RAZOV *et al.*, 2017; CAN CENGİZ *et al.*, 2018; CAN *et al.*, 2019; EFETOV *et al.*, 2010, 2011, 2015b, 2019; TARMANN *et al.*, 2019; VRENOZI *et al.*, 2019).

In this study, the fauna of Zygaenidae in Kazakhstan was investigated using the attractant EFETOV-2 (2-butyl 2-dodecenoate). Its property as a sex attractant for the males of some Procrinae species has been proved in our field experiments earlier (EFETOV *et al.*, 2016, 2018, 2019, 2020; EFETOV & KUCHERENKO, 2020).

## Materials and methods

Field observations were carried out in eastern and northern Kazakhstan in VI-VII-2018. To prepare the baits, the substance EFETOV-2 (2-butyl 2-dodecenoate) was used. This attractant was synthesised in the Crimean Federal University as described in EFETOV *et al.* (2014c). For this study, the sex attractant was applied onto grey rubber vial caps in dose of 200 mcl. The baits were prepared in 2014, wrapped singly in aluminium foil and stored in a freezer for the first year and then at room temperature until the date of the fieldwork. In study biotopes the lures were hung on bushes at a height 0.5-1.0 m above the ground. The attracted specimens were collected by netting them near the lure. All moths were determined by genitalia examination.

The map (Fig. 1) is compiled by the BioOffice software, Tiroler Landesmuseen, Ferdinandeum, Hall in Tirol, Austria.

## Results and discussion

*Jordanita (Solaniterna) subsolana* (Staudinger, 1862) is the type species of the subgenus *Solaniterna* Efetov, 2004. In 2018 this species was collected for the first time in Kazakhstan. This became possible due to an application of the sex attractant EFETOV-2. One male of this species was attracted to the bait with EFETOV-2 in Zaysan District, 19 km S of Zaysan, Saur Mountains, 47°17'50,07" N, 84°48'28,54" E, 1100 m, 1-2-VII-2018, S. Knyazev leg. (Figs 1-4). This species was recorded (EFETOV, 1990) from the Russian Altai (Fig. 1), but until now, there was no information about the presence of *J. (S.) subsolana* in eastern Kazakhstan based on original material. The known distribution of *J. (S.) subsolana* outside of Kazakhstan: Morocco (Rif), Spain, France, Belgium, Germany, Switzerland, Austria, Italy (including Sicily), Slovenia, Croatia, Bosnia and Herzegovina, Serbia, Montenegro, Albania, North Macedonia, Greece, Czech Republic, Slovakia, Hungary, Romania, Bulgaria, Ukraine, Moldova, Russia (European part including Crimea, Northern Caucasus, southern Siberia), Georgia, Armenia, Azerbaijan, and Turkey (EFETOV, 2004; BioOffice database, Tiroler Landesmuseen, Ferdinandeum, Hall in Tirol, Austria). Moreover, as this species is now found on the border of Kazakhstan and China, most probably that *J. (S.) subsolana* is also present in China.

The list of the Zygaenidae fauna of Kazakhstan includes 26 species which are listed below.

### Procrinae:

1. *Rhagades (Rhagades) pruni* ([Denis & Schiffermüller], 1775)
2. *Adscita (Procriterna) subtristis* (Staudinger, 1887)
3. *Adscita (Procriterna) subdolosa* (Staudinger, 1887) (EFETOV, 1995)
4. *Adscita (Adscita) statices* (Linnaeus, 1758)
5. *Jordanita (Roccia) budensis* (Speyer & Speyer, 1858)
6. *Jordanita (Roccia) paupera* (Christoph, 1887)
7. *Jordanita (Roccia) volgensis* (Möschler, 1862)
8. *Jordanita (Roccia) naufocki* (Alberti, 1937)
9. *Jordanita (Roccia) almatiensis* Mollet, 2008
10. *Jordanita (Tremewania) splendens* (Staudinger, 1887)
11. *Jordanita (Jordanita) chloros* (Hübner, 1813) (EFETOV, 1997)

12. *Jordanita (Solaniterna) subsolana* (Staudinger, 1862)
13. *Jordanita (Solaniterna) solana* (Staudinger, 1887) (EFETOV, 1995)

#### Zygaeninae:

14. *Zygaena (Mesembrynus) purpuralis* (Brünnich, 1763)
15. *Zygaena (Mesembrynus) cynarae* (Esper, 1789)
16. *Zygaena (Mesembrynus) centaureae* Fischer von Waldheim, 1832
17. *Zygaena (Mesembrynus) laeta* (Hübner, 1790)
18. *Zygaena (Agrumenia) truchmena* Eversmann, 1854
19. *Zygaena (Agrumenia) sogdiana* Erschoff, 1874
20. *Zygaena (Agrumenia) separata* Staudinger, 1887
21. *Zygaena (Agrumenia) carniolica* (Scopoli, 1763)
22. *Zygaena (Agrumenia) exulans* (Hohenwarth, 1792)
23. *Zygaena (Agrumenia) viciae* ([Denis & Schiffermüller], 1775)
24. *Zygaena (Lictoria) loti* ([Denis & Schiffermüller], 1775)
25. *Zygaena (Zygaena) osterodensis* Reiss, 1921
26. *Zygaena (Zygaena) lonicerae* (Scheven, 1777)

The systematic position of two taxa, viz. *Jordanita (Roccia) scintillosa* Zolotuhin, 2020, and *Jordanita (Roccia) smaragdonna* Zolotuhin, 2020, both mentioned recently for Kazakhstan (ZOLOTUHHIN, 2020), needs verification. *J. scintillosa* and *J. smaragdonna* have no genitalia differences from *J. paupera*. Moreover, the colour and shape of wings and colour and length of proboscis mentioned by Zolotuhin (2020) as important characters are variable even in *J. paupera* in Turkmenistan, from where the latter species had been described. In this publication we consider *J. scintillosa* and *J. smaragdonna* as the synonyms (**syn. n.**) of *J. paupera*. These two taxa of Zolotuhin will be discussed in a separate publication (EFETOV & TARMANN, in prep.).

It is known that (2*R*)-butyl 2-dodecenoate, named as EFETOV-S-2, is a very effective sex attractant for the detection of *J. (S.) subsolana* in different biotopes (EFETOV *et al.*, 2020). However, males of this species also actively respond to the attractant EFETOV-2 (2-butyl 2-dodecenoate). Previously, this was shown in the Crimea (Russia) (EFETOV *et al.*, 2016) and in the Thrace Region (European Turkey) (CAN CENGİZ *et al.*, 2018). It is interesting that the baits with the attractant EFETOV-2 used in this study were prepared in 2014. This means that EFETOV-2 retains its attractive properties for several years. Of course, evaporation from rubber caps occurs, but the remaining dose is sufficient to attract and accordingly to detect *J. (S.) subsolana* in biotopes. We reported about a long duration of the activity of EFETOV-2 lures, but only during one season (three months) (VRENOZI *et al.*, 2019). Now we have the proof for a much longer activity. This is convenient because we can use attractive baits for a long time without the need to change dispensers. Moreover, it is cost-effective.

In addition, one more species, viz. *Adscita (Adscita) statice* (Linnaeus, 1758), was attracted to EFETOV-2 lures in Kazakhstan by the second author in 2018. Totally 17 males of *A. (A.) statice* were caught: 1 ♂ - eastern Kazakhstan, Katon-Karagay District, Southern Altai Ridge, bank of Kara-Kaba River, 49°01'42.96" N, 86°01'04.44" E, 1653 m, 28-VI-2018; 11 ♂♂ - eastern Kazakhstan, Markakol District, 1.5 km N of village Urunkhaika, eastern bank of Markakol Lake, 48°48'24.83" N, 86°02'00.19" E, 1450 m, 29-VI-2018; 5 ♂♂ - northern Kazakhstan, North Kazakhstan Region, Taiynshinsky District, 6 km. SE of village Akkudyk, 53°37'58.64" N, 70°42'47.80" E, 145 m, 08-VII-2018. This result was expected and confirmed our data obtained in Turkey (CAN CENGİZ *et al.*, 2018). Besides this, *A. (A.) statice* males were also attracted to the sex attractant EFETOV-S-2 (mentioned above) in Russia (EFETOV & GORBUNOV, 2016) and Italy (EFETOV *et al.*, 2020).

#### Conclusions

The males of two Procridinae species, viz. *A. (A.) statice* and *J. (S.) subsolana*, were attracted to

the lures with the sex attractant EFETOV-2 in Kazakhstan. *J. (S.) subsolana* is reported for Kazakhstan for the first time.

## Acknowledgements

We are indebted to Dr M. Y. Baevsky and Mr A. I. Poddubov (Simferopol, Russia) for help in preparing sex attractants. We also thank Prof. Dr G. M. Tarmann (Innsbruck, Austria) for the compilation of the map and fruitful discussions. The second author thanks Mrs Aliya M. Gabdullina (Katon-Karagai, Kazakhstan) and Mr Vadim V. Ivonin (Novosibirsk, Russia) for their help and good company during the expedition in Kazakhstan in 2018. The study of the second author was funded by RFBR, project number 20-34-90105.

## BIBLIOGRAPHY

- CAN, F., EFETOV, K. A., BURMAN, J., KAYA, K., KUCHERENKO, E. E., ULAŞLI, B. & TARMANN, G. M., 2019.– A study of the Zygaenidae (Lepidoptera) fauna of Central Anatolia, Turkey.– *Turkish Journal of Entomology*, **43**(2): 189-199. Available from <https://doi.org/10.16970/entoted.512580>.
- CAN CENGİZ, F., EFETOV, K. A., KAYA, K., KUCHERENKO, E. E., OKYAR, Z. & TARMANN, G. M., 2018.– Zygaenidae (Lepidoptera) of Thrace Region of Turkey.– *Nota lepidopterologica*, **41**(1): 23-36. Available from <https://doi.org/10.3897/nl.41.21065>.
- EFETOV, K. A., 1990.– Family Zygaenidae, pp. 84-85.– In K. A. EFETOV & YU. I. BUDASHKIN. *Lepidoptera of the Crimea*: 112 pp., 40 pls. Tavriya, Simferopol.
- EFETOV, K. A., 1995.– New foresters (Lepidoptera, Zygaenidae) for the fauna of Kazakhstan.– *Vestnik zoologii*, **29**(1): 32.
- EFETOV, K. A., 1997.– *Jordanita (Jordanita) chloros* (Hübner, [1813]) (Lepidoptera, Zygaenidae) - a new forester for the fauna of Kazakhstan and Asian part of Russia.– *Vestnik zoologii*, **31**(3): 85.
- EFETOV, K. A., 1999.– *Inouela* gen. n. from Japan and Taiwan (Lepidoptera: Zygaenidae, Chalcosiinae).– *Entomologist's Gazette*, **50**(2): 91-95.
- EFETOV, K. A., 2001.– *Jordanita (Jordanita) chloros* (Lepidoptera, Zygaenidae) - a new forester for the fauna of Kazakhstan and Asian part of Russia.– *Vestnik zoologii*, **35**(3): 45.
- EFETOV, K. A., 2004.– *Forester and Burnet Moths (Lepidoptera: Zygaenidae). The genera Theresimima Strand, 1917, Rhagades Wallengren, 1863, Zygaenoprocris Hampson, 1900, Adscita Retzius, 1783, Jordanita Verity, 1946 (Procridae), and Zygaena Fabricius, 1775 (Zygaeninae)*: 272 pp. CSMU Press, Simferopol.
- EFETOV, K. A., CAN, F., TOSHOVA, T. B. & SUBCHEV, M., 2010.– New sex attractant for *Jordanita anatolica* (Naufock) (Lepidoptera: Zygaenidae: Procridae).– *Acta Zoologica Bulgarica*, **62**(3): 315-319.
- EFETOV, K. A. & GORBUNOV, O. G., 2016.– Attraction of the males of *Adscita statices* (Linnaeus, 1758) (Lepidoptera: Zygaenidae, Procridae) by synthetic sex attractant in Moscow Region.– *Tavrisheskiy Mediko-biologicheskii Vestnik*, **19**(3): 40-46.
- EFETOV, K. A., HOFMANN, A. & TARMANN, G. M., 2014a.– Application of two molecular approaches (use of sex attractants and DNA barcoding) allowed to rediscover *Zygaenoprocris eberti* (Alberti, 1968) (Lepidoptera, Zygaenidae, Procridae), hitherto known only from the female holotype.– *Nota lepidopterologica*, **37**(2): 151-160. Available from <https://doi.org/10.3897/nl.37.7871>.
- EFETOV, K. A., HOFMANN, A., TARMANN, G. M. & TREMEWAN, W. G., 2014b.– Taxonomic comments on the treatment of the Zygaenidae (Lepidoptera) in volume 3 of *Moths of Europe*, Zygaenids, Pyralids 1 and Brachodids (2012).– *Nota lepidopterologica*, **37**(2): 123-133. Available from <https://doi.org/10.3897/nl.37.7940>.
- EFETOV, K. A. & KNYAZEV, S. A., 2014.– New records of *Jordanita (Roccia) volgensis* (Möschler, 1862) (Lepidoptera: Zygaenidae, Procridae) from Siberia (Russia) and Ukraine.– *Entomologist's Gazette*, **65**(3): 175-178.
- EFETOV, K. A., KOSHIO, C. & KUCHERENKO, E. E., 2018.– A new synthetic sex attractant for males of *Illiberis (Primilliberis) pruni* Dyar, 1905 (Lepidoptera: Zygaenidae, Procridae).– *SHILAP Revista de lepidopterología*, **46**(182): 263-270.
- EFETOV, K. A. & KUCHERENKO, E. E., 2020.– Structural analysis of sex pheromones and attractants in Zygaenidae (Insecta, Lepidoptera): biochemical and evolutionary aspects.– *Journal of Evolutionary Biochemistry and Physiology*, **56**(5): 381-395. Available from <https://doi.org/10.1134/S0022093020050014>.

- EFETOV, K. A., KUCHERENKO, E. E., PARSHKOVA, E. V. & TARMANN, G. M., 2016.– 2-butyl 2-dodecenoate, a new sex attractant for *Jordanita (Tremewania) notata* (Zeller, 1847) and some other Procridinae species (Lepidoptera: Zygaenidae).– *SHILAP Revista de lepidopterología*, **44**(175): 519-527.
- EFETOV, K. A., KUCHERENKO, E. E. & TARMANN, G. M., 2019.– New synthetic sex attractants for the males of two endemic Iberian Procridinae species (Lepidoptera: Zygaenidae).– *SHILAP Revista de lepidopterología*, **47**(186): 307-315.
- EFETOV, K. A., KUCHERENKO, E. E. & TARMANN, G. M., 2020.– An application of the synthetic sex attractants from the series “EFETOV-2” for studying Procridinae in Italy (Lepidoptera: Zygaenidae).– *SHILAP Revista de lepidopterología*, **48**(192): 733-749.
- EFETOV, K. A., PARSHKOVA, E. V., BAEVSKY, M. Y. & PODDUBOV, A. I., 2014c.– *Sec*-butyl ester of dodecenoate: synthesis and attractive properties.– *Ukrainian Biochemical Journal*, **86**(6): 175-182. Available from <https://doi.org/10.15407/ubj86.06.175>.
- EFETOV, K. A., PARSHKOVA, E. V., TARASOVA, L. G. & TARMANN, G. M., 2015a.– The karyotypes of Procridinae (Lepidoptera: Zygaenidae), with the first record of the karyotype of *Pollanisus commoni* Tarmann, 2004, a representative of the tribe Artonini.– *Entomologist's Gazette*, **66**(2): 121-125.
- EFETOV, K. A. & SAVCHUK, V. V., 2009.– The first record of *Jordanita (Roccia) volgensis* (Möschler, 1862) (Lepidoptera: Zygaenidae, Procridinae) from the Crimea.– *Entomologist's Gazette*, **60**(3): 155-158.
- EFETOV, K. A., SUBCHEV, M. A., TOSHOVA, T. B. & KISELEV, V. M., 2011.– Attraction of *Zygaenoprocris taftana* (Alberti, 1939) and *Jordanita horni* (Alberti, 1937) (Lepidoptera: Zygaenidae, Procridinae) by synthetic sex pheromones in Armenia.– *Entomologist's Gazette*, **62**(2): 113-121.
- EFETOV, K. A. & TARMANN, G. M., 1999.– *Forester Moths. The genera Theresimima Strand, 1917, Rhagades Wallengren, 1863, Jordanita Verity, 1946, and Adscita Retzius, 1783* (Lepidoptera: Zygaenidae, Procridinae): 192 pp. Apollo Books, Stenstrup.
- EFETOV, K. A. & TARMANN, G. M., 2014.– A new European species, *Adscita dujardini* sp. nov. (Lepidoptera: Zygaenidae, Procridinae) confirmed by DNA analysis.– *Entomologist's Gazette*, **65**(3): 179-200.
- EFETOV, K. A. & TARMANN, G. M., 2016.– *Pseudophacusa multidentata* Efetov & Tarmann, a new genus and species of Procridini from Myanmar, China and Laos (Lepidoptera: Zygaenidae, Procridinae).– *SHILAP Revista de lepidopterología*, **44**(173): 81-89.
- EFETOV, K. A. & TARMANN, G. M., 2017a.– The hypothetical ground plan of the Zygaenidae, with a review of the possible autapomorphies of the Procridinae and the description of the Inouelinae subfam. nov.– *Journal of the Lepidopterists' Society*, **71**(1): 20-49. Available from <https://doi.org/10.18473/lepi.v71i1.a5>.
- EFETOV, K. A. & TARMANN, G. M., 2017b.– *Thibetana keili* Efetov & Tarmann, a new species of the genus *Thibetana* Efetov & Tarmann, 1995, from Tibet (Lepidoptera: Zygaenidae, Procridinae, Artonini).– *SHILAP Revista de lepidopterología*, **45**(180): 581-587.
- EFETOV, K. A., TARMANN, G. M., HAYASHI, E. & PARSHKOVA, E. V., 2006.– New data on the chaetotaxy of the first instar larvae of Procridini and Artonini (Lepidoptera: Zygaenidae, Procridinae).– *Entomologist's Gazette*, **57**(4): 229-233.
- EFETOV, K. A., TARMANN, G. M., TOSHOVA, T. B. & SUBCHEV, M. A., 2015b.– Enantiomers of 2-butyl 7Z-dodecenoate are sex attractants for males of *Adscita mannii* (Lederer, 1853), *A. geryon* (Hübner, 1813), and *Jordanita notata* (Zeller, 1847) (Lepidoptera: Zygaenidae, Procridinae) in Italy.– *Nota lepidopterologica*, **38**(2): 161-169. Available from <https://doi.org/10.3897/nl.38.6312>.
- HOFMANN, A. & TREMEWAN, W. G., 1996.– *A Systematic Catalogue of the Zygaeninae* (Lepidoptera: Zygaenidae): 251 pp. Harley Books, Colchester.
- KNYAZEV, S. A., EFETOV, K. A. & PONOMARYOV, K. B., 2015a.– Zygaenidae (Lepidoptera) from Omsk Region.– *Zoologicheskii Zhurnal*, **94**(11): 1297-1302. Available from <https://doi.org/10.7868/S0044513415110069>.
- KNYAZEV, S. A., EFETOV, K. A. & PONOMARYOV, K. B., 2015b.– Zygaenidae (Lepidoptera) of Omsk Province.– *Entomological Review*, **95**(8): 1106-1111. Available from <https://doi.org/10.1134/S0013873815080175>.
- MOLLET, B., 2008.– *Jordanita (Roccia) almatiensis* sp. nov. from Kazakhstan (Lepidoptera, Zygaenidae, Procridinae).– *Entomologist's Gazette*, **59**(1): 57-62.
- RAZOV, J., EFETOV, K. A., FRANIN, K., TOSHOVA, T. B. & SUBCHEV, M. A., 2017.– The application of sex pheromone traps for recording the Procridinae fauna (Lepidoptera: Zygaenidae) in Croatia.– *Entomologist's Gazette*, **68**(1): 49-53.
- SUBCHEV, M. A., EFETOV, K. A., TOSHOVA, T. B. & KOSHIO, C., 2016.– Sex pheromones as isolating mechanisms in two closely related *Illiberis* species – *I. (Primilliberis) rotundata* Jordan, 1907, and *I. (P.) pruni* Dyar, 1905 (Lepidoptera: Zygaenidae, Procridinae).– *Entomologist's Gazette*, **67**(1): 51-57.
- SUBCHEV, M. A., KOSHIO, C., TOSHOVA, T. B. & EFETOV, K. A., 2012.– *Illiberis (Primilliberis) rotundata* Jordan



- (Lepidoptera: Zygaenidae: Procridinae) male sex attractant: Optimization and use for seasonal monitoring.– *Entomological Science*, **15**: 137-139. Available from <https://doi.org/10.1111/j.1479-8298.2011.00485.x>.
- SUBCHEV, M., KOSHIO, C., TOSHOVA, T., EFETOV, K. A. & FRANCKE, W., 2013.– (2R)-butyl (7Z)-dodecenoate, a main sex pheromone component of *Illiberis (Primilliberis) pruni* Dyar (Lepidoptera: Zygaenidae: Procridinae)?– *Acta Zoologica Bulgarica*, **65**(3): 391-396.
- TARMANN, G. M., EFETOV, K. A. & KUCHERENKO, E. E., 2019.– A second generation of *Theresimima ampellophaga* (Bayle-Barelle, 1808) (Lepidoptera: Zygaenidae, Procridinae) discovered by using the sex attractant EFETOV-2 on the Kassandra peninsula (Halkidiki) in Greece.– *Entomologist's Gazette*, **70**(1): 19-26.
- VRENOZI, B., TOSHOVA, T. B., EFETOV, K. A., KUCHERENKO, E. E., RREDHI, A. & TARMANN, G. M., 2019.– The first well-documented record of the vine bud moth *Theresimima ampellophaga* (Bayle-Barelle, 1808) in Albania established by field screening of sex pheromone and sex attractant traps (Lepidoptera: Zygaenidae, Procridinae).– *SHILAP Revista de lepidopterología*, **47**(187): 567-576.
- ZOLOTUHIN, V. V., 2020.– What is *Jordanita paupera* (Christoph, 1887) in the Lower Volga Region? (Lepidoptera: Zygaenidae).– *Eversmannia*, **62**: 3-11.

\*K. A. E.

Department of Biochemistry and Laboratory of Biotechnology  
V. I. Vernadsky Crimean Federal University  
RU-295051 Simferopol (Crimea)  
RUSSIA / RUSSIA  
E-mail: shysh1981@mail.ru  
E-mail: efetov.konst@gmail.com  
<https://orcid.org/0000-0003-1468-7264>

S. A. K.

Russian Entomological Society  
Irtyskaya Naberezhnaya, 14-16,  
RU-644042 Omsk  
RUSSIA / RUSSIA  
E-mail: konungomsk@yandex.ru  
<https://orcid.org/0000-0002-3887-0971>

y / and

Altai State University  
Lenina Str., 61  
RU-656049 Barnaul  
RUSSIA / RUSSIA

E. E. K.

Department of Biochemistry and Laboratory of Biotechnology  
V. I. Vernadsky Crimean Federal University  
RU-295051 Simferopol (Crimea)  
RUSSIA / RUSSIA  
E-mail: shysh1981@mail.ru  
<https://orcid.org/0000-0003-2567-8785>

\*Autor para la correspondencia / *Corresponding author*

(Recibido para publicación / *Received for publication* 3-V-2021)

(Revisado y aceptado / *Revised and accepted* 7-VI-2021)

(Publicado / *Published* 30-VI-2022)



**Figures 1-4.** **1.** Map of Kazakhstan with locality (red dot) of *Jordanita subsolana*: eastern Kazakhstan, Zaysan District, 19 km S of Zaysan, Saur Mountains. *Jordanita subsolana* is also known from Altai Republic (Russia), Ongudayskiy District (yellow dot) (EFETOV, 1990). **2.** Habitat of *Jordanita subsolana* in Kazakhstan: eastern Kazakhstan, Zaysan District, 19 km S of Zaysan, Saur Mountains. **3.** Male of *Jordanita subsolana* attracted to the lure EFETOV-2 on 1-2-VII-2018, 19 km S of Zaysan. **4.** Its genitalia.