

Report of a New Larval Host Plant of *Galleria mellonella* (Linnaeus, 1758) from Kolkata, West Bengal, India (Lepidoptera: Pyralidae, Galleriinae)

Srinjoy Das & Kaustav Chakraborty

Abstract

In this study, we report for the first time a new larval host plant *Cassia fistula* L. for *Galleria mellonella* (Linnaeus, 1758) from Kolkata, West Bengal, India. *Galleria mellonella* is an established pest of beehives and till date, its larvae were known to feed only on the wax comb of *Apis* sp., and destroy it. For this reason, beekeeping enterprises face high economic losses. Our finding opens a scope for extensive surveys in areas surrounding apiaries in order to quantify infestations in *C. fistula* which might provide a new insight in the pest management practices for *G. mellonella*.

Keywords: Lepidoptera, Pyralidae, Galleriinae, new host plant, *Galleria*, Kolkata, India.

**Informe de una nueva planta alimenticia de las larvas de *Galleria mellonella* (Linnaeus, 1758) en Calcuta, Bengala Occidental, India
(Lepidoptera: Pyralidae, Galleriinae)**

Resumen

En este estudio presentamos por primera vez una nueva planta alimenticia de las larvas de *Galleria mellonella* (Linnaeus, 1758), *Cassia fistula* L., de Calcuta, Bengala Occidental, India. *Galleria mellonella* es una plaga establecida de las colmenas y, hasta la fecha, se sabía que sus larvas sólo se alimentaban del panal de cera de *Apis* sp. y lo destruían. Por esta razón, las empresas apícolas se enfrentan a elevadas pérdidas económicas. Nuestro hallazgo abre la posibilidad de realizar estudios exhaustivos en las zonas circundantes a los colmenares para cuantificar las infestaciones de *C. fistula*, lo que podría aportar una nueva perspectiva sobre las prácticas de control de plagas de *G. mellonella*.

Palabras clave: Lepidoptera, Pyralidae, Galleriinae, nueva planta alimenticia, *Galleria*, Calcuta, India.

Introduction

The term wax moth is a common name which represents different species of moths that usually invade, attack and damage honeybee colonies and hive products. The list of wax moths are *G. mellonella* (Linnaeus, 1758), *Achroia grisella* (Fabricius, 1794), *Plodia interpunctella* (Hübner, [1813]) *Aphomia sociella* (Linnaeus, 1758) and *Ephestia kuehniella* Zeller, 1879. *G. mellonella*, the pest was classified *Tinea cerella* by Fabricius (1775), *Vindana obliquella* by Walker (1866) and later reclassified and named *G. melonella* (Linnaeus, 1758). Subfamily of *Galleria mellonella* is Galleriinae and its family is Pyralidae of order Lepidoptera. The species is well known for its parasitization of honeybees and their hives (Kwadha et al. 2017). *G.*

mellonella undergoes four developmental stages in its life cycle (egg, larva, pupa, and adult), so it is a typical holometabolous insect (Wojda et al. 2020). The eggs of this particular species vary in size (average length and width 0.478 mm and 0.394 mm), spheroidal shape with interspersed wavy lines which gives it a rough texture. Difference between male and female is not possible at larval stage, because Sex specific external morphological characters are absent. Polypod larvae with six legs on the thorax. The head of larva is composed of 3 well developed apical teeth but lacks sub-apical teeth with cream white body with sclerotized body parts. Male pupae are smaller in size than females. Pupa is white to yellow and gradually changes to dark brown. Sexual differentiation is possible in pupa. The adult moth shows sexual dimorphism. Males are generally smaller and less dark than females. The forewings for both sexes show varying intensities of pigmentation. The anterior two-third of the forewing is covered by scales that give it a uniformly darker pigmentation with comparison to the posterior one-third which is a mixture of stripes of darker and lighter pigment (Kwadha et al. 2017).

Female moths of *Galleria mellonella* (Linnaeus, 1758) lay eggs in honeybee comb cracks and crevices. The larvae feed honey, pollen, and brood, destroying the comb structure. Sometimes they show aggregation and cannibalism when there is a shortage of food (Kwadha et al. 2017). It was first reported in colonies of Asian honeybee, *Apis cerana* Fabricius, 1793. Williams (1997) and Shimanuki (1980), later designated this species as a pest as it is ubiquitously distributed everywhere beekeeping is practiced (Kwadha et al. 2017). After hatching, larvae move from cracks and crevices to the honeycomb where they begin to feed and build protective silken tubes, destroying the honeycomb structure in the process. This directional movement and feeding are stimulated chemically (Paddock, 1918; Nielson & Brister, 1979; Wojda et al. 2020). Worldwide record of larval host species of *Galleria melonella* is *Apis* sp. (Paddock, 1918; Nielsen & Briester, 1979; Shimanuki, 1980; Williams, 1997; Hosamani et al. 2017; Kwadha et al. 2017; Wojda et al. 2020).

Beekeeping has taken the shape of a promising enterprise. It is a very popular and highly economically beneficial integrated farming. India has a great scope for increasing bee farming. Bee-farming has two beneficial sides - firstly, the wax, honey, and different medical products, all have great market demands globally and secondly pollination services. But in India have some problems with beekeeping. One of the major problems is their natural enemies (Hosamani et al. 2017).

The *Galleria's* larval stage destroys *Apis* sp. For this reason, beekeepers face heavy economic loss, approximately 60-70% worldwide (Ambaw et al. 2020). The larvae can also damage the box hive frames in highly infected hives, it can cause a huge amount of economic loss for beekeepers (Ambaw et al. 2020).

Result and Discussion

The larvae are feed mainly honeybee comb substances, but we observed a new type of food is consumed by these larvae. On 28-VIII-2022, we collected a fruit of *Cassia fistula* L. from a place nearby the Mahanayak Uttam Kumar metro station, Tollygunge, Southern part of the Kolkata, West Bengal, India. Kolkata, the eastern part of India is a warm, humid place. High temperature along with enough humidity is one the characteristics of this place during summer. In winter season, supposed to be in December and January, the average high temperature remains around 27°C and low temperature remains around 13.8°C along with 11-17 mm average rainfall. During monsoon in July and August (to some extent the September month), this place accounts high rainfall. In July average rainfall is 411 mm and in August 349 mm respectively. This specimen is collected by us in the month of August 2022 when the average temperature was 26°C -32°C (Mitra, 2019) (Figure 1). It is a metropolitan city with Latitude 22.4986°N and Longitude 88.3454°E.

The next day we suddenly observed a hole in the fruit, cut around the hole, and found in the area the pupa shell of a moth covered in a silk cotton-like material and surrounded by old excreta left by moth's larvae (Figure 2). Hoping to know what came out of the hole; we placed the fruit

in a clean transparent container and covered its open mouth with a clean cloth. The next morning, we noticed that there were three more holes in the fruit as before, but this time the organisms that came out through the holes to stay covered could not escape. We took the opportunity to take photos of them and the photos were identified by one of the experts in the field as adult *Galleria melonella*.




Figure 1. Collection location of the *C. fistula* fruit - Mahanayak Uttam Kumar metro station, Tollygunge, Kolkata, West Bengal, India

In *Galleria melonella* sexual dimorphism is observable. *Galleria melonella* moth's body length is 15-20 mm and wingspan are 31 mm, darker grey in color. In *Galleria melonella*, the labial palps of female projecting forward which looks like a beak in the head region (Kwadha et al. 2017). In males semi-lunar notch present in outer margin of front wing (Hosamani et al. 2017). All these morphological characters are able to separate this moth from the close congeners.

This study suggests a new larval host plant for the larvae of *Galleria mellonella*, which is not previously reported elsewhere. Based on our observation, we can suggest that planting this host plant *Cassia fistula* around the beekeeping area may reduce the damage to the bee hives by this moth larvae. However, further studies are warranted for the outcome whether prefer this new host plants more than the wax hives or not. If so, this will give both-way benefits. It is very known to us that the ability to feed on multiple host species permits them to avail more resources. The larvae of the greater wax moth will get options and not depend fully on beehives. On the other hand, the practice of beekeeping or the apiary will be benefited to some extent and, so the apiculture industry. In the future, we have plans to explore more on it.

Acknowledgement

Authors are supported by their respective institutes. The first author would like to thank his mother Mrs. Shyamali Das and his friend Mr. Sahanur Mondal for maintenance and proper scientific caring.

References

- Ambaw, M., Teklehaimanot, T., & Workye, M., (2020). The prevalence of wax moth and associated risk factors in selected districts of Arsi Zone. *Journal of Entomology and Zoology Studies*, 8(1), 201-205.
- Hosamani, V., Hanumantha Swamy, B. C., Kattimani, K. N., & Kalibavi, C. M. (2017). Studies on Biology of greater wax moth (*Galleria mellonella* L.). *International journal of current microbiology and applied science*, 6(11), 3811-3815. <https://doi.org/10.20546/ijcmas.2017.611.447>
- Kwadha, C. A., Ong'amo, G. O., Ndegwa, P. N., Raina, S. K., & Fombong, A. T. (2017). The Biology and Control of the Greater Wax Moth, *Galleria mellonella*. *Insects*, 8(2), 61. <https://doi.org/10.3390/insects8020061>
- Mitra, S. (2019). Examining The Thermal Performance of Old Indigenous Architecture of Kolkata to Attain Sustainable Development. *International Journal of Advanced Research in Engineering & Technology*, 10(2). <https://doi.org/10.34218/IJARET.10.2.2019.039>
- Nielsen, R. A., & Briester, C. (1979). Greater wax moth: Behaviour of larvae. *Annals of the Entomological Society of America*, 72, 811-815. <https://doi.org/10.1093/aesa/72.6.811>
- Paddock, F. B. (1918). The Bee moth or Wax worm. Texas Agricultural Experiment Stations; College Station. TX, USA 217-234.
- Shimanuki, H. (1980). Diseases and Pests of Honey Bees. In *Beekeeping in the United States* (1980) (pp. 118-128). United States Department of Agriculture, Science and Education Administration.
- Williams, J. L. (1997). Insects: Lepidoptera (moths). In Morse R., Flottum K. (Ed.) (1997). *Honey Bees Pests, Predators, and Diseases* (pp. 121-141). AI Root Company.
- Wojda, I., Staniec, B., Sulek, M., & Kordaczu, J. (2020). The greater wax moth *Galleria mellonella*: biology and use in immune studies. *Pathogens and Disease*, 78(9), ftaa057. <https://doi.org/10.1093/femspd/ftaa057>

Srinjoy Das
Department of Zoology
Bidhannagar College
West Bengal State University
Kolkata - 700064, West Bengal
INDIA / INDIA
E-mail: srinjoydas1998@gmail.com
<https://orcid.org/0000-0002-1924-0499>

*Kaustav Chakraborty
Department of Zoology
S. B. S. Government College, Hili
University of Gour Banga
Dakshin Dinajpur - 733126, West Bengal
INDIA / INDIA
E-mail: kaustavc17@gmail.com
<https://orcid.org/0000-0002-6842-6193>

*Autor para la correspondencia / *Corresponding author*

(Recibido para publicación / *Received for publication* 14-VI-2023)

(Revisado y aceptado / *Revised and accepted* 18-X-2023)

(Publicado / *Published* 30-VI-2024)

Derechos de autor: El autor(es). Este es un artículo de acceso abierto distribuido bajo los términos de la Licencia de Reconocimiento 4.0 Internacional de Creative Commons (CC BY 4.0), que permite el uso, distribución y reproducción sin restricciones en cualquier medio, siempre que se cite al autor original y la fuente. / **Copyright:** The author(s). This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.




Figure 2. New larval host plant, *C. fistula* for *Galleria mellonella*. **A)** Adult moth. **B)** Exit hole of adult moth on the surface of *C. fistula* fruit. **C)** A silk-like cover of pupa of *G. mellonella* (L.) inside the fruit of *C. fistula*. **D)** Exuviae of a pupa inside silk-like cover. **E)** Excreta of larvae inside the plant fruit.



Available in:

<https://www.redalyc.org/articulo.oa?id=45579522007>

How to cite

Complete issue

More information about this article

Journal's webpage in redalyc.org

Scientific Information System Redalyc
Diamond Open Access scientific journal network
Non-commercial open infrastructure owned by academia

Srinjoy Das, Kaustav Chakraborty

Report of a New Larval Host Plant of *Galleria mellonella* (Linnaeus, 1758) from Kolkata, West Bengal, India (Lepidoptera: Pyralidae, Galleriinae)

Informe de una nueva planta alimenticia de las larvas de *Galleria mellonella* (Linnaeus, 1758) en Calcuta, Bengala Occidental, India (Lepidoptera: Pyralidae, Galleriinae)

Shilap Revista de Lepidopterología

vol. 52, no. 206, p. 255 - 259, 2024

Sociedad Hispano-Luso-Americana de Lepidopterología,

ISSN: 0300-5267

ISSN-E: 2340-4078

DOI: <https://doi.org/10.57065/shilap.901>