

Contribution to the biology of *Coleophora namaqua* Baldizzone & van der Wolf, 2015 from South Africa (Lepidoptera: Coleophoridae)

Thomas Kaltenbach & Giorgio Baldizzone¹

Abstract

Larvae of *Coleophora namaqua* Baldizzone & van der Wolf, 2015 were collected on their foodplant and reared to imagines. It is the first documented breeding of a *Coleophora* Hübner, 1822 species from South Africa. The larval case is described, and COI sequences were obtained. Further, variations in male and female genitalia are briefly discussed and illustrated.

Keywords: Lepidoptera, Coleophoridae, COI, *Coleophora namaqua*, foodplant, larval case, South Africa.

Contribución a la biología de *Coleophora namaqua* Baldizzone & van der Wolf, 2015 de Sudáfrica (Lepidoptera: Coleophoridae)

Resumen

Se recogieron larvas de *Coleophora namaqua* Baldizzone & van der Wolf, 2015, en su planta nutricia y se criaron hasta imagos. Se trata de la primera cría documentada de una especie de *Coleophora* Hübner, 1822 de Sudáfrica. Se describe el estuche larvario y se obtuvieron secuencias COI. Además, se discuten e ilustran brevemente las variaciones en la genitalia del macho y de la hembra.

Palabras clave: Lepidoptera, Coleophoridae, COI, *Coleophora namaqua*, planta nutricia, estuche larvario, Sudáfrica.

Introduction

During the last decades, lots of collection efforts for Microlepidoptera were done in Southern Africa. Amongst other groups, focus was also put on the genus *Coleophora* Hübner, 1822 (Coleophoridae), which turned out to be very diverse in that region, with more than 80 species reported so far (e.g. Baldizzone, 2021, 2022a, 2022b; Baldizzone & van der Wolf, 2011, 2015, 2020). *Coleophora* is one of the largest genera of Lepidoptera with more than 1550 species described worldwide. The big majority is distributed in the Holarctic region, but a substantial number is also known from tropical and subtropical regions, mainly in Africa (Baldizzone et al. 2006; Baldizzone, 2021). The imagines of *Coleophora namaqua* Baldizzone & van der Wolf, 2015, have a wingspan of 13-16 mm and brownish forewings with a distinct white line along the costa (Figure 1a), a habitus which they share with a number of other Afrotropical species and also with Palearctic species of the 9th species group of Toll's system (Baldizzone & van der Wolf, 2015; Toll, 1953). The differentiation of

¹ Contribution to the knowledge of Coleophoridae CLVIII

these species can be done with the morphology of the genitalia, the most similar species is *C. aarviki* Baldizzone & van der Wolf, 2011 (Baldizzone & van der Wolf, 2015). Here, we describe the larval case and report the foodplant of *C. namaqua*, both hitherto unknown, based on the first rearing of a *Coleophora* species in South Africa.

Materials and methods

Eight larvae were found on their foodplant on 03-XI-2014 during a search specifically for *Coleophora* larvae in the Cedar Mountains and kept in a carton container covered with gauze. For two days, twigs with leaves were added into the container, as two larvae were still feeding. Thereafter, all larvae remained in the same container during pupation, first for about ten days in South Africa and thereafter at room temperature in Switzerland.

For the genitalia, the abdomens were at first treated with heated KOH, the dissection was done in Cellosolve (2-Ethoxyethanol) with subsequent mounting on slides with Euparal liquid, using an Olympus SZX7 stereomicroscope.

The DNA of part of the specimens was extracted using a hindleg of two different specimens. We amplified a 658 bp fragment of the mitochondrial gene cytochrome oxidase subunit 1 (COI) using the primers LCO 1490 and HCO 2198 (Folmer et al. 1994). Sequencing was done with Sanger's method (Sanger et al. 1977). The genetic variability between two specimens was estimated using Kimura-2-parameter distances (K2P; Kimura, 1980), calculated with the program MEGA 11 (Tamura et al. 2021, <http://www.megasoftware.net>).

Photograph of an imago in toto was taken using a Canon EOS 6D camera and processed with the programs Adobe Photoshop Lightroom (<http://www.adobe.com>) and Helicon Focus version 5.3 (<http://www.heliconsoft.com>). Pictures of parts of genitalia were taken with an Olympus BX43 microscope, processed with the program Olympus Cell Sense v. 3.2. Photographs were subsequently enhanced with Adobe Photoshop Elements 13.

Results

Material examined: SOUTH AFRICA, Cedar Mountains, near Kagga Kamma Lodge; 32°44'45''S, 19°33'46''E; 1.073 m; 1 ♂, 2 ♀; 03-XI-2014; e. 1. *Wiborgia obcordata* 02-03-XII-2014; leg. Th. Kaltenbach; genitalia preparations GBIFCH01221758, GBIFCH01221757.

Remarks to variations in the genitalia: The male genitalia show variations in the sacculus and the cornuti (Baldizzone & van der Wolf, 2015, figs 35-38). The male specimen in this study has genitalia identical to Figure 37 in Baldizzone & van der Wolf (2015), but cornuti, which are different in number and orientation (Figure 1e). The female genitalia of a specimen in this study are as shown in the original description (Baldizzone and van der Wolf, 2015, figs 40-41). However, the signum has a small spine (Figure 1d), which was also discovered in the genitalia of a paratype (GP BLDZ 15005).

Description of larval case and remarks to life-history: Eight larval cases in final stage were found on their foodplant *Wiborgia obcordata* (P. J. Bergius) Thunb. (Fabaceae). Two of the larvae were still feeding by mining the very small leaves from beneath. Abandoned mines in the leaves were also observed in the neighborhood.

One case of the young larva (Figure 1b) was found nearby a feeding mature larva, attached to a twig. It is made of two segments of leaf cuticle, flattened laterally, length ca. 4 mm, color yellow brown. The anal opening is bivalve, the oral opening is set at ca. 45° to the long axis. We assume that the first case initially is made from one segment of leaf cuticle and later completed with a second segment. However, it remains unclear, if the first case is used by the larva for the construction of the final case or just abandoned and left behind during the construction of the final case.

The case in its final form (Figure 1c) is elongate tubular, slightly flattened laterally. It is composed of mostly three segments of leaf cuticle; anal segment conical, partially with a ventral keel consisting of the leaf cuticle margin; length 9-11 mm, color yellow brown to brown. The anal opening is bivalve, the

oral opening is set at ca. 40° to the long axis. Pupation as usually in the case, attached to twigs of the foodplant or adjacent places.

The habitat where the specimens were collected in the Cedar Mountains above 1000 m is a semi desert scrub with sandstone rock formations and sandy soil of reddish color (Figure 1f). Nearby are areas of Mediterranean fynbos, typical for this region in South Africa. It is hot and arid during summer and cold with some rainfall during winter. The vegetation is loose, and drought adapted with shrubs like *Wiborgia obcordata* as highest plants.

Genetics: The two COI sequences obtained from two specimens of *C. namaqua* from the same location have a genetic distance of 0% (K2P), as it is expected in such a case. The sequences are accessible on GenBank with the following numbers: PP297496, PP297497.

Discussion

Not much is known about the larval cases and the foodplants of Afrotropical species of *Coleophora* in general. Apart from *C. namaqua* treated in this study, only the larval case and foodplant of *C. enchitis* Meyrick, 1920 from Kenya are known (Baldizzone & van der Wolf, 2015). The foodplant of *C. namaqua*, *Wiborgia obcordata*, belongs to the family Fabaceae. Interestingly, the known larvae of species belonging to the 9th group of Toll (Toll, 1953), to which *C. namaqua* has affinities, also feed on Fabaceae. The knowledge on larval cases, foodplants and life-history of this megadiverse genus is an important prerequisite to better understand the phylogeny of this group in future studies (see Bauer et al. 2012). Therefore, it is recommended to also focus on the collection and rearing of *Coleophora* larvae during future field trips in the Afrotropical region.

Acknowledgements

We sincerely thank Céline Stoffel (MZL, UNIL) for her support with lab work and preparation of the COI barcodes. Further we are very grateful to Dr. Antonio Vives for the translation of the abstract into Spanish language. We are also thankful to the reviewers for their valuable comments.

References

- Baldizzone, G. (2021). On the taxonomy of Afrotropical Coleophoridae (VI). New species of the genus *Coleophora* Hübner, 1822 from South Africa (Lepidoptera, Coleophoridae). *Zootaxa*, 5071, 167-205. <https://doi.org/10.11646/zootaxa.5071.2.1> PMID:34810673
- Baldizzone, G. (2022a). On the taxonomy of Afrotropical Coleophoridae (V). New species of the genus *Coleophora* from Namibia (Lepidoptera, Coleophoridae). *Revue Suisse de Zoologie*, 129, 19-49. <https://doi.org/10.35929/RSZ.0060>
- Baldizzone, G. (2022b). On the taxonomy of Afrotropical Coleophoridae (VII). New species of genus *Coleophora* Hübner, 1822 from Central, Southern Africa and Oman (Lepidoptera, Coleophoridae). *Zootaxa*, 5133, 431-442. <https://doi.org/10.11646/zootaxa.5133.3.7> PMID:36101090
- Baldizzone, G., & van der Wolf, H. W. (2011). On Afrotropical Coleophoridae (I) (Lepidoptera: Coleophoridae). *SHILAP Revista de lepidopterología*, 39(156), 351-377. <http://www.redalyc.org/articulo.oa?id=45522548003>
- Baldizzone, G., & van der Wolf, H. W. (2015). On the taxonomy of Afrotropical Coleophoridae (II) (Lepidoptera, Coleophoridae). *Zootaxa*, 4000, 335-362. <http://dx.doi.org/10.11646/zootaxa.4000.3.2> PMID:26623620
- Baldizzone, G., & van der Wolf, H. W. (2020). On the taxonomy of Afrotropical Coleophoridae (IV). New or little-known species from South Africa (Lepidoptera, Coleophoridae). *Zootaxa*, 4816, 151-170. <https://doi.org/10.11646/zootaxa.4816.2.1> PMID:33055701
- Baldizzone, G., van der Wolf, H. W., & Landry, J.-F. (2006). Coleophoridae, Coleophorinae (Lepidoptera). In B. Landry (Ed.). *World Catalogue of Insects* (Vol. 8). Apollo Books. <https://doi.org/10.1163/9789004475403>
- Bauer, F., Stübner, A., Neinhuis, C., & Nuss, M. (2012). Molecular phylogeny, larval case architecture, host-plant associations, and classification of European Coleophoridae (Lepidoptera). *Zoologica Scripta*, 41, 248-265. <https://doi.org/10.1111/j.1463-6409.2012.00532.x>

- Folmer, O., Black, M., Hoeh, W., Lutz, R., & Vrijenhoek, R. (1994). DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from divers' metazoan invertebrates. *Molecular Marine Biology and Biotechnology*, 3, 294-299. http://www.mbari.org/staff/vrijen/PDFS/Folmer_94MMBB.pdf
- Kimura, M. (1980). A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. *Journal of Molecular Evolution*, 16, 111-120. <https://doi.org/10.1007/BF01731581> PMID:7463489
- Sanger, F., Nicklen, S., & Coulson, A. R. (1977). DNA sequencing with chain-terminating inhibitors. *Proceedings of the National Academy of Sciences U.S.A.*, 74, 5463-5467. <https://doi.org/10.1073/pnas.74.12.5463> PMID:271968 PMCID:PMC431765
- Tamura, K., Stecher, G., & Kumar, S. (2021). MEGA 11: molecular evolutionary genetics analysis version 11. *Molecular Biology and Evolution*, 38, 3022-3027. <https://doi.org/10.1093/molbev/msab120> PMID:33892491 PMCID:PMC8233496
- Toll, S. (1953). Rodzina Eupistidae Polski. *Documenta Physiographica Poloniae*, 32, 1-293, 38 pls. [1952].

*Thomas Kaltenbach
Muséum cantonal des Sciences Naturelles
Département de Zoologie
Palais de Rumine, Place Riponne, 6
CH-1005 Lausanne
SUIZA / SWITZERLAND
E-mail: thomas.kaltenbach@bluewin.ch
<https://orcid.org/0000-0001-8052-0388>

Giorgio Baldizzone
Via Manzoni, 24
I-14100 Asti
ITALIA / ITALY
E-mail: baldizzonegiorgio@gmail.com
<https://orcid.org/0000-0001-8127-0843>

y / and

y / and

University of Lausanne (UNIL)
Department of Ecology and Evolution
CH-1015 Lausanne
SUIZA / SWITZERLAND

Museum d'Histoire Naturelle de Genève
C.P. 6434
CH-1211 Geneve 6
SUIZA / SWITZERLAND
(Corresponding member)

*Autor para la correspondencia / *Corresponding author*

(Recibido para publicación / *Received for publication* 9-II-2024)

(Revisado y aceptado / *Revised and accepted* 19-III-2024)

(Publicado / *Published* 30-XII-2024)

Derechos de autor: El autor(es). Este es un artículo de acceso abierto distribuido bajo los términos de la Licencia de Reconocimiento 4.0 Internacional de Creative Commons (CC BY 4.0), que permite el uso, distribución y reproducción sin restricciones en cualquier medio, siempre que se cite al autor original y la fuente. / **Copyright:** The author(s). This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.




Figure 1. *Coleophora namaqua*: **a.** female. **b.** first larval case. **c.** final larval case. **d.** signum of female genitalia (arrow: small spine) **e** cornuti of male genitalia **f** habitat in Cedar Mountains, with shrubs of *Wiborgia obcordata* **g** *Wiborgia obcordata*. Scale bars: 1 mm (a, b, c), 20 µm (d, e).



Available in:

<https://www.redalyc.org/articulo.oa?id=45581999002>

How to cite

Complete issue

More information about this article

Journal's webpage in redalyc.org

Scientific Information System Redalyc
Diamond Open Access scientific journal network
Non-commercial open infrastructure owned by academia

Thomas Kaltenbach, Giorgio Baldizzone
**Contribution to the biology of *Coleophora namaqua*
Baldizzone & van der Wolf, 2015 from South Africa
(Lepidoptera: Coleophoridae)**
**Contribución a la biología de *Coleophora namaqua*
Baldizzone & van der Wolf, 2015 de Sudáfrica
(Lepidoptera: Coleophoridae)**

Shilap Revista de Lepidopterología
vol. 52, no. 208, p. 677 - 681, 2024
Sociedad Hispano-Luso-Americana de Lepidopterología,
ISSN: 0300-5267
ISSN-E: 2340-4078

DOI: <https://doi.org/10.57065/shilap.1010>