



DYNA

ISSN: 0012-7353

ISSN: 2346-2183

Universidad Nacional de Colombia

Marín-Arango, Zaira T.; Cortes-Rodríguez, Misael;
Montoya-Campuzano, Olga I.; Arango-Tobón, Julio C.
Viability of *Lactobacillus casei* ATCC 393 and properties in Andean
blackberry suspensions with probiotic and prebiotic characteristics
DYNA, vol. 86, no. 210, 2019, July-September, pp. 179-186
Universidad Nacional de Colombia

DOI: <https://doi.org/10.15446/dyna.v86n210.72929>

Available in: <https://www.redalyc.org/articulo.oa?id=49662789023>

- How to cite
- Complete issue
- More information about this article
- Journal's webpage in redalyc.org

UNAM

Scientific Information System Redalyc
Network of Scientific Journals from Latin America and the Caribbean, Spain and
Portugal

Project academic non-profit, developed under the open access initiative

Viability of *Lactobacillus casei* ATCC 393 and properties in Andean blackberry suspensions with probiotic and prebiotic characteristics

Zaira T. Marín-Arango ^a, Misael Cortes-Rodríguez ^b, Olga I. Montoya-Campuzano ^c & Julio C. Arango-Tobón ^b

^a Facultad de Ciencias Agroindustriales, Universidad del Quindío, Armenia, Colombia. ztmarin@uniquindio.edu.co

^b Facultad de Ciencias Agrarias, Universidad Nacional de Colombia, Sede Medellín, Medellín, Colombia. mcortesro@unal.edu.co, jcarango@unal.edu.co

^c Facultad de Ciencias, Universidad Nacional de Colombia, Sede Medellín, Medellín, Colombia. oimontoy@unal.edu.co

Received: June 18th, 2018. Received in revised form: September 21th, 2018. Accepted: November 25th, 2018.

Abstract

The food industry has grown in recent years due to the supply of functional foods with probiotics and prebiotics. The viability of *Lactobacillus casei* and the properties of blackberry suspensions added with inulin were evaluated using a central design composed of the following factors: mass fraction of total solid blackberry concentrate (X_{TSBC}) (0.103-0.120), [McFarland] (6-10), inulin (2.78-3.68%), and mass blackberry concentrate (BC)/mass inoculum (R) (5-20); and the dependent variables: moisture (M), pH, acidity, °Brix, total solids suspension (TSs), viscosity (μ), zeta potential (ζ), and Log CFU/g. Results of the experimental optimization of multiple responses were: X_{TSBC} (0.103), [McFarland] (10), inulin (3.12%w/w), and R (18.5); being the dependent variables: $M=86.7\pm0.0\%$, $pH=2.74\pm0.01$, $acidez=3.1\pm0.0\%$, $^{\circ}Brix=11.0\pm0.1\%$, $TSs=13.3\pm0.0\%$, $\mu=1288.0\pm20.1$ cP, $\zeta=-13.3\pm0.3$ mV and Log CFU/g=7.8±0.5. Probiotic microorganism *Lb. casei* ATCC 393 presents adaptability in suspensions based on BC and inulin, which guarantee its viability and possible use for obtaining an innovative spray-dried product.

Keywords: *Rubus glaucus* Benth; functional food; inulin; colloids.

Viabilidad de *Lactobacillus casei* ATCC 393 y propiedades en suspensiones de mora de castilla con características probióticas y prebióticas

Resumen


La industria alimentaria en los últimos años ha crecido principalmente por la oferta de alimentos funcionales con probióticos y prebióticos. Se evaluó la viabilidad del *Lactobacillus casei* y las propiedades de suspensiones de mora adicionadas con inulina, utilizando un diseño central compuesto con los factores: fracción másica sólidos totales concentrado de mora (X_{STCM}) (0,103-0,120), [McFarland] (6-10), inulina (2,78-3,68%) y masa (CM)/masa inóculo (R) (5-20) y las variables dependientes: humedad (H), pH, acidez, °Brix, sólidos totales suspensión (STs), viscosidad (μ) potencial zeta (ζ) y LogUFC/g. Los resultados de la optimización experimental de múltiples respuestas fueron: X_{STCM} (0,103), [McFarland] (10), Inulina (3,12 %p/p) y R (18,5), siendo las variables dependientes: $H=86,7\pm0,0\%$, $pH=2,74\pm0,01$, $acidez=3,1\pm0,0\%$, $^{\circ}Brix=11,0\pm0,1\%$, $STs=13,3\pm0,0\%$, $\mu=1288,0\pm20,1$ cP, $\zeta=-13,3\pm0,3$ mV y LogUFC/g=7,8±0,5. El microorganismo probiótico *Lb. casei* ATCC 393 presenta una adaptabilidad en suspensiones a base de CM e inulina, que garantizan su viabilidad y un posible uso para la obtención de un producto innovador secado por aspersión.

Palabras clave: *Rubus glaucus* Benth; alimento funcional; inulina; coloides.

1. Introduction

Blackberry (*Rubus glaucus* Benth) is a berry rich in phenolic components such as anthocyanins [1] and

How to cite: Marín-Arango, Z.T., Cortes-Rodríguez, M., Montoya-Campuzano, O.I. and Arango-Tobón, J.C., Viability of *Lactobacillus casei* ATCC 393 and properties in Andean blackberry suspensions with probiotic and prebiotic characteristics. DYNA, 86(210), pp. 179-186, July - September, 2019.

© The author; licensee Universidad Nacional de Colombia. 
Revista DYNA, 86(210), pp. 179-186, July - September, 2019, ISSN 0012-7353
DOI: <http://doi.org/10.15446/dyna.v86n210.72929>

ellagitannins [2] which are related to health, the decrease of cancer cells [1,3], protection against liver damage, blood pressure reduction, anti-inflammatory and antimicrobial activities [1], among others. This fruit has a high respiratory rate, which makes it a perishable product and, therefore, it has a short postharvest life and limits its use as fresh fruit [1]; for this purpose, some research has reported technological alternatives which allow delaying respiration rate and moisture loss: hypobaric storage, protective films [4], refrigeration, traditional and rapid freezing, spray drying, modified atmospheres and irradiation [1,5]. Its flavor and functional properties make blackberry a much-desired fruit all around the world [6]. The United States is the largest exporter of blackberry in the world, followed by Spain and Mexico [7], 90% of production is converted into processed products such as nectars, juices, frozen pulps, concentrates, jams, jellies, and dehydrated powders, and the remaining 10% is destined to fresh consumption [6].

Nowadays, world market demands specific attributes on functional foods, which are those that, besides providing basic nutrients, contain components with physiological activity or active components which contribute to the prevention of some diseases [3]. In this context, one of the components most widely used in food industry is *Lactobacillus casei*, a gram-positive microaerophilic, auxotroph bacillus, part of the microbiota of fruits, vegetables, meat, milk, wine, intestinal and vaginal mucous, able to reach the human intestine and multiply without damaging the host [8-10]. In addition, it has a series of healthy properties: it prevents the alteration of intestinal microbiota, helps to eliminate cholesterol and reduces osteoporosis risk, among others [11]. Therefore, it is considered a probiotic microorganism widely used in the development of functional foods [12]. By regulation, a functional food with probiotic characteristics must contain a number $\geq 1 \times 10^6$ CFU/g or mL until the end of its useful life [9,13-15].

Literature does not report viability studies of probiotic microorganisms in blackberry-based colloidal. However, there is interest in using different dietary matrices to maintain the viability of these probiotics and thus offer the modern consumer innovative, stable and healthy products within the range of functional foods [16]. Many of these foods range from suspensions to dry products, where the guarantee of microorganism viability is essential at any stage of the process. For example, pear juice obtained levels of 10^7 CFU/mL of *Lb. casei* inoculated at 2% [17], Cape gooseberry pulp inoculating a [McFarland] of 0.5 *Lb. casei* ATCC 393 obtained 5.4×10^8 CFU/mL [18], while guava juice 8 Log CFU/mL units of encapsulated *Lb. casei* 431 were inoculated obtaining 7.06 Log CFU/g [19]. On the other hand, characteristics of foods can influence strains survival, so it is convenient to protect probiotics by reducing their physical exposure during the passage through the gastrointestinal tract, either through encapsulants or with the addition of prebiotics that helps them survive during the passage through the stomach [20].

The aims of this investigation was to evaluate microbial cell viability of the commercial strain *Lactobacillus casei* ATCC 393 and the properties in blackberry suspensions

added with inulin as a prebiotic agent.

2. Materials and methods

2.1. Materials

Blackberry concentrate (BC) (Nutrium SAS) was used with specifications according to the technical data of °Brix: 13-15%, citric acidity: 3.1-5.5, and pH: 2.5-3.5. The commercial strain was *Lactobacillus casei* ATCC393. It was activated in 5 mL of MRS broth (Man, Rogosa and Sharpe) (Scharlau) at 37 °C for 72 h in anaerobic conditions, and added in 200 mL of the same broth. Then, it was incubated under the same conditions to generate biomass. Finally, Fibruline S20 inulin (Cosucra, Groupe Warcoing SA) was used as a prebiotic agent.

2.2. Preparation of the inoculum

The microorganism suspended in MRS broth was centrifuged at 2000 rpm for 10 min at 25 °C, separating the biomass used to prepare McFarland standards 6, 8 and 10 by optical density. The microorganism was diluted in sterile water and taken to spectrophotometry for reading at 625 nm [18].

2.3. Experimental design and preparation of suspensions

It was carried out using response surface methodology with a composite central design, face-centered ($\alpha = 1$) with 21 experiments (Table 1), considering the independent variables: mass fraction of TS contributed by BC (X_{TSBC}) (0.103 - 0.120), [McFarland] (6 - 10), inulin (2.78 - 3.68%), and R (5 - 20); and the dependent variables: moisture (M), pH, acidity, °Brix, TS in final suspension (TS_s), viscosity (μ), zeta potential (ζ), and Log CFU/g. The valuations were made at times 0, 24 and 48 hours (T_0 , T_{24} , T_{48}). The quantities of each component were determined from material balances for batches of 250g; initially, a suspension of BC was prepared to X_{TSBC} of design, using a Silverson homogenizer L5M-A at 10,000 rpm, then inulin and finally inoculum biomass with desired [McFarland] were added. This suspension was refrigerated at 4 °C and stored at times of 0, 24, and 48 h.

2.4. Viability of *Lb. casei*

It was determined by plate count of microorganisms. Initially, 10 g of suspension were homogenized with 90 mL of universal peptone (0.1% w/v) using 80 Lab Blender stomacher, then successive dilutions were made up to 10^6 with universal peptone in a ratio of 1:9, which were inoculated 0.1 mL in petri dishes with MRS agar by the planting on surface method; finally, they were incubated at 37 °C for 72 h under anaerobic conditions.

2.5. Characterization of suspensions

Acidity tests were performed according to AOAC method 942.15/2012 expressed as malic acid, by titration with 0.1N

NaOH, and using phenolphthalein as indicator [21]; pH by AOAC method 981.12/2012 with a Schott potentiometer CG840B [21]; °Brix according to AOAC method 932.12/2012 with a Leica auto ABBE refractometer [21]; M following AOAC method 20.013/2012 with vacuum oven at 60 °C until constant weight [21]; TSs was determined by difference with M in the final suspension; μ was measured with a Brookfield viscometer DV-III Ultra at 100 rpm, RV-5 spindle at 25 °C [22]; water activity (a_w) according to AOAC method 978.18/2012 with a dew point hygrometer Aqualab 4TE, Decagon Devices USA at 25 °C [21]; ζ was determined on a Zetasizer Nano ZS90 (Malvern Instruments Ltda.) using a capillary electrophoresis cell [23]; and optical density with a Thermo Scientific Evolution 60 UV/VIS spectrophotometer [24].

2.6. Data analysis

Results were analyzed using Statgraphics Centurion XVI® software, by means of the optimization of multiple responses and the analysis of variance (ANOVA) at a confidence level of 95%. All analyzes were performed in triplicate.

3. Results and analysis

The characterization of BC presented values of M: $83.97 \pm 0.02\%$ and a_w : 0.990 ± 0.002 , which makes it a product with characteristics suitable for the growth of beneficial microorganisms, alterants and pathogens [25]. TS values: $16.03 \pm 0.02\%$ and °Brix: $12.10 \pm 0.14\%$ (soluble in liquid fraction), which affect the osmotic pressure of food system and at the same time the viability of microorganisms of interest [26]; acidity values: $5.85 \pm 0.16\%$ and pH: 2.74 ± 0.00 , which frames it within acidic foods; ζ values : -11.17 ± 0.60 mV which identifies BC as a colloidal system with repulsive electric forces of negative charge in the proximity of coions layer, formed around the interphase of the suspended particles [27] and for this case, negative ions obtained from the dissociation of salts present in BC [28]. Blackberry pulp has been identified as a non-Newtonian, pseudoplastic liquid [29]; while, BC is a much more viscous product (μ : 5093.3 ± 201.7 cP), which could affect the mobility of suspended particles, contributing to the colloidal system stability.

Some authors have reported different results in food matrices based in blackberry: [28] in BC (citric acidity: 1.67%, pH: 2.89 and °Brix: 65.0%), and in blackberry pulp (citric acidity: 1.65%, pH: 2.86 and °Brix: 8.9%); [1] in blackberry pulp (citric acidity: 0.76%, pH: 3.31, °Brix: 6.77% and M: 91.96%); [30] in blackberry pulp (citric acidity: 1.88%, pH: 3.03, Brix: 9.06, M: 83.5% and a_w : 0.984) and finally, [31] in blackberry pulp (pH: 3.24; Brix: 8.5%, M: 87.7% and a_w : 0.981). A variability is observed when compared with the results obtained in BC of the present investigation, which is attributable on the one hand to production conditions and concentration levels of BC; on the other hand, the differences between pulps for the same variety of fruit, which is affected by other pre-harvest factors: climatic, solar radiation, soil, rain, among others [32], and

post-harvest factors such as maintenance of the cold chain [33].

Table 1 shows the mean values and standard deviation of Log CFU/g of blackberry inoculated with *Lb. casei* and inulin suspensions evaluated at times 0, 24, and 48 h (T_0 , T_{24} , T_{48}) and the physical and physicochemical properties of suspensions at T_0 . A very definite tendency is observed in suspensions, where Log CFU/g units decrease with storage time; therefore, T_0 was selected for its highest content of probiotic microorganisms. This situation is attributable to different factors, on the one hand, the effect of osmotic, thermal and oxidative stress on *Lactobacillus* cell wall; on the other hand, to the effect of the pH of BC itself. The pH has an effect on growth and viability of microbial cells, because intracellular pH of bacteria remains close to neutrality and the decrease of extracellular pH can lead to the destruction of cellular macromolecules, since these are sensitive to acids, especially weak acids that enter cells and dissociate (H^+) in the cytoplasm, decreasing the ability of cells to generate energy [34]. A sudden increase in external osmolarity results in the elimination of water from the cell to the outside, producing loss of turgor, as well as changes in intracellular solute concentration and cell volume, decreasing metabolic activity and therefore lower viability [35]. These types of stress, by subjecting the microorganism to a prolonged exposure over time can lead cells to microbial death; however, if microorganisms are subjected to an acid stress with pH<4.5 in short times, heat shock proteins can be produced in the microorganism, helping them to generate resistance to heat stress [36].

Fig. 1 shows response surface graphs of dependent variables with respect to independent variables; while Table 2 presents the ANOVA that identifies dependent variables showing significant differences ($p < 0.05$) with respect to independent variables, with linear and quadratic interactions. The average values obtained from plate count varied between 5.74 and 7.79 Log CFU/g, equivalent to 5.5×10^5 and 6.2×10^7 CFU/g, which indicates that some values do not meet the minimum required by Colombian and international regulations (10^6 CFU/g) [12, 37].

The behavior of Log CFU increases mainly when [McFarland] is higher than 9, being enhanced when R is minimum ($R = 5$) and [McFarland] = 10, where logarithmic units per gram reached their highest values: 7.79 ± 0.03 (inulin: 2.78% and X_{TSBC} : 0.103) and 7.79 ± 0.05 (inulin: 3.68% and X_{TSBC} : 0.120). This situation is attributed to the fact that inoculation is done at the highest microbial concentration and with the highest mass of inoculum, which results in a good microorganism recovery.

Some authors have investigated the viability of different microorganisms in various food matrices and prebiotic agents: in yogurt and orange juice complemented with inulin and galacto-oligosaccharides (GOS). *Lb. acidophilus* and *Lb. casei* 01 were evaluated during storage at 4°C for 4 weeks, increasing probiotic protection and growth as a result of implementing GOS (1.5%) ($>10^7$ CFU/g or mL) compared to a control without GOS; obtaining counts for yogurt above control in logarithmic units of 1.1 and 0.4 for *Lb. acidophilus* and *Lb. casei* 01 respectively [38].

Table 1.
Log CFU/g of blackberry suspensions inoculated with *Lb. casei* and inulin at times of 0, 24 and 48 hours..

Run	Independent variables			Dependent variables										
	X _{TSBC}	R	[McFarland] Inulin (%)	T ₀		T ₂₄		T ₄₈		T ₀				
				Log CFU/g	Log CFU/g	Log CFU/g	Log CFU/g	M (%)	TS _s (%)	°Brix (%)	Acidity (%)	pH	ζ(mV)	μ(cP)
1	0.112	12.5	10	3.23	7.01 ± 0.08	6.16 ± 0.30	5.72 ± 0.28	85.65 ± 0.44	14.35 ± 0.44	12.05 ± 0.06	4.21 ± 0.44	2.68 ± 0.01	-12.03 ± 0.33	2713 ± 52.50
2	0.112	12.5	8	3.68	5.95 ± 0.03	5.38 ± 0.04	4.78 ± 0.20	83.95 ± 1.62	16.05 ± 1.62	12.15 ± 0.06	3.50 ± 0.23	2.84 ± 0.00	-14.25 ± 0.94	938 ± 13.66
3	0.112	20.0	8	3.23	5.75 ± 0.05	5.35 ± 0.14	4.15 ± 0.17	85.75 ± 0.10	14.25 ± 0.10	12.10 ± 0.08	4.07 ± 0.42	2.79 ± 0.01	-13.23 ± 0.19	1286 ± 20.78
4	0.103	20.0	10	3.68	7.12 ± 0.12	6.37 ± 0.03	6.04 ± 0.30	86.00 ± 0.00	14.00 ± 0.00	11.90 ± 0.18	3.52 ± 0.03	2.90 ± 0.07	-13.53 ± 0.91	773 ± 11.94
5	0.112	12.5	8	3.23	5.74 ± 0.01	5.06 ± 0.07	4.50 ± 0.23	85.75 ± 0.21	14.25 ± 0.21	12.23 ± 0.05	3.85 ± 0.04	2.79 ± 0.01	-12.30 ± 0.64	2013 ± 169.15
6	0.112	5.0	8	3.23	6.23 ± 0.10	6.33 ± 0.10	6.06 ± 0.07	85.93 ± 0.05	14.08 ± 0.05	12.15 ± 0.06	4.16 ± 0.08	2.79 ± 0.01	-10.70 ± 0.61	1798 ± 15.49
7	0.103	20.0	6	3.68	7.49 ± 0.08	6.87 ± 0.15	7.13 ± 0.22	86.45 ± 0.10	13.55 ± 0.10	11.55 ± 0.35	3.37 ± 0.12	2.72 ± 0.00	-12.75 ± 0.99	1303 ± 16.12
8	0.112	12.5	8	3.23	5.91 ± 0.09	5.08 ± 0.28	4.15 ± 0.17	85.38 ± 0.05	14.63 ± 0.05	11.93 ± 0.10	3.88 ± 0.22	2.68 ± 0.01	-12.03 ± 0.84	1617 ± 8.25
9	0.120	20.0	6	2.78	6.37 ± 0.05	6.35 ± 0.01	6.23 ± 0.04	85.30 ± 0.08	14.70 ± 0.08	12.33 ± 0.05	4.32 ± 0.06	2.72 ± 0.01	-11.55 ± 0.29	2384 ± 5.66
10	0.120	12.5	8	3.23	6.07 ± 0.14	5.89 ± 0.04	5.36 ± 0.15	84.90 ± 0.08	15.10 ± 0.08	12.20 ± 0.08	4.18 ± 0.01	2.72 ± 0.01	-11.48 ± 0.19	2306 ± 13.27
11	0.112	12.5	6	3.23	6.39 ± 0.04	6.28 ± 0.03	6.25 ± 0.02	85.73 ± 0.05	14.28 ± 0.05	12.08 ± 0.05	4.27 ± 0.13	2.73 ± 0.00	-11.83 ± 0.34	1797 ± 3.83
12	0.103	5.0	6	2.78	7.10 ± 0.04	6.78 ± 0.37	6.99 ± 0.23	86.88 ± 0.05	13.13 ± 0.05	10.95 ± 0.21	4.27 ± 0.10	2.72 ± 0.00	-11.65 ± 0.13	1496 ± 16.00
13	0.112	12.5	8	3.23	6.43 ± 0.17	6.09 ± 0.02	6.14 ± 0.13	85.65 ± 0.06	14.35 ± 0.06	11.95 ± 0.17	3.32 ± 0.16	2.89 ± 0.01	-13.88 ± 0.30	962 ± 14.79
14	0.112	12.5	8	2.78	6.02 ± 0.07	6.09 ± 0.23	4.78 ± 0.00	89.25 ± 3.18	10.75 ± 3.18	11.83 ± 0.24	2.09 ± 0.19	2.91 ± 0.13	-13.10 ± 0.22	893 ± 27.40
15	0.103	12.5	8	3.23	6.48 ± 0.18	6.40 ± 0.14	6.15 ± 0.12	88.55 ± 2.41	11.45 ± 2.41	11.60 ± 0.29	1.88 ± 0.03	2.85 ± 0.01	-13.25 ± 0.69	651 ± 22.24
16	0.120	5.0	6	3.68	7.08 ± 0.01	6.75 ± 0.01	5.30 ± 0.35	84.75 ± 0.47	15.25 ± 0.47	13.25 ± 0.13	4.85 ± 0.12	2.70 ± 0.01	-11.35 ± 0.42	2185 ± 12.81
17	0.103	5.0	10	2.78	7.79 ± 0.03	6.54 ± 0.07	6.51 ± 0.04	86.53 ± 0.05	13.48 ± 0.05	10.98 ± 0.15	3.91 ± 0.12	2.70 ± 0.01	-11.80 ± 0.54	1404 ± 27.71
18	0.120	20.0	10	2.78	6.14 ± 0.04	5.78 ± 0.01	5.79 ± 0.05	85.20 ± 0.08	14.80 ± 0.08	12.15 ± 0.06	4.24 ± 0.12	2.82 ± 0.01	-12.40 ± 1.02	1479 ± 25.38
19	0.120	5.0	10	3.68	7.79 ± 0.05	6.10 ± 0.00	6.68 ± 0.05	84.38 ± 0.05	15.63 ± 0.05	12.88 ± 0.53	2.21 ± 0.07	2.85 ± 0.02	-11.83 ± 1.04	1131 ± 12.38
20	0.112	12.5	8	3.23	6.69 ± 0.18	6.35 ± 0.01	5.72 ± 0.49	85.83 ± 0.10	14.18 ± 0.10	12.15 ± 0.06	3.74 ± 0.07	2.73 ± 0.00	-11.75 ± 0.24	1799 ± 10.00
21	0.112	12.5	8	3.23	7.50 ± 0.07	5.35 ± 0.17	3.39 ± 0.10	85.95 ± 0.06	14.05 ± 0.06	12.75 ± 0.64	3.89 ± 0.26	2.76 ± 0.01	-11.18 ± 0.99	1611 ± 43.50

Source: The Authors.

Table 2.

ANOVA of blackberry suspensions inoculated with *Lb. casei* and inulin to T0.

Font	p-Value					
	Log CFU/g	M (%)	TS (%)	μ (cP)	°Brix	Acidity (%)
A: X _{TSBC}	0.1505	0.0000*	0.0000*	0.0000*	0.0036*	0.0000*
B: R (g/g)	0.1214	0.8797	0.8797	0.0173*	0.9112	0.2539
C: [McFarland]	0.0258*	0.3699	0.3699	0.0012*	0.6766	0.0000*
D: Inulin (%)	0.7850	0.0000*	0.0000*	0.7427	0.1096	0.0000*
AA	0.0677	0.2614	0.2614	0.7047	0.1277	0.0000*
AB	0.0624	0.0000*	0.0000*	0.0956	0.0396*	0.0000*
AC	0.7239	0.8029	0.8029	0.0052*	0.0233*	0.0000*
AD	0.4991	0.9961	0.9961	0.0615	0.9685	0.2034
BB	0.8060	0.1405	0.1405	0.8202	0.4213	0.0000*
BC	0.0007*	0.8967	0.8967	0.5109	0.1916	0.0000*
BD	0.7046	0.0068*	0.0068*	0.0000*	0.0022*	0.0000*
CC	0.0000*	0.0499*	0.0499*	0.0000*	0.9211	0.0000*
CD	0.8184	0.7808	0.7808	0.1839	0.7545	0.0000*
DD	0.7299	0.2859	0.2859	0.0000*	0.4546	0.0000*

Source: The Authors.




Figure 1. Response surface graph of BC suspensions inoculated with *Lb. casei* and inulin for a time of 0 hours (T_0).
Source: The Authors.

On the other hand, in cape gooseberry pulp and in glucose solution, *Lb. plantarum* and *Lb. casei* ATCC 393 were inoculated, reporting counts at 15 days and 4 °C of *Lb. plantarum* of 3.23×10^9 and 1.64×10^9 CFU/mL respectively; while *Lb. casei* ATCC 393 obtained counts 5.4×10^8 and 7.34×10^8 respectively [18]; and in mango, lulo and strawberry juices inoculating 10^{10} CFU/mL of *Lb. plantarum*, a viability of 10^8 CFU/mL was reported after 30 days of storage at 8 °C [39].

Average values of M fluctuated between 83.95% and 89.25%, and for TS_s between 10.75% and 16.05%, observing coherence between them. M was higher when the suspension formulation showed lower X_{TSBC}, higher R and lower inulin contents, and TS_s levels decreased under these compositions. On the other hand, these parameters are also related to the μ of suspension, which is a relevant property in spray drying processes and which average values were between 651 and 2713 cP. An important relationship of μ with X_{TSBC} was evident, getting higher when it increases, which is consistent due to higher content of TS_s.

The effect of [McFarland] became more relevant to the scale of 8, where μ decreases to higher X_{TSBC}; whereas, the influence of R is more noticeable when suspension has $\mu < 1000$ cP, which increases as R decreases (< presence of BC) and X_{TSBC} is smaller. These interactions influence the rheological response of suspension and at the same time in microbial population, due to the effect of TS_s on the medium's osmotic pressure, which causes cell plasmolysis, resulting in possible inhibition of microorganisms and therefore in its viability [35,40].

There isn't much literature regarding the rheological properties of blackberry suspensions, seen from the pulp itself to concentrates. Blackberry pulp is reported for viscosity values of 2061.8 cP and enzymatically treated 150 cP [41].

Average °Brix suspension values fluctuated between 10.95% and 13.25%, which are available for strain maintenance. A decrease was observed, mainly at lower X_{TSBC}, improving when it interacts at low values of R and [McFarland], which is consistent due to the greater mass of inoculum (>water content) and lower content of microorganism in the inoculum. On the other hand, there is an important effect of inulin-R interaction, which occurs at low values of R and inulin, attributed to an effect similar to that previously mentioned. Literature does not report values of soluble solids in blackberry suspensions inoculated with microorganisms, they are reported mainly in fruit with significant variability, due to maturity degree, climatology, soils, solar radiation, among others [32]; 7.74% with degree of maturity of 5 [42], 12.07 and 13.53% [43]. On the other hand, other authors report values in blackberry pulp of 6.77% [1] and in soft drinks 9.54% and 10.53% [44].

Average pH values fluctuated between 2.7 - 2.9, which indicates that there is no marked influence of independent variables considered on this variable. Blackberry and blackberry products are known to have an acidic pH, literature shows values between 2.65 and 2.93 [43,45] for fruit, and values of 3.31 [1] for pulp. Acidity results fluctuated in a higher proportion than that of pH: 1.9% and 4.8%, with a noticeable increase as R and [McFarland]

decrease, along with an increase of X_{TSBC} and inulin. However, major changes (> acidity) happen at high R values and their interactions with X_{TSBC} and inulin, which is attributed to a higher content of BC (malic acid: 5.85%) and inulin, with little inoculum mass (> water content), which favors acid concentration in suspension. For Andean blackberry, reported values of acidity fluctuate: 3.66% [42], 2.30% [43] and 2.83% [45] and 2.62-2.74%; while, for blackberry pulp a value of 0.76% is reported and expressed as citric acid. Literary sources do not mention growth for blackberry-based substrates, some work is highlighted in cashew juice where a greater viability of *Lb. casei* at pH of 6.4 was obtained [46]; on the other hand, *Lb. plantarum* survival of 68.7% at pH 2.0, inoculated in fruit juices was reported [39].

ζ values showed low fluctuation: -14.25 and -10.70 mV, which denotes a negative predominant electric charge in the outer coion layers, which contributes to low repulsive forces in the colloidal system ($|\zeta| > 25$ mV). However, there is an alleged synergy with suspension viscosity that makes it stable, since no separation phase or precipitates were seen with time [47]. ζ is favored mainly by the effect of R-X_{TSBC} interactions, with repulsive forces being greater (ζ more negative) when the colloidal system has a higher concentration of BC and with higher R (more BC and less water), which causes a higher amount of anions adsorbed in coion layers. Inulin and inoculum represent another important aspect affecting the colloidal system stability, which individually reached values of -5.18 and -3.46 mV respectively.

Model optimization and validation: optimization parameters of BC suspension inoculated with *Lb. casei* and inulin, were determined under the criteria of promoting microorganism viability and compliance with design criteria operability of pilot Vibrasec PSALAB1.5 spray dryer ($\mu \approx 1000$ cP and TS $\leq 45\%$). Weights and impacts considered and optimization results are listed in Table 3.

Multiple optimization results showed a desirability of 69.1%, defining suspension independent variables or formulation as follows: X_{TSBC}: 0.103, R: 15.6, McFarland Scale: 10% inulin: 3.1%. Most dependent variables showed an acceptable residual error between the predicted value of

Table 3.
Results of experimental optimization of BC suspension inoculated with *Lb. casei* and inulin.

Response	Criterion	Weight	Impact	Predicted value	Experimental value	Residual error
μ (cP)	1000	3.0	5.0	1000.0	1288.0 \pm 20.1	28.8
ζ (mV)	Minimize	1.0	3.0	-13.3	-13.3 \pm 0.26	0
pH	Maximize	1.0	3.0	2.82	2.74 \pm 0.01	2.8
Acidity (%)	Minimize	1.0	3.0	2.3	3.1 \pm 0.004	34.7
TS _s (%)	Maximize	0.1	1.0	10.7	13.3 \pm 0.0	24.3
M (%)	Maximize	0.1	1.0	89.2	86.7 \pm 0.0	2.8
°Brix	Maximize	0.1	1.0	11.9	11.0 \pm 0.1	7.6
Log CFU/g	Maximize	3	5.0	7.3	7.8 \pm 0.5	6.8

Source: The Authors.

mathematical model and the experimental value obtained from 3 replicas of this formulation. The ones displaying a greater residual errors are attributed because the response surface generates trends that are expected to coincide with experimental values.

4. Conclusions

Lb. casei ATCC 393 probiotic microorganism showed an adaptability in BC based and inulin suspensions, which guarantee its viability. The independent variables that are believed to affect microorganism viability the most are BC, the amount of inoculum and [McFarland]. Counts obtained from microorganism could identify colloidal suspension as a functional food with probiotic characteristics. For an effective use of suspension in a spray drying process, an inoculation time of 0 hours was considered adequate, since microorganism viability decreases with time and the effect of prebiotic was not evident.

References

- [1] Ferrari, C., Germer, S. and De Aguirre, J., Effects of spray-drying conditions on the physicochemical properties of blackberry powder, *Dry Technology*, 30(2), pp. 154-163, 2012. DOI: 10.1080/07373937.2011.628429
- [2] Oszmiański, J., Nowicka, P., Teleszko, M., Wojdyło, A., Cebulak, T. and Oklejewicz, K., Analysis of phenolic compounds and antioxidant activity in wild blackberry fruits. *Molecular Diversity Preservation International*, 16(7), pp. 14540-14553, 2015. DOI: 10.3390/ijms160714540
- [3] Szajdek, A. and Borowska, E., Bioactive compounds and health-promoting properties of Berry fruits: a review, *Plant Foods for Human Nutrition*, 63(4), pp. 147-53, 2008. DOI: 10.1007/s11130-008-0097-5
- [4] Galus, S. and Kadzińska, J., Food applications of emulsion-based edible films and coatings. *Trends in Food Science & Technology*, 45(2), pp.273-283, 2015. DOI: 10.1016/j.tifs.2015.07.011
- [5] Ramírez, J., Aristizábal, I. y Restrepo, J., Conservación de mora de castilla mediante la aplicación de un recubrimiento comestible de gel de mucilago de penca de sábila. *Vitae* [Online], 20(3), pp. 172-183, 2013. [Date of reference: January 30th, 2018]. Available at: <http://www.redalyc.org/articulo.oa?id=169829162003>. ISSN 0121-4004
- [6] Kaume, L., Howard, L.R. and Devareddy, L., The blackberry fruit: a review on its composition and chemistry, metabolism and bioavailability, and health benefits, *Journal of Agricultural and Food Chemistry*, 60(23), pp. 5716-5727, 2012. DOI: 10.1021/jf203318p
- [7] Ministerio de Agricultura, Colombia. Cadena productiva nacional de la mora: cifras sectoriales, 2015, 51 P.
- [8] Páez, R., Lavari, L., Audero, G., Cuatrin, A., Zaritzky, N., Reinheimer, J. and Vinderola, G., Study of the effects of spray-drying on the functionality of probiotic lactobacilli, *International Journal of Dairy Technology*, 66(2), pp. 155-161, 2013. DOI: 10.1111/1471-0307.12038
- [9] Liao, L.K., Wei, X.Y., Gong, X., Li, J.H., Huang, T. and Xiong, T., Microencapsulation of *Lactobacillus casei* LK-1 by spray drying related to its stability and in vitro digestion, *LWT - Food Science and Technology*, 82, pp. 82-89, 2017. DOI: 10.1016/j.lwt.2017.03.065
- [10] Kerry, R., Patra, J.K., Gouda, S., Park, Y., Shin, H.-S. and Das, G., Benefaction of probiotics for human health: A review, *Journal of Food and Drug Analysis*, 30, pp. 1-13, 2018. DOI: 10.1016/j.jfda.2018.01.002
- [11] Dimitrellou, D., Kandyli, P. and Kourkoutas, Y., Effect of cooling rate, freeze-drying, and storage on survival of free and immobilized *Lactobacillus casei* ATCC 393, *LWT - Food Science and Technology*, 69, pp. 468-473, 2016. DOI: 10.1016/j.lwt.2016.01.063
- [12] Rodríguez-Barona, S., Giraldo, G.I. and Zuluaga, Y., Evaluación de la incorporación de fibra prebiótica sobre la viabilidad de *Lactobacillus casei* impregnado en matrices de mora (*Rubus glaucus*), *Información Tecnológica*, 26(5), pp. 25-34, 2015. DOI: 10.4067/S0718-07642015000500005
- [13] Ministerio de la protección Social, Colombia. Rotulado nutricional de alimentos - Resolución 333 de 2011.
- [14] Ministerio de la protección Social, Colombia. Suplementos - Resolución 3096 de 2007.
- [15] Instituto Colombiano de normas técnicas y certificación (ICONTEC), NTC 805. Productos lácteos, leches fermentadas, 2005, 9 P.
- [16] Vijaya, B., Vijayendra, S. and Reddy, O., Trends in dairy and non-dairy probiotic products - a review, *Journal of Food Science and Technology*, 52(10), pp. 6112-24 2015. DOI: 10.1007/s13197-015-1795-2
- [17] Vergara, A., Estudio de la viabilidad de *Lactobacillus casei* en jugo de pera, Tesis, Universidad Austral de Chile, Valdivia, Chile, 2015.
- [18] Marín, Z., Cortés, M. y Montoya, C. Evaluación de la viabilidad de crecimiento de la cepa nativa *Lactobacillus plantarum* LPBM10 y la cepa comercial *Lactobacillus casei* ATCC 393 en pulpa de uchuva y en solución isotónica de glucosa, *Vitae*, [Online], 16(2), pp. 210-17, 2009. [Date of reference: January 30th of 2018]. Available at: http://www.scielo.org.co/scielo.php?script=sci_arttext&pid=S0121-40042009000200005&lng=en&nrm=iso. ISSN 0121-4004
- [19] De la Cruz, A. y Terán, A., Evaluación de la viabilidad de *Lactobacillus casei* libre encapsulado en alginato sódico como probiótico en jugo de guayaba, Tesis de grado, Escuela Agrícola Panamericana, San Antonio de Oriente, Honduras, 2013.
- [20] Reid, G., Probiotics: definition, scope and mechanisms of action, *Best Pract Res Clin Gastroenterol*, 30(1), pp.17-25, 2016. DOI: 10.1016/j.bpg.2015.12.001
- [21] AOAC. Official methods of analysis. Association of Official Analytical Chemists, Arlington, 19th edition, Arlington, Virginia (USA), 2012, 684 P.
- [22] Wardy, W., Pujols-Martínez, D., Xu, Z., Kyoona, H. and Prinyawiwatkul, W., Viscosity changes of chitosan solution affect physico-functional properties and consumer perception of coated eggs during storage, *LWT - Food Science and Technology*, 55, pp. 67-73, 2014. DOI: 10.1016/j.lwt.2013.07.013
- [23] Niu, F., Zhang, Y., Chang, C., Pan, W., Sun, W., Su, Y. and Yang, Y., Influence of the preparation method on the structure formed by ovalbumin/gum arabic to observe the stability of oil-in-water emulsion, *Food Hydrocolloids*, 63, pp. 602-610, 2017. DOI: 10.1016/j.foodhyd.2016.10.007
- [24] Koneman, E., Koneman's color atlas and textbook of diagnostic microbiology. 6^a Ed., Lippincott Williams & Wilkins, 2006.
- [25] Badui, S., Química de los alimentos, Editorial Pearson, 5^a Ed., Ciudad de México, 2013, 723 P.
- [26] Shori, A., Influence of food matrix on the viability of probiotic bacteria: a review based on dairy and non-dairy beverages, *Food Bioscience*, 13, pp. 1-8, 2016. DOI: 10.1016/j.fbio.2015.11.001
- [27] Drapala, P., Mulvihill, M. and Mahony, A., A review of the analytical approaches used for studying the structure, interactions and stability of emulsions in nutritional beverage systems, *Food Structure*, 16, pp. 27-42, 2018. DOI: 10.1016/j.foostr.2018.01.004
- [28] Wang, W. and Xu, S., Degradation kinetics of anthocyanins in blackberry juice and concentrate. *Journal of Food Engineering*, 82, pp. 271-275, 2007. DOI: 10.1016/j.jfoodeng.2007.01.018
- [29] Haminiuk, C., Sierakowski, D. and Masson, M., Rheological characterization of blackberry pulp. *Brazilian Journal of Food Technology*, [Online], 9(4), pp. 291-296, 2006. [Date of reference: February 3rd of 2018]. Available at: <http://www.ital.sp.gov.br/bj/artigos/bjft/2006/p06262.pdf>
- [30] Franceschini, L., Salvatori, D., Sosa, N. and Schebor, C., Physical and functional properties of blackberry freeze- and spray-dried powders. *Drying Technology*, 32(2), pp. 197-207, 2014. DOI: 10.1080/07373937.2013.814664
- [31] Rodríguez, B., Zuluaga-Pava, Y. y Cruz-Rios, D., Producto potencialmente simbiótico a partir de mora de castilla (*Rubus glaucus*) aplicando impregnación a vacío, *Scientia Agropecuaria*, [Online], 3, pp. 273-278, 2012. [Date of reference: February 3rd of 2018]. Available at: <http://www.redalyc.org/articulo.oa?id=357633704001> ISSN 2077-

- 9917
- [32] Moreno, B. y Deaquiz, Y., Caracterización de parámetros fisicoquímicos en frutos de mora (*Rubus alpinus* Macfad). Acta Agronómica, 25, pp. 130-136, 2016. DOI: 10.15446/acag.v65n2.45587
 - [33] Kaume, L., Howard, L.R. and Devareddy, L., The blackberry fruit: a review on its composition and chemistry, metabolism and bioavailability, and health benefits. J Agric Food Chem, 60(23) pp. 5716-5727, 2012. DOI: 10.1021/jf203318p
 - [34] Ray, B. y Bhunia, A., Fundamentos de microbiología de los alimentos. 4ª Ed., Mc Graw Hill, 2010, 352 P.
 - [35] Huang, S., Vignolles, M., Chen, D., Le Loir, Y., Jan, G., Schuck, P., Spray drying of probiotics and other food-grade bacteria: a review. Trends Food Sci Technol, 63, pp. 1-17, 2017. DOI: 10.1016/j.tifs.2017.02.007
 - [36] Páez, R., Lavari, L., Audero, G., Cuatrin, A., Zaritzky, N., Reinheimer, J. and Vinderola, G., Study of the effects of spray-drying on the functionality of probiotic lactobacilli. International Journal of Dairy Technology, 66(2), pp. 155-161, 2013. DOI: 10.1111/1471-0307.12038
 - [37] FAO/WHO., Expert consultation. Human vitamin and mineral requirement. Report of a joint, Bangkok, 2001. 303 P.
 - [38] Krasaekoopt, W. and Watcharapoka, S., Effect of addition of inulin and galactooligosaccharide on the survival of microencapsulated probiotics in alginate beads coated with chitosan in simulated digestive system, yogurt and fruit juice, LWT - Food Science and Technology, 57, pp. 761-766, 2014. DOI: 10.1016/j.lwt.2014.01.037
 - [39] Serna, J., Elaboración de jugos de fruta con adición de bacterias ácido lácticas con potencial probiótico. Tesis, Universidad de la Sabana, Bogotá, Colombia, 2012.
 - [40] Piuri, M., Sanchez-Rivas, C., Ruzal, S.M., Cell wall modifications during osmotic stress in *Lactobacillus casei*, J Appl Microbiol, 98(1), pp. 84-95, 2005. DOI: 10.1111/j.1365-2672.2004.02428.x
 - [41] Agudelo, A., Lucas, J. y Quintero, V., Evaluación de los cambios de viscosidad en pulpa de mora (*Rubus glaucus*) tratada enzimáticamente, Agronomía Colombiana, 34(1Supl), pp. S626-S629, 2016. DOI: 10.15446/agron.colomb.v34n1supl.58408
 - [42] Sora, A., Gerhard, F. y Flórez, R., Almacenamiento refrigerado de frutos de mora de Castilla (*Rubus glaucus* Benth.) en empaques con atmósfera modificada, Agronomía Colombiana, [Online], 24(2), pp. 306-316, 2006. [Date of reference: March 2nd of 2018]. Available at: <https://revistas.unal.edu.co/index.php/agrocol/article/view/20044>
 - [43] Farinango, M., Estudio de la fisiología postcosecha de la mora de Castilla (*Rubus glaucus* Benth) y de la mora variedad brazos (*Rubus* spp.), Tesis, Escuela Politécnica Nacional, Quito, Ecuador, 2010. 142 P.
 - [44] Galvis, B., Estudio de la durabilidad de la pulpa de mora de Castilla y mora San Antonio (*Rubus glaucus*), Tesis, Universidad Nacional de Colombia, 2003, 79 P.
 - [45] Ayala, C., Valenzuela, P. y Bohórquez, Y., Caracterización fisicoquímica de mora de castilla (*Rubus glaucus* Benth) en seis estados de madurez. Biotecnol en el Sect Agropecu y Agroindustrial, [Online], 11(2) pp. 10-8, 2013. [Date of reference: March 2nd of 2018]. Available at: <http://revistas.unicauca.edu.co/index.php/biotecnologia/article/view/306/503>
 - [46] Pereira, A., Maciel, T. y Rodrigues, S., Probiotic beverage from cashew apple juice fermented with *Lactobacillus casei*. Food Res Int, 44(5), pp.1276-83, 2011. DOI: 10.1016/j.foodres.2010.11.035
 - [47] Genovese, B. and Lozano, J., The effect of hydrocolloids on the stability and viscosity of cloudy apple juices, Food Colloids, 15, pp. 1-7, 2001. DOI: 10.1016/S0268-005X(00)00053-9

of 14 articles published in scientific journals. Her research interests include: food engineering, biotechnology, functional foods and probiotics. ORCID ID: 0000-0003-1734-6414

M. Cortés-Rodríguez, received the BSc. in Eng in Chemical Engineering in 1989 and the PhD. in Food Engineering in 2004. His research interests include: food engineering, science and technology, functional foods. Currently, he is a titular professor in the Agricultural and Food Engineering Department, Faculty of Agricultural Sciences, Universidad Nacional de Colombia. Author of 3 patents, 93 articles published in scientific journals and 3 book chapters. Participation in 39 research projects; director of 5 extension projects; director of 4 Dr. Thesis and 33 MSc. Thesis; director of 7 Dr Thesis and 5 MSc. Thesis in development. Speaker in 95 national and international events, national industry consultant, evaluator of research projects, scientific articles of national and international scientific journals; member of editorial and scientific committees of national and international scientific journals. ORCID: 0000-0003-3407-1635

O.I. Montoya-Campuzano, received the BSc. in Biologist in 1983, from the Universidad de Antioquia, Food Engineer in 1991, from the Corporación Universitaria Lasallista. MSc. in Microbiology in 1997, from the Universidad Nacional de Colombia Bogotá, professor at the Universidad Nacional de Colombia, Medellín in the Faculty of Sciences and the School of Biosciences, since 1991. I have taught the subjects of general microbiology, food microbiology, the undergraduate course in Biological Engineering, and Microbiology and Food Hygiene in the postgraduate course in Food Science and Technology. Especial emphasis on probiotics. Director of the research group of at "Colciencias": Probiotics Functional Prospection and Metabolites, assigned since 2017. Director of degree works, special academic practices, biological engineering students, master's thesis at students of biotechnology, agricultural sciences, food science and technology. I have participated in the tutorial committees of PhD students of biotechnology, agricultural sciences and food science and technology. ORCID ID: 0000-0002-4820-0381

J.C. Arango-Tobón, received the BSc. in Agricultural Engineer, MSc. in Exploitation of Hydraulic Resources. Is titular professor in the Universidad Nacional de Colombia, Department of Agricultural Engineering and Food, professional experience for 11 years at Colombian Institute of Hydrology Meteorology and Land Adaptation, and 27 years in teaching, research and extension, in areas like: study auditor, design and construction of works for land adaptation; irrigation principles; irrigation engineering; administration, operation and conservation of irrigation districts; erosion control engineering, management and operation of irrigation districts. Researcher of the Agricultural Engineering Research Group, ranked in Colciencias as A1 group; director of 45 degree projects; author of 73 publications of academic material, with participation in 51 national and international events; Production of 7 didactic videos on irrigation, mango and soursop crops; national consultant; evaluator of teaching calls from Colombian universities. ORCID ID: 0000-0003-1625-0541

Z.T. Marín-Arango, is BSc. in Food Engineering in 2003, MSc. in Food Science and Technology in 2009, currently PhD candidate in Food Science and Technology; student in Agricultural Sciences, all of them in the Universidad Nacional de Colombia. She is also in the study commission of the University of Quindío, where she is working since 2014 to full time in the Faculty of Agroindustrial Sciences and in the assistant category. Author