



REVISTA DE INGENIERIA DE LA FACULTAD DE INGENIERIA, UNIVERSIDAD NACIONAL DE COLOMBIA - BOGOTÁ

DYNA

ISSN: 0012-7353

ISSN: 2346-2183

Universidad Nacional de Colombia

Dias Batista, Pedro Henrique; Pontes de Almeida, Gledson Luiz; Bezerra da Silva, Jhon Lennon; Pandorfi, Héilton; Vinícius da Silva, Marcos; Batista da Silva, Rodes Angelo; Neves de Melo, Maria Vitória; Abraão Costa Lins, Frederico; Florentino, José Jairo

Short-term grazing and its impacts on soil and pasture degradation

DYNA, vol. 87, no. 213, 2020, April-June, pp. 123-128

Universidad Nacional de Colombia

DOI: <https://doi.org/10.15446/dyna.v87n213.81853>

Available in: <https://www.redalyc.org/articulo.oa?id=49664596017>

- How to cite
- Complete issue
- More information about this article
- Journal's webpage in redalyc.org

UNEN
redalyc.org

Scientific Information System Redalyc

Network of Scientific Journals from Latin America and the Caribbean, Spain and Portugal

Project academic non-profit, developed under the open access initiative

Short-term grazing and its impacts on soil and pasture degradation

Pedro Henrique Dias Batista ^a, Gledson Luiz Pontes de Almeida ^a, Jhon Lennon Bezerra da Silva ^a, Héilton Pandorfi ^a, Marcos Vinícius da Silva ^a, Rodes Angelo Batista da Silva ^a, Maria Vitória Neves de Melo ^a, Frederico Abraão Costa Lins ^a & José Jairo Florentino Cordeiro Junior ^b

^a Department of Agricultural Engineering, Federal Rural University of Pernambuco, Recife-PE, Brazil. giga_pedro@hotmail.com; gledson81@hotmail.com; jhonlennoigt@hotmail.com; hpandorf@hotmail.com; marcolino_114@hotmail.com; rodesangel@gmail.com; vitorianevevss@outlook.com; frederico_acl@hotmail.com

^b Federal University of Sergipe, Nossa Senhora da Glória-SE, Brazil. jairofcordeiro@hotmail.com

Received: August 23th, 2019. Received in revised form: March 17th, 2020. Accepted: March 30th, 2020.

Abstract

The objective was to verify possible modifications of the soil structure and the pattern of the spectral response of pasture vegetation cover to animal trampling. The study was carried out on a farm in the Agreste region of Pernambuco, Brazil in an area with continuous grazing by heifers. Soil samples were collected at 36 regular points, before and after the grazing period, where the physical properties of the soil were determined at a 0.00–0.10 m depth. Before and after grazing, images of the Sentinel-2A satellite were also obtained to observe the pasture vegetation response pattern over time through Vegetation Indexes. The soil attribute data were submitted to multivariate factorial analysis. The vegetation index maps were evaluated for spatial variability. The results showed that after the grazing, there was a significant change in soil attributes and pasture, which can indicate possible degradation processes.

Keywords: animal trampling; dairy cattle; soil physics; vegetation indices; remote sensing.

Pastoreo a corto plazo y sus impactos en la degradación del suelo y los pastos

Resumen

El objetivo fue verificar las posibles modificaciones de la estructura del suelo y el patrón de la respuesta espectral de la cubierta vegetal de los pastizales en función del pisoteo de los animales. El estudio se llevó a cabo en una granja en la región Agreste de Pernambuco, Brasil. En un área manejada bajo pastoreo continuo de vaquillas. Se recolectaron muestras de suelo en 36 puntos regulares, antes y después del período de pastoreo, donde se determinaron las propiedades físicas del suelo en la profundidad de 0.00-0.10 m. Antes y después del pastoreo, también se obtuvieron imágenes del satélite Sentinel-2A para observar el patrón de respuesta de la vegetación del pasto a lo largo del tiempo a través de los Índices de Vegetación. Los datos de los atributos del suelo se sometieron a análisis factorial multivariado. Los mapas del índice de vegetación se evaluaron para determinar la variabilidad espacial. Los resultados mostraron que después del pastoreo, hubo un cambio significativo en los atributos del suelo y el pasto, que puede indicar posibles procesos de degradación.

Palabras clave: pisoteo animal; ganado lechero; física del suelo; índices de vegetación; teledetección.

1. Introduction

In Brazil, about 159 million hectares of agricultural land is occupied by pastures, and the country has established in most of these areas the largest herd of cattle in the world with 221.81 million head [1].

However, high stocking rates, soil compaction and lack

of management for nutrient replacement negatively affect the quality of pastures and soil, becoming precursors to degradation [26]. Thus, the use of new technologies for the correct management of these areas is important to prevent the degradation of new areas, since the use of pastures represents a practical alternative to feed ruminants and, concomitantly, to produce meat and milk [12].

How to cite: Batista, P.H.D., Almeida, G.L.P., Silva, J.L.B., Pandorfi, H., Silva, M.V., Silva, R.A.B., Melo, M.V.N., Lins, F.A.C. and Cordeiro Junior, J.J.F., Short-term grazing and its impacts on soil and pasture degradation. DYNA, 87(213), pp. 123-128, April - June, 2020.

© The author; licensee Universidad Nacional de Colombia.
Revista DYNA, 87(213), pp. 123-128, April - June, 2020, ISSN 0012-7353
DOI: <http://doi.org/10.15446/dyna.v87n213.81853>



Therefore, degradation in pasture areas can also be a consequence of overgrazing, so animal trampling, with the high intensity of vegetation removal by grazing, decreases the infiltration rate, increases soil density and reduces root growth of plants. Thus, the impact of trampling and high stocking rates can cause soil compaction and biological degradation of pastures. However, these possible phenomena are mainly dependent on the soil class, the soil's edaphoclimatic conditions, its moisture content, the animal stocking rate, the available forage mass and the forage species used in the system [15,26].

Orbital satellite images following the study of soil physical properties have been used to determine degraded pasture areas [23,25]. Therefore, through the vegetation indices that can be obtained in digital images by specific software, it is possible to monitor and document variables that cause degradation in areas with grazing as well as areas that are arid or without vegetation, etc., in large or small fields and at different time scales [19,21].

Digital image processing with the application of the Surface Energy Balance Algorithm for Land (SEBAL), for example, allows obtaining thematic maps of the energy balance components, with analyses in space and time practically and efficiently with low operational cost, determining biophysical parameters such as the vegetation indices Soil Adjusted Vegetation Index (SAVI) and Leaf Area Index (LAI). Changes in the cover and soil use can be observed directly by the reduction of the SAVI and LAI indices [5,18,21].

Given this context and given the need of the environmental monitoring of the vegetation cover of pasture in semiarid soil, the objective was to verify possible modifications of the soil structure and the pattern of the spectral response of pasture vegetation cover in the function of animal trampling. This was done with the assistance of multivariate factorial analysis of soil physical-hydric attributes and the use of the geoprocessing in the application of remote sensing techniques as well as satellite images to determine biophysical parameters, such as the vegetation indices, before and after grazing in a Brazilian semiarid region.

2. Material and methods

The study area is located in the municipality of Capoeiras (Fig. 1), which is in the Agreste region. This region is in the dairy basin. The sites are in the State of Pernambuco, which has Brazilian semiarid climate, and is located between the parallels of 08°37'10.27" S (UTM: 9046392 m) and 08°37'11.82" S (UTM: 9046344 m), and between the meridians of 36°36'46.50" W (UTM: 762715 m) and 36°36'47.98" W (UTM: 762670 m), with an altitude of 800 m, being west of the meridian of Greenwich. According to the climatic classification of Köppen, the climate of the region is characterized as semiarid (Bsh), and the average annual precipitation of the region is 588 mm. The soil of the study area was classified as Planosol according to the Agroecological Zoning of Pernambuco [31]. Soil

characterization was performed according to the methods described by [6] in Table 1.

The research was conducted in the spring season, from 16/10 to 05/11/2016, totalling 21 consecutive days. The grazing field had a size of 40 × 40 m (Fig. 1) and an enclosure for the animals, measuring 8 × 10 m, containing a drinking fountain and a shaded area of approximately 16 m². The soil of the study area was prepared with a harrow before planting of the grass; however, no fertilization or soil corrections were used. The area also had negligible slope. Management was adopted in a continuous grazing system, in soil with grazing of *Brachiaria decumbens*, under rainfed conditions. Three Girolando heifers with an average weight of 300 kg were used, which started the grazing period with a mean forage height of 0.90 m and it was suspended when a mean height of 0.23 m was reached, as indicated by [11].

In the grazing area, 36 points were defined in a 6 × 6 m regular mesh (Fig. 1), where resistance to penetration, moisture, bulk density, total porosity and soil particle density were determined at depths of 0.00–0.10 m in addition to the hydraulic conductivity in saturated soil and the organic matter in the soil surface. An Uhland-type sampler was used for the undeformed collection of soil.

The undeformed soil samples were weighed to determine the wet soil mass and then oven-dried at 105 °C for 24 h to obtain the dry soil mass. Soil density (SD) was calculated from the weight of the oven-dried soil and the total volume of the soil cores. Gravimetric water content (w) at sampling was then calculated as the difference between the wet weight of each soil core at sampling and then after oven drying. The particle density (PD) was determined by the volumetric flask method. The total soil porosity (TP) was quantified using the ratio of SD to PD. The soil organic matter content (OM) was measured via oxidation with potassium dichromate in sulfuric acid, and the fractions of sand, silt and clay were determined by the hydrometer method. All procedures were performed according to [6].




Figure 1. Spatial location map of the municipality of Capoeiras-PE (A), study area (B), grazing field before (C), during (D) and after (E) the grazing period. The image was georeferenced by the Universal Transverse Mercator (UTM), with a spatial resolution of 10 m.

Source: The Authors.

Table 1.

Physical characterization of the study area, a commercial property of dairy cattle in the municipality of Capoeiras-PE.

Characteristics	Pasture field
Layer	0.00–0.10 m
Declivity (%)	0
Textural class	sandy-loam
Sand (g.kg ⁻¹)	817.82
Silt (g.kg ⁻¹)	178.78
Clay (g.kg ⁻¹)	4.0
Particle density (g.cm ⁻³)	2.64
Organic matter (g.kg ⁻¹)	39.57

Source: The Authors.

An impact penetrometer, model IAA/Planalsucar/Stolf, was used to determine soil penetration resistance (PR), with the number of impacts dm⁻¹ transformed into dynamic resistance (MPa) utilizing eq. (1) as proposed by [24].

$$PR = \frac{M + m + \left(\frac{M}{M + m} \times \frac{M \times h}{X} \right)}{A} \quad (1)$$

where: PR—Soil penetration resistance, kgf.cm⁻² (kgf.cm⁻² × 0.098 = MPa); M—Mass of piston, 1.6 kg; m—Mass of machine without piston, 1.5 kg; h—Height of piston fall, 54 cm; X—Penetration of machine stick, cm by impact; A—Area of cone, 1.35 cm².

To characterize soil water infiltration, the Beerkan method was used, based mainly on simplified infiltration tests and soil particle size distribution analysis [14]. For the determination of the hydraulic conductivity of the saturated soil, the methodology developed by [4] according to eq. (2).

$$K_0 = \frac{b}{0.467 \left(\frac{2.92}{r \times \alpha} + 1 \right)} \quad (2)$$

where: K₀ soil water infiltration, the Beerkan method Hydraulic conductivity in saturated soil, mm.s⁻¹; b—Slope of an equation linearized in the function of cumulative infiltration over time; r—Ring radius, 75 mm; α—adopted 0.012, for practices in permeameters and infiltrometers in soils that range from thick sand to compacted clay.

Remote sensing techniques were introduced to aid and consolidate terrestrial information using two images of the Sentinel-2 satellite (processing level 2A) Multispectral Instrument (MSI) sensor and with a spatial resolution of 10 m. The mapping was carried out of the pasture field of the study area. The images were made available by the European Space Agency (ESA) via the EOS/Land Viewer (Earth Observing System) platform, dated 08/10/2016 (before grazing) and 07/11/2016 (after grazing). The digital and semi-automated processing of the images was performed using the ERDAS IMAGINE® 9.1 Software, in which some steps of the SEBAL (Surface Energy Balance Algorithm for Land) algorithm, responsible for the mathematical operations of image processing, was implemented (2002). The algorithm is based on the satellite's multispectral bands, and is used to obtain thematic maps of the terrestrial surface [5,2]. From the level 2A product of reflectance (corrected atmospheric

product, including cloud tracking and adjacency/slope effects correction), the multispectral bands of the Sentinel-2A images are a baseline for the geophysical parameters [7,9,10]. The images were georeferenced by the Universal Transverse Mercator (UTM) Map Projection System, DATUM: WGS84—Zone 24 L.

From the multispectral reflective bands of satellite images, the vegetation indexes were determined: SAVI (Soil Adjusted Vegetation Index) and LAI (Leaf Area Index). The SAVI was determined according to eq. (3) [3,10,21].

$$SAVI = \frac{(1 + L) \times (B_{08} - B_{04})}{(L + B_{08} + B_{04})} \quad (3)$$

where: SAVI—Soil Adjusted Vegetation Index; B₀₈ e B₀₄—near-infrared multispectral bands (NIR) and red of the Sentinel-2A satellite respectively [9]. L is the soil adjustment factor, which varies between 0 and 1, according to the vegetation cover density of the study area. An adjustment factor of 0.5 was used, which indicates an area with intermediate vegetation cover [16,2]. The chosen factor refers to the study area, which is located in the Brazilian semiarid and has Caatinga vegetation [22,21].

With the determination of the Soil Adjusted Vegetation Index (SAVI), the Leaf Area Index (LAI, m² m⁻²) was determined as an indicator of the biomass amount according to eq. (4) [2, 19].

$$LAI = - \frac{\ln \left(\frac{0.69 - SAVI}{0.59} \right)}{0.91} \quad (4)$$

The results of the measured soil attributes were subjected to a multivariate factorial analysis, an exploration that makes it possible to ascertain the greatest amount of original information contained in the variables, and projects the attributes that better explain the impact of animal trampling under the soil during the study period.

The results of the SAVI and LAI vegetation indexes were analysed by descriptive statistics, central tendency (mean) and dispersion (minimum, maximum, standard deviation—SD and coefficient of variation—CV). The SAVI and LAI thematic maps were evaluated for spatial variability, by CV values (%), according to [28] criteria, which were being of low variability—CV<12%; mean variability—12%<CV<60% and high variability—CV>60%.

3.. Results and discussion

The soil physical attributes promoted patterns of important responses to the pasture field, based on multivariate statistical analyses, which allowed observing results over the level of soil compaction and the vulnerability to degradation due to grazing area management over time.




The spatial monitoring of the pasture vegetation management through the SAVI and LAI vegetation indices was possible by the development of thematic maps of the terrestrial surface, which showed different patterns of

The multivariate factorial analysis explained 59.21% of the total information in Factor 1 and Factor 2 (Fig. 2A) for the time before grazing, and, 57.61% of the two main factors explained the system for the period after grazing. [29] who evaluated the impact of pasture degradation on soil carbon and nitrogen fractions, through multivariate analysis and observed values of total variance greater than 50%, similarly to the present study.

Similar results were observed by [17], who reported that after animal trampling, the soil's physical-hydraulic properties underwent some type of modification, mainly with increased density and reduced infiltration rate, the authors also pointed out that clear signs of vegetation degradation were observed, manifested by a loss in species diversity and vegetation cover.

Fig. 3 illustrates the behaviour pattern of the SAVI index through thematic maps of the terrestrial surface in the study area.

The monitoring of the pasture area by satellite images via remote sensing confirmed the decrease in vegetation, that is, the reduction of the leaf canopy and the amount of biomass



It is interesting to observe the conditions of changes in grazing, which were well supported quantitatively by the variation of the minimum and maximum values and in the averages of the spatial analysis. Thematic maps of the SAVI and LAI indices showed a pattern of the spatial distribution of low variability, with CV ranging between 10.48 and 11.87% (Table 2).

Table 2.

Statistical parameters and quantitative variability of the SAVI and LAI vegetation indices in the study area, before and after grazing.

Satellite	Date	Vegetation Index—SAVI				
		Min	Max	¹ A	² SD	³ CV
Sentinel-2A	10/08/2016	0.27	0.46	0.38	0.04	10.53
	11/07/2016	0.13	0.22	0.18	0.02	11.11
		Leaf Area Index—LAI (m ² .m ⁻²)				
	10/08/2016	1.64	2.73	2.19	0.26	11.87
	11/07/2016	0.83	1.29	1.05	0.11	10.48

¹Average, ²Standard deviation, ³Coefficient of variation (%).

Source: The Authors.

environmental degradation of the soil in the semiarid region over time (Table 2).

The SAVI was a good indicator of the conditions of the pasture in the study. [31,26], highlight that the use of NDVI is a good indicator of pasture conditions, showing values below 0.6 and 0.5, respectively, as the beginning of pasture degradation. As SAVI is an improved NDVI index, it is a better indicator for identifying degraded areas, with values below 0.4 describing a possible degradation of pasture in the present study.

Because of this aspect, the application of remote sensing techniques to identify pasture degradation areas proved a promising tool, combined with vegetation indexes, which in turn, expressed the conditions of the study area [8,20].

4. Conclusion

The intensive management of the grazing, with the bovine trampling pressure altered the soil structure, thus increasing soil density and soil penetration resistance, favouring soil compaction and, consequently, hindering regrowth and renewal of the pasture and reducing the productivity.

The SAVI and LAI vegetation indices presented thematic maps of low variability and were sensitive to changes in the vegetation cover of pasture before and after grazing.

Remote orbital sensing through vegetation indexes was efficient in determining the response pattern of pasture vegetation in the semiarid region.

Acknowledgments

We would like to thank the Fundação de Amparo à Ciência e Tecnologia do Estado de Pernambuco (FACEPE APQ-0465-5.03/14) for financial support for the development of this research and Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES, Brazil) —Finance Code 001 by the granting of masters and doctoral scholarships. The European Space Agency (ESA) via the EOS/Land Viewer (Earth Observing System) platform by the free availability of the satellite images.

References

- [1] ABIEC - Associação Brasileira das Indústrias Exportadoras de Carnes. Perfil da pecuária no Brasil: Relatório anual [online]. 2018. [accessed: November 26th of 2018]. Available at: <http://abiec.siteoficial.ws/>

- [2] Allen, R.G., Tasumi, M. and Trezza, R., Satellite-Based energy balance for mapping evapotranspiration with internalized calibration (METRIC) – Model. Journal of Irrigation and Drainage Engineering, 133(4), pp. 380-394, 2007. DOI: 10.1061/(ASCE)0733-9437(2007)133:4(380)
- [3] Allen, R.G., Tasumi, M., Trezza, R., Waters, R. and Bastiaanssen, W., SEBAL (surface Energy Balance Algorithms for Land). Advance training and users manual—Idaho implementation, version 1, 2002, 97 P.
- [4] Bagarello, V., Di Prima, S., Iovino, M. and Provenzano, G., Estimating field-saturated soil hydraulic conductivity by a simplified Beerkan infiltration experiment. Hydrological Processes, 28(3), pp. 1095-1103. DOI: 10.1002/hyp.9649
- [5] Bastiaanssen, W.G., SEBAL-based sensible and latent heat fluxes in the irrigated Gediz Basin, Turkey. Journal of hydrology, 229(1-2), pp. 87-100, 2000. DOI: 10.1016/S0022-1694(99)00202-4
- [6] Donagema, G.K., Campos, D.B., Calderano, S.B., Teixeira, W.G. and Viana, J.M., Manual de métodos de análise de solo. Embrapa Solos-Documentos (INFOTEC-A-E), 2011.
- [7] Drusch, M., Del Bello, U., Carlier, S., Colin, O., Fernandez, V., Gascon, F., Hoersch, B., Isola, C., Laberinti, P., Martimort, P., Meygret, A., Spoto, F., Sy, O., Marchese, F. and Bargellini, P., Sentinel-2: ESA's optical high-resolution mission for GMES operational services. Remote Sensing of Environment, 120, pp. 25-36, 2012. DOI: 10.1016/j.rse.2011.11.026
- [8] Eddy, I.M., Gergel, S.E., Coops, N.C., Henebry, G.M., Levine, J., Zeriffi, H. and Shibkov, E., Integrating remote sensing and local ecological knowledge to monitor rangeland dynamics. Ecological Indicators, 82, pp. 106-116, 2017. DOI: 10.1016/j.ecolind.2017.06.033
- [9] ESA - European Space Agency. Sentinel-2 User Handbook, [online]. 2015. [accessed: June 02nd of 2019]. Available at: https://sentinels.copernicus.eu/documents/247904/685211/Sentinel-2_User_Handbook
- [10] Fernández-Manso, A., Fernández-Manso, O. and Quintano, C., SENTINEL-2A red-edge spectral indices suitability for discriminating burn severity. International Journal of Applied Earth Observation and Geoinformation, 50, pp. 170-175, 2016. DOI: 10.1016/j.jag.2016.03.005
- [11] Fidalski, J. and Alves, S.J., Grazing height of brachiaria and stocking rate limited by soil physical properties three years into an integrated soybean crop-livestock system. Revista Brasileira de Ciência do Solo, 39(3), pp. 864-870, 2015. DOI: 10.1590/01000683rbcs20140667
- [12] Figueiredo, E.B., Jayasundara, S., de Oliveira Bordonal, R., Berchielli, T.T., Reis, R.A., Wagner-Riddle, C. and La Scala Júnior, N., Greenhouse gas balance and carbon footprint of beef cattle in three contrasting pasture-management systems in Brazil. Journal of Cleaner Production, 142, pp. 420-431, 2017. DOI: 10.1016/j.jclepro.2016.03.132
- [13] Ganiyu, S.A., Evaluation of soil hydraulic properties under different non-agricultural land use patterns in a basement complex area using multivariate statistical analysis. Environmental monitoring and assessment, 190(10), pp. 595-610, 2018. DOI: 10.1007/s10661-018-6959-x
- [14] Haverkamp, R., Arrúe, J.L., Vandervaere, J.P., Braud, I., Boulet, G., Laurent, J.P., Taha, A., Ross, P.J. and Angulo-Jaramillo, R., Hydrological and thermal behaviour of the vadose zone in the area of Barrax and Tomelloso (Spain): experimental study, analysis and modeling. Project UE n. EV5C-CT, 92, pp. 00-90, 1996.
- [15] Hoffmann, C., Giese, M., Dickhoefer, U., Wan, H., Bai, Y., Steffens, M., Liu, C., Butterbach-Bahl, K. and Han, X., Effects of grazing and climate variability on grassland ecosystem functions in Inner Mongolia: synthesis of a 6-year grazing experiment. Journal of Arid Environments, 135, pp. 50-63, 2016. DOI: 10.1016/j.jaridenv.2016.08.003
- [16] Huete, A.R., A soil adjusted vegetation index (SAVI). Remote Sensing of Environment, 25, pp. 295-309, 1988. DOI: 10.1016/0034-4257(88)90106-X
- [17] Morón, N.A.C., Manzano, M.G. and Mata-González, R., Soil hydrology and vegetation as impacted by goat grazing in Vertisols and Regosols in semi-arid shrublands of northern Mexico. The

- Rangeland Journal, 39(4), pp. 363-373, 2017. DOI: 10.1071/RJ17061
- [18] Oliveira, L.M.M., Montenegro, S.M.G.L., Antonino, A.C.D., Silva, B.B., Machado, C.C.C. and Galvêncio, J.D., Análise quantitativa de parâmetros biofísicos de bacia hidrográfica obtidos por sensoriamento remoto. *Pesquisa Agropecuária Brasileira*, 47(9), pp. 1209-1217, 2012. DOI: 10.1590/S0100-204X2012000900005
- [19] Serrano, J., Shahidian, S. and Marques da Silva, J., Evaluation of normalized difference water index as a tool for monitoring pasture seasonal and inter-annual variability in a Mediterranean agro-silvo-pastoral system. *Water*, 11(1), pp. 62-82, 2019. DOI: 10.3390/w11010062
- [20] Serrano, J., Shahidian, S. and Marques da Silva, J., Monitoring seasonal pasture quality degradation in the Mediterranean Montado ecosystem: proximal versus remote sensing. *Water*, 10(10), pp. 1422, 2018. DOI: 10.3390/w10101422
- [21] Silva, J.L.B., Moura, G.B.A., França e Silva, Ê.F., Lopes, P.M.O., Silva, T.T.F., Lins, F.A.C., Silva, D.A.O. and Ortiz, P.F.S., Spatial-temporal dynamics of the Caatinga vegetation cover by remote sensing in municipality of the Brazilian semi-arid. *Revista Brasileira de Ciências Agrárias (Agrária)*, 14(2), pp. 1-10, 2019. DOI: 10.5039/agraria.v14i4a7128
- [22] Silva, L.G., Galvêncio, J.D. Análise comparativa da variação nos índices NDVI e SAVI no Sítio PELD – 22, em Petrolina – PE, na primeira década do século XXI. *Revista Brasileira de Geografia Física*, 5(6), pp. 1446-1456, 2012. DOI: 10.26848/rbgf.v5.6.p1446-1456
- [23] Sha, Z., Wang, Y., Bai, Y., Zhao, Y., Jin, H., Na, Y. and Meng, X., Comparison of leaf area index inversion for grassland vegetation through remotely sensed spectra by unmanned aerial vehicle and field-based spectroradiometer. *Journal of Plant Ecology*, 12(3), pp. 395-408, 2019. DOI: 10.1093/jpe/rtz036
- [24] Stolf, R., Teoria e teste experimental de fórmulas de transformação dos dados de penetrômetro de impacto em resistência de solo. *Revista brasileira de ciência do solo*, 15(3), pp. 229-235, 1991.
- [25] Tepanosyan, G.H., Asmaryan, S.G., Muradyan, V.S. and Saghatelian, A.K. Mapping man-induced soil degradation in Armenia's high mountain pastures through remote sensing methods: a case study. *Remote Sensing Applications: Society and Environment*, 8, pp. 105-113, 2017. DOI: 10.1016/j.rsase.2017.08.006
- [26] Valle Júnior, R. F., Siqueira, H. E., Valera, C. A., Oliveira, C. F., Fernandes, L.F.S., Moura, J.P. and Pacheco, F.A.L. Diagnosis of degraded pastures using an improved NDVI-based remote sensing approach: an application to the Environmental protection area of Uberaba River Basin (Minas Gerais, Brazil). *Remote Sensing Applications: Society and Environment*, 14, pp. 20-33, 2019. DOI: 10.1016/j.rsase.2019.02.001
- [27] Vasques, I.C., Souza, A.A., Morais, E.G., Benevenuto, P.A., da Silva, L.D.C., Homem, B.G., Casagrande, R.D. and Silva, B.M., Improved management increases carrying capacity of Brazilian pastures. *Agriculture, Ecosystems & Environment*, 282, pp. 30-39, 2019. DOI: 10.1016/j.agee.2019.05.017
- [28] Warrick, A.W., Nielsen, D.R., Spatial variability of soil physical properties in the field. In: D. Hillel, Ed., *Applications of soil physics Academic D.*, New York, US, 1980, pp. 319-344.
- [29] Yuan, Z.Q., Wu, Q.B., Song, X., Jiang, X.J., Gao, S.R., Wang, Q.F. and Li, G.Y., Pasture degradation impact on soil carbon and nitrogen fractions of alpine meadow in a Tibetan permafrost region. *Journal of Soils and Sediments*, pp. 1-13, 2020. DOI: 10.1007/s11368-020-02596-1
- [30] Zape Digital. Zoneamento agroecológico de Pernambuco. Embrapa Solos, Documentos, (35). Embrapa Solos, Recife, PE, Brasil, 2011.
- [31] Zhumanova, M., Mönnig, C., Hergarten, C., Darr, D. and Wrage-Mönnig, N., Assessment of vegetation degradation in mountainous pastures of the Western Tien-Shan, Kyrgyzstan, using eMODIS NDVI. *Ecological Indicators*, 95, pp. 527-543, 2018. DOI: 10.1016/j.ecolind.2018.07.060
- P.H.D. Batista**, is Irrigation and Drainage Technologist in 2015 from the Federal Institute of Ceará (IFCE), Brazil, a MSc. in 2017 and PhD. student in Agricultural Engineering at the Federal Rural University of Pernambuco (UFRPE), Brazil.
ORCID: 0000-0002-8710-5678
- G.L.P. Almeida**, has a BSc. in Agricultural Engineering in 2007, from the Federal Rural University of Pernambuco (UFRPE), Brazil, and a PhD. in Agricultural Engineering from the Federal Rural University of Pernambuco (UFRPE), Brazil. He is currently an adjunct professor at the Federal Rural University of Pernambuco (UFRPE), Brazil.
ORCID: 0000-0002-5813-4517
- J.L.B. Silva**, is Irrigation and Drainage Technologist in 2014 from the Federal Institute of Ceará (IFCE), Brazil, an MSc. in 2016 and is a PhD. student in Agricultural Engineering at the Federal Rural University of Pernambuco (UFRPE), Brazil.
ORCID: 0000-0002-2611-4036
- H. Pandorfi**, has a BSc. in Agronomy Engineering in 1999, from the Luiz de Queiroz Higher School of Agriculture (ESALQ), Brazil, and a PhD. in Physics of the Agricultural Environment from the Luiz de Queiroz Higher School of Agriculture (ESALQ), Brazil. He is currently an adjunct professor at the Federal Rural University of Pernambuco (UFRPE), Brazil.
ORCID: 0000-0002-2037-8639
- M.V. Silva**, has a BSc. in Agricultural Engineering in 2016 from the State University of Goiás (UEG), Brazil, a MSc. in 2019 and is a PhD. student in Agricultural Engineering at the Federal Rural University of Pernambuco (UFRPE), Brazil.
ORCID: 0000-0002-1318-2320
- R.A.B. Silva**, has a BSc. in Agricultural Engineering in 2016 from the Federal Rural University of Pernambuco (UFRPE), Brazil, an MSc. in 2019 and a PhD. student in Agricultural Engineering at the Federal Rural University of Pernambuco (UFRPE), Brazil.
ORCID: 0000-0003-3263-9194
- M.V.N. Melo**, BSc student in Agricultural Engineering at the Federal Rural University of Pernambuco (UFRPE), Brazil.
ORCID: 0000-0002-9590-4381
- F.A.C. Lins**, has a BSc. Agronomy Engineering in 2015 from the Federal Rural University of Pernambuco (UFRPE), Brazil, a MSc. in 2018 and is a PhD. student in Agricultural Engineering at the Federal Rural University of Pernambuco (UFRPE), Brazil.
ORCID: 0000-0001-6498-9805
- J.J.F. Cordeiro Junior**, has a BSc. in Agricultural Engineering in 2011, from the Federal Rural University of Pernambuco (UFRPE), Brazil and a PhD. in Agricultural Engineering from the Federal Rural University of Pernambuco (UFRPE), Brazil. He is currently an adjunct professor at the Federal University of Sergipe (UFS), Brazil.
ORCID: 0000-0002-1138-8309