

Odovtos International Journal of Dental Sciences

ISSN: 1659-1046 ISSN: 2215-3411

Facultad de Odontología. Universidad de Costa Rica

Genaro, Luis Eduardo; Marconato, José Victor; Hanai, Daniel; Gobbo-Pawloski, Carolina Luisa; Capote, Ticiana Sidonko de Oliveira Virtual Reality in Oral Hygiene Instruction: an Immersive Approach Odovtos International Journal of Dental Sciences, vol. 24, no. 1, 2022, January-April, pp. 177-187 Facultad de Odontología. Universidad de Costa Rica

DOI: https://doi.org/10.15517/IJDS.2022.49283

Available in: https://www.redalyc.org/articulo.oa?id=499570300017



Complete issue

More information about this article

Journal's webpage in redalyc.org



Scientific Information System Redalyc

Network of Scientific Journals from Latin America and the Caribbean, Spain and Portugal

Project academic non-profit, developed under the open access initiative



International Journal of Dental Sciences

https://revistas.ucr.ac.cr/index.php/Odontos | ISSN: 2215-3411

DOI: 10.15517/IJDS.2022.49283

CLINICAL RESEARCH

Received: 1-X-2021

Virtual Reality in Oral Hygiene Instruction: an Immersive Approach

Accepted:

17-XI-2021

Realidad virtual en la enseñanza de la higiene bucal:

Published Online: 26-XI-2021

un enfoque inmersivo

Luis Eduardo Genaro¹; José Victor Marconato²; Daniel Hanai³; Carolina Luisa Gobbo Pawloski⁴, Ticiana Sidonko de Oliveira Capote⁵

1. São Paulo State University (UNESP), School of Dentistry, Department of Morpholog, Genetics, Orthodontic and Pediatric Dentistry, Araraquara, São Paulo, Brazil. https://orcid.org/0000-0003-4206-2974

2. Central Paulista University Center (UNICEP), School of Dentistry, São Carlos, São Paulo, Brazil. https://orcid.org/0000-0002-5495-5860

3. Social Service of Commerce (SESC), Araraquara, São Paulo, Brazil.

https://orcid.org/0000-0002-4335-0233

4. Social Service of Commerce (SESC), Araraquara, São Paulo, Brazil. https://orcid.org/0000-0002-1658-5547

5. São Paulo State University (UNESP), School of Dentistry, Department of Morpholog, Genetics, Orthodontic and Pediatric Dentistry, Araraquara, São Paulo, Brazil. https://orcid.org/0000-0003-0979-5826

Correspondence to: Dr. Luis Eduardo Genaro - luis-genaro@outlook.com

ABSTRACT: Currently, oral health issues are increasingly been addressed and virtual reality (VR) is being used as an aid in health areas. However, there is little evidence of VR being used in preventing oral health problems. To evaluate the effectiveness of VR in oral hygiene education. Three hundred pupils, aged between 9 and 12 years, received instructions in oral hygiene by immersion VR and they gave their opinions about the performed activities. Chi-square and Fisher's exact tests were applied, with a significance level of 5%. Part perception of oral hygiene showed no statistically significant difference (P > 0.550). There was an increase in the use of dental floss (P<0.001), tongue brushing (P<0.001), and tooth brushing (P<0.001). Both drawings and comments made by participants related to oral hygiene were positive regarding the project. Conclusions: The use of VR in oral hygiene instruction was effective. Most participants increased brushing and flossing frequency and produced drawings related to oral hygiene. VR is effective and should be used in dentistry for teaching and prevention, especially with children, as it presents an interactive instructional space that can be approached in a playful way. This study presents dentists with a promising alternative for changing oral hygiene habits.

KEYWORDS: Virtual reality; Dentistry; Oral hygiene.

RESUMEN: Actualmente, los problemas de salud bucal se abordan cada vez más y la realidad virtual (RV) se utiliza como avuda en las áreas de salud. Sin embargo, hav poca evidencia de que la RV se utilice para prevenir problemas de salud bucal. Evaluar la efectividad de la RV en la educación en higiene bucal. Trescientos alumnos, con edades comprendidas entre los 9 y los 12 años, recibieron instrucciones en higiene bucal por inmersión VR y opinaron sobre las actividades realizadas. Se aplicaron las pruebas de Chi-cuadrado y exacta de Fisher, con un nivel de significancia del 5%. La percepción parcial de la higiene bucal no mostró diferencias estadísticamente significativas (P > 0.550). Hubo un aumento en el uso de hilo dental (P < 0.001), cepillado de la lengua (P < 0.001) y cepillado de dientes (P < 0.001). Tanto los dibujos como los comentarios de los participantes relacionados con la higiene bucal fueron positivos con respecto al provecto. Conclusiones: El uso de la RV en la instrucción de higiene bucal fue eficaz. La mayoría de los participantes aumentaron la frecuencia de cepillado y uso del hilo dental y produjeron dibujos relacionados con la higiene bucal. La realidad virtual es eficaz y debe utilizarse en odontología para la enseñanza y la prevención, especialmente con los niños, ya que presenta un espacio educativo interactivo que se puede abordar de manera lúdica. Este estudio presenta a los odontólogos una alternativa prometedora para cambiar los hábitos de higiene bucal.

PALABRAS CLAVE: Realidad virtual; Odontología; Higiene oral.

INTRODUCTION

With population growth, oral health issues are increasingly important to society. Oral diseases are still an important public health problem. According to WHO (World Health Organization) statistics, more than 60% of children worldwide have dental caries (1).

Advances in technology influence different areas of society (2) and can be applied in different areas, such as health care (3). Standing out in health focused technologies, virtual reality (VR) immerses users in a three-dimensional, interactive, computer-generated environment that simulates the characteristics of the real world, spaces and imaginary objects (3-7). It originated in the 20th century with great advances in the 1980s (3).

This technology provides a subjective experience of being in one place or environment,

even when physically located elsewhere. Another effect is immersion, in which sensory and perceptual information can offer a therapeutic environment (7).

VR is used as a professional learning tool (2) where educational programs based on VR technology can provide pupils with instant and consistent feedback (8). There is growing interest and evidence for the benefit of using interactive computer-based technologies to transform education and increase the autonomy and quality of life of individuals with VR (9).

Through VR, it is possible to create a learning environment which allows participants to practice social skills in a non-threatening, synthetic, but realistic, motivating, controllable and diversifiable environment during learning (10). There is evidence that VR technologies can, and do represent authentic and plausible social settings

and scenarios, reflecting and supporting real-world conventions, understanding and behaviors for a variety of different user groups (8,11).

VR has been used in neuropsychological assessment to diagnose and treat various mental illnesses. Therapy through VR permits the use of a realistic and involving environment, adapted to the needs of the individual and allows repetitive, consistent and systematic training. The capabilities of VR make it useful for patients with mental health problems. In the field of pediatric psychiatry, VR is used for attention training, social skills, cognitive rehabilitation and education. Previous studies have shown that children displayed more interest in participating in training through VR; one explanation being that in children and adolescents, neurological development makes the VR environment appear more vivid than in adults. Therefore, the VR experience can have a stronger impact on children and adolescents (12,13).

Dentistry is one of areas where VR applications can be used in various oral health related topics. Childhood, especially the first years of school life, is a time of discovering sensations (14); it is also when dental eruption of the first permanent teeth occurs (15), making it a propitious time for oral hygiene focused learning.

The benefits of using VR to reduce dental anxiety and pain levels during dental procedures have been widely addressed in scientific literature (16-21), and its usefulness as a distraction tool is receiving increasing attention in medical settings (20). During aversive experiences, VR can improve pain management and reduce the perceived duration of a procedure (21). Some studies have shown that, for both pain and anxiety, the use of VR is more effective in children than adults. One possible reason for this may be that VR is especially attractive to children as they become more involved in capricious

thoughts and are fascinated by imaginative play (19,20,21).

Health education programs in the school environment which focus on healthy habits and behavior, are able to improve the level of knowledge children have in relation to the health-disease process, and are thus considered an effective option. In addition, children in a certain age group and who have contact with the construction of healthy habits, can perpetuate such habits during adulthood.

Using VR, we can create simulators that allow the improvement of skills, such as simulating virtual patients with certain clinical conditions, so that the user can train on interventions, and observe the effects and reactions in the virtual organism (5).

Although studies show positive results with VR, there is little evidence related to oral health, especially with a focus on prevention. Thus, the objective of this study was to evaluate the effectiveness of virtual reality in oral hygiene education. The study is based on assessing whether VR will be able to change the habits of participants. Our hypothesis is that VR will be effective in improving oral health habits.

MATERIALS AND METHODS

This study was approved by the by the Ethics Committee of São Paulo State University (UNESP), School of Dentistry, Araraquara (CAEE: 24730219.9.0000.5416). Children aged 9 to 12 years, pupils from public schools in the city of Araraquara, São Paulo, Brazil were invited to participate in the study. To participate in the study, free and informed written consent was obtained prior to intervention from guardians. Participants were informed about their free choice of whether or not to participate in the study. In the first stage

of the study, 350 pupils took part, but 50 of these did not participate in the second stage. Thus, a final sample size of 300 participants was used in the statistical analysis.

We employed systematic probabilistic sampling. For inclusion criteria, participants were defined as being 9 to 12 years old without any psychological handicap. For exclusion criteria, participants were defined as those who refused to participate in the study, those who were younger or older, and those with some psychological disability.

This project was undertaken in partnership with the Social Service of Commerce of Araraquara City (SESC Araraquara). SESC Araraquara developed the project "Inside the Mouth", which consisted of visiting some public schools in the city and using VR glasses to show themes related to oral health.

Oral hygiene questionnaires were applied before and after (one week apart) using VR glasses, where oral hygiene instructions were demonstrated (Figure 1 and Figure 2). The questionnaires were pre-tested by means of a pilot evaluation, in order to establish consistency and effectiveness. Forty

virtual reality glasses (Oculus Go, Facebook Technologies, Auburn, USA) were purchased by SESC Araraquara and used in the study. Participants were encouraged to write or make drawings related to the activity. After an interval of one week, the participants answered the oral hygiene questionnaire again.



Figure 1. Pupils using VR glasses and receiving oral hygiene information.




Figure 2. A) Representation of the structures of the oral cavity and dental biofilm. B) Representation of mixed dentures. C) Representation of the toothbrushing movement. D) Representation of tongue brushing. E) Representation of the use of dental floss. F) Representation of a healthy mouth, with a clean tongue and teeth, after oral hygiene.

VR glasses allowed the development of a three-dimensional virtual environment of a mouth by a video lasting eight minutes in which issues related to oral hygiene are addressed in a playful way. In addition, this methodology is consistent with reality and is suitable for target audience understanding (children aged 9 to 12 years).

In this "Inside the Mouth" journey, participants visualized oral cavity structures, such as the tongue, teeth and cheek, and were able to observe curiosities and the care necessary for maintaining oral hygiene.

In the oral cavity, participants observed the dental biofilm, composed of bacteria, food debris and saliva, demonstrating that the dental biofilm is a dynamic structure, as bacteria are active organisms in the mouth. It showed that by using a toothbrush and dental floss, it is possible to remove or disorganize this biofilm, reducing the possibility of oral diseases appearing.

Pupils visualized the existence and difference between primary and permanent teeth. The focal point was the first permanent molar, showing that this tooth had just erupted, and being below the occlusal plane was subjected to greater plaque accumulation, thus stressing its importance for the life of the individual.

The virtual animation also presented brushing movement, using a toothbrush with properly positioned bristles, effectively removing the dental biofilm; and tongue brushing was also introduced. The correct use of dental floss was also demonstrated with correct positioning between the proximal surfaces of the teeth reinforcing its importance in removing food residues from areas that the toothbrush cannot reach.

A healthy mouth with a clean tongue and teeth was shown after oral hygiene, once again emphasizing the importance of oral health for a healthy body as a whole.

The journey ended with the appearance of an educational book, with stories, tips, and playful games. Those monitoring the session ended the educational activity by punctuating the main topics addressed by the dental games, and participants wrote or made drawings related to the activities.

Data were analyzed using the chi-squared and Fisher's exact tests with a significance level set at 5%.

RESULTS

Table 1 presents the results of oral hygiene issues before and after the project activities. Issues such as proper oral hygiene, flossing and tongue brushing. The data are presented as percentages and total values.

There was no difference regarding oral hygiene knowledge before and after the project

activities (P=0.550, Fisher's exact test). However, after the VR project, a greater number of participants started to use dental floss (P<0.001, Chi-square test), and brush their tongue (P<0.001, Chi-square test).

Table 2 shows the daily toothbrushing frequency before and after participating in the project activities. The data are presented as percentages and total values.

An increase in daily toothbrushing frequency can be seen after participation in the VR project (P<0.001, Fisher's exact test).

Figure 3 shows some drawings made by participants after immersion in VR.

Two evaluators performed the descriptive evaluation, Kappa analysis was performed to measure the degree of interrater agreement on the results. The level of agreement was acceptable.

Analyzing the pupils drawings, we verified that 180 drawings were related to oral cavity structures, and depicted products used in oral hygiene, such as toothbrushes, toothpaste and dental floss; 30 drawings were related to VR glasses. Ninety messages were written by participants, most of them reported "I learned how to take care of my teeth", "it was really cool", "now I know how to brush my teeth", "I learned how to use dental floss" and "I didn't know I needed to clean my tongue".

Table 1. Results of oral hygiene issues before and after the project activities.

	Before the activities		After the	Statistical	
_	Yes	No	Yes	No	analysis
Care related to oral hygiene	256 (85.4%)	44 (14.6%)	295 (98.3%)	5 (1.7%)	<i>P</i> =0.550
Use of dental floss	140 (46.6%)	160 (53.4%)	250 (83.4%)	50 (16.6%)	<i>P</i> <0.001
Tongue brushing	236 (78.6%)	64 (21.4%)	290 (96.4%)	10 (3.3%)	<i>P</i> <0.001

Table 2. Daily frequency of toothbrushing before and after project activities.

	None	Once	Twice	Three times	Four times	Statistical analysis
Before the activities	5 (1.6%)	44 (14.7%)	128 (42.6%)	110 (36.7%)	13 (4.4%)	<i>P</i> <0.001
After the activities	0	4 (1.3%)	57 (19%)	158 (52.7%)	81 (27%)	



Figure 3. Drawings made by participants after immersion in VR.

DISCUSSION

Due to the COVID-19 pandemic our study was only able to perform assessments in the week after the activity. Technically, our study would evaluate other periods such at one week, and one, three and six months, as well as clinically evaluate plaque score; unfortunately, we had to therefore adapt our study to the situation we are experiencing. Thus, we were only able to evaluate effectiveness based on oral hygiene measures reported by the pupils, which is a limitation of the study. Even so, we were able to observe from the questionnaires that participants increased the frequency of toothbrushing, flossing and tongue brushing.

VR has allowed an advance in the teaching-learning process, improving the population's quality of life (2,6,7). In recent decades, there has been a huge advance in technology aimed at VR (24) creating new immersive educational experiences, such as human anatomy (25) training in oral surgery (26,27) and even entertaining children during dental treatment (28). Modern 3D virtual systems have established the way for new approaches in medical imaging and education (6).

Pediatric dentistry also follows the high technology path, highlighting the importance of

establishing habits aimed at oral hygiene. Oral education goes through recent developments and extraordinary research (29).

This present work is innovative, as no other studies were found analyzing the effectiveness of VR in oral hygiene instruction. It demonstrated an interactive, immersive, instructional and motivational environment in oral health using VR resources that allow effective interaction with the target audience.

This research reached children from low-income families who were pupils at public schools located in city suburbs. In many cases, parents do not have an education level to ensure adequate oral hygiene guidance, as there is a correlation between parent educational level and child oral hygiene (30).

Although there was no significant difference regarding oral hygiene care knowledge (Table 1), at the beginning of the study children reported having regular or good oral hygiene and, after motivation they reported good to excellent oral hygiene, according to brushing habits and dental flossing. Our results were similar to those of Markeviciute & Narbutaite (31) who when analyzing orphaned children after oral hygiene motivation, observed a significant improvement in tooth brushing frequency.

Most people require motivation in oral hygiene instruction to acquire the habit of regular tooth brushing and dental flossing (32,33). We observed an improvement in oral health care using VR motivation and instruction in young pupils.

Oral hygiene is extremely important for the prevention of oral diseases, especially caries and periodontal disease (34). Tooth brushing is one of the main habits that constitute oral hygiene. In this study, after receiving guidance and immersion in VR, most participants reported performing three brushings a day. Zotti *et al.* (30) when evaluating

children who used technological applications to motivate oral hygiene also observed an improvement in oral hygiene, with a change in tooth brushing habits.

The results of our study corroborate with data found in literature, which show an improved adherence to oral hygiene performance in children when applying technological support (29-31,34). These digital tools are a good option for instigating behavioral change in oral hygiene, as they meet the needs while at the same time offer children motivation through fun (29).

As an educational method, VR allows the use of videos containing instructions related to oral hygiene; an additional advantage is that a video can be used repeatedly without any extra effort (35).

Above all, health promotion that seeks to address socioeconomic inequalities is extremely important since low-income children miss more school days due to dental diseases than children from high-income families (34).

At the beginning of the study, some children reported not having dental floss in their homes or not knowing how to use it. It is therefore necessary to provide instructions and support for the habitual use of dental floss, as only 10% of the population habitually use it, and it is indispensable for adequately cleaning the proximal surfaces when considering satisfactory oral hygiene (33).

Blake *et al.* (34) also found a significant increase in the level of dental floss use in children after oral health intervention, demonstrating that oral hygiene behavior changes after motivation.

They emphasized that promoting oral health can generate positive changes in knowledge and behaviors about oral hygiene in children.

The drawings and messages made by participants after participating in the VR project were all related to instructions in the video; most drawings presented oral cavity structures with the presence of dirt, and after brushing, sanitized structures (Figure 2) as well as the positive messages. We suggest that instruction by VR was considered fun as the children were committed and enthusiastic.

These instructed and analyzed hygiene habits are of great importance, as they are acquired in childhood by an educational and instructional process (33). Thus, the daily practice of oral hygiene is of paramount importance in educational programs which are aimed at preventing oral diseases.

This study presented an improvement of oral hygiene reports with the assistance of VR as a method of teaching good oral hygiene practices. One limitation of the study is the verification and quantification of bacterial biofilm before and after participation in the VR project. Future studies are needed to evaluate this variable.

CONCLUSION

The use of VR for oral hygiene instruction is effective and has a positive impact. It is an interactive, motivational and playful environment for oral health education. Most participants increased their brushing, flossing and tongue brushing frequency. VR presents innovative perspectives that can assist oral health education for young pupils.

AUTHOR CONTRIBUTION STATEMENT

Conceptualization: L.E.G. and T.S.O.C.

Methodology: L.E.G. Formal analysis: C.L.G.P. Investigation: J.V.M. he g. Resources: D.H. and C.L.G.P.

Writing-preparation of the original draft: L.E.G.

and D.H.

Writing-proofreading and editing: L.E.G. and C.L.G.P.

Visualization: J.V.M., D.H. and T.S.O.C.

Supervision: T.S.O.C.

ACKNOWLEDGMENTS

We would like to thank the Social Service for Commerce-SESC, for the tech-nical support and financial support.

REFERENCES

- 1. Huang T.K., Yang C.H., Hsieh Y.H., Wang J.C., Hung C.C. Augmented reality (AR) and virtual reality (VR) applied in dentistry. Kaohsiung J Med Sci. 2018; 34 (4), 243-48. doi: 10.1016/j.kjms.2018.01.009
- 2. Bezerra T.F. & de Souza V.L.V. The use of Virtual Reality as an occupational therapeutic resource in the neurological rehabilitation of children and adolescents. Rev Interinst Bras Ter Ocup. 2018; 2 (2), 272-91. doi: 10.47222/2526-3544.rbto12739
- 3. O'Connor S. Virtual Reality and Avatars in Health care. Clin Nurs Res. 2019; 28 (5), 523-28, doi: 10.1177/1054773819845824
- 4. Aminabadi N.A., Erfanparast L., Sohrabi A., Oskouei S.G., Naghili A. The impact of virtual reality distraction on pain and anxiety during dental treatment in 4-6-year-old children: a

- randomized controlled clinical trial. J Dent Res Dent Clin Dent Prospectspects. 2012; 6 (4), 117. doi: 10.5681/joddd.2012.025
- 5. Paiva P.V.F., dos Santos Machado L., dos Santos S.R., Romero R.O.G. Health Education capabilities Based on Virtual Reality. Rev Elet Com, Infor e Inov em Saúde. 2013; 7 (3). doi: 10.3395/reciis.v7i3.540
- 6. Moro C., Stromberga Z., Raikos A., Stirling A. The effectiveness of virtual and augmented reality in health sciences and medical anatomy. Anat Sci Educ. 2017; 10 (6), 549-59. doi: 10.1002/ase.1696
- 7. Burrai F., Othman S., Brioni E., Silingardi M., Micheluzzi V., Luppi M., La Manna G. Virtual reality in dialysis: a new perspective on care. J Ren Care. 2018; 44 (4), 195-96. doi: 10.1111/jorc.12264
- 8. Ke F., Moon J., Sokolikj Z. Virtual reality-based social skills training for children with autism spectrum disorder. J Spec Educ Tech. 2020. doi: 10.1177/0162643420945603
- 9. Walsh E., Holloway J., McCoy A., Lydon H. Technology-aided interventions for employment skills in adults with autism spectrum disorder: A systematic review. J Autism Dev Disord. 2017; 4 (1), 12-25. doi: 10.1007/s40489-016-0093-x
- 10. Bellani M., Fornasari L., Chittaro L., Brambilla P. Virtual reality in autism: State of the art. Epidemiol Psychiatr Sci. 2011; 20 (3), 235-38. doi: 10.1017/S204579601 1000448
- 11. Parsons S. Authenticity in virtual reality for assessment and intervention in autism: A conceptual review. Educ Res Rev. 2016; 19, 138-157. doi: 10.1016/j.edurev.2016.08.001
- 12. Bailey J.O. & Bailenson J.N. Immersive virtual reality and the developing child. In Cognitive development in digital contexts.

- Academic Press. 2017, 181-200. doi: 10.1016/B978-0-12-809481-5.00009-2
- 13. Kim S. & Kim E. The use of virtual reality in psychiatry: a review. J Korean Acad Child Adolesc Psychiatry. 2020; 31 (1), 26. doi: 10.5765/jkacap.190037
- 14. Madeira M.C.; Rizzolo R.J.C. Anatomia do Dente. 8 ed. São Paulo: Sarvier. 2016.
- 15. Mills A.C. Creating Web-based, multimedia, and interactive courses for distance learning. Comput Nurs. 2000; 18 (3): 125-31. doi: 10835811
- 16. Wiederhold MD, Gao K, Wiederhold BK. Clinical use of virtual reality distraction system to reduce anxiety and pain in dental procedures. Cyberpsychol Behav Soc Netw. 2014; 17:359-65. doi: 10.1089/ cyber.2014.0203
- 17. Raghav K., Van Wijk A.J., Abdullah F., Islam M.N., Bernatchez M, De Jongh A. Efficacy of virtual reality exposure therapy for treatment of dental phobia: A randomized control trial. BMC Oral Health. 2016; 27: 25. doi: 10.1186/s12903-016-0186-z
- 18. Eijlers R., Utens E., Staals L.M., de Nijs P.F.A., Berghmans J.M., Wijnen R.M.H., Hillegers M.H.J., Dierckx B., Legerstee J.S. Systematic Review and Meta-analysis of Virtual Reality in Pediatrics: Effects on Pain and Anxiety. Anesth Analg. 2019; 129:1344-53. doi: 10.1213/ANE.00000000000004165
- 19. Gujjar K.R., van Wijk A., Kumar R., de Jongh A. Efficacy of virtual reality exposure therapy for the treatment of dental phobia in adults: A randomized controlled trial. J Anxiety Disord. 2019; 62:100-108. doi: 10.1016/j.janxdis.2018.12.001
- 20. Ahmadpour N., Keep M., Janssen A., Rouf A.S., Marthick M. Design strategies for virtual reality interventions for managing pain and anxiety in children and adolescents: scoping review. JMIR serious games. 2020; 8 (1). doi: 10.2196/14565

- 21. López-Valverde N., Muriel Fernández J., López-Valverde A., Valero Juan L.F., Ramírez J.M., Flores Fraile J., & Bravo M. (2020). Use of virtual reality for the Management of anxiety and pain in dental treatments: Systematic review and meta-analysis. J Clin Med, 9 (4), 1025. doi: 10.3390/jcm9041025
- 22. Marques I.R., Marin H.D.F. Nursing on the web: the creation and validation process of a coronary heart disease web site. Rev Lat Am Enfermagem. 2002; 10 (3): 298-07. doi: 10.1590/S0104-11692002000300005
- 23. Amantini S.N.S.R., Montilha A.A.P., Antonelli B.C., Leite K.T.M., Rios D., Cruvinel T., Machado M.A.A.M. Using Augmented Reality to Motivate Oral Hygiene Practice in Children: Protocol for the Development of a Serious Game. JMIR Research Protocols. 2020; 9 (1). doi: 10.2196/10987
- 24. Rizzo A., Hartholt A., Grimani M., Leeds A., Liewer M. Virtual reality exposure therapy for combat-related posttraumatic stress disorder. Computer. 2014; 47: 31-7. doi: 10.1109/MC.2014.199
- 25. Messier E., Wilcox J., Dawson-Elli A., Diaz G., Linte C.A. An interactive 3D virtual anatomy puzzle for learning and simulation Initial demonstration and evaluation. Stud Health Technol Inform. 2016; 220: 233-40. doi: 10.3233/978-1-61499-625-5-233
- 26. Ayoub A., Pulijala Y. The application of virtual reality and augmented reality in Oral & Maxillofacial Surgery. BMC Oral Health. 2019; 19 (1) 238. doi: 10.1186/s12903-019-0937-8
- 27. Bartella A.K., Kamal M., Scholl I., Schiffer S., Steegmann J., Ketelsen D., Lethaus B. Virtual reality in preoperative imaging in maxillofacial surgery: implementation of "the next level"? Br J Oral Maxillofac Surg. 2019; 57 (7), 644-48. doi: 10.1016/j. bjoms.2019.02.014

- 28. Koticha P., Katge F., Shetty S., Patil D.P. Effectiveness of Virtual Reality Eyeglasses as a Distraction Aid to Reduce Anxiety among 6-10-year-old Children Undergoing Dental Extraction Procedure. J Clin Pediatr Dent. 2019; 12 (4), 297. doi: 10.5005/jp-journals-10005-1640
- 29. Hotwani K., Sharma K., Nagpal D., Lamba G., Chaudhari P. Smartphones and tooth brushing: content analysis of the current available mobile health apps for motivation and training. Eur Arch Paediatr Dent. 2020; 21(1), 103-08. doi: 10.1007/s40368-019-00457-1
- Zotti F., Pietrobelli A., Malchiodi L., Nocini P.F., Albanese M. Apps for oral hygiene in children 4 to 7 years: Fun and effectiveness. J Clin Exp Dent. 2019; 11 (9), 795. doi: 10.4317/jced.55686
- 31. Markeviciute G & Narbutaite J. Effectiveness of a motivation and practical skills development methods on the oral hygiene of orphans children in Kaunas, Lithuania. J Oral

- Maxillofac Surg. 2015; 6 (3). doi: 10.5037/jomr.2015.6302
- 32. Ciancio S. Improving oral health: current considerations. J Clin Periodontol. 2003; 30:4-6. doi: 10.1034/j.1600-051X.30.s5.2.x
- 33. Kubo F.M.M., Mialhe F.L. Dental floss: from difficulty to success in the removal of interproximal biofilms. Arq Odontol. 2011; 47 (1), 51-5. doi: 725232.
- 34. Blake H., Dawett B., Leighton P., Rose-Brady L., Deery C. School-based educational intervention to improve children's oral health-related knowledge. Health Promot Pract. 2015; 16 (4), 571-82. doi: 10.1177/1524839914560568
- 35. Baliga S., Deshpande M.A., Thosar N., Rathi N., Bane S., Deulkar P. Comparison of impact of oral hygiene instructions given via sign language and validated customized oral health education skit video on oral hygiene status of children with hearing impairment. J Indian Soc Pedod Prev Dent. 2020; 38 (1), 20. doi: 10.4103/JISPPD.JISPPD 37 20