



Geologica Acta: an international earth science journal

ISSN: 1695-6133

ISSN: 1696-5728

geologica-acta@ictja.csic.es

Universitat de Barcelona

España

Fatka, O.; Budil, P.

Digestive structures in Middle Ordovician trilobite *Prionocheilus*
ROUAULT, 1847 from the Barrandian area of Czech Republic

Geologica Acta: an international earth science journal,
vol. 16, no. 1, 2018, January-March, pp. 65-73

Universitat de Barcelona
España

DOI: <https://doi.org/10.1344/GeologicaActa2018.16.1.4>

Available in: <https://www.redalyc.org/articulo.oa?id=50558534004>

- How to cite
- Complete issue
- More information about this article
- Journal's webpage in [redalyc.org](https://www.redalyc.org)

redalyc.org

Scientific Information System Redalyc

Network of Scientific Journals from Latin America and the Caribbean, Spain and
Portugal

Project academic non-profit, developed under the open access initiative

Digestive structures in the Middle Ordovician trilobite *Prionocheilus* ROUAULT 1847, from the Barrandian area of Czech Republic

O. FATKA¹ P. BUDIL²

¹Charles University, Faculty of Sciences, Institute of Geology and Palaeontology

Albertov 6, 128 43, Prague, Czech Republic. E-mail: fatka@natur.cuni.cz

²Czech Geological Survey

Klárov 3, 118 21, Prague, Czech Republic. E-mail: petr.budil@geology.cz

ABSTRACT

Remains of a digestive system from a slightly damaged articulated specimen of the comparatively rare bathycheilid trilobite *Prionocheilus vokovicensis* (ŠNAJDR, 1956) are described for the first time. The specimen comes from the Middle Ordovician Šárka Formation of the Prague Basin and contains the midgut region of the digestive system preserved through the axial region of glabella and six anterior thoracic segments. The anterior-most part of the digestive system is unknown as the anterior glabellar lobes are not preserved in the studied specimen. In the cephalic shield, the remains of two pairs of gut diverticulae are seen in the posterior region of the glabella. Remains of five pairs of small cavities developed in the axis of the first six thoracic segments represent the remains of thoracic gut diverticulae. The discussed specimen possess the first undoubted remain of digestive structures established within the family *Bathycheilidae* (PŘIBYL, 1953).

KEYWORDS | *Prionocheilus*. Digestive tract. Ordovician. Barrandian area. Czech Republic.

INTRODUCTION

Nearly one thousand trilobite species have been described from the Cambrian to Devonian aged rocks of the Barrandian area of the Czech Republic (Valíček and Vaněk, 2001; Vaněk and Valíček, 2002) but remains of soft parts are extremely rare (*e.g.* Šnajdr, 1990; Fatka *et al.*, 2014). About twenty exceptionally preserved articulated trilobite exoskeletons with remains of digestive system have been collected from Cambrian Buchava and Jince formations (fms.) of the Skryje-Týřovice and Příbram-Jince basins (Jaekel, 1901; Fatka *et al.*, 2014). Similar number of trilobite specimens with undoubted remains of gut, cephalic and thoracic diverticulae and/or crop were recently documented in Ordovician specimens from the Mílina (Tremadocian), Šárka (Darriwilian), Letná (Sandbian),

and Bohdalec (Katian) fms. of the Prague Basin (for summary see Fatka *et al.*, 2014) (Fig. 1C).

Here, we describe the morphology of the digestive system in the rare bathycheilid trilobite *Prionocheilus vokovicensis* (ŠNAJDR, 1956). A slightly damaged internal mould of an originally complete and well articulated exoskeleton was recently identified by the junior author in collections housed in the National Museum of Prague (housed under the inventory number NM L35774). The external mould of the studied specimens is not stored in the National Museum collections. This extraordinary specimen was collected near the town of Rokycany, particularly in the fields of the Osek locality. In these fields, diverse Middle Ordovician fossils, preserved in silicified nodules of the Šárka and Dobrotivá fms., have been collected since the middle of the 19th century (Lajblová and Kraft, 2014).




FIGURE 1. Maps showing the location of the discovery site and Ordovician stratigraphy in the Barrandian area: A) Map of the Czech Republic and the Bohemian Massif with the Ordovician rocks of the Barrandian area; B) Ordovician of the Prague Basin showing the location of the Osek locality at which the studied specimen was found. C) Chart showing the correlation between global series, stages, stage slices, Time Slices (TS), Time Units (TU), and the regional chronostratigraphic and lithostratigraphic units recognized in the Ordovician of the Prague Basin with marked ranges of the species of *Prionocheilus* (stratigraphy modified from Bergström *et al.*, 2008; Fatka *et al.*, 2013). Ranges of the species of *Prionocheilus* based on Šnajdr (1956), Vaněk (1965), Budil and Kraft (2003), Mergl *et al.* (2008), Peršín and Budil (2009), Vokáč *et al.* (2015).

GEOLOGICAL SETTING

The Šárka Formation

The Šárka Formation (Fm.) contains a highly diverse skeletal fauna (*e.g.* Havlíček and Vaněk, 1966, 1990; Fatka and Mergl, 2009). The thickness of this formation ranges from several metres in the marginal parts of the basin to nearly 300 metres in segments of supposed rapid synsedimentary subsidence (Havlíček, 1981). The stratigraphy and depositional setting of the Šárka Fm. have been discussed by numerous authors (*e.g.* Kukul, 1962; Havlíček and Vaněk, 1966; Kraft and Kraft, 1992, 1999; Havlíček and Fatka, 1992; Havlíček, 1998; Servais *et al.*, 2008). In earlier schemes, the Šárka Fm. was correlated with the British Llanvirn Series (*e.g.* Havlíček and Vaněk, 1966); later, it was supposed to correspond to the late Arenig to early Llanvirn interval (*e.g.* Kraft *et al.*, 2001). Fatka *et al.* (2013) correlated the Šárka Fm. with the Oretanian Regional Stage which is equal to the middle Darriwilian (Bergström *et al.*, 2008). Based on the restricted ranges of graptolites, the Šárka Fm. has been divided into two biozones: the older *Corymbograptus retroflexus* and the younger *Didymograptus clavulus* biozones (Kraft *et al.*, 2001). The Šárka Fm. typically consist of poorly fossiliferous

dark shales. Our knowledge about fauna is generally based on specimens collected from siliceous nodules, which have attracted the attention of amateur collectors. However, such loose nodules do not provide information about the stratigraphic position and original palaeoenvironment (Budil *et al.*, 2007; Mergl *et al.*, 2008).

Fossil associations

Abundant remains of brachiopods, gastropods, trilobites, agnostids, phyllocarids, ostracods, bivalves, hyoliths, cephalopods, echinoderms, and conulariids associated with graptolites and ichnofossils have been studied for more than 150 years (Barrande, 1872; Havlíček and Vaněk, 1966; Chlupáč, 1970, 2003; Mikuláš, 1991; Horný, 2001; Mergl, 2002; Kraft and Kraft, 2003; Budil *et al.*, 2007; Manda, 2008; Mergl *et al.*, 2008; Fatka and Mergl, 2009; Steinová, 2012; Polechová, 2013; Aubrechtová, 2015). The intensively studied and diverse fauna made possible to distinguish several trilobite and non-trilobite associations of the Šárka Fm. (Havlíček, 1982, 1998; Havlíček and Vaněk, 1990; Mergl, 2002; Lefebvre, 2007). Mikuláš (1991, 1998) studied assemblages of ichnofossils, which he assigned to a transition from *Cruziana* to *Zoophycos* ichnofacies.

Trilobite associations

More than 50 species of trilobites have been described from the Šárka Fm. (Havlíček and Vaněk, 1966; Budil *et al.*, 2007; Mergl *et al.*, 2007). Remains of trilobite exoskeletons together with gastropod, brachiopod and bivalve shells are common elements in fossil associations. Trilobite associations have been discussed by Havlíček and Vaněk (1990), Havlíček (1998), Bruthansová (2003), Budil *et al.* (2007), Mergl *et al.* (2008), Fatka and Mergl (2009) and Mergl and Kozák (2016). All earlier proposals of trilobite associations were recently summarized by Fatka *et al.* (2015) and are shown in Figure 2.

The discussed damaged internal mould of an originally complete exoskeleton of *Prionocheilus vokovicensis* is preserved in a siliceous nodule that is known from the shallower part of the *Placoparia* Association of the Šárka Fm. (Fig. 2).

METHODS

Methods used to analyse the trilobite specimen included standard light microscopy of the external

surface of the internal mould (Microscope NIKON SMZ 1500, Leica S8APO); photographs were taken using digital cameras (a NIKON D 300 and an Olympus SZX-ILLB200) after coating with ammonium chloride. The drawing was made from a photograph using Corel Draw X3 and Adobe Photoshop CS5. The terminology used follows that proposed by Whittington and Kelly (1997), including the following abbreviations: sag. (sagittal), tr. (transverse). The specimen is housed in the National Museum of Prague, under the inventory number NML 35774.

SYSTEMATIC PALAEONTOLOGY

Family: Bathycheilidae PŘIBYL, 1953

Subfamily: Pharostomatinae HUPÉ, 1953

GENUS *Prionocheilus* ROUAULT, 1847

Type of species. *Prionocheilus verneuili* (ROUAULT, 1847), Sandbian/Katian, Formation de Riadan in the Martigné-Ferchaud South of Rennes, Caradocian, Brittany, France (by original designation).




FIGURE 2. Sketch representing the distribution of the major biofacies associated with the late Darriwilian Šárka Fm. and occurrence of *Prionocheilus*. The shallowest part of the basin was inhabited by the poor orthid brachiopod association, in the offshore areas the *Placoparia* Association rich with trilobites, brachiopods and other skeletal fauna; in the offshore, downslope, it gradually passes into the poor atheloptic trilobite association (not represented in this figure), which includes also the poor benthic dendroid 'gardens'. The water column was inhabited by poor planktonic graptolites and taxa of the poorly diverse *Caryocaris* and *Cyclopygid* biofacies. Poorly oxygenated black shales in central parts of the basin are dominated by the *Paterula* Association. Modified from Fatka and Mergl (2009, fig. 11D) and Fatka *et al.* (2015, fig. 2).




FIGURE 3. Morphology of the preserved parts of the digestive system in the cephalic and thoracic regions of *Prionocheilus* from the Šárka Fm. (Middle Ordovician, Darriwilian= Oretanian Regional Stage). A) Internal mould of *Prionocheilus vokovicensis* (ŠNAJDR, 1956) in dorsal view; National Museum of Prague, NML 35774; fields North of Rokycany (Osek locality). B) Interpretative sketch. Dc: cephalic gut diverticulae; Dt: thoracic gut diverticula; Ar: ichnofossils *Arachnostega*.

Remarks. The validity of *Prionocheilus* (ROUAULT, 1847) and *Pharostoma* (HAWLE AND CORDA, 1847) has been discussed for a long time. Šnajdr (1956), Vaněk (1965), Whittington (1965), and Siveter (1973, 1976) placed *Prionocheilus* in the synonymy of *Pharostoma*. However, Dean (1964) showed that the contribution of Rouault (1847) predates that of Hawle and Corda (1847). Ingham (1977) followed the proposal of Dean (1964) bringing the argumentation that *Prionocheilus* was used in zoological publications. However, in an addendum, Siveter (1976, pg.: 393) stressed the information about decision published in the Bulletin of Zoological Nomenclature bearing on the question of the usage of *Prionocheilus* instead of *Pharostoma*. In agreement with the scientific revisors in Horný and Bastl (1970), Hammann and Henry (1978), Hammann (1983), Zhou *et al.* (1984), Rábano (1989) and other authors, we follow the use of *Prionocheilus*. The systematic revision of *Prionocheilus* is, however, out of the scope of this paper.

Species. Siveter (1973, pg.: 2) assigned 19 species to *Pharostoma* (syn. of *Prionocheilus*). Sixteen years later, Rábano (1989, pg.: 93) listed 18 species classified under *Prionocheilus*. In comparison, Lemke (unpublished) provided the most complete list of 27 species assigned to *Prionocheilus*. Evaluation of their lists and detailed discussion is, however, out of the scope of this contribution.

Distribution. *Prionocheilus* has been established in West Gondwana (Argentina, Mendoza area; Morocco), European peri-Gondwana (France, Iberian Peninsula, Czech Republic, Great Britain, Ireland), Baltica (Estonia), Laurentia (Canada), Kazakhstania, SW China (Guizhou Province).

Prionocheilus ranges from the Tremadocian of France and Spain [*P. languedocensis* (COURTESSOLE AND PILLET, 1975)]

to the Ashgillian of Canada [*P. rarus* (COOPER AND KINDLE, 1936)], Ireland [*P. obtusum* (MCCOY, 1846)], and Kazakhstan [*P. solitus* (APOLLONOV, 1974)].

In Ordovician sequence of the Prague Basin, five species have been assigned to *Prionocheilus*: *P. borni* (VANĚK, 1995), *P. derceto* (VANĚK, 1995), *P. mendax* (VANĚK, 1965), *P. pulcher* (BARRANDE, 1846), and *P. vokovicensis* (ŠNAJDR, 1956). Stratigraphically, they range from the Middle Ordovician Šárka Fm. (Darriwilian) to the Upper Ordovician Bohdalec Fm. (Late Berounian Regional Stage = mid Katian) (Fig. 1C);

Prionocheilus vokovicensis (ŠNAJDR, 1956) (Figs. 3, 4 and 5).

Description. The studied specimen is preserved in a hard siliceous nodule, which represents one of lithotypes of the Middle Ordovician Šárka Fm. (Kukal, 1962; Chvátal, 2003; Drost *et al.*, 2003). It is a slightly incomplete, partly damaged, internal mould of posterior part of cephalon associated with a nearly complete articulated thorax of a middle-sized holaspis specimen.

The exoskeleton is preserved in a prone attitude; the preserved part is 28mm long (sag.) and reaches 24mm in maximum width (tr.). The width of the axial region of thoracic segments ranges from 10mm in the first segment to 7mm in the eleventh (=posterior-most preserved) segment.

The anterior border and preglabellar field were situated outside the siliceous nodule and, as a result, are not preserved. The surface of the fixigena and the anterior part of the glabella are strongly weathered. The pygidium and

the posterior-most thoracic segment were situated outside the siliceous nodule and are not preserved.

Two, quite deep, lobate cavities are developed in the left posterior part of the glabella; one narrow, sagittally oriented cavity is seen in the right posterior part of the glabella. Similar paired and quite deep cavities are developed also in the axial part of the third and fourth thoracic segment; unpaired, morphologically comparable cavities are seen also in the left part of the second segment axis as well as in the right part of the axis in the fifth and sixth thoracic segment.

Comparatively narrow, centrally placed and nearly parallel-sided remains of the dorsal exoskeleton are visible in several segments of the axial part of the thorax. These remains extend from the second to the sixth thoracic segment.

Interpretation. The above described cavities in the glabella and in the anterior part of the thorax recall the earlier described remains of trilobite digestive system (Lerosey-Aubril *et al.*, 2011, 2012; Fatka *et al.*, 2013, 2015). The placement inside the axial part of the cephalon and thorax combined with an apparent arrangement in pairs makes it possible to interpret these structures as paired gut diverticulae supposedly associated with a centrally placed gut (most probably not preserved in this specimen, the original position of the gut was under the parallel-sided and narrow area seen between paired cavities).

The alimentary canal, preserved in the posterior part of the glabella, is associated with two larger, separated, lobate imprints (left) and one longer and narrow cavity. All of these imprints are interpreted as two pairs of cephalic gut diverticulae (Dc2-3 in Fig. 3B). The small cavities developed on either side of the gut in the axial region of the second to sixth thoracic segments belong to partially preserved, paired thoracic gut diverticulae (Dt2 to Dt6 in Fig. 3B). The slight shift of thoracic diverticulae on the left side is, most probably, caused by decay of soft tissue during early diagenesis.

The thin fissure developed ventrally in the axis of the fourth and fifth segments, separates dorsally positioned, a sagittally elongated body which could represent the remains of a tubular gut (arrow in Fig. 4).

Remarks. The cephalon and all preserved thoracic segments are well articulated and the remains lack any signs of disarticulation in the posterior part of the thorax. Therefore, the disarticulation along the posterior part of the thorax in an unspecific location, *sensu* Daley and Drage (2016), could be excluded. Consequently, this specimen

represents partly preserved internal mould of an originally complete carcass.

Based on comparison with other complete specimens of *Prionocheilus vokovicense*, sag. length of the herein described specimen could be estimated to range from 35 to 40mm; it apparently represents a late holaspid stage.

Fine, branched and curved tunnels of the ichnospecies *Arachnostega gastrochaenae* BERTLING, 1992, are developed directly below the exoskeletal surface in different parts of the mould, especially in the eighth to eleventh thoracic segment (Ar in Fig. 3B). *Arachnostega* tunnels associated with body fossils have been reported from numerous Ordovician areas, including the Šárka Fm. of the Barrandian area (for summary see Fatka *et al.*, 2011).

Similarly, in specimens of *Colpocoryphe bohémica* described from the Šárka Fm. of the Barrandian area, as well as in the herein studied specimen of *Prionocheilus*, it is possible to suppose that the missing parts of the exoskeleton were situated outside of the nodule and were destroyed during diagenetic processes (compare Fatka *et al.*, 2015). Such a type of preservation agrees well with the very early diagenetic formation of siliceous nodules (for detailed discussion see Dabbard and Loi, 2012 and the following section of this paper).




FIGURE 4. Morphology of the preserved parts of the digestive system in the cephalic and thoracic regions of *Prionocheilus* from the Šárka Fm. (Middle Ordovician, Darrivilian=Oretanian Regional Stage). Internal mould of *Prionocheilus vokovicensis* (ŠNAJDR, 1956) in dorso-lateral view; National Museum of Prague, NML 35774; fields North of Rokycany (Osek locality). Ce: cephalic shield; or: occipital ring; 1 to 12: thoracic segments; arrow: sagittally elongated body which could represent the remains of a tubular gut.




FIGURE 5. Reconstruction of the morphology of the digestive system in *Prionocheilus* in dorsal view.

DISCUSSION

Decay of soft parts

Babcock and Chang (1997) and Babcock *et al.* (2000) showed, that the decay of internal soft parts started only a few hours after death in modern *Limulus* and the decay did not last more than one month. Also, other decay experiments have shown a rapid destruction of internal organs in marine arthropods by the activity of microbes associated with phosphatization of internal tissues (Butler *et al.*, 2015; Strang *et al.*, 2016). Similarly, as in other trilobites with preserved soft parts, the processes leading to preservation of the delicate remains of soft tissue should be initiated very shortly after the entombment of carcasses. This agrees with the model of early diagenetic phosphogenesis linked to Carbonate Fluor-Apatite (CFA) precipitation in upper levels of the sediment (Dabbard and Loi, 2012). Such type of preservation of soft parts could be explained by a rapid mineralisation of the gut associated with the early diagenetic formation of the siliceous nodule. This process was later on followed by a diagenetic removal of the original gut mineral phase, possibly due to weathering. The exoskeleton may have been dissolved at the same time.

Morphology of alimentary tract

Two major morphological types of alimentary tract have been distinguished by Lerosey-Aubril *et al.* (2011): i) simple tube with a crop, and ii) simple tube with metamERICALLY paired caeca (=gut diverticulae). Both morphologies have

been observed in Ordovician trilobites of the Prague Basin (Fatka *et al.*, 2014). Recently, Gutiérrez-Marco *et al.* (2017) described a third type of alimentary tract characterised by a co-occurrence of both crop and metamERICALLY paired caeca in large asaphid *Megistaspis (Ekeraspis) hammondi* from the Fezouata Lagerstätte of Morocco.

In *Prionocheilus*, similarly as in the Cambrian specimen of *Jiumenia anhuiensis* (Zhu *et al.*, 2014), the blind *Conocoryphe* (Budil and Fatka, 2008) and the small-eyed Middle Ordovician *Colpocoryphe* (Fatka *et al.*, 2015), a bell-shaped, anteriorly narrowing glabella is developed. Due to this glabellar morphology, metamERICALLY paired caeca could be expected. We also agree with Zhu *et al.* (2014), who explained the more abundant preservation of guts as sediment-like infilling (Šnajdr, 1990), while more rarely preserved metamERICALLY paired digestive caeca is connected with a suspected phosphatisation of this part of the digestive system (for discussion see Lerosey-Aubril *et al.*, 2012 and Fatka *et al.*, 2013).

Palaeoecology

Based on general morphology of exoskeleton and hypostome, Hammann (1983, pg.: 34, text-fig. 12) suggested an epibenthic mode of life for *Prionocheilus*; this is generally accepted (*i.e.* Mergl *et al.*, 2008, pg.: 279) (Fig. 2). Budil *et al.* (2007, pg.: 68) classified *P. vokovicensis* as a scavenger-predator. The occurrence of paired diverticulae, however, implies a long-standing digestive process of ingested particles, which is in agreement with a detritus feeding habit.

CONCLUSIONS

i) Remains of the digestive system are, for the first time, documented in an Ordovician bathycheilid trilobite.

ii) The studied carcass of *Prionocheilus vokovicensis* shows an apparent presence of two cephalic and at least six thoracic diverticulae (Fig. 5). In the studied specimen, preservation of the gut tract is questionable.

iii) The alimentary tract observed in *Prionocheilus* is known in trilobites with diverse visual abilities. Comparable tracts have been reported in the contemporaneous genus *Colpocoryphe*, as well as in the Cambrian genera *Conocoryphe* and *Jiumenia*.

ACKNOWLEDGMENTS

We thank Thomas Hegna (Western Illinois University, U.S.A.) and an anonymous reviewer for their helpful reviews and

the linguistic improvements made on our text. This study was supported by PROGRES Q45 of the Ministry of Education, Youth and Sports of Czech Republic and the Czech Geological Survey internal project No 339900. This paper is a contribution to the project IGCP 653 “Filling the gap between Cambrian Explosion and the GOBE”.

REFERENCES

- Apollonov, M.K. 1974. Ashgillskie trilobity Kazachstana [Ashgill trilobites from Kazakhstan]. Izdatel'stvo Nauka Kasachskoj ASSR, Alma-Ata. 136pp. [in Russian]
- Aubrechtová, M., 2015. A revision of the Ordovician cephalopod *Bactrites sandbergeri* BARRANDE: Systematic position and palaeobiogeography of *Bactroceras*. *Geobios*, 48(3), 193-211. DOI: 10.1016/j.geobios.2015.03.002
- Babcock, L.E., Chang, W.T., 1997. Comparative taphonomy of two nonmineralized arthropods: *Naraoia* (Nektaspida; Early Cambrian, Chengjiang Biota, China) and *Limulus* (Xiphosurida; Holocene, Atlantic Ocean). *Bulletin of the National Museum of Natural Science*, 10, 233-250.
- Babcock, L.E., Merriam, D.F., West, R.R., 2000. *Paleolimulus*, an early limuline (Xiphosurida), from Pennsylvanian-Permian Lagerstätten of Kansas, and taphonomic comparison with modern *Limulus*. *Lethaia*, 33(3), 129-141.
- Barrande, J., 1846. Notice préliminaire sur le Système Silurien et les trilobites de Bohême. C.L. Hirschfeld, libraire. Leipzig, 97pp. DOI: 10.5962/bhl.title.9141
- Barrande, J., 1872. Système Silurien du centre de la Bohême. I, Supplement au vol. 1. Trilobites, crustacés divers et poissons. Prague and Paris, 647pp.
- Bergström, S.M., Chen, X., Gutiérrez-Marco, J.C., Dronov, A., 2008. The new chronostratigraphic classification of the Ordovician System and its relations to major regional series and stages and to $\delta^{13}\text{C}$ chemostratigraphy. *Lethaia*, 42(1), 97-107. DOI: 10.1111/j.1502-3931.2008.00136.x
- Bertling, M., 1992. *Arachnostega* n. ichnog. - burrowing traces in internal moulds of boring bivalves (Late Jurassic, Northern Germany). *Paläontologische Zeitschrift*, 66(1-2), 177-185.
- Bruthansová, J., 2003. The trilobite Family Illaenidae HAWLE ET CORDA, 1847 from the Ordovician of the Prague Basin (Czech Republic). *Transactions of the Royal Society Edinburgh, Earth Sciences*, 93(2), 167-190.
- Budil, P., Fatka, O., 2008. Bohemian and Moravian Trilobites and their relatives. Czech Geological Survey, Prague, 47pp. ISBN: 978-80-7075-711-6.
- Budil, P., Kraft, P., Kraft, J., Fatka, O., 2007. Faunal associations of the Šárka Formation (Middle Ordovician, Darriwilian, Prague Basin, Czech Republic). *Acta Palaeontologica Sinica*, 46(Supplement), 64-70.
- Butler, A.D., Cunningham, J.A., Budd, G.E., Donoghue, P.C.J., 2015. Experimental taphonomy of *Artemia* reveals the role of endogenous microbes in mediating decay and fossilization. *Proceedings of the Royal Society Biological Sciences*, 282(1808), 20150476. DOI: 10.1098/rspb.2015.0476
- Chlupáč, I., 1970. Phyllocarid crustaceans of the Bohemian Ordovician. *Sborník Geologických Věd, Paleontologie*, 12, 41-77.
- Chlupáč, I., 2003. Phyllocarid crustaceans from the Middle Ordovician Šárka Formation at Praha-Vokovice. *Bulletin of Geosciences*, 78(1), 107-111.
- Chvátal, M., 2003. Mineral assemblage of the Červený vrch locality. *Bulletin of Geosciences*, 78(2), 103-106.
- Cooper, G.A., Kindle, C.H. 1936. New brachiopods and trilobites from the Upper Ordovician of Percé, Quebec. *Journal of Paleontology* 10, 348-372.
- Courtesole, R., Pillet, J. 1975. Contribution à l'étude des faunes trilobitiques de l'Ordovicien inférieur de la Montagne Noire. Les Eulominae et les Nileidae. *Annales de la Société Géologique du Nord*, 95, 251-272.
- Dabbard, M.-P., Loi, A., 2012. Environmental control on concretion-forming processes: Examples from Paleozoic terrigenous sediments of the North Gondwana margin, Armorican Massif (Middle Ordovician and Middle Devonian) and SW Sardinia (Late Ordovician). *Sedimentary Geology*, 267-268, 93-103. DOI: 10.1016/j.sedgeo.2012.05.013
- Daley, A.C., Drage, H.B., 2016. The fossil record of ecdysis, and trends in the moulting behaviour of trilobites. *Arthropod Structure & Development*, 45, 71-96. DOI: 10.1016/j.asd.2015.09.004
- Dean, W.T. 1964. The status of the Ordovician trilobite genera *Prionocheilus* and *Polyeres*. *Geological Magazine*, 101(1), 95-96.
- Drost, K., Linnemann, U., Wemmer, K., Budil, P., Kraft, P., Fatka, O., Marek, J., 2003. Provenance, geotectonic setting, and early diagenetic processes of the Šárka Formation at Praha – Červený vrch (Ordovician, Barrandian, Czech Republic). *Bulletin of Geosciences*, 78(1), 147-156.
- Fatka, O., Mergl, M., 2009. The “microcontinent” Perunica: status and story 15 years after conception. In: Bassett, M.G. (ed.). *Early Palaeozoic Peri-Gondwanan Terranes: New Insights from Tectonics and Biogeography*. London, Geological Society, Special Publications, 325, 65-102. DOI: 10.1144/SP325.4
- Fatka, O., Mikuláš, R., Szabad, M., Micka, V., Valent, M., 2011. *Arachnostega* BERTLING, 1992, in Cambrian of the Barrandian area (Czech Republic). *Acta Geologica Polonica*, 61(4), 367-381.
- Fatka, O., Lerosey-Aubril, R., Budil, P., Rak, Š., 2013. Fossilised guts in trilobites from the Upper Ordovician Letná Formation (Prague Basin, Czech Republic). *Bulletin of Geosciences*, 88(1), 95-104. DOI: 10.3140/bull.geosci.1329
- Fatka, O., Budil, P., David, M., Kozák, V., Micka, V., Szabad, M., 2014. Digestive structures in Cambrian and Ordovician trilobites from the Barrandian area (Czech Republic). In: Zhan, R.B., Huang, B. (eds.). *IGCP 591 Field Workshop 2014, Kunming China 12-21 August 2014*. Nanjing University Press, Extended Summary, 49-51.

- Fatka, O., Budil, P., David, M., 2015. Digestive structures in Ordovician trilobites *Colpocoryphe* and *Flexicalymene* from the Barrandian area of Czech Republic. *Estonian Journal of Earth Sciences*, 64(4), 255-266. DOI: 10.3176/earth.2015.32
- Gutiérrez-Marco, J.C., García-Bellido, D.C., Rábano, I., Sá, A.A., 2017. Digestive and appendicular soft-parts, with behavioural implications, in large Ordovician trilobite from the Fezouata Lagerstätte, Morocco. *Scientific Reports*, 7, 39728. DOI:10.1038/srep39728
- Hammann, W., 1983. Calymenacea (Trilobita) aus dem Ordovizium von Spanien; ihre Biostratigraphie, Ökologie und Systematik. *Abhandlungen der Senckenbergischen naturforschenden Gesellschaft*, 542, 1-177. [in German, with English abstract]
- Hammann, W., Henry, J.L., 1978. Quelques espèces de *Calymenella*, *Eohomalonotus* et *Kerfornella* (Trilobita, Ptychopariidae de l'Ordovicien du Massif Armoricaire et de la Péninsule Ibérique). *Senckenbergiana Lethaea*, 59(4/6), 401-429. [in French, with English abstract]
- Havlíček, V., 1981. Development of a linear sedimentary depression exemplified by the Prague Basin (Ordovician–Middle Devonian; Barrandian area – central Bohemia). *Sborník Geologických Věd, Geologie*, 35, 7-48.
- Havlíček, V., 1982. Ordovician of Bohemia: development of the Prague Basin and its benthic communities. *Sborník Geologických Věd, Geologie*, 37, 103-136.
- Havlíček, V., 1998. Ordovician. In: Chlupáč, I., Havlíček, V., Kříž, J., Kukal, Z., Štorch, P. (eds.). *Palaeozoic of the Barrandian*. Czech Geological Survey, Prague, 149-164.
- Havlíček, V., Fatka, O., 1992. Ordovician of the Prague Basin (Barrandian area, Czechoslovakia). In: Webby, B.D., Laurie, J.R. (eds.). *Global perspectives on Ordovician geology*. Balkema, Rotterdam, 461-472.
- Havlíček, V., Vaněk, J., 1966. The Biostratigraphy of the Ordovician of Bohemia. *Sborník Geologických Věd, Paleontologie*, 8, 7-69.
- Havlíček, V., Vaněk, J., 1990. Ordovician invertebrate communities in black-shale lithofacies (Prague Basin, Czechoslovakia). *Věstník Ústředního Ústavu Geologického*, 65(4), 223-236.
- Hawle, I., Corda, A.J.C., 1847. *Prodrom einer Monographie der böhmischen Trilobiten*. *Abhandlungen der königlichen böhmischen Gesellschaft der Wissenschaften*, 5, 119-292.
- Horný, R., 2001. Ordovician Tergomya and isostrophic Gastropoda (Mollusca) of Bohemia: Types and referred specimens in the collections of the National Museum, Prague, Czech Republic. *Acta Musei Nationalis Pragae, Series B – Historia Naturalis*, 57(3-4), 69-102.
- Horný, R., Bastl, F., 1970. Type specimens of fossils in the National Museum of Prague. Volume 1, Trilobita; [Scientific revisors: Chlupáč, I., Marek, L., Příbyl, A., Šnajdr, M.]. Prague, Museum of Natural History, 354pp.
- Hupé, P., 1953. Classification des Trilobites. *Annales de Paléontologie*, 39, 61-168.
- Ingham, J.K., 1977. A monograph of the upper Ordovician trilobites from the Cautley and Dent districts of Westmorland and Yorkshire. *Palaeontographical Society Monograph*, 3, 89-121.
- Jaekel, O., 1901. Ueber die Organisation der Trilobiten. *Zeitschrift der Deutschen Geologischen Gesellschaft*, 53(1), 133-171. [in German]
- Kraft, J., Kraft, P., 1992. Biostratigraphy of the Klabava and Šárka Formations (Bohemia, Lower Ordovician) – a brief overview of new investigations. *Acta Universitatis Carolinae, Geologica*, 33(1-2), 23-29.
- Kraft, J., Kraft, P., 1999. Graptolite biozones of the Bohemian Lower and Middle Ordovician and their historical development. *Journal of the Czech Geological Society*, 44(1-2), 53-62.
- Kraft, P., Kraft, J., 2003. Middle Ordovician graptolite fauna from Praha – Červený vrch (Prague Basin, Czech Republic). *Bulletin of Geosciences*, 78(1), 129-139.
- Kraft, P., Kraft, J., Prokop, R.J., 2001. A possible hydroid from the Lower and Middle Ordovician of Bohemia. *Alcheringa*, 25(2), 143-154.
- Kukal, Z., 1962. Petrografický výzkum šareckých vrstev barrandienského ordoviku [Petrographical investigation of the Ordovician Šárka beds in the Barrandian area]. *Sborník Ústředního Ústavu Geologického, Odd. Geologický*, 27, 175-214. [in Czech, with English abstract]
- Lajblová, K., Kraft, P., 2014. The earliest ostracods from the Ordovician of the Prague Basin, Czech Republic. *Acta Geologica Polonica*, 64(4), 367-392.
- Lefebvre, B., 2007. Early Palaeozoic palaeobiogeography and palaeoecology of stylophoran echinoderms. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 245, 156-199. DOI: 10.1016/j.palaeo.2006.02.021
- Lemke, U., 2016. Katalog verfügbarer Namen der Trilobiten-Taxa. Unpublished, 503pp. [Available at: https://www.researchgate.net/publication/297418418_Katalog_verfuegbarer_Namen_der_Trilobiten-Taxa_Stand_Marz_2016]
- Lerosey-Aubril, R., Hegna, T.A., Olive, S., 2011. Inferring internal anatomy from the trilobite exoskeleton: the relationship between frontal auxiliary impressions and the digestive system. *Lethaia*, 44(2), 166-184. DOI: 10.1111/j.1502-3931.2010.00233.x
- Lerosey-Aubril, R., Hegna, T.A., Kier, C., Bonino, E., Habersetzer, J., Carré, M., 2012. Controls on gut phosphatisation: The Trilobites from the Weeks Formation Lagerstätte (Cambrian; Utah). *PLOS ONE*, 7(3), e32934. DOI:10.1371/journal.pone.0032934
- Manda, Š., 2008. *Trocholites* CONRAD, 1838 (Nautiloidea, Tarphycerida) in the Middle Ordovician of the Prague Basin and its palaeobiogeographical significance. *Bulletin of Geosciences*, 83(3), 327-334. DOI: 10.3140/bull.geosci.2008.03.327
- McCoy, F. 1846. A synopsis of the Silurian fossils of Ireland. University Press, M.H. Gill, Dublin, 72pp.

- Mergl, M., 2002. Linguliformean and craniiformean brachiopods of the Ordovician (Třenice to Dobrotivá formations) of the Barrandian, Bohemia. *Acta Musei Nationalis Pragae, Series B*, 58, 1-82.
- Mergl, M., Kozák, V., 2016. The effaced Ordovician trilobite *Svobodapeltis* and the earliest history of illaenimorph trilobites in the Ordovician of the Prague Basin (Bohemia). *Bulletin of Geosciences*, 91(1), 155-168. DOI: 10.3140/bull.geosci.1516
- Mergl, M., Fatka, O., Budil, P., 2007. Lower and early Middle Ordovician trilobite associations of the Prague Basin (Perunica, Czech Republic). *Acta Palaeontologica Sinica*, 6 (Supplement), 320-327.
- Mergl, M., Fatka, O., Budil, P., 2008. Lower and Middle Ordovician trilobite associations of Perunica: from shoreface endemicity to offshore uniformity (Prague Basin, Czech Republic). In: Rábano, I., Gozalo, R., García-Bellido, D. (eds.). *Advances in trilobite research*. Instituto Geológico y Minero de España, Cuadernos del Museo Geominero, 9, 275-282.
- Mikuláš, R., 1991. Trace fossils from siliceous concretions in the Šárka and Dobrotivá Formations (Ordovician, central Bohemia). *Časopis pro Mineralogii a Geologii*, 36(1), 29-38.
- Mikuláš, R., 1998. Ordovician of the Barrandian area: Reconstruction of the sedimentary basin, its benthic communities and ichnofacies. *Journal of the Czech Geological Society*, 43(3), 143-159.
- Peršín, J., Budil, P., 2009. Nové poznatky ze šáreckého a dobrotivského souvrství (ordovik, stupeň darriwil) v severozápadní a severní části Prahy [New observations on Šárka and Dobrotivá Formation (Ordovician, Darriwilian Stage) in northwestern and northern surroundings of Prague]. *Český Kras*, 35, 25-35. [in Czech, with English abstract]
- Polechová, M., 2013. Bivalves from the Middle Ordovician Šárka Formation (Prague Basin, Czech Republic). *Bulletin of Geosciences*, 88(2), 427-461. DOI: 10.3140/bull.geosci.1426
- Příbyl, A., 1953. Seznam eských trilobitových rodů. [Index of trilobite genera in Bohemia]. *Knihovna Ústředního Ústavu Geologického*, 25, 1-80. [in Czech and English]
- Rábano, I., 1989. Trilobites del Ordovicio Medio del sector meridional de la Zona Centroibérica Española. III. Calymenina y Cheirurina. *Boletín Geológico y Minero*, 100(5), 767-841.
- Rouault, M., 1847. Extrait du Mémoire sur les Trilobites du Département d'Ille-et-Villaine. *Bulletin de la Société Géologique de France*, 4, 309-328.
- Servais, T., Dzik, J., Fatka, O., Heuse, T., Vecoli, M., Verniers, J., 2008. Ordovician. In: McCann, T., (ed.). *Geology of Central Europe*. London, Geological Society, 203-248.
- Siveter, D.J., 1973. Trilobites of the family Calymenidae from the Ordovician, Silurian and Devonian systems of North-West Europe. PhD Thesis, University of Leicester, 380pp.
- Siveter, D.J., 1976. The Middle Ordovician of the Oslo Region, Norway. 27. Trilobites of the family Calymenidae. *Norsk Geologisk Tidsskrift*, 56, 335-396.
- Šnajdr, M., 1956. Trilobiti drabovských a letenských vrstev českého ordoviku. [The trilobites from the Drabov and Letná Beds of the Ordovician of Bohemia]. *Sborník Ústředního Ústavu Geologického*, 22, 477-533. [in Czech, with English abstract]
- Šnajdr, M., 1990. Bohemian trilobites. Prague, Czech Geological Survey, 265pp.
- Steinová, M., 2012. Probable ancestral type of actinodont hinge in the Ordovician bivalve *Pseudocyrtodonta* Pfab, 1934. *Bulletin of Geosciences*, 87(2), 333-346. DOI: 10.3140/bull.geosci.1330
- Strang, K.M., Armstrong, H.A., Harper, D.A.T., 2016. Minerals in the gut: scoping a Cambrian digestive system. *Royal Society of Open Science*, 3, 160420. DOI: 10.1098/rsos.160420
- Valíček, J., Vaněk, J., 2001. New index of the genera, subgenera, and species of Barrandian trilobites. Part A-B (Cambrian and Ordovician). *Palaeontologia Bohemiae*, 7, 1-49.
- Vaněk, J., 1965. New species of the suborder Calymenina Swinneton, 1915 (Trilobita) from the Barrandian area. *Sborník Geologických Věd, Paleontologie*, 6, 21-37.
- Vaněk, J., 1995. New deeper water trilobites in the Ordovician of the Prague Basin, Czech Republic. *Palaeontologia Bohemiae*, 5, 1-12.
- Vaněk, J., Valíček, J., 2002. New index of the genera, subgenera, and species of Barrandian trilobites. Part C-D (Silurian and Devonian). *Palaeontologia Bohemiae*, 8, 1-74.
- Vokáč, V., Hartl, F., David, M., Pavlovi, Doubrava, M., Kozák, V., Grigar, L., 2015. Významné nálezy trilobitů z ordoviku (daping-sandby) pražské pánve (Barrandien, Česká Republika) (Notable findings of trilobites from the Ordovician (Dapingian-Sandbian) of the Prague Basin (Barrandian area, Czech Republic). *Erica*, 22, 141-157. [in Czech, with English abstract]
- Whittington, H.B., 1965. A monograph of the Ordovician trilobites of the Bala area, Merioneth. *Palaeontographical Society Monograph*, 2, 33-62.
- Whittington, H.B., Kelly, S.R.A., 1997. Morphological terms applied to trilobites. In: Moore, R.C., Kaesler, R.L. (eds.). *Treatise on Invertebrate Paleontology, Part O Arthropoda 1 Trilobita, Revised 1*. Kansas, Boulder, Colorado and Lawrence, O313-O329.
- Zhou, Z.Y., Yin, G.Z., Tripp, R.P., 1984. Trilobites from the Ordovician Shihtzupu Formation, Zunyi, Guizhou Province, China. *Earth and Environmental Science Transactions of the Royal Society of Edinburgh*, 75(1), 13-36. DOI: 10.1017/S0263593300009755
- Zhu, X.J., Lerosey-Aubril, R., Esteve, J., 2014. Gut content fossilization and evidence for detritus feeding habits in an enrolled trilobite from the Cambrian of China. *Lethaia*, 47(1), 66-76. DOI: 10.1111/let.12041

**Manuscript received March 2017;
revision accepted June 2017;
published Online October 2017.**