



Journal of Negative and No Positive Results

ISSN: 2529-850X

Research and Science S.L.

Baltasar, Aniceto; Bou, Rafael; Bengochea, Marcelo; Perez, Nieves
Beginnings of bariatric and metabolic surgery in Spain
Journal of Negative and No Positive Results, vol. 5, no. 4, 2020, pp. 448-467
Research and Science S.L.

DOI: <https://doi.org/10.19230/jonnpr.3324>

Available in: <https://www.redalyc.org/articulo.oa?id=564564530007>

- How to cite
- Complete issue
- More information about this article
- Journal's webpage in redalyc.org

redalyc.org

Scientific Information System Redalyc

Network of Scientific Journals from Latin America and the Caribbean, Spain and Portugal

Project academic non-profit, developed under the open access initiative



HISTORICAL CORNER (English version)

Beginnings of bariatric and metabolic surgery in Spain

Inicios de la cirugía bariátrica y metabólica en España

Aniceto Baltasar, Rafael Bou, Marcelo Bengochea, Nieves Pérez

Hospital Comarcal de Alcoy. España

* Corresponding Author.
e-mail: abaltasar@coma.es (Aniceto Baltasar).

Received 19 October 2019; accepted 30 October 2019.

How to cite this paper:

Baltasar A, Bou R, Bengochea M, Pérez N. Beginnings of bariatric and metabolic surgery in Spain. JONNPR. 2020;5(4):448-67. DOI: 10.19230/jonnpr.3324

Cómo citar este artículo:

Baltasar A, Bou R, Bengochea M, Pérez N. Inicios de la cirugía bariátrica y metabólica en España. JONNPR. 2020;5(4):448-67. DOI: 10.19230/jonnpr.3324



This work is licensed under a Creative Commons
Attribution-NonCommercial-ShareAlike 4.0 International License
La revista no cobra tasas por el envío de trabajos,
ni tampoco cuotas por la publicación de sus artículos.

Abstract

Bariatric surgery (BS) from the Greek *bari* = weight and *iatrein* = cure) treats obesity and began in Spain in 1973. Its greatest development occurs after the founding of SECO (Spanish Society of Obesity Surgery) in 1997. The purpose of this work is to reflect the changes that have occurred in these 22 years.

Keywords

Morbid obesity; Bariatric surgery; SECO

Resumen

La cirugía bariátrica (**CB**) de *bari* = peso y *iatrein* = cura) trata la obesidad y comienza en España en 1973. Su mayor desarrollo ocurre tras la fundación de SECO (Sociedad Española de Cirugía de la obesidad) en 1997. La finalidad de este trabajo es reflejar los cambios que han ocurrido en éstos 42 años.



Palabras clave

Obesidad mórbida; Cirugía bariátrica; SECO

Nomenclature

Add L for laparoscopic

English

Vertical Banded Gastroplasty
Gastric Banding
Roux-en-Y gastric bypass
One Anastomosis GBP /mini GBP
Bilio-pancreatic Diversion
Intestinal diversion
Duodenal Switch
Single Anastomosis Duodenal-Ileostomy
Sleeve-Forming Gastrectomy
Adolescent Bariatric Surgery
Revision/Conversion of prior surgery
Morbid Obesity
Weight loss
% Excess Weight Loss
% Expected BMI Loss

Abbreviated

VBG
GB
RY-GBP
OAGB
BPD
ID
DS
SADI
SFG
ABS
REV
MO
WL
%EWL
%ExBMIL

Obesity is a multifactorial epidemic ailment of environmental origin, affecting subjects from all countries, and whose origins are not in the stomach or intestine. It represents a unique case of surgery to operate healthy organs, which are not the cause of the disease and do not improve after the operation.

Obesity is a multifactorial epidemic ailment of environmental origin, affecting subjects from all countries, and whose origins are not in the stomach or intestine. It represents a unique case of surgery to operate healthy organs, which are not the cause of the disease and do not improve after the operation. Payne⁽³⁾ and Scott⁽⁴⁾ developed these ID techniques in the 1960s leaving only 14-4 inches (35-10 cm) as an absorptive zone and those were abandoned in the 1970s because of their serious metabolic (malnutrition) and hepatic (liver failure) complications.

Buchwald⁽⁵⁾ initiated the ID of the last third of the intestine for hypercholesterinemia (POSCH) and showed its protective role at 25 years in the development of atherosclerosis. Now it has also been abandoned, not because of lack of effectiveness, but because of the development of nystatin in the medical control of cholesterol. Dr. Henry Buchwald remains active 67 years later, and in 2012 Barcelona was appointed as *Honorary Member of the Spanish Society of Obesity Surgery (SECO)* and he will participate in Madrid-IFSO 2019. Baltasar⁽⁶⁾ published in 1991 the only three ID in Spain for hypercholesterinemia.



First Spanish experience

Prof. Sebastián García Díaz of Seville carried out the 1st Scott-type Jejunum-ileal diversion (JID) in the Virgen Macarena Hospital on 11.19.1973. He began bariatric surgery in Spain with 12 cases⁽⁸⁻⁹⁾ and then published 20 more, the 1st work in English by a Spanish author⁽⁷⁾ in the World Journal of Surgery in 1981. For this 2nd work he received the award by the Seville *Hospital of the Five Sores* in 1979 (Figure 2). His work went unnoticed for 40 years until we rescued them in 2013⁽¹¹⁾.




Figure 1-1. Genealogical tree of bariatric surgery.

Figure 1. Bariatric techniques



There are two Spanish JID done by Drs. Sánchez/Masdevall in Barcelona Bellvitge Hospital on March 1976 and another by Dr. Martínez, Zaragoza Clinic Hospital on July 12, 1978.

Buchwald⁽¹²⁾ describe (Figure 1) the tree of the development of this surgery in the 1970s with a big malabsorptive and restrictive operation. The evolution of malabsorptive surgery is on the left, and restrictive surgery and gastric bypass (**GBP**) were reported.

In 1964 Mason⁽¹³⁾ initiates the GBP. It was a total change in strategy. Capella⁽¹⁴⁾, Álvarez-Cordero⁽¹⁵⁾ (both first SECO Honorary Members) and Fobi⁽¹⁶⁾ made very important contributions to this technique and many other authors in the 1970s.

Baltasar AB⁽¹⁷⁾ performed in *June 17, 1977* the first Mason-type GBP in Spain. Vara⁽¹⁸⁾, De la Cruz⁽¹⁹⁾ and Sitges⁽²⁰⁾ wrote in Spain about ID in hypercholesterinemia and GBP in obesity.

Scopinaro⁽²¹⁾, a *tireless researcher and clinician*, initiated experimental and clinical gastrectomy with biliopancreatic diversion (**BPD**) in 1976^(22,23). He is the leader and "*father of European bariatric surgery*" and participated in multiple congresses and publications. His combined mixed a technique of gastrectomy plus BPD as the most effective for treating obesity⁽²⁴⁾. He is an Honorary Member of SECO and the only foreign Outstanding Achievement Award winner (OAAW) of the American Association of Metabolic and Bariatric Surgery (**ASMBS**).

DBP became a very popular and was a commonly used technique in Spain. Larrad and Sánchez⁽²⁵⁻²⁸⁾ published extensively on a modification of their own and other authors used this technique in Zaragoza⁽²⁹⁻³²⁾, Santander^(33,34) and Barcelona.

Mason⁽³⁵⁾ "*father of BS*" published in 1982, 18 patients on vertical banded gastroplasty (**VBG**) and it was the 1st great revolution by making OS "easy and affordable".



AB (36) in 1984 broadcasts in RTVE the *1st Spanish documentary* in MO entitled "*JJ Soriano more moral than the Alcoy man*". with IMC-52 and that blocked the telephone lines of the hospital. Laporte⁽³⁷⁾ published the *1st Spanish VBG experience* in 9 cases.

Reopening of the vertical staple-line was a serious disadvantage of the VBG because it cancels out the restrictive effect of the operation. Baltasar⁽³⁸⁾ in 1989 described the separation with staples between gastric tube and major curvature and did not have a single recanalization in 100 cases. McLean⁽³⁹⁾ used the same technique three years later. Alcoy's Andreo⁽⁴⁰⁾ describes the typical radiological "*peanut deformity*" of the VBG.

Many Spanish surgeons performed VBG^(41,43). Baltasar⁽⁴⁴⁾ published his *first* 100 cases in 1990 with excellent results, *but* 5 years later he reviewed the same patients and describes it as a "*frustrated hope*"^(45,46). And two years later this technique was abandoned.

Belachew^(47,48) in Belgium initiated on 1/09/1993 laparoscopy the 2nd revolution in bariatrics with the *1st laparoscopic adjustable gastric band (LAGB)* operation.




Figura 2. H. Buchwald S. García Díaz E. Mason N. Scopinaro

Favretti⁽⁴⁹⁾ performed on Sept 27.1995 the *1st LAGB* operation in Spain at Madrid "La Paz" Hospital (Figure 3) assisted by Masdevall / Baltasar




Figure 3. Favretti 1ª BGLA española. Hospital "La Paz" Madrid. 1995



Carbajo⁽⁵⁰⁾ in 1986 made the 1st 12 BGL in Spain and Alastrué⁽⁵¹⁾ compares VBG with BGLA. Thousands of BGLA were done all over the world and then it was abandoned. *Laparoscopy* changed the way of doing surgery not only in BS but also in all general surgery. Advances in bariatric laparoscopy, being repetitive operations and performed on healthy organs were the greatest advance in XX century surgery.

In the 90's the 1st *national bariatric societies* were created, starting with the American Society of Bariatric Surgery [ASBS] by Edward Mason on June 3, 1983 in Iowa City, IO⁽⁵²⁾, the town he worked all his life. Deitel created *Obesity Surgery*, as the 1st *journal of obesity* in 1990⁽⁵²⁾.

In 1995 the International Federation of Surgery for Obesity (IFSO) is founded and standards⁽⁵³⁾ devised for reporting results.




Figure 4. M. Deitel M. Belachew A. Wittgrove W. Clark

The Spanish Society of Obesity Surgery (**SECO**) was founded at Residencia Pérez Mateos, San Juan, Alicante by 26 surgeons on December 12, 1976 (Figure 5).




Figure 5. SECO 1976 SECO 2015 Baltasar 2011 AMSBS-OAAW

SECO joined IFSO in 1998 and AB became *IFSO-2002 president*. He organizes and preside IFSO-Salamanca-2003. Dr. Antonio Torres, 2nd *Spanish president* of IFSO organizes IFSO-2019-Madrid, the world's largest bariatric meeting. Spain will be the 2nd *country* that organizes this congress twice (Figure 6 and Figure 7).



SECO founding members

December 12. 1976

San Juan. Alicante, España

1. Aniceto Baltasar Alcoy
2. Juan Pujol Barcelona
3. Miguel A. Carbajo Valladolid
4. Santiago Tamames Madrid
5. Carlos Escalante Santander
6. Horacio Urquijo Madrid
7. Cándido Martínez Vitoria
8. Francisco Arlandis Alcoy
9. Rafael Bou Alcoy
10. Miguel A Calvo Bilbao
11. Antonio Alastrué Badalona
12. Eugenio Urquijo Madrid
13. Carlos Cerquella Madrid
14. Felipe de la Cruz Madrid
15. Mario García Madrid
16. Luis García Vallejo Santiago
17. Federico Leruite Granada
18. Juan Machuca La Coruña
19. José M^a Recio Barcelona
20. Carlos Masdevall Barcelona
21. Salvador Serrano Burgos
22. Tomeu Feliú Gerona
23. Antonio Soro Mallorca
24. Antonio Martín Madrid
25. Mariano Martínez Zaragoza

Figure 6. SECO Founding members




1. Dec 1997 Aniceto Baltasar Torrejón
2. May 2001 Cándido Martínez Blázquez
3. Oct 2004 José C. Fernández-Escalante
4. April 2007 Antonio José Torres García
5. April 2011 Carles Masdevall Noguera
6. April 2013 Felipe de la Cruz Vigo
7. May 2015 Juan C. Ruiz Adana Belbel
8. March 2017 José A. Ramírez Felipe
9. Sept 2019 Raquel Sánchez-Santos

Figure 7. SECO Presidents

Wittgrove and Clark in *October 27, 2003*⁽⁵⁵⁻⁵⁷⁾ made the *most significant step in performing the 1st laparoscopic gastric bypass (LGBP)* in San Diego. This is the *3rd revolution of the BS*. AB was the *1st visitor* in September 2007 in San Diego, and to *our proposal* Wittgrove⁽⁵⁸⁾ dropped the huge circular #33 port and use the *circular stapler without trocar*, a very important step in those early times.

We performed the *1st LGBP* in Spain on *1.14.1997*⁽⁵⁹⁾ and presented it⁽⁶⁰⁾ at Bruges IFSO-1998 the *1st European to report it* on video. Serra⁽⁶¹⁾ published in 1999 the *1st world hernia after LGBP*. Higa⁽⁶²⁾ made fortunately the *1st LGBPs* with manual sutures.

Baltasar⁽⁶³⁾ published the *1st book* in Spanish in OS in 2000. Martínez⁽⁶⁴⁾ from Vitoria published a bilingual BS book (Spanish and Basque) on 2001. De la Cruz⁽⁶⁵⁾ published in 2006 the *1st book* on LGBP in Spain and carried out the *1st surgical session* in León. García-Caballero published an English book on diabetes surgery (Figure 8).



Baltasar-2000 C. Martínez F. de la Cruz Diabetes de Caballero
Figure 8. Books published by SECO members

In 1988, Hess⁽⁶⁶⁾ and Marceau⁽⁶⁷⁾ initiated the duodenal switch (**DS**), a *Sleeve-forming Vertical Gastrectomy (SFVG)* plus BPD and Baltasar⁽⁶⁸⁾ started it on *3.17.1994*.

Ren/Gagner⁽⁶⁹⁾ performed the *1st world LDS* in October 1999 and Baltasar on *5.10.2000*⁽⁷⁰⁾ the *1st LDS in Europe*. (Figure 9). This difficult and controversial technique by *the Switchers* surgeons, is rarely used today, in less than 1%.

Baltasar⁽⁷¹⁾ on 950 patients had a low mortality of 0.4% with LDS and is *the most effective technique to lose weight*.




Figure 9. Switchers logo DS AB/ Hess/ Marceau M. Gagner

There have been many technical variations in laparoscopy. In general, all viscera division is done with auto sutures. The anastomosis is done either with auto sutures or manual ones. We advocate the manual suture starting always with the *sliding, self-locking sliding knot* Resa and Solano⁽⁸²⁾.

SECO members reported BS surgeries in the early years of OS and SOARD. Spain was the 2nd country with the highest number of publications after the USA in 2005 and 2006 and in 2013 was the 5th country with the highest number of publications in SOARD. (Figure 10 and Figure 11)

Papers in OBESITY SURGERY Jan.-Sept. 2005			
Argentina	1	Korea	1
Australia	7	Mexico	1
Austria	5	New Zealand	2
Belgium	6	Netherlands	3
Brazil	20	Poland	4
Canada	6	Saudi Arabia	3
Chile	4	Spain	33
France	20	Sweden	11
Germany	3	Switzerland	6
Greece	8	Taiwan	1
India	1	Turkey	2
Israel	6	United Kingdom	4
Italy	20	USA	84
Total		280	

Table 1. Papers in OBESITY SURGERY by Country, January – December 2006					
Argentina	3	Germany	10	Russia	1
Australia	11	Greece	12	Saudi Arabia	2
Austria	13	Israel	10	Singapore	2
Belgium	12	Italy	25	Spain	33
Brazil	27	Korea	3	Sweden	5
Canada	8	Kuwait	1	Switzerland	12
Chile	5	Lebanon	2	Taiwan	7
China	3	Mexico	2	Turkey	3
Cyprus	1	New Zealand	4	United Kingdom	7
Czech Republic	2	Netherlands	9	USA	106
Finland	1	Poland	5		
France	24	Portugal	3	Total	374

© FD-Communications Inc.

Obesity Surgery, 16, 2006 1553

Figure 10. Publicaciones españolas de SECO en Obes. Surg. 2005-6

SURGERY FOR OBESITY AND RELATED DISEASES OFFICIAL JOURNAL OF THE AMERICAN SOCIETY FOR METABOLIC AND BARIATRIC SURGERY	
United States	590
United Kingdom	23
Canada	23
Italy	19
Spain	13

Figure 11. Spanish publications in SOARD 2014

SECO members have been very active publishing in OS and SOARD (Figure 8 and Figure 9). Among the SECO founding members there has been a high bibliographic volume



(Alcoy-124, Carbajo-76, Belvitge-38, Alastrué-30, Martín-Duce-27, Martínez C.-22, Ballesta-9, Zaragoza-5). In 2003 SECO made the Salamanca Declaration on BS⁽⁷⁴⁾.

Laparoscopic Sleeve-forming gastrectomy (LSFG), the restriction part of DS, was so described by several authors in 2005. Baltasar LSFG⁽⁷⁵⁾, according to Ahmad⁽⁷⁶⁾ is the 61st most cited article in all the bariatric literature. Angrisani⁽⁷⁷⁾ claims that the LSFG is today the most commonly performed operation in the world.

We start the gastrectomy at the pylorus and suture the gastric anterior and posterior serosa, covering the staples, to prevent rotation of the sleeve and avoid leaks.

Rutledge described 1,274 cases of *mini-gastric bypass* in 2001⁽⁷⁸⁾ and Spanish authors have developed two popular techniques. Carbajo⁽⁷⁹⁾ performs since 2004 the *lesser curvature reservoir without gastric resection*, the *one anastomosis* OAGB, a GBP with latero-lateral diversion to an intestinal loop. He presented more than 3,500 cases at the 1919s World Congress in Valladolid and is today the fastest growing technique in the world. (Figure 12)



Figure 12. OAGP R. Rutledge OAGBP MA. Carbajo

Sánchez and Torres⁽⁸⁰⁾ at Madrid Clinic Hospital, describe in 2005 the *one anastomosis* DS or SADI in English. There is gastric resection in the form of SFG and the BPD is done end-to-side at the duodenum (D1). The operation is becoming very popular all over the world. Currently they have more than 350 cases. (Figure 13)




Figure 13. A. Sánchez SADI A. Torres

BS at Barcelona Bellvitge hospital has organized 16 consecutive bariatric courses and Torres/Sánchez another 16 annual courses at Madrid Clinic Hospital.

We founded *BMI-Latina Journal* (Iberoamerican Bariatric and Metabolic) www.bmilatina.com in 2011 as an online magazine published in Spanish, English and Portuguese. Adopted by SECO in 2015, it should play a valuable role in Spanish and



Portuguese speaking BS societies *but* since 2015 only the abstract is published in English, and that may restrict its dissemination.

In 2009, Scopinaro, Melissas, Fried and AB create the IFSO European Chapter of the Centers of Excellence Program (ECEP). Currently, several Spanish centers and surgeons use this prestigious quality control program.

SECO members have participated in numerous local, regional, national, and international meetings and congresses, especially in Latin American societies where SECO has always been very kindly well received. In May 1998 we organized a table at the meeting of the prestigious course of Prof. Moreno González, with Drs. Cowan, Fobi, Scopinaro, Clark & F.de la Cruz.

In 2003, AB inaugurated the Spring (NYC) and Fall (Chicago) two keynote lectures at the American Congress of Surgery with on DS.

AB received the *IFSO-Lifetime Membership Award* in June 2011 and was a *finalist* for the prestigious *ASMBS-Outstanding Achievement Award* at the ASMBS meeting in Orlando, Florida. (Figure 4)

Diabetes Surgery

Part of the BS is dedicated to diabetes as **metabolic surgery**. Baltasar⁽⁸⁰⁾ publishes a successful *1st intervention* in 2004 *BPD-without-SFG*.

Resa and Solano⁽⁸²⁾ describe 65 cases of gastro-ileal bypass as the simplest, quickest and safest technique we know for treating obesity. And then Resa⁽⁸³⁾ again publishes 1512 more cases. Alhambra⁽⁸⁴⁾, Vidal⁽⁸⁵⁾, Vilarrasa⁽⁸⁶⁾, García⁽⁸⁷⁾, Cruz⁽⁸⁸⁾ and Torres⁽⁸⁹⁾ have published on the same subject.

Laparoscopic OS in Children and Adolescents (ABS)

OS is increasingly used in children. Baltasar⁽⁹⁰⁻⁹¹⁾ published the *1st national SFG* in 2004 on a 10-year-old boy with excellent results 10 years later.

Carbajo⁽⁹²⁾ has a case with 5 years follow-up and then in 2019 again⁽⁹³⁾ reports the more extensive experience in CLONA with 39 patients treated with OAGBP with excellent results.



Vilallonga⁽⁹⁴⁾ reports that the overweight rate in 4–24 years-old children has increased by approximately 10% in the last 20 years. It is estimated that today, 20% of boys and teenagers and 15% of girls are overweight.

Robotics bariatric surgery

Cadiere and Favretti performed the *world's* 1st robotic bariatric operation at a distance in 1998. Díez and Blázquez perform the *first 12 robotic* Spanish bariatric surgeries in Vitoria-2013. Vilallonga (*1st accredited robotics surgeon in Spain*) and Fort from 2010 in the Hospital Vall d'Hebron in Barcelona develop robotic surgery⁽⁹⁴⁾ and performed more than 540 cases with the da Vinci (Fig.13) Surgical System® (Intuitive Surgical, Sunnyvale, CA) at the beginning with GV and then DG robotics⁽⁹⁵⁾.

Morales⁽⁹⁶⁾ performs in Seville the complex single port surgery of since 2012 and today leads the European surgery⁽⁹⁷⁾. (Figure 14)



Figure 14. Da Vinci R. Vilallonga S. Morales A. Lacy

A. Lacy initiated AIS-Channel as a pioneering worldwide on-line TV transmission and made the *1st BS remote operation by cellphone G5* from Barcelona Clinic Hospital on 4.14.2019 at the WORLD-VIDEO Forum Barcelona-2019.

Surgeries performed in Spain 2018

1st-Surgeries: 5.952; **2nd-Revision:** 343:

Total Complications: LGBP-3.7% and LSFG-3.6%; LGBP-Leaks-1.2%, Bleeding-2.3%, PET-0.1% **Re-op**-2.3% Exitus-0.1% and in **SFG**-Leaks-1.5%, Bleeding-1.9% PET-0.1% 2.1% **Deaths**-0.1%. Extraordinary good results!

Private centers report multiple operations over the years with different techniques: Valladolid-4.255, Teknon-3.000, De la Cruz-2.493, Zaragoza-2.649, Alcoy-1.729.

We should publish all serious complications such as malnutrition⁽⁹⁷⁻⁹⁸⁾, leaks^(98,100), total gastrectomy's⁽¹⁰¹⁾, liver failure⁽¹⁰²⁻¹⁰⁵⁾, stapling of the bougie⁽¹⁰⁶⁾, emergency tracheostomy⁽¹⁰⁷⁾



and mistakes as the Journal of negatives or non-positive results (JONNPR)⁽¹⁰⁸⁾, since it is more likely to learn from negative results than positive ones.

BS is performed in public centers in all regions of Spain with very low leakage and complication rates. But... *it is not yet performed on an outpatient basis.*

If OM is an epidemic pathology and if CB is the best solution, it should be accessible to more subjects. Duncan⁽¹⁰⁹⁾ performs ambulatory BS, and this will be the 4th great bariatric revolution. His A-team uses 2 operating rooms, 2 anesthesiologist nurses, 2 scrub nurses and an operating room technician as an assistant. Operative times of 22'. The surgeon passes to OR-2 with the patient already anesthetized. He changes gloves and do 5 patients in total in the morning. In the afternoon, surgeon-B operates while team-A have consultations. Total 10 patients per day. 50 cases per week, 2,2000 cases per year. No overnight stay. All morning patients are discharged before 14 hours. There is no hospitalization.

Referencias

1. Henrikson V.: Kantunn farmsresektion forsvaras som terapi mot fett sot? Nordisk Medicin 1952; 47: 744
2. Kremen AJ, Linner JH, Nelson CH. An experimental evaluation of the nutritional importance of proximal and Distal small intestine. Ann Surg 1954; 140:439–48.
3. Payne JH, de Wind L, Schab CE, Kern WH. Surgical treatment of morbid obesity. Sixteen years of experience. Arch Surg. 1973; 106:431–7.
4. Scott HW, Law DH, Sandstead H. et al: Jejunoileal shunt in surgical treatment of morbid obesity. Ann Surg.1971; 171:770–81.
5. Buchwald H, Rucker R.: The History of Metabolic Surgery for Morbid Obesity and a Commentary. World J. Surg. 1981; 5, 781-787
6. Buchwald H, Williams SE, Watts SP et al.: Overall mortality in the program of the surgical control of the hyperlipidemias (POSCH). J Am Coll Surg. 2002; 195:327–316.
7. Baltasar A, Marcote E, Bou R, Bengochea M, Arlandis F.: Exclusión ileal parcial en la hiper colesterinemia. A propósito de tres operaciones. Cir. Esp 1991; 6:4475–8
8. García Díaz S. Indicaciones del tratamiento quirúrgico en la obesidad mórbida. Rev. Quir. Española 1975; 2:165-72
9. García Díaz S., Pera C, Loscertales J et al. Trabajos originales. Estado actual del tratamiento de cirugía de la obesidad. Rev Esp Enf del Ap. Digestivo 1977; 51:1-25



10. García Díaz S., Pera C, Loscertales J et al. Avances en cirugía. Barcelona. Editorial Salvat 1979:209-24
11. García Díaz S, García G.: Medical and surgical indications for treatment of morbid obesity. World Journal of Surgery 1981. 5:795-9
12. Baltasar A. Domínguez-Adame E.: Inicios de la cirugía bariátrica y metabólica en España. Cir. Esp 2013.; 91:407-416
13. Mason EE, Ito C.: Gastric bypass in obesity. Surg Clin N Amer 1967; 47:1345.
14. Capella R, Capella J, Mandac H.: Vertical Banded Gastroplasty-Gastric Bypass: preliminary report. Obesity Surgery 1991; 1, 389-395
15. Álvarez-Cordero R, Aragon V. Post-operative Complications in a Series of Gastric Bypass Patients. Obes Surg 1992; 2, 87-89
16. Fobi MAL. Why silastic ring vertical banded gastric bypass for treatment of obesity? Obes Surg 1991; 1: 423-6.
17. Baltasar A, Del Río J, Bengochea A.: Bypass gástrico en la obesidad mórbida. Revista clínica española 1980; 157: 395-9
18. Vara López R.: Tratamiento quirúrgico de la obesidad. Real Academia de Medicina. Anales XCI. 1974
19. De la Cruz Caro F.: Mesa redonda sobre obesidad. Mayo 1975
20. Sitges Creus A, Laporte E: Tratamiento quirúrgico de la obesidad masiva. Práctica quirúrgica 1982.
21. Scopinaro N, Gianetta E, Civalleri D, et al. Biliopancreatic bypass for obesity: II. Initial experience in man. Br J Surg 1979; 66: 619-20.
22. Scopinaro N, Gianetta E, Civalleri D, et al. Two years of clinical experience with biliopancreatic bypass for obesity. Am j Clin Nutr 1980; 33: 50614.
23. Scopinaro N, Gianetta E, Civalleri D, et al. Partial and total biliopancreatic bypass in the surgical treatment of obesity. Inf J Obesity 1981; 5: 421-9.
24. Scopinaro N: Why the Operation I Prefer is Biliopancreatic Diversion (BPD). Obes Surg 1991; 1:307-309
25. Sánchez C, Larrad N.: Analysis of Weight Loss with the Biliopancreatic Diversion of Larrad: Absolute Failures or Relative Successes? Obes Surg 2002; 12, 249-252
26. Larrad A, Sánchez C.: Indicadores de calidad en cirugía bariátrica y criterios de éxito a largo plazo. Quality indicators in bariatric surgery and criteria for long-term success Cir Esp 2004; 75: 5, 2 301-304



27. Larrad A, Sánchez C, Cuadros P et al. Course of Metabolic Syndrome following the Biliopancreatic Diversion of Larrad. *Obes Surg*, 2007, 17: 202-210
28. Larrad A, Sánchez C, Cuadros P: Short-, Mid- and Long-Term Results of Larrad Biliopancreatic Diversion. *Obes Surg*, 2007; 17, 202-210
29. Elía M, Ariba's D, Gracia J, Artigas C. Results of bilio-pancreatic diversion after five years. *Obes Surg* 2004; 14:766-772.
30. Gracia J, Martínez M, Aguilera V et al. Postoperative morbidity of biliopancreatic diversion depending on common limb length. *Obes Surg* 2007; 117:1306-1311
31. Aguilera V, Royo P, Jiménez A. et al. Obesity surgery results depending on technique performed: long-term outcome. *Obes Surg* 2009; 19:4323-438
32. Gracia JA, Elía M, Aguilera V et al: Metabolic syndrome after bariatric surgery. Results depending on the technique performed. *Obes Surg* 2011; 21:179-185
33. Domínguez A, Olmedo F, Ingelmo A, Gómez M, Escalante C. Bypass bilio-pancreático. *Cir Esp* 2004; 75:251-6. 10.1016/S0009-739X(04)72313-1
34. Palomar R, Fernández G, Domínguez A. Effects of Weight Loss after Biliopancreatic Diversion on Metabolism and Cardiovascular Profile *Obes Surg* 2005, 15: 794–798
35. Mason E. Vertical banded gastroplasty for obesity. *Arch. Surg* 1982; 117 :191-4
36. Laporte E, Badosa F, Masdevall C.: La gastro plastia para el tratamiento de la obesidad. *Cir. Esp.* 1985; 38:621–5.
37. Baltasar A.: "JJ Soriano más moral que el Alcoyano. Vivir cada día. RTVE 1984. https://www.youtube.com/watch?v=pQtW3H9B_kg
38. Baltasar A.: Modified vertical gastroplasty. Technique with vertical division and serosal patch. *Acta Chir. Scand* 1989; 155:107.112
39. McLean LD, Rhode BM, Sampalis J, et al. Results of the surgical treatment of obesity. *Am J Surg* 1993; 165:155-62.
40. Andreo L, Lasieria R, Baltasar A, et al. Evaluación radiológica de la gastroplastia vertical anillada. *Radiología.* 1988; 30:303–8.
41. Alastrué A, Rull M, Casas D et al: Gastroplastia vertical anillada. Experiencia de un grupo multidisciplinario en 65 pacientes. Metodología y técnica quirúrgica. Resultados antropométricos. *Cir Esp* 1991; 50 50-58
42. Alastrué A, Formiguera J, Rull M, Casas D, Escudero LE, Del Moral P et al. Cirugía bariátrica: gastroplastia vertical anillada. Obesidad mórbida y super obesidad. Resultados antropométricos. Complicaciones técnicas y reintervenciones a largo plazo. *Rev Endocrinol* 1992; 38:365-375



43. Arribas D, Martínez M, Elía M et al.: Vertical banded gastroplasty: Is it a durable operation for morbid obesity? *Obes Surg* 2004; 14:536-538
44. Baltasar A, Tomás, Marcote et al. Cirugía bariátrica. Experiencia con 100 pacientes operados. *Cir Esp* 1991; 50: 271.5
45. Baltasar A, Bou R, Del Río J, et al. Cirugía bariátrica: resultados a largo plazo de la gastroplastia vertical anillada. ¿Una esperanza frustrada? *Cir Esp* 1997; 62:175-9. 21.
46. Baltasar A, Bou R, Arlandis F, et al. Vertical banded gastroplasty at more than five years. *Obes Surg* 1998; 8:29-34.
47. Belachew M, Lagrard TH, Defrebreuren M.: Laparoscopic adjustable silicone gastric banding in the treatment of morbid obesity. *Endoscopy* 1994; 8:1354-6
48. Belachew M; Legrand M; Vincent V. Laparoscopic Placement of Adjustable Silicone Gastric Band in the Treatment of Morbid Obesity: How to Do It. *Modern Surgery: Technical Innovations. Obesity Surg* 1995; 5: 66-70
49. F. Favretti F; Cadiere GB; Segato G; Bruyns G; Laparoscopic Adjustable Silicone Gastric Banding: Technique and Results *Obes Surg*, 1995; 5:364-371
50. Carbajo MA, Martín JC, Blanco J.: Banda ajustable por laparoscopia (Lapband) en el tratamiento de la obesidad grave. Experiencia preliminar *Cir Esp* 1998; 63:185-8
51. Alastrué A, M Rull, D Casas, et al. Evaluación de dos técnicas restrictivas en cirugía bariátrica. Gastroplastia vertical anillada (GVA) vs bandeleta hinchable (Lapband). Valoración a los dos años. *Cir Esp* 1996; 60: 181-187,
52. Scott HW.: A tribute to Edward E Mason. The 1st Annual Edward E. Mason Founders lecture. *Obes Surg* 1991; 1: 13-19
53. Blommers T. The ASBS first nine years. *Obes Surg* 1992; 2: 115-7
54. Mason E, Amaral J, Cowan G, Deitel E.: Standards for reporting results. *Obes Surg* 1994; 4:56-65
55. Wittgrove A, Clark W, Tremblay L. Laparoscopic gastric bypass. Roux-en-Y. Preliminary report. *Obes Surg* 2004; 4:353-7.
56. Wittgrove A, Clark W. Laparoscopic gastric bypass. Roux-en Y. Experience with 27 cases with 3-18 months follow up. *Obes Surg*. 2006; 6:54-7.
57. Wittgrove A, Clark W, Schubert K. Laparoscopic gastric bypass. Roux-en Y. Technique and results in 75 patients with 3-30 months follow up. *Obes Surg*. 2006; 6:500-4.
58. Wittgrove A. Chapter 21: Laparoscopic gastric bypass. Circular stapler technique. In: *Obesity Surgery: Principles and Practice*. Pitombo C, Jones KB, Higa KD, Pareja JC). Albany, New York: McGraw Hill Medical 2007, 221-224.



59. Baltasar A, Bou R, Miró J, Serra C, et al. Bypass gástrico por laparoscopia. Cir Esp. 2000; 67:556–9.
60. Baltasar A, Bou R, Bengochea M et al: Lap Roux-Y gastric bypass. V-18. Obes Surg. 1998; 8:393.
61. Serra C, Baltasar A. Internal hernias and gastric perforation after laparoscopic gastric bypass. Obes Surg. 1999; 9:546
62. Higa K, Boone K, Ho T.: Complications of the Laparoscopic Roux-en-Y Gastric Bypass: 1,040 Patients - What Have We Learned? Obes Surg, 2000; 10:509-513
63. Baltasar A.: Obesidad y cirugía. Cómo dejar de ser obeso. Libro de Editorial Arán 2000.
64. Martínez C. Martínez G. Bypass gástrico laparoscópico. Vitoria. 2001
65. De la Cruz F, la Cruz JL Bypass Gástrico por Laparoscopia. Laparoscopic Gastric Bypass. Obes Surg, 2006;16, 112
66. Hess DS, Hess DW. Biliopancreatic diversion with a duodenal switch. Obes Surg. 1988; 8:267–82.
67. Marceau P, Biron S, MD, Bourque A.: Biliopancreatic Diversion with a New Type of Gastrectomy Obes Surg 1993; 3: 29-35
68. Baltasar A, Bou R, Marcote E, et al. Hybrid bariatric surgery: Biliopancreatic diversion and duodenal switch. Obes Surg. 1995; 5:419–23.
69. Ren CJ, Patterson E, Gagner M. Early results of laparoscopic biliopancreatic diversion with duodenal switch: a case series of 40 consecutive patients. Obes Surg 2000; 10:514-523.
70. Baltasar A, Bou R, Miró J. et al.: Avances en técnica quirúrgica. Cruce duodenal por laparoscopia en el tratamiento de la obesidad mórbida: técnica y estudio preliminar. Cir. Esp. 2001; 70:102-104. www.youtube.com/watch?v=GSfzgYYxZJ8.
71. Baltasar A, Pérez N, Bou R et al. A switch to the Duodenal Switch. Glob J Obes Diabetes Metab Syndr 2019, 6: 001-009. DOI: <http://doi.org/10.17352/2455-8583.000035>
72. Serra C, Pérez N, Bou R, Baltasar A.: Sliding Self-Locking First Stitch and Aberdeen Knots in Suture Reinforcement with Omentoplasty of the Laparoscopic Gastric Sleeve Staple line. Obes Surg 2014; 24:1739–1740. DOI10.1007/s11695-014-1352-5 <https://www.youtube.com/watch?v=i4kNc8PJcEk>
73. Baltasar A, Bou R, Serra R, Bengochea M and Pérez N: Use of self-locking knots in running intestinal bariatric sutures. Glob Surg, 2015; 2:100-101. doi:10.15761/GOS.1000132



74. Recomendaciones de la SECO para la práctica de la cirugía bariátrica (Declaración de Salamanca). *Cir Esp* 2004;75(5):312-4
75. Baltasar A, Serra C, Pérez N, et al. Laparoscopic sleeve gastrectomy. A multi-purpose operation. *Obes Surg*. 2005; 15:1124–8.
76. Ahmad SS, Ahmad SS, Kohl S. The Hundred Most Cited Articles in Bariatric Surgery. *Obes Surg*. 2015; 25:900–9.
77. Angrisani L, Santonicola A, Iovino P et al. IFSO Worldwide Survey 2016: Primary, Endoluminal, and Revisional Procedures. *Obes Surg* 2018; 28:3783–3794
<https://doi.org/10.1007/s11695-018-3450-2>
78. Rutledge R.: The Mini-Gastric Bypass: Experience with the First 1,274 Cases Obesity Surgery, 11, 276-280
79. Carbajo M, García M, Toledano M et al. One anastomosis gastric bypass by laparoscopy: results of the first 209 patients. *Obes Surg* 2004; 15: 398-404.
80. Sánchez-Pernaute A, Miguel Rubio MA, Torres A. et al: Proximal Duodenal–Ileal End-to-Side Bypass with Sleeve Gastrectomy: Proposed Technique. *Obes Surg* 2007; 17:1614–1618 DOI 10.1007/s11695-007-9287-8
81. Baltasar A. Historical note: First diabetes metabolic operation in Spain *Integr Obesity Diabetes*, 2015. doi: 10.15761/IOD.1000140
82. Resa J, Solano J, Fatás A et al. Laparoscopic Biliopancreatic Diversion: Technical Aspects and Results of our Protocol. *Obes Surg* 2004; 14, 329-333
83. Resa J, Lagos J, Pérez A, et al.: Laparoscopic Gastro ileal Bypass with Single Anastomosis: Analysis of the First 1512 Patients. *Laparoscopic Gastroileal Bypass with Single Anastomosis*. *Obes Surg* 2019. <https://doi.org/10.1007/s11695-019-03788-6>
84. Alhambra M, Molina M, Prior M, et al Variations in diabetes remission rates after bariatric surgery in Spanish adults according to the use of different diagnostic criteria for diabetes. *BMC Endocrine Disorders* 2017; 17:51 DOI 10.1186/s12902-017-0201-7
85. Vidal J, Ibarzabal A, Romero F, et al. Type 2 diabetes mellitus and the metabolic syndrome following sleeve gastrectomy in severely obese subjects. *Obes Surg*. 2008;18(9):1077-1082.
86. Vilarrasa N, Rubio MA, Miñambres I et al. Long-term outcomes in patients with morbid obesity and Type 1 Diabetes Undergoing Bariatric Surgery *Obes Surg* 2017; 27:856–863 DOI 10.1007/s11695-016-2390-y
87. García M; Tinahones FG; Cohen R.: McGraw-Hill Interamericana de España S.L. 1ª ed., 1ª imp (10/04/2011) ISBN: 8448175409 ISBN-13: 9788448175405



88. Cruz M, Rubio MA, Calle L. Remisión de diabetes mellitus tipo 2 a largo plazo tras cirugía bariátrica Tesis. Hospital Clínico. 2017.
89. Torres A, Sánchez A, Rubio MA et al. Metabolic surgery for obesity a critical account ISSN 1575-0922, Vol. 60, Nº. Extra 1, 2013
90. Baltasar A, Serra C, Bou R et al. Sleeve Gastrectomy in a 10-year-old Child. *Obes Surg* 2008; 18:733–736 DOI 10.1007/s11695-007-9328-3
91. Baltasar A.: Nine years follow up of the 1st Adolescent Bariatric Surgery in Spain Nine years follow up of the 1st Adolescent Bariatric Surgery in Spain. *J J Obesity*. 2017. 3(1): 026.
92. Carbajo M, Vázquez R, Aparicio R.:12-year old adolescent with super morbid obesity, treated with laparoscopic one anastomosis gastric bypass (LOAGB/BAGUA): A case report after 5-year follow-up. *Nutr Hosp*. 2015; 31:2327-2332
93. Carbajo MA, González G, Jiménez JM. Et al. A 5-Year Follow-up in Children and Adolescents Undergoing One-Anastomosis Gastric Bypass (OAGB) at a European IFSO Excellence Center (EAC-BS). *Obesity Surgery* <https://doi.org/10.1007/s11695-019-03908-2>
94. Vilallonga R, Moreno JM, Yeste D et al.: Initial Approach to Childhood Obesity in Spain. A Multisociety Expert Panel Assessment. *Obes Surg* 2017; 27:997–1006 DOI 10.1007/s11695-016-2413-8
95. Vilallonga R, Fort JM, Caubet E, González O, Armengol M. Robotic Sleeve Gastrectomy Versus Laparoscopic Sleeve Gastrectomy: a Comparative Study with 200 Patients. *Obes Surg* 2013; 23:1501–1507 DOI 10.1007/s11695-013-1039-3
96. Morales S. Bypass gástrico por puerto único: detalles técnicos. De un nuevo abordaje. *Cir Es* 2015; 90:6:100-6
97. Morales-Conde S, Peeters A, Meyer YM, et al. European association for endoscopic surgery (EAES) consensus statement on single-incision endoscopic surgery. *Surg Endosc*. 2019 Apr;33(4): 996-1019. doi: 10.1007/s00464-019-06693-2.
98. Baltasar A, Bou R, Bengochea M, Serra C.: Malnutrición calórico-proteica. Tres tipos de alargamiento de asa común. *BMI-Latina* 2011; 5:96-97.
99. Baltasar A, Bou R, Bengochea M: Fatal perforations in laparoscopic bowel lengthening operations for malnutrition. *Surg Obes Relat Dis* 2010; 6:572–574
100. Bueno J, Pérez N, Serra C et al. Fistula pancreato-cutánea secundaria a pancreatitis postoperatoria tras cruce duodenal laparoscópico. *Cir. Esp* 2004; 76:184–6.



101. Serra C, Baltasar A, Andreo L. et al: Treatment of Gastric Leaks with Coated Self-expanding Stents after Sleeve Gastrectomy. *Obes Surg* 2007; 17:866-872
102. Serra C, Baltasar A, Pérez N. et al: Total Gastrectomy for Complications of the Duodenal Switch, with Reversal. *Obes Surg* 2006; 16:1082-1086
103. Castillo J, Fábrega F, Escalante C: Liver Transplantation in a Case of Steatohepatitis and Subacute Hepatic Failure after Biliopancreatic Diversion for Morbid Obesity. *Obes Surg* 2001; 11: 640–642 |
104. Baltasar A.: Liver failure and transplantation after duodenal switch. *Surg Obes Relat Dis* 2014; 10: c93-c96.
105. Mayo MA, Pacheco JM, Pérez D et al. Caso clínico Fallo hepático fulminante a largo plazo en pacientes sometidos a bypass gástrico por obesidad mórbida *Nutr Hosp.* 2015;32(1):430-434
106. Baltasar A, Bou R, 2, Serra C, et al.: Grapado de la sonda en gastrectomía vertical laparoscópica. *BMI-latina* 2013; 3:4-7
107. Baltasar A, Bou R, Bengochea M Intubación Dificil y Traqueotomía en el obeso mórbido Baltasar A.: *BMI-2013*; 3.:4.
108. Baltasar A.: Prevenir y Deshacer entuertos en cirugía. *Journal of negative or non-positive results.* 2019; 4: 252-64 DOI: 10.19230/jonnpr.2889
109. Duncan T, Tuggle K, Larry Hobson L et al. PL-107. Feasibility of laparoscopic gastric bypass performed on an outpatient basis. *Surg Obes Relat Dis* 2011; 7: 339–35