



Atmósfera

ISSN: 0187-6236

editor@atmosfera.unam.mx

Universidad Nacional Autónoma de México  
México

Pérez-García, Ismael; Aguilar-Sierra, Alejandro; Hernández-Alfaro, Jaime Steven

**Connections in the development of tropical cyclone  
Mitch (1998): A tribute to the human loss in this calamity**

Atmósfera, vol. 31, no. 3, 2018, July-September, pp. 285-292

Universidad Nacional Autónoma de México  
México

DOI: <https://doi.org/10.20937/ATM.2018.31.03.05>

Available in: <https://www.redalyc.org/articulo.oa?id=56557023009>

- How to cite
- Complete issue
- More information about this article
- Journal's webpage in redalyc.org

UNAM  redalyc.org

Scientific Information System Redalyc

Network of Scientific Journals from Latin America and the Caribbean, Spain and  
Portugal

Project academic non-profit, developed under the open access initiative

## **Connections in the development of tropical cyclone Mitch (1998): A tribute to the human loss in this calamity**

Ismael PÉREZ-GARCÍA<sup>†</sup>, Alejandro AGUILAR-SIERRA\* and Jaime Steven HERNÁNDEZ-ALFARO

*Centro de Ciencias de la Atmósfera, Universidad Nacional Autónoma de México, Circuito de la Investigación Científica s/n, Ciudad Universitaria, 04510 Ciudad de México, México*

Corresponding author; email: [asierra@unam.mx](mailto:asierra@unam.mx)

Received: April 6, 2017; accepted: May 9, 2018

<sup>†</sup>Dr. Ismael Pérez-García, a great meteorologist, scientist, and teacher passed away on April 20, a few days before the acceptance of this paper, in which he worked during his last days. This paper is a tribute to the victims of Mitch, but also to the life and work of Ismael Pérez-García.

### **RESUMEN**

Se estudia el ciclón tropical Mitch, que ocurrió en el mar Caribe en octubre de 1998. Mitch afectó a varios países de Centroamérica, donde causó grandes pérdidas humanas y económicas, y los autores consideran que la comunidad científica no le ha otorgado la atención que requiere. Dos patrones dipolares con orientación NE-SE asociados con sus zonas de convergencia-divergencia contribuyeron significativamente a su formación e intensificación, las cuales se analizan desde el punto de vista sinóptico observacional. Las características más notables de este fenómeno en el campo medio ocurrieron entre el 18 de octubre y el 4 de noviembre de 1998, cuando se formó otro patrón dipolar con dirección NO-SE, conformado por un anticiclón de Norteamérica de niveles altos (HLA, por sus siglas en inglés) y un anticiclón de Sudamérica de niveles altos (HLC, por sus siglas en inglés). Para discernir estos patrones se utilizaron dos fuentes de datos: el reanálisis NCAR-ds627.0 (ERA-Interim) y el de NCEP/NCAR. La idealización más simple de un vórtice dipolar NO-SE es un modón tropical, que es una solución exacta de la ecuación de vorticidad barotrópica (BVE, por sus siglas en inglés) en una esfera en rotación. Se explora el papel del HLA y el HLC en la excitación de las ondas tropicales utilizando un método espectral numérico para estudiar el modo de estabilidad normal de las soluciones exactas de la BVE. La iniciación de la onda tropical en niveles bajos podría estar asociada con perturbaciones en niveles altos del modo normal más inestable.

### **ABSTRACT**

Tropical cyclone Mitch, which occurred in October 1998 on the Caribbean Sea, is studied. Mitch affected countries in Central America, where it caused great economic and human losses, and the authors believe that the scientific community has not paid enough attention to this phenomenon. Two NE-SE oriented dipole patterns associated with their convergence-divergence zones contributed significantly to its formation and intensification, which are analyzed from the observational synoptic point of view. The most remarkable characteristics of this phenomenon in the mean field occurred between October 18 and November 4, 1998 when another NW-SE dipolar pattern was formed, consisting of a high-level anticyclone from North America (HLA) and a high-level anticyclone from South America (HLC). To discern these patterns, two data sources were used: the NCAR-ds627.0 (ERA-Interim) reanalysis and the NCEP/NCAR reanalysis. The simplest idealization of the NW-SE dipolar vortex is a tropical modon, which is an exact solution of the barotropic

vorticity equation (BVE) on a rotating sphere. The role of the HLA and the HLC in the excitation of tropical waves is explored using a numerical spectral method for studying the normal mode stability of exact solutions of the BVE. The initiation of the tropical wave at low levels could be associated with disturbances at high levels of the most unstable normal mode.

**Keywords:** Tropical cyclone Mitch, La Niña, transient connection patterns, tropical wave, barotropic vorticity equation, Verkley's modon solution, normal mode instability.

## 1. Introduction

The coasts of Mexico and Central America are affected by tropical cyclones of the north Atlantic and the northeast Pacific. The relationship between El Niño-Southern Oscillation (ENSO) and the tropical cyclones of the Northeast Pacific and the North Atlantic has been studied by many authors (Landsea, 1993; Demaria and Kaplan, 1994; Goldenberg et al., 2001; Vitart and Anderson, 2001; Amador et al., 2006; Camargo et al., 2007; Wood and Ritchie, 2013). Gray (1984) and Shapiro (1987) point out that the smaller vertical shear that accompanies La Niña phase directly contributes to increasing the number of tropical cyclones in the Atlantic. Camargo et al. (2007) noted that tropical cyclone activity decreased in the Atlantic during El Niño years.

In this paper, we show that large horizontal convergences are generated at higher levels during La Niña phase near the central-eastern Pacific, which can suppress the activity of tropical cyclones in this region due to the vertical shear. In section 2, we look more closely at how the tropical waves are amplified by the tropical upper tropospheric trough (TUTT).

Gray (1984) suggests that El Niño can suppress hurricane activity in tropical Atlantic regions due to its influence in the upper troposphere; however, he also indicates that the summer TUTT will act as a process to decrease the inhibitory influences of the vertical shear and the ventilation of higher levels over the center of a disturbance. In section two, we will look more closely at the way in which tropical waves are amplified by the TUTT. It is not clear which physical mechanisms are responsible for the greater number and intensity of hurricanes during La Niña years, although it has been observed that the trajectories of tropical cyclones are mainly towards the west during these periods, tending to recurve during El Niño years.

The first part of this study, presented in section 2, focuses on the La Niña period of 1998, with a

particular emphasis on the study of the development of tropical cyclone Mitch in the Caribbean sea. Our basic idea is based on the fact that during a La Niña phase, a large-scale convergence can be created near Central America. Mitch's general movement was slow, resulting in a large amount of rainfall, mainly in Central America, where it caused catastrophic losses and devastation, especially in Honduras, El Salvador, Guatemala and Nicaragua (Hellin and Haigh, 1999; Lidy et al., 2001; Pasch et al., 2001). Although tropical cyclone Mitch caused widespread damage, the scientific community has not studied it extensively.

The relationships observed between low pressure and the number of tropical cyclones in the Caribbean Sea show peaks in June and October (Inoue et al., 2002), and in the last month there is a weak easterly wind and a sea level pressure (SLP) minimum (Wang, 2007). In the lower troposphere, tropical waves appear as an inverted "V" and move along with the trade winds across the Atlantic Ocean (Frank, 1968). They begin to appear from April/May and continue until October/November (Burpee, 1972). It has been recognized that around 60% of Atlantic tropical storms originate from the eastern waves and that almost all tropical cyclones that occur in the eastern Pacific Ocean can also be traced from Africa (Burpee, 1972; Avila and Pasch, 1995). Burpee (1972) documented a mechanism for the origin of these waves, the instability of the east African jet. Although other characteristics were proposed (Krishnamurti et al., 2013), here we suggest that tropical systems are disturbed by the instability of a dipolar vortex in the upper troposphere, which is first examined by means of observations and then by the analysis of eigenvalues of the non-divergent linearized barotropic vorticity equation. So, some of the basic aspects of the development and movement of tropical cyclone Mitch can be illustrated in terms of the barotropic vorticity equation (BVE) theory over the sphere.

The present work provides an observational study of a dipolar vortex in the upper troposphere associated with tropical cyclone Mitch; these dipolar vortices have not been documented in the literature (Pérez-García et al., 2017). Tropical cyclone Mitch was not an isolated solitary vortex but it was pushed by a directional flow generated within a dipolar vortex of the upper troposphere. Our challenge is to understand in how many ways the environment interacted with Mitch. The dipolar vortex could have helped to intensify the cyclone and move it towards the SW (Pérez-García et al., 2017). This environment is the dipolar vortex of the upper troposphere on a large scale and can be analytically represented by a Gill solution (Gill, 1980) or an exact solution of the non-linear BVE. The structure is very similar to a modon trapped near the equator (Verkley, 1984; Pérez-García and Skiba, 1999; Skiba and Pérez-García, 2009). In the context of the BVE exact solutions on the rotating unit sphere, a dipolar vortex of the upper troposphere is constructed and is called the tropical modon. The modons are localized solutions of nonlinear dynamic equations. They are stable against interactions with external fields (Leith, 1981) and are obtained without the existence of a local source of warming.

Despite its simplicity, this BVE contains the main elements that describe the complexity of atmospheric behavior (Simmons et al., 1983; Pérez-García et al., 2016) and provides information on tropical circulation. This has not been previously examined in the literature. Our modest goal is to fill this gap for the equatorial waves of non-linear basic dynamics. This is first presented observationally in section 2 and then analytically in section 3. The concluding remarks are given in section 4.

## 2. Transient connection pattern and tropical circulation during tropical cyclone Mitch, 1998

Many tropical disturbances drive large amounts of clouds by low-level air convergence; therefore, the wind field provides more information about tropical circulation than pressure. For this investigation, NCAR-ds627.0 (ECMWF, 2009) and NCEP/NCAR reanalysis data (Kalnay et al., 1996) were used. In particular, we used the relative humidity and the zonal ( $u$ ) and meridional ( $v$ ) components of the wind field at different pressure levels. These wind components

were used to calculate the divergence  $\delta$ , the vertical component of the relative vorticity  $\zeta$ , the stretch deformation  $E$  and the shear deformation  $F$ . To obtain a clear view of the entire circulation over the tropical region, we took the whole sphere as domain.

The term  $(u, v)$  is decomposed into a non-divergent flow and an irrotational flow. The non-divergent flow is described by the stream function  $\psi$ , which satisfies  $\nabla^2\psi = \zeta$ , and the irrotational flow satisfies  $\nabla^2\chi = \delta$ , where  $\chi$  is the velocity potential. These elliptic equations are solved using the numerical spectral method (Sardeshmukh and Hoskins, 1987) where we use a grid of  $128 \times 64$  and a truncation of T31.

Figure 1a shows the relative humidity in the 700–925 hPa layer and the average streamlines for the upper layer (200–300 hPa) on October 16, 1998 at 12 UTC. The interesting feature in this field that passes over the tropical American region is the dipolar envelope that contains two vortices; one is the high-level anticyclone from South America (HLC), and the other is the high-level anticyclone from North America (HLA) (Reed, 1933). Two trough lines referred to as TUTT and small cyclone (C1) and anticyclone (A1) can also be seen in Figure 1a.

Figure 1b shows the average streamlines for the lower layer (925–1000 hPa) and the relative humidity (for the 700–925 hPa layer, shaded) for October 17, 1998 at 00 UTC. A longitudinal band of eastern zonal wind was located at the south of the Gulf of Mexico, induced by a transient anticyclone labeled as TA and a zonal wind traced to the central Atlantic caused by the southern edge of the North Atlantic Subtropical High (NASH). As the eastern zonal wind continued to the Americas, some streamlines are directed toward Mexico-Central America and other streamlines to South America. As the eastern wind continued to flow into the Caribbean Sea, a tropical wave associated to the origin of Mitch is not present in the streamlines.

Figure 2a shows the GOES-IR satellite images for the same date (October 17) at 00 UTC, where we can see that a cloud cluster associated with the development of the tropical wave was not yet noticeable, only a cover of clouds to the east is observed. On October 18 at 23:45 UTC, a cloud cluster was seen in the vicinity of 60 W, 15 N (Fig. 2b). Comparing Figure 1b with Figure 2a, 2b, that cloud cluster embedded in an easterly wind is easier to track than a tropical wave (Sikdar and Hentz, 1980).






Fig. 1. (a) Mean streamlines for the upper layer (200-300 hPa) and mean relative humidity (shaded) in the 700-925 hPa layer for October 16, 1998 at 12 UTC. The high-level anticyclone of North America (HLA), the high-level anticyclone (HLC) of South America, the cyclone C1 and the small anticyclone A1 are designed. The bold arrows (red and blue) indicate the small jet cores and the tropical upper tropospheric troughs (TUTT) are designated by a thin blue arrow. (b) Streamlines and winds of the 925-1000 hPa layer and mean RH of 700-925 hPa (shaded) for October 19, 1998 at 00 UTC. HR areas greater than 50% in (a) are indicated by darker shades, the contour interval for 50-100% is 10, white shading indicates values below 50%. The maximum wind intensity (m/s) is labeled on the lower right corner. Transient anticyclones are labeled as TA, northeast and southeast Pacific subtropical highs as NPSH and SPSH, respectively, as well as NASH and SASH for North Atlantic and South Atlantic subtropical highs.




Fig. 2. GOES8-IR satellite images for (a) October 17, 1998 at 00:00 UTC and (b) October 18, 1998 at 23:45 UTC.

We must go back days before the tropical wave approached Jamaica, to understand the characteristics of the phenomenon that occurred. By October 15 at 00 UTC, at an average of 200-300 hPa, two more trough

lines originated: the first, oriented N-S, was located to the east side of HLA and the second (NE-SE) to the east side of the HLC. As the NW-SE trough to the NE of the HLC was longer, it quickly became a

small A1 anticyclone on October 15 at 06 UTC. By October 16, the upper trough on the east side of the HLA was transformed into a small C1 cyclone at high levels and the A1 was amplified to the northeast of the HLC, allowing a SW-NE flow between C1 and A1, as shown in Figure 1a.

By October 17 at 00 UTC, a new tropical upper troposphere trough (TUTT) line oriented SW-NE was generated on the east side of the HLA. After the TUTT was transformed into a C2 cyclone of high levels on day 18 at 00 UTC (not shown), it allowed the formation of a small A2 anticyclone of high levels between C1 and C2. The presence of the HLA and this C2 cyclone of high levels disturbed the tropical flow to the southeast, creating a small chain of disturbances or waveguide composed of small cyclones and anticyclones, decreasing in amplitude toward the southeast (Fig. 3a). It was then that the tropical flow was disturbed at lower levels (Fig. 3b).

Under this condition, there was a flow to the NE between these small cyclones and anticyclones and the cloud cluster appeared, as shown in Figure 2b. In this place the tropical wave gained amplitude, which originated the tropical cyclone Mitch (Fig. 3b). The amplitude of A2 increased at the expense of the loss of energy and decrease in amplitude of the HLA. Later on, the C2 cyclonic circulation moved between the HLA and the HLC, and around the HLA. Because C2 was much closer to the HLA, it was guided by the circulation of the HLA and on October 22 it was located over Mexico (Fig. 5a).

By October 21 at 00:00 UTC, this horizontal distribution of the streamlines showed a hyperbolic deformation field, which developed on the east side of the dipolar vortex between the island of Puerto Rico and Colombia. The kinematics or motion geometry within the upper troposphere dipolar vortex, the conditions for La Niña event and the beginning of the tropical Mitch perturbation were examined by the potential function  $\chi$  for the divergent average wind in the 200-300 hPa and 925-1000 hPa layers. We will see how another mechanism can disturb a tropical wave.

The intense warming of the warm pool of the eastern Pacific, the warm pool of the Caribbean, the Sierra Madre Occidental and the deserts of Chihuahua and Sonora cause the air to rise, which in turn generates a low surface pressure (centered near the coast NW-SW

of Mexico and Central America), named as the maritime-continental thermal low (Pérez-García et al., 2017). Therefore, large horizontal convergences are generated at lower levels between the eastern Pacific about 100° W and the eastern Caribbean about 70° W.

Figure 3c shows the convergent/divergent average wind and its corresponding potential function in the 925-1000 hPa layer and the average relative humidity (gray shades) in the 700-925 hPa layer for October 18, at 00 UTC. A notable feature in this figure is the strong convergence and high relative humidity throughout Central America, Mexico and the northwest of South America, since the Caribbean Sea, the eastern Pacific and the Gulf of Mexico serve as sources of atmospheric humidity for rainfall in the Americas (Hales, 1974; Ting and Wang, 1997; Barlow et al., 1998; Klimczak, 2000; Wang, 2007). Near the northeastern equatorial Pacific, the trade winds southwest of Baja California become wet toward Mexico and Central America (Fig. 3b).

A wide area of convergence extends between Mexico, Central America and just north of South America, while around it we can find the four primary centers of action, namely the Northeastern Pacific Subtropical High (NPSH), the Southeastern Pacific Subtropical High (SPSH), the Azores High or North Atlantic Subtropical High (NASH), and the South Atlantic Subtropical High (SASH). Diverging winds generate each one of these centers of action and contribute to the converging winds over America in the tropical region, as can be seen in Figure 3c. A transient anticyclone, indicated by the letters "TA", is also part of the convergence zone in tropical America.

In October, when the NASH is weak and extends to the NE, the SST in the Caribbean and NE Pacific near Mexico and Central America are abnormally warm (Wang, 2007). The SST anomaly chart of October 20, 1998, shows negative values in most of the eastern-central Pacific domain (30° S-30° N, 180°-115° W) and minimum values along the equator, as corresponding to La Niña phase (<http://www.ospo.noaa.gov/data/sst/anomaly/1998/anom-night.10.20.1998.gif>).

In Figure 3d we also observe the trajectories (blue) of these primary centers of action during most of October 1998. NASH and SASH remain almost stationary. From October 16 to 20 the trajectory of a migratory anticyclone (traced in red) labeled as TA1






Fig. 3. (a) Same as in Fig. 1a, but for October 19, 1998 at 00 UTC. (b) Streamlines at 800 hPa and mean relative humidity for the 700-925 hPa layer for October 19, 1998 at 00 UTC. The axis of the eastern waves (EW and EW\*) is marked by bars. Tropical cyclone Mitch originated from EW\* and EW is related to the cloud cluster located over the western Caribbean Sea. (c) Mean convergent/divergent wind and its corresponding potential function contours ( $\times 10^5$ ) in the 925-1000 hPa layer and mean relative humidity (shaded) in the 700-925 hPa layer for October 18, 1998, 00 UTC. (d) Same as in Fig. 3c, but for October 19, 1998, 00 UTC. Blue lines designate the trajectory of the primary centers of action and the red lines, transient anticyclones (in mid-latitudes), for the better part of October 1998; numbers indicate the initial and final day for each case. The transient connection pattern to La Niña is traced in green. (e) Same as Figure 3c for layer 200-300 hPa. The trajectories in red and orange designate the greatest horizontal convergences for some transient anticyclones, during October 1998. The transient connection pattern to SPSH is traced in green for October 19, 1998, 00 UTC. (f) Same as in Fig. 3d, the connection pattern to SPSH and the transient connection pattern to the east of USA are traced with green for October 23, 1998, 00 UTC.

can be observed, reaching northwest of the Atlantic Ocean by October 20. On October 23, the SPSH fluctuated zonally and suffered a TA2 dispersion that moved on October 24 over the southern part of South America. A third transient anticyclone TA3 was located near the equator of the central Pacific.

In Figure 3d, we track these transient anticyclones (red) from October 16 to 20 for TA1, from 18 to 21 for TA2 and from 17 to 19 for TA3 (not shown). We point out (orange) the centers of the dry air intrusion, with a W-E elongation generated near the equatorial central Pacific that which could be more related to a local air-ocean heat exchange process (La Niña phase) (Lindzen and Nigam, 1987) in the east-center of the cold water pool of the Pacific. This SST anomaly in the central-eastern Pacific seems to explain the dry air intrusion and the zone of divergence.

In the 925-1000 hPa mean layer of the lower troposphere (Fig. 3d), we find the convergence on tropical America that connects (represented by a green line) with two other divergent centers of TA3 (or SPSH) and TA1 action; moreover, TA1 may have been connected to the northeast with a convergent center. TA1, TA3, NASH, SPSH and the wide area in the eastern central Pacific (the area of 30° S-30° N and 160°-120° W) had a direct influence on the low level convergence in the tropical region of America.

The greatest horizontal divergences were generated in the middle and upper levels of the troposphere (Fig. 3e) and the convergences were generated in the lower levels (Fig. 3c and 3d), favoring the different stages of development of the tropical system in the Caribbean Sea, as shown above with more details. The warm air rises and cools in a low pressure zone and the areas of descending air produce high pressure zones. In the region between the southeastern Pacific and Central America, convergent-divergent cells are formed at low to high levels. The zone of convergence located between Mexico and the North Atlantic, seen in Figure 3d, is associated with the position of the migratory anticyclone TA1.

The connection between TA1 and TA3 is called Transient Connection Pattern to La Niña (TCPNa). It describes the convection cells of northeastern and southeastern Central America. TA1 is connected to the northeast with a convergent center located in the North Atlantic. However, this convergent field can also be connected to a divergent field located in

Greenland as shown in Figure 3d. TCPNa is an alternate pattern between pressures in the central Pacific Ocean and various centers of action in Central America and the northwest Atlantic Ocean. Another route for this connection is through the divergent eastern Pacific centers of action and with North America as in the Pacific/North American (PNA) Classic Pattern (Barnston and Livezey, 1987).

The four primary centers of action and the transient anticyclone are associated with the convergent/divergent fields in the 200-300 hPa layers as seen in Figure 3e. An intensified elongation of the NW-SE direction in the divergence field appears over Central America and a divergence is also observed with a more marked asymmetry over the north Atlantic. A stronger and more southern convergence was limited to the southeast Pacific, while a weaker convergence was observed over the northwest north Atlantic, thus creating a NW-SE feedback system. The wide field of divergence that appears over Mexico, Central America and the northwest of the Southern Hemisphere unites them from above.

On the SPSH center of action on October 19, 1998, in the upper average atmosphere of 200-300 hPa, there is a strong convergence that connects the three different convergent, divergent and convergent centers of action, as shown in Figure 3e, which we call the connection pattern to SPSH. According to the classification of teleconnection weather patterns by Barnston and Livezey (1987), two of these patterns are the North Atlantic Oscillation (NAO) and the PNA. The NAO is a north-south oscillation on the path of storms and depressions across the north Atlantic and flows into Europe. It has been observed that the positive phase of the PNA is strongly related to the ENSO phenomenon, while the negative phase tends to be associated with La Niña. In this work, the SPSH and TCPNa connections are not statistical patterns of low frequency variability, because their occurrence lasts only a few hours or a few days at most.

Figure 4 shows schematically the relative humidity in the 700-925 hPa layer and the average field of streamlines for the upper layer (200-300 hPa) from October 17 to November 4. The most interesting feature of this mean field of streamlines that passes over the tropical America region is a dipolar vortex formed by the HLC and the HLA. This dipolar vortex, which has a Gill-Matsuno wave configuration (Pérez-García






Fig. 4. Mean streamlines at 200-300 hPa, mean RH at 700-925 hPa and trajectories (blue lines) of the HLC and HLA vortices from October 17 to November 4, 1998. The red line represents Hurricane Mitch trajectory from October 22 to November 4, 1998.

et al., 2017), is the main research topic in this work. This may have contributed to the movement of tropical cyclone Mitch to the SW, as shown below. Figure 4 also shows the trajectories of the HLC, HLA and Mitch vortices, respectively, and it is observed how this tropical cyclone moved between the HLC and the HLA.

When comparing Figures 3c and 4, we noticed that the dipolar vortex and the tropical wave that would give rise to tropical cyclone Mitch were trapped within a convergence zone below (Fig. 3c) and a divergence zone above. To understand a tropical dipole circulation formed by an anticyclone at the upper level, it is necessary first to understand how an HLA is formed, where rainfall occurs and how it relates to the development of tropical cyclones within that dynamic system.

Following the streamlines analysis maps in Figure 4, we can see the movement of tropical cyclone Mitch within a dipolar vortex as a dynamic system. We examine below the resulting stationary perturbation in the linear barotropic model, using a numerical spectral method (Skiba and Pérez-García, 2009) that is tested for an isolated stationary equatorial modon similar to that constructed by Verkley (1984). As the northeast wind flowed to the eastern Pacific (October 21 00 UTC, 200-300 hPa layer), a longitudinal band of strong northeasterly winds (located between the HLC and the HLA in the eastern Pacific) intensified,

forming a diffluent jet (between Panama and Nicaragua) that can form a line of divergence and induce a convergence zone on the surface, which is why a convergence nucleus was observed at the 925-1000 hPa level in the southwest region of the Caribbean Sea. Therefore, frontogenesis (Keyser et al., 1988; Pérez-García, 1993) was driven by the confluence to form a higher level front and two jet cores oriented SW-NE and directed to opposite sides. The south-eastern core might have intensified the disturbance at the lower levels or to the tropical wave.

A new TUTT oriented SW-NE was developed towards the east side of the HLA, then it was transformed into an upper-level low C2 on October 21 at 18 UTC (not shown). This allowed the formation of an upper-level high A2 east of the C2, thus the condition of October 19 (Fig. 3a) was replicated on October 22 at 18 UTC (Fig. 5a). A jet core located between HLA and the C2 oriented NE-SW and directed towards the western Caribbean Sea makes the low level tropical disturbance unstable, which would give origin to Mitch (Fig. 5c).

The TUTT and the upper-level low C2 appeared when the HLA was of greater amplitude and with an almost symmetric circulation. In addition, the HLC was more elongated towards the west. Thus, as a result of the external mechanisms of tropical cyclone formation (Montgomery and Farrell, 1993), the tropical depression that gave rise to Mitch was formed below the line of divergence between the HLC and the C2 on October 22 at 18 UTC, about 76° W, 12° N. A line of divergence in the upper troposphere (Fig. 5a) induces a zone of convergence at lower levels (Fig. 5b), and may become a mechanism giving rise to tropical disturbances (see Fig. 5c and Pérez-García et al., 2016). This area of convergence is reinforced by the connections established SW of the eastern Pacific and to the east of the United States (Fig. 5b).

Early on October 24, Mitch became a hurricane. Later that day, as it moved westward, it began to intensify rapidly. In about 24 hours, its central pressure dropped to 924 hPa on the afternoon of October 25. The center of the high divergence zone associated with the dipolar vortex (200-300 hPa) was located on the southwest of the Caribbean Sea on October 25 (not shown). Mitch continued to strengthen and its central pressure reached a minimum of 905 hPa some