



Atmósfera

ISSN: 0187-6236

editor@atmosfera.unam.mx

Universidad Nacional Autónoma de México
México

Montero-Martínez, Guillermo; Rivera-Arellano, Joel; Roy, Priyadarsi D.; Rosado-Abón, Anielka;
Hernández-Nagay, Déborah Patricia; Mendoza-Trejo, Angélica; Andraca-Ayala, Gema Luz

**Content and composition of dissolved organic carbon
in precipitation at the southern part of Mexico City**

Atmósfera, vol. 31, no. 4, 2018, October-December, pp. 331-346

Universidad Nacional Autónoma de México

México

DOI: <https://doi.org/10.20937/ATM.2018.31.04.02>

Available in: <https://www.redalyc.org/articulo.oa?id=56557569002>

- How to cite
- Complete issue
- More information about this article
- Journal's webpage in redalyc.org

UNAM  redalyc.org

Scientific Information System Redalyc

Network of Scientific Journals from Latin America and the Caribbean, Spain and
Portugal

Project academic non-profit, developed under the open access initiative

Content and composition of dissolved organic carbon in precipitation at the southern part of Mexico City

Guillermo MONTERO-MARTÍNEZ^{1*}, Joel RIVERA-ARELLANO¹, Priyadarsi D. ROY²,
Anielka ROSADO-ABÓN³, Déborah Patricia HERNÁNDEZ-NAGAY¹, Angélica MENDOZA-TREJO¹
and Gema Luz ANDRACA-AYALA¹

¹ Centro de Ciencias de la Atmósfera, Universidad Nacional Autónoma de México, Circuito de la Investigación Científica s/n, Ciudad Universitaria, 04510 Ciudad de México, México.

² Instituto de Geología, Universidad Nacional Autónoma de México, Circuito de la Investigación Científica s/n, Ciudad Universitaria, 04510 Ciudad de México, México.

³ Instituto de Química, Universidad Nacional Autónoma de México, Circuito de la Investigación Científica s/n, Ciudad Universitaria, 04510 Ciudad de México, México.

*Corresponding author; email: gmontero@atmosfera.unam.mx

Received: August 28, 2017; accepted: June 15, 2018

RESUMEN

Se presentan las concentraciones de carbono total disuelto (TC) y carbono orgánico (OC) en muestras de agua de lluvia recolectadas durante 2016 en el sur de la Ciudad de México. El TC varió de 10 a 25 mgC L⁻¹ y el OC, reportado como carbón orgánico no purgable (NPOC, por sus siglas en inglés), tuvo un rango de 2.5 a 20 mgC L⁻¹. Los valores más altos de NPOC se midieron en las muestras recolectadas entre mayo y julio, y se detectó un cambio en la razón de NPOC/TC para la segunda parte de la campaña (es decir, agosto-octubre). El análisis por ¹H RMN sugiere que aldehídos, ácidos carboxílicos y otros hidrocarburos oxigenados constituyen la fracción orgánica disuelta en las muestras de precipitación. La evaluación de las trayectorias de las masas de aire no encontró ninguna relación entre los contenidos de compuestos orgánicos y las direcciones del viento dominante para la formación de las tormentas, lo que sugiere que los compuestos orgánicos disueltos en el agua de lluvia tienen su origen en fuentes dentro de la cuenca de México. Además, los valores de pH más altos en las muestras estudiadas indican que la neutralización de la lluvia ocurre cuando las tormentas provienen del norte de la ciudad, donde se encuentran las fuentes de los agentes que producen este efecto.

ABSTRACT

We present concentrations of dissolved total carbon (TC) and organic carbon (OC) in rainwater samples collected in 2016 in the southern part of Mexico City in a preliminary study. TC ranged from 10 to 25 mgC L⁻¹ and OC – reported as non-purgable organic carbon (NPOC) – varied between 2.5 and 20 mgC L⁻¹. NPOC concentrations were higher in samples collected between May and July and a change in the ratio of NPOC/TC was detected for the second part of the campaign (i.e., August-October). ¹H NMR analysis indicated presence of aldehydes, carboxylic and other oxygenated hydrocarbons in the precipitation. An evaluation of air mass back-trajectories suggested no relationship between the organic contents and wind directions. Carbon contents were originated from sources within the basin of Mexico and higher pH of rainwater during the passage of air masses indicates that the sources causing rain neutralization are located in northern parts of the city.

Keywords: Water-soluble organic compounds, rainwater characterization, rain neutralization, urban pollution, Mexico City.

1. Introduction

Carbon is present as gaseous and condensed phases in the atmosphere and both of them are associated with anthropogenic and biogenic sources (Siudek et al., 2015). Atmospheric aerosols contain inorganic (IC) and organic carbon (OC). More than 2300 Tg of OC are emitted annually into the atmosphere and a significant fraction remains in the condensed aerosol phase (Yan and Kim, 2012; Nozière et al., 2015), affecting air quality and the radiative balance of the atmosphere (Gilardoni et al., 2016), both in urban and rural locations (Putaud et al., 2004; Zhang et al., 2007). The organic fraction includes saturated and unsaturated aliphatic compounds, aromatics, alcohols, ketones, aldehydes, carboxylic acids, amines, sugars, polyols, and organic sulphur compounds (Seinfeld and Pandis, 2006). OC is ubiquitous in urban locations (e.g., Zhang et al., 2007) and constitutes a major component (20-90% of the total mass) of the sub-micron atmospheric aerosol particles (Kanakidou et al., 2005; Moffet et al., 2010). Furthermore, it has been shown that water-soluble organic compounds (WSOC) represent 20-70% of the total organic aerosol mass (Saxena and Hildemann, 1996; Facchini et al., 1999) and can reach a considerable fraction (~50%) of the total water-soluble materials in the size range of the cloud condensation nuclei (Huang et al., 2006).

Wet deposition processes (e.g., rainout and washout) remove particulate material and pollutants from the atmosphere (Guidotti et al., 2000; Kawamura et al., 2001; Seinfeld and Pandis, 2006; Balla et al., 2014). Therefore, OC is a typical constituent of rainwater (Balla et al., 2014) and it can be used as an initial screening of air quality in different urban and/or industrial areas (Guidotti et al., 2000), and to infer some of its effects on ecosystems (Santos et al., 2009). WSOC are poorly understood due to their inherent complexity and difficulties in chemical analysis, even though they are essential components of atmospheric aerosols. The organic fraction consists of low-molecular weight (MW) polar compounds (Saxena and Hildemann, 1996; Kawamura et al., 2001) as well as high-MW, polydisperse macromolecules, termed as humic-like substances (HULIS) due to their similarities to terrestrial humic acids (Graber and Rudich, 2006). During the last two decades, OC present in the atmosphere was characterized using

chromatographic techniques, mass and UV spectrometry, and nuclear magnetic resonance, among others; however, there remain many challenges in this research field (Nozière et al., 2015). Studies of OC in precipitation are scarce compared to those for other atmospheric substrates, but results indicate that OC in rainwater is lower compared to concentrations observed in snow and fog (Iavorivska et al., 2016 and references therein). Some of the previous reports have focused on the measurement of total WSOC concentrations (Coehlo et al., 2008; Pan et al., 2010; Yan and Kim, 2012; Siudek et al., 2015) and others have characterized the different fractions by using gas, liquid, and UV-visible chromatographies; molecular fluorescence; FTIR-ATR, and ^1H NMR spectroscopies (Santos et al., 2009; Xu et al., 2008; Guidotti et al., 2000; Čosović et al., 2007; Chon et al., 2015).

Previous studies on rainwater characterization in samples from Mexico City were conducted by analyzing major inorganic ions and trace elements (Báez et al., 2006, 2007; 2009; García et al., 2006). Báez et al. (1989, 1993, 2008) measured organic compounds in rainwater samples but focused only on carbonyl compounds. Balla et al. (2014, and references therein) reported that aldehydes constitute up to 16% of dissolved OC in rainwater samples. This study aims to measure the total organic content and characterize the organic compounds present in rainwater samples collected in an urban area of Mexico City during 2016. As rain collects air pollutants in gaseous and liquid phase, the determination and characterization of OC could be used as a screening of air quality in the area and for the study of surface-active effects (a key factor in the collision-coalescence process) in fog and cloud droplets due to the presence of these organic compounds.

2. Experimental

Mexico City's metropolitan area (MCMA) is located to the southwest of the basin of Mexico. Except for the northeast, this basin is surrounded by mountains, some of them with altitudes higher than 4500 masl. Due to the subtropical latitude and the mean altitude (2250 m) of the basin, the intense solar radiation promotes the formation of photochemical pollutants in the area (Salcedo et al., 2006). The main rainy

season – in terms of accumulated rain – occurs from May to October (summer and autumn), and it is characterized by the development of convective clouds with warm bases and cold tops (Kucieńska et al., 2010) with predominant winds coming from north-northeast (Jáuregui, 2000). Furthermore, the passage of cold fronts during the winter produces a second rainy season, although its contribution in terms of precipitated water quantity is much lower (Montero et al., 2011).

The sampling site is located on the roof of the Centro de Ciencias de la Atmósfera (Atmospheric Sciences Center, CCA) building (19.33° N, 99.18° W, 2280 masl) of the National Autonomous University of Mexico (Fig. 1), in the southwestern part of Mexico City. Instrumentation includes diverse sensors for the measurement of common meteorological parameters (temperature, pressure, wind direction and intensity, etc.) and evaluation of air quality. A Texas Electronics TR-525M tipping bucket rain gauge records rainfall intensity with a resolution of 0.1 mm and an accuracy of 1%. Rainwater samples were gathered from May to October 2016 on an event basis. Sample collection was made with a stainless steel funnel in order to avoid problems of absorption of organic compounds

on the surface (Decesari et al., 2000) and amber glass bottles (washed with both deionized and grade HPLC water and combusted at 280 °C during 24 h to remove volatile organic compounds). All the samples were filtered and stored in amber bottles – sealed with an aluminum foil baked at 280 °C – and kept in refrigeration at 4 °C after collection until analyzed. Filtration was done using filters TITAN3 (manufactured by Thermo Scientific) with nylon membrane and a diameter pore of 0.45 mm previously washed with deionized water.

2.1 Characterization of organic compounds

Proton nuclear magnetic resonance (^1H NMR) spectroscopic analysis was done to obtain preliminary information about the organic matter dissolved in rainwater and aerosol samples. Some important characteristics of this technique include that good results – spectra – can be acquired to identify the functional groups of organics in one sample and the small amount (about 5 to 30 mg) necessary to perform the assays (Suzuki et al., 2001). Rain samples were dried using a rotary evaporator with a high purity nitrogen flow at ambient temperature (ca. 20 °C). ^1H NMR spectra were recorded in deuterated water (~0.6 mL) with an Eclipse JEOL 300 spectrometer.

2.2 Carbon analysis

Concentrations of total carbon (TC) and organic carbon (OC) dissolved in the rainwater samples were estimated by high temperature non-catalytic temperature using a Thermo Scientific HiperTOC carbon analyzer with an auto-sampler. The HiperTOC uses a dual NDIR approach (15.2 and 38.1 cm sample cells) to detect CO_2 levels. Both detectors are calibrated to 1% CO_2 full scale and operate reliably in the 0–1000 ppm CO_2 gas range (Thermo Scientific, 2007). All results were gathered from the detector with the longest cell (low bench detector). Total OC was measured as Non-purgeable organic carbon (NPOC), which estimates the non-volatile OC (Saarikoski et al., 2008). The procedure involves removal of inorganic carbon (IC) by treatment with acid ($\text{Na}_2\text{S}_2\text{O}_8$, also known as persulfate) prior to measuring the OC (Bisutti et al., 2004). The reference solutions for OC determinations were prepared from reagent grade anhydrous potassium hydrogen phthalate (KHP), while a mixture of anhydrous sodium carbonate and




Fig. 1. Geographical location of the basin of Mexico, its main orographic features as well as the locations mentioned in the text.

anhydrous sodium hydrogen carbonate was used for TC measurements. To perform carbon analyses, 0.25 mL of solution (sample) was injected into the analyzer, which oxidizes carbon at 1000 °C without a catalyst. In particular, for NPOC measurements, the acidified samples were sparged with CO₂ free carrier gas (UHP oxygen [99.8% purity] with < 0.5 ppm of total hydrocarbons) at a flow rate of 200 mL/min for 5 min to remove inorganic carbon prior to the actual analysis. Total IC can be estimated as the difference between TC and NPOC.

2.3 Back trajectory analysis

Backward trajectories of the air masses were calculated based on the hybrid single-particle Lagrangian integrated trajectory model – NOAA HYSPLIT (Stein et al., 2015) – to study a potential association between air masses and the organic contents in rainwater. The input parameters were as follows: meteorological database, NARR; trajectory duration, 48-h; vertical motion based on model vertical velocity, and starting heights, 500 and 1500 m above the ground level. Both height levels are within the air layer below the cloud base of the storms that occur in the area during the sampling period (Montero et al., 2011). The lowest level aims to facilitate studying the differences in the air mass trajectories closest to the superficial pollution sources with respect to those at higher levels.

The Red Automática de Monitoreo Ambiental (Automated Environmental Monitoring Network, RAMA) (<http://www.aire.cdmx.gob.mx>) employs 29 stations distributed throughout Mexico City to provide information about the pollutant levels. The hourly air pollution data from a RAMA station installed in the sampling site (CCA) was used in this study.

3. Results and discussion

3.1 Characterization of organics by ¹H NMR

Based on the results of previous reports for urban areas worldwide (a few dozens of mg of carbon per liter of rainwater) and volumes of the collected samples during the first part of this study (less than 500 mL), it was decided to mix a pair of samples in order to get enough material for the ¹H NMR analysis. The selected samples correspond to events occurred

during the first part of the rainy season. The mixture of rainwater samples was injected into the ¹H NMR instrument to obtain a proxy of the diversity of soluble compounds contained in the precipitation over the study area. Fig. 2 shows the gathered spectrum for the solution prepared after mixture two rainwater samples. Four or five categories of functional groups have usually been identified in aerosol and fog water samples (Decesari et al., 2000; Graham et al., 2002; Cavalli et al., 2006; Duarte and Duarte, 2011). The peak at 9.67 ppm can be attributed to aldehyde protons. Chemical shifts in the spectral region between 7.25 and 7.9 ppm suggest protons in aromatics with electron-attractor substituents such as -CHO and -COOH. On the other hand, the shifts at about 6.8 ppm may be attributed to hydroxyl- and methoxyl-substituents in the aromatic hydrocarbons. Signals between 3.59 and 4.36 ppm correspond to aliphatic protons adjacent to alcohols and ethers. In addition, there are signals between 2.02 and 2.77 ppm corresponding to aliphatic protons adjacent to aromatic rings or carbonyl groups. Aliphatic protons adjacent to methylene groups and/or alcohols are abundant in the 1.25–1.5 ppm spectral region. Finally, the presence of methyl groups is evidenced by the 0.93 ppm signal (Decesari et al., 2000; Suzuki et al., 2001; Pretsch et al., 2009).

The results from the ¹H NMR spectrum show that the composition of the water-soluble organic fraction in rainwater samples is in agreement with other previously published reports about the composition of aerosols in Mexico City, with aliphatic saturated and carboxylic acids groups as major components in sub-micron particles (Gilardoni et al., 2009) and other atmospheric pollutants. This suggests that composition of the water-soluble organic fraction in rainwater over the sampling site is due to the occurrence of wet deposition processes.

Báez et al. (1989, 1993, 2008) detected carbonyl compounds, especially formaldehyde, both in rainwater and particulate samples from Mexico City and other near places. They associated the presence of these compounds with anthropogenic emissions (motor vehicles exhaust). Aromatic and aliphatic acids have also been detected in rainwater samples from urban areas around the world (Chon et al., 2015). It is known that gaseous aliphatic and aromatic anthropogenic volatile organic compounds (VOCs) can produce particulate




Fig. 2. ^1H NMR spectrum at 300 MHz for the rainwater extract. The reported chemical shifts are referred to deuterated water used to dissolve the remnants after the extraction.

aliphatic and aromatic carboxylic acids and aldehydes via secondary organic aerosols (SOA) formation, but these organic acids and aldehydes can also be emitted directly by motor vehicles (Sullivan et al., 2006 and references therein) and cigarette emissions (Chon et al., 2015). Low molecular weight carboxylic acids mainly originate from vehicular exhaust but can also be produced by photo-oxidation of terpenes emitted from biological organisms (Sun and Ariya, 2006). Aliphatic acids have their origin from both primary natural (microbial activity and vegetation) as well as anthropogenic sources (biomass burning and traffic exhaust). Finally, the organic carbon in aerosol particles produced from biomass burning has been shown to be predominately aliphatic, composed mainly of oxygenated compounds with a minor content of aromatic compounds containing poly acidic compounds and phenols (Graham et al., 2002; Sullivan et al., 2006). This analysis and the report of Tsimpidi et al. (2011) indicate that the main sources of organic compounds in Mexico City are anthropogenic pollution (motor vehicles, including diesel and gasoline) and biomass burning.

3.2 TC and NPOC results

Former storage experiments at 4 °C refer that rainwater dissolved organic carbon (DOC) concentrations were stable (3%) for a week (Willey et al., 2000). On the other hand, Campos et al. (2007) reported that DOC loss in rainwater samples may be smooth or even not significant in samples with storage periods up to 21 days under the same conditions. In this study, the maximum variability for the analysis performed of a pair of samples with four weeks storage was of 17 and 25% for TC and NPOC, respectively. In the case of a sample with up to six weeks of storage, the depletion of organics (NPOC) amount was of 38%. Our results for loss of organics are comparable to those from Campos et al. (2007). Table I presents the contents of TC and NPOC in rainwater samples with a storage period up to three weeks at 4 °C. Carbon concentrations are expressed as $(\bar{x} \pm \text{unc})$, where (\bar{x}) is the mean value of the instrument measurements and *unc* is the standard deviation of the mean (SDOM) – calculated by dividing the standard deviation (σ_x) with squared root of the number of injections (\sqrt{N}) – plus the variability due to storage. For those rainwater samples with storage period up to

Table I. TC and NPOC concentrations, pH and accumulated rain data for samples collected during the field campaign. Accumulated rain is expressed in mm while carbon contents are reported in mgC L^{-1} (mean \pm *unc*). N_1 and N_2 correspond to the number of analysis performed by the analyzer.

Date	R_{ac} (mm)	pH	N_1	TC (mgC L^{-1})	N_2	NPOC (mgC L^{-1})
26/05/2016	2.9	5.3	3	24.8 ± 4.3	3	16.6 ± 4.2
02/06/2016	1.8	6.2	3	20.3 ± 3.5	3	15.1 ± 4.0
07/06/2016	7.2	4.2	3	20.6 ± 3.5	3	11.9 ± 0.2
14/06/2016	1.6	6.7	3	24.0 ± 4.2	3	15.4 ± 4.1
20/06/2016	3.3	5.1	3	21.5 ± 3.7	3	19.4 ± 4.9
08/08/2016	1.4	5.7	3	8.9 ± 1.6	3	3.5 ± 0.9
10/08/2016	4.7	5.5	3	9.1 ± 1.7	3	3.5 ± 0.9
16/08/2016	16.7	6.2	2	13.4 ± 2.4	1	3.3
24/08/2016	23.8	5.2	3	13.3 ± 2.5	3	3.4 ± 0.9
06/09/2016	2.9	7.3	3	11.8 ± 1.2	2	6.6 ± 1.9
12/09/2016	14.6	6.3	3	14.4 ± 0.6	3	3.9 ± 0.9
13/09/2016	0.4	6.8	3	14.9 ± 1.5	3	4.3 ± 0.4
19/09/2016	1	6.7	3	15.5 ± 2.7	3	7.8 ± 1.9
28/09/2016	6.9	4.4	3	11 ± 2.2	3	3.2 ± 0.8
12/10/2016	10.8	7.1	3	10.4 ± 0.5^a	3	2.5 ± 0.1

^aThis value is just below the limit of detection for TC; *unc*: sum of the standard deviation of the mean plus the variability due to sample storage.

one week, *unc* does not include the storage variability. The limit of detection for TC and NPOC are 10.6 and 2.40 mgC L^{-1} , respectively. The accumulated rain data (R_{ac}) correspond to the amount of precipitation event estimated by the TR-525M rain gauge.

Concentrations of TC and NPOC divide the samples into two different groups. The first one corresponds to samples collected between May and July with higher mean values of TC (22.2 mgC L^{-1}) and NPOC (15.7 mgC L^{-1}) compared to the samples obtained for the second period between August and October (TC, 15.7 mgC L^{-1} ; NPOC, 4.2 mgC L^{-1}). The difference is more evident in the case of NPOC, since concentrations are reduced significantly during the second period. A similar behavior between spring and summer is observed in Pan et al. (2010). Although the authors do not mention a clear justification for their observations during this particular period, they refer that emission changes from diverse sources during the study as well as variability in rain amounts may be responsible for the organic concentration differences. Moreover, the results for NPOC in Table I are higher than the continental rain (1.9 mgC L^{-1} ;

Willey et al., 2000), indicating that the occurrence of rain is associated with terrestrial and local features.

The relation NPOC/TC suggests that contribution of organic carbon to total carbon in rainwater fluctuated along the year. The proportion NPOC/TC was 0.7 on average during the first part of the campaign and it is comparable to the observed by Pan et al. (2010) but lower than the value reported by Siudek et al. (2015). Organic levels in our sampling site are comparable to those reported for other urban areas in China and Poland, although the larger concentrations gathered in Mexico City were not so large compared to those reported in the mentioned studies. The NPOC/TC ratio in the sampling site decreases to 0.3 during the August-October period; plausible explanations regarding this difference can be that air masses were more photochemically processed during the first part of the campaign (Kondo et al., 2007) or that there was a change in the source contributions for WSOC during the sampling period (Saarikoski et al., 2008). It is worthy of note that the amount of inorganic carbon, reported as water-soluble inorganic carbon ($WSIC = TC - NPOC$) in the rainwater samples for Mexico City remained similar for the whole campaign. Mexico City's

Table II. Organic concentrations in rainwater from this study compared with other studies performed at urban areas of different parts of the world. Data show the minimum and maximum values obtained for the collected samples.

Location	DOC range (mgC L ⁻¹)	Reference
Tokio, Japan	15.4 – 1.2	Sempere and Kawamura (1994)
Los Angeles, USA	18 – 2.0	Kawamura et al. (2001)
Guangzhou, China	8.3 – 0.93	Xu et al. (2008)
Northern China	44.1 – 0.3	Pan et al. (2010)
Zagreb, Croatia	7.84 – 0.19	Orlović-Leko et al. (2010)
Seoul, Korea	9.36 – 0.18	Yan and Kim (2012)
Thessaloniki, Greece	21.3 – 0.46	Balla et al. (2014)
Poznań, Poland	48.3 – 0.1	Siudek et al. (2015)
Gwangju, Korea	6.9 – 0.15	Chon et al. (2015)
Mexico City	19.5 – 2.4	This study

rainwater has an average of 7.6 ± 3.5 mgC L⁻¹ of WSIC, which is higher than the values reported for Poland and China and explains the larger NPOC/TC values for Poland (Siudek et al., 2015) respect to those found in our sampling site. The WSIC content found in this study is higher than the continental rain expected from Henry's Law and carbonic dissociation (0.2 mgC L⁻¹; Willey et al., 2000) and global average (1.9 mgC L⁻¹; Liu et al., 2004), but comparable to levels reported for urban and polluted areas (up to 10 mgC L⁻¹; Pan et al., 2010). It suggests that local sources contribute dominantly to this carbonaceous fraction.

A comparison of the results obtained in this study with organic concentrations measured in rainwater from several urban areas around the world is presented in Table II. Except for Los Angeles (USA), most of the studies used TOC analyzers for the measurements of carbon. Kawamura et al. (2001) oxidized the samples from Los Angeles with K₂Cr₂O₄ and determined the resultant CO₂ by a coulometric detector. Organic carbon (NPOC) content in rainwater samples from Mexico City shows a range comparable to dissolved organic concentrations reported from Tokyo (Japan), Los Angeles and Thessaloniki (Greece). However, the higher values of our study are higher compared to urban areas located at Guangzhou (southern China), Zagreb (Croatia), Seoul and Gwangju (Korea). Some of the rainwater samples from northern China and Poznań (Poland) exhibited much higher DOC concentrations (above 44 and 48 mg L⁻¹, respectively) than in Mexico City.

Table III shows maximum concentrations of tropospheric O₃ in the atmosphere for the sampling period of this study. It also has the average concentration of PM_{2.5} three hours before the rain event (i.e., PM_{2.5}[-3h]) – calculated from the measurements registered by the RAMA station installed at Atmospheric Sciences Center (CCA) – and the scavenging efficiencies

Table III. Maximum concentrations of O₃ (ppb) in the atmosphere, as well as 3-h mean levels of PM_{2.5} registered by the CCA station (source, RAMA website) before precipitation in the sampling site. Scavenging efficiencies estimates for particulate matter and ozone are also shown.

Date	O ₃ (ppb)	PM _{2.5} (-3h) μg m ⁻³	η _{PM_{2.5}} (%)	η _{O₃} (%)
26/05/2016	123	31.3	56.7	40.4
02/06/2016	104	24.7	64.7	29.4
07/06/2016	140	44.7	43.5	43.6
14/06/2016	98	13.7	26.7	9.7
20/06/2016	47	24.3	46.2	36.8
08/08/2016	55	7.3	25.0	39.2
10/08/2016	81	15.0	27.8	55.6
16/08/2016	62	12.0	42.9	38.2
24/08/2016	93	21.3	28.6	63.9
06/09/2016	95	15.7	14.3	25.0
12/09/2016	111	37.7	54.5	40.0
13/09/2016	91	25.3	47.6	17.6
19/09/2016	67	14.0	28.6	13.3
28/09/2016	50	20.3	44.0	42.9
12/10/2016	NA	39.3	66.1	NA

NA: not available.

(η_m) for ozone and particulate matter estimated according to Collett et al. (2008) as, $1 - (Conc_{m, \text{interstitial}}/Conc_{m, \text{pre-rain}})$ where the prefix *m* refers to O_3 and particulate matter. Reported concentrations of O_3 during the months of May and first half of June – corresponding to some of the highest pollution levels for 2016 – have a similar behavior to TC and NPOC results. Moreover, it is found that O_3 is well correlated ($R^2 = 0.69$) with $PM_{2.5}$ concentrations.

It can be observed that concentrations of the organic fraction in rainwater samples decrease as precipitation increases (Fig. 3). This behavior is similar to that reported by Pan et al. (2010) and Yan and Kim (2012), indicating that organics are washed out effectively during the initial stages of rain events. Precipitation is among the key factors controlling the pollution levels in the study area since it is considered to be the dominant scavenging mechanism in the atmosphere (Yan and Kim, 2012); therefore, it should represent an important cleansing mechanism. Associations between O_3 and $PM_{2.5}$ scavenging efficiencies with the amount of precipitation are shown in

Fig. 3b, c. Displayed results in these panels indicate higher efficiencies for O_3 and $PM_{2.5}$ scavenging as the amount of rain from an event increases, which is in agreement with the expected role of precipitation as an atmospheric cleansing process. However, there seems to be an apparent limit for the scavenging efficiency value, but the obtained data are insufficient to confirm this observation. The correlation between $PM_{2.5}$ and rainfall, however, is not clear (Fig. 3c). This weak association may be due to a continuous supply of particulate matter during rain events (Kieber et al., 2002; Yan and Kim, 2012).

Figure 4 exhibits the association of NPOC observations with sulfur dioxide (SO_2) and carbon monoxide (CO). SO_2 emissions from anthropogenic and natural sources are oxidized quickly in the atmosphere, leading to aerosol formation and acid deposition through sulfuric acid (Lee et al., 2011; He et al., 2012). Although SO_2 emission have been reduced since 2006 due to changes in fuel formulations, the main sources of SO_2 in the Mexico City area are industrial activities – especially power plants




Fig. 3. (a) NPOC concentration and O_3 and (b, c) $PM_{2.5}$ scavenging factors as function of rain amount for precipitation events collected in the sampling site.




Fig. 4. Concentrations of sulfur dioxide (red diamonds) and carbon monoxide (blue squares) vs. NPOC for the studied rain events.

and the oil refinery in Tula, Hidalgo (northwest of the city). On the other hand, carbon monoxide (CO) is emitted from methane and non-methane hydrocarbon oxidation, from fossil fuel combustion (associated with car traffic, industry and domestic heating), and from vegetation burning for agricultural purposes or from wildfires (Clerbaux et al., 2008). CO in Mexico is mainly emitted from vehicles (Benítez et al., 2014). Data displayed in Figure 4 show a trend to larger NPOC concentrations as the amount of these pollutants increases. The association between NPOC and CO is more evident compared to the relation between NPOC and SO₂, suggesting that the presence of water-soluble organic matter in the area is more dependent on local sources (vehicle emissions).

Meteorological and geographical conditions affect the evolution of pollutants (Sun et al., 2013; You et al., 2015) over the MCMA. The 48-h trajectories of air masses for 500 and 1500 m above the ground level for the studied rainstorms are shown in Figure 5. The trajectories evidence that air masses were located almost all the time over the continent, as suggested by the obtained results of organic and inorganic carbon contents in rainwater samples – which are higher than those representative for continental rain (Willey et al., 2000). With the aid of trajectory analyses, we tried to find a trend for the organic levels and pH values respect to the origin of air masses. The trajectory corresponding to 500 m above the ground was intended to look into the relation between the nearest (local) sources and

the rainwater parameters studied in this work. The motivation for the corresponding trajectories at 1500 m above the ground was to study the influence of transport of pollutant sources located away from the sampling site. Panels (a) and (b) in Fig. 6 exhibit the NPOC results (concentric circles) for the rainstorms studied in this report in function of prevalent wind from the HYSPLIT diagrams. Although most of the storms were advected from the north-northeast, as expected according to Jáuregui (2000), data for the 500 and 1500 m levels do not show a clear relationship between the air masses' origin and those cases with high ($> 10 \text{ mgC L}^{-1}$) carbon contents, since some of them arrived from north but others from other directions.

The pH varied from 4.2 to 7.3 and it is similar to the observations of Báez et al. (2007). Panels (c) and (d) in Fig. 6 display the pH values for the rainwater samples collected during the field campaign in function of wind direction for the selected height levels in the back-trajectory diagrams. Due to limited data, it is only possible to speculate about the location of pollutants sources near or within Mexico City that control the constituent levels in rainwater. A meticulous analysis of the pH and the mass air origin shows that most of the rainwater samples had pH greater than 5.6 (pH of rainwater in equilibrium with atmospheric CO₂; Shukla and Sharma, 2010) when the wind pattern had a dominant northern or northeastern component. On the other hand, there is not any specific prevailing wind direction for those rainwater samples that had pH in the acidic range.

Physical and chemical interactions between aerosols and gases with cloud droplets and raindrops represent many uncertainties for the prediction of pH in rainwater. Even though there are important sources of SO₂ in Tula, located north of the city, the pH of rainwater was not affected by these sources. Our results and other previous studies suggest that sources situated north of the sampling site have more influence on the rain neutralization processes reflected by decreasing pH. Báez et al. (2007) indicated that particles with NH₄⁺ and Ca²⁺ were among the main agents for rain neutralization in the basin of Mexico. NH₄⁺ is derived from cleaning items, industry, excrements and pet urine, and agricultural activities, whilst Ca²⁺ is sourced from soils located northeast of Mexico City. Furthermore, another plausible element for rain neutralization




Fig. 5. (a) Air-mass back trajectories randomly selected for the field campaign observed during the rainy season of 2016. (b) Continuation of (a).




Fig. 5. (a) Air-mass back trajectories randomly selected for the field campaign observed during the rainy season of 2016. (b) Continuation of (a).

may be the scavenging of ammonia emitted by more than 4.8 million vehicles, most of them with catalytic converters (Perrino et al., 2002; INEGI, 2015).

4. Summary and conclusions

The present study reports water-soluble total and organic carbon contents in rainwater samples collected from the southwestern part of Mexico City during 2016 (May–October) and their chemical characteristics through a preliminary study. ^1H NMR analysis indicates the presence of aldehydes, carboxylic and other oxygenated groups present in aromatic and aliphatic hydrocarbons. Carbon analyses were

performed using a Thermo Scientific HiPerTOC analyzer to gather data of total as well as organic carbon, measured as NPOC. Results show notable changes in concentrations of both TC and OC during May–June (first part) and August–October (second part). Mean values for TC (22.2 mgC L^{-1}) and NPOC (15.7 mgC L^{-1}) in the first part were higher compared to the second part (TC, 12.3 mgC L^{-1} ; NPOC, 4.2 mgC L^{-1}). The chemical characteristics and concentration values are comparable to organic content observed in other urban areas throughout the world. Furthermore, the NPOC/TC ratio for rainwater samples decreased during the second part of the campaign (the amount of organics render the 71% of the total carbon detected




Fig. 6. Measurements of NPOC concentrations and pH (in concentric circles) for the rainwater samples collected during the field campaign in function of wind direction for the 500 (red diamonds) and 1500 m (blue diamonds) above the ground. Vertical and horizontal lines refer to the cardinal coordinates. The wind direction corresponds to the component 6 h before rain occurrence. The grey circles in pH panels indicate the limit for neutral pH of atmospheric water with CO_2 equilibrium. Points outside this circle identify rainwater samples from alkaline events.

during the first part of the campaign but it decreased to 34% during the last one). However, the inorganic carbon concentration was 7.6 mgC L^{-1} in average and it is higher than the continental rain, suggesting that local sources contribute to this carbonaceous fraction. Analysis of air trajectories for 500 and 1500 m above the ground level did not show an association between carbon contents in rainwater and the air mass origin. It suggests that most of the carbon detected

in rain samples was produced by local sources. We observed that rainwater pH tends to be alkaline when the wind direction had a dominant north or northeast component. Similar to the observations of Báez et al. (2007), the industrial emissions and particles from sediments and soils from the northern parts of the city could be the neutralizing agents. Moreover, the scavenging of ammonia emitted by vehicles was another plausible element for rain neutralization. The organic

compounds (aromatic and alkenes) control production of daytime ozone in the MCMA (e.g., Tie et al., 2007) and we observed some of the highest NPOC levels and higher concentrations of O₃ during May and the first half of June. More data, however, is required to relate the organic concentrations in rainwater with the pollution levels in the basin.

Acknowledgments

This research was funded by the UNAM-DGA-PA-PAPIIT (IA100715) program. The authors acknowledge Ana Rosa Flores Márquez, Rocío García Martínez, José Manuel Hernández, and Alfredo Rodríguez Manjarrez for support during the field campaign, and Bertha Eugenia Mar Morales and Víctor Zarraluqui for the elaboration of the figures. The authors are also grateful to an anonymous reviewer for his valuable comments.

References

- Báez A., Belmont R., González O.G. and Rosas I.P., 1989. Formaldehyde levels in air and wet precipitation at Mexico City, Mexico. *Environ. Pollut.* 62, 153-169. DOI: 10.1016/0269-7491(89)90184-X
- Báez A., Padilla H. and Belmont R., 1993. Scavenging of atmospheric formaldehyde by wet precipitation. *Environ. Pollut.* 79, 271-275. DOI: 10.1016/0269-7491(93)90100-3
- Báez A., Belmont R., García R., Torres M.C. and Padilla H.G., 2006. Rainwater chemical composition at two sites in Central Mexico. *Atmos. Res.* 80, 67-85. DOI: 10.1016/j.atmosres.2005.06.008
- Báez A., Belmont R., García R., Padilla H. and Torres M.C., 2007. Chemical composition of rainwater collected at a southwest site of Mexico City, Mexico. *Atmos. Res.* 86, 61-75. DOI: 10.1016/j.atmosres.2007.03.005
- Báez A., Torres M.C., García R., Padilla H. and Belmont R., 2008. Measurement of carbonyls in three urban zones of the Mexico City metropolitan area and one rural zone. *Open Atmos. Sci. J.* 2, 61-67. DOI: 10.2174/1874282300802010061
- Báez A., Belmont R., García R., Padilla H.G. and Torres M.C., 2009. Trends in chemical composition of wet precipitation in Mexico City, Mexico: 1992-2007. *Open Atmos. Sci. J.* 3, 187-195. DOI: 10.2174/1874282300903010187
- Balla D., Papageorgiou A. and Voutsas D., 2014. Carbonyl compounds and dissolved organic carbon in rainwater of an urban atmosphere. *Environ. Sci. Pollut. Res.* 21, 12062-12073. DOI: 10.1007/s11356-014-3080-8
- Benítez-García S.E., Kanda I., Wakamatsu S., Okazaki Y. and Kawano M., 2014. Analysis of criteria air pollutant trends in three Mexican metropolitan areas. *Atmosphere* 5, 806-829. DOI: 10.3390/atmos5040806
- Bisutti I., Hilke I. and Raessler M., 2004. Determination of total organic carbon – An overview of current methods. *Trends Anal. Chem.* 23, 716-726. DOI: 10.1016/j.trac.2004.09.003
- Campos M.L.A.M., Nogueira R.F.P., Dametto P.R., Francisco J.G. and Coelho C.H., 2007. Dissolved organic carbon in rainwater: Glassware decontamination and sample preservation and volatile organic carbon. *Atmos. Environ.* 41, 8924-8931. DOI: 10.1016/j.atmosenv.2007.08.017
- Cavalli F., Facchini M.C., Decesari S., Emblico L., Mircea M., Jensen N.R. and Fuzzi S., 2006. Size-segregated aerosol chemical composition at a boreal site in southern Finland, during the QUEST project. *Atmos. Chem. Phys.* 6, 993-1002. DOI: 10.5194/acp-6-993-2006
- Chon K., Kim Y., Bae D.H. and Cho J., 2015. Confirming anthropogenic influences on the major organic and inorganic constituents of rainwater in an urban area. *Drink. Water Eng. Sci.* 8, 35-48. DOI: 10.5194/dwes-8-35-2015
- Clerbaux C., Edwards D.P., Deeter M., Emmons L., Lamarque J.-F., Tie X.X., Massie S.T. and Gille J., 2008. Carbon monoxide pollution from cities and urban areas observed by the Terra/MOPITT mission. *Geophys. Res. Lett.* 35, L03817. DOI: 10.1029/2007GL032300
- Coelho C.H., Francisco J.G., Nogueira R.F.P. and Campos M.L.A.M., 2008. Dissolved organic carbon in rainwater from areas heavily impacted by sugar cane burning. *Atmos. Environ.* 42, 7115-7121. DOI: 10.1016/j.atmosenv.2008.05.072
- Collett Jr. J.L., Herckes P., Youngster S. and Lee T., 2008. Processing of atmospheric organic matter by California radiation fogs. *Atmos. Res.* 87, 232 - 241. DOI: 10.1016/j.atmosres.2007.11.005
- Ćosović B., Orlović-Leko P. and Kozarac Z., 2007. Rainwater dissolved organic carbon: characterization of surface active substances by electrochemical method. *Electroanalysis* 19, 2077-2084. DOI: 10.1002/elan.200703907

- Decesari S., Facchini M.C., Fuzzi S. and Tagliavini E., 2000. Characterization of water-soluble organic compounds in atmospheric aerosol: A new approach. *J. Geophys. Res.* 105, 1481-1489. DOI: 10.1029/1999JD900950
- Duarte R.M. and Duarte A.C., 2011. A critical review of advanced analytical techniques for water-soluble organic matter from atmospheric aerosols. *Trends Anal. Chem.* 30, 1659-1671. DOI: 10.1016/j.trac.2011.04.020
- Facchini M.C., Mircea M., Fuzzi S. and Charlson R.J., 1999. Cloud albedo enhancement by surface-active organic solutes in growing droplets. *Nature* 401, 257-259. DOI: 10.1038/45758
- García R., Torres M.C., Padilla H., Belmont R., Azpra E., Arcega-Cabrera F. and Báez A., 2006. Measurement of chemical elements in rain from Rancho Viejo, a rural wooded area in the State of Mexico, Mexico. *Atmos. Environ.* 40, 6088-6100. DOI: 10.1016/j.atmosenv.2006.05.048
- Gilardoni S., Liu S., Takahama S., Russell L.M., Allan J.D., Steinbrecher R., Jimenez J.L., de Carlo P.F., Dunlea E.J. and Baumgardner D., 2006. Characterization of organic ambient aerosol during MIRAGE 2006 on three platforms. *Atmos. Chem. Phys.* 9, 5417-5432. DOI: 10.5194/acp-9-5417-2009
- Gilardoni S., Massoli P., Paglione M., Giulianelli L., Carbone C., Rinaldi M., Decesari S., Sandrini S., Costabile F., Gobbi G.P., Pietrogrande M.C., Visentin M., Scotto F., Fuzzi S. and Facchini M.C., 2016. Direct observation of aqueous secondary organic aerosol from biomass-burning emissions. *P. Natl. Acad. Sci. USA* 113, 10013-10018. DOI: 10.1073/pnas.1602212113
- Graber E.R. and Rudich Y., 2006. Atmospheric HULIS: How humic-like are they? A comprehensive and critical review. *Atmos. Chem. Phys.* 6, 729-753. DOI: 10.5194/acp-6-729-2006
- Graham B., Mayol-Bracero O.L., Guyon P., Roberts G.C., Decesari S., Facchini M.C., Artaxo P., Maenhaut W., Köll P. and Andreae M.O., 2002. Water-soluble organic compounds in biomass burning aerosols over Amazonia 1. Characterization by NMR and GC-MS. *J. Geophys. Res.* 107, 8047. DOI: 10.1029/2001JD000336
- Guidotti M., Giovinazzo R., Cedrone O. and Vitali M., 2000. Determination of organic micropollutants in rain water for laboratory screening of air quality in urban environment. *Environ. Int.* 26, 23-28. DOI: 10.1016/S0160-4120(00)00074-X
- He H., Li C., Loughner C.P., Li Z., Krotkov N.A., Yang K., Wang L., Zheng Y., Bao X., Zhao G. and Dickerson R.R., 2012. SO₂ over central China: Measurements, numerical simulations and the tropospheric sulfur budget. *J. Geophys. Res.* 117, D00K37. DOI: 10.1029/2011JD016473
- Huang X.-F., Yu J.Z., He L.-Y. and Yuan Z., 2006. Water-soluble organic carbon and oxalate in aerosols at a coastal urban site in China: Size distribution characteristics sources and formation mechanisms. *J. Geophys. Res. Atmos.* 111, D22212. DOI: 10.1029/2006JD007408
- Iavorivska L., Boyer E.W. and DeWalle D.R., 2016. Atmospheric deposition of organic carbon in precipitation. *Atmos. Environ.* 146, 153-163. DOI: 10.1016/j.atmosenv.2016.06.006
- INEGI, 2015. Instituto Nacional de Estadística y Geografía. Available at: <http://www.beta.inegi.org.mx/app/buscador/default.html?q=veh%C3%ADculos#tabMCCollapse-Indicadores> (last accessed on July 2017)
- Jáuregui E., 2000. El clima de la ciudad de México. Instituto de Geografía, UNAM/Plaza y Valdés, México, 131 pp.
- Kanakidou M., Seinfeld J., Pandis S., Barnes I., Dentener F., Facchini M.C., van Dingenen R., Ervens B., Nenes A., Nielsen C.J., Swietlicki E., Putaud J.P., Balkanski Y., Fuzzi S., Horth J., Moortgat G., Winterhalter R., Myhre C., Tsigaridis K., Vignati E., Stephanou E. and Wilson J., 2005. Organic aerosol and climate modeling: a review. *Atmos. Chem. Phys.* 5, 1053-1123. DOI: 10.5194/ACP-5-1053-2005
- Kawamura K., Steinberg S., Ng L. and Kaplan I.R., 2001. Wet deposition of low molecular weight mono- and di-carboxylic acids, aldehydes and inorganic species in Los Angeles. *Atmos. Environ.* 35, 3917-3926. DOI: 10.1016/S1352-2310(01)00207-2
- Kieber R.B., Peake B., Willey J.D. and Avery G.B., 2002. Dissolved organic carbon and organic acids in coastal New Zealand rainwater. *Atmos. Environ.* 36, 3557-3563. DOI: 10.1016/S1352-2310(02)00273-X
- Kondo Y., Miyazaki Y., Takegawa N., Miyakawa T., Weber R.J., Jiménez J.L., Zhang Q. and Worsnop D.R., 2007. Oxygenated and water-soluble organic aerosols in Tokyo. *J. Geophys. Res.* 112, D01203. DOI: 10.1029/2006JD007056
- Kucieńska B., Montero-Martínez G. and García-García F., 2010. A simulation of the influence of organic and inorganic pollutants on the formation and development of warm clouds over Mexico City. *Atmos. Res.* 95,

- 487-495. DOI: 10.1016/j.atmosres.2009.11.011
- Lee C., Martin R.V., van Donkelaar A., Lee H., Dickerson R.R., Hains J.C., Krotkov N., Richter A., Vinnikov K. and Schwab J.J., 2011. SO₂ emissions and lifetimes: Estimates from inverse modeling using in situ and global, space-based (SCIAMACHY and OMI) observations. *J. Geophys. Res.* 116, D06304. DOI: 10.1029/2010JD014758
- Liu C.J., Hannu I., Werner K., Ma X.Q., Carl J.W. and Pekka K., 2004. An estimate on the rainout of atmospheric CO₂. *J. Environ. Sci.* 1, 86-89
- Moffet R.C., Henn T.R., Tivanski A.V., Hopkins R.J., Desyaterik Y., Kilcoyne A.L.D., Tylliszczak T., Fast J., Barnard J., Shutthanandan V., Cliff S.S., Perry K.D., Laskin A. and Gilles M.K., 2010. Microscopic characterization of carbonaceous aerosol particle aging in the outflow from Mexico City. *Atmos. Chem. Phys.* 10, 961-976. DOI: 10.5194/acp-10-961-2010
- Montero-Martínez G., Zarraluqui-Such V., Caetano E.S. and García-García F., 2011. Hydrometeor vertical characterization of precipitating clouds over the Mexico Basin. *Int. J. Remote Sens.* 32, 6367-6378. DOI: 10.1080/01431161.2010.510489
- Nozière B., Kalberer M., Claeys M., Allan J., D'Anna B., Decesari S., Finessi E., Glasius M., Grgic I., Hamilton J.F., Hoffmann T., Iinuma Y., Jaoui M., Kahnt A., Kampf C.J., Kourtchev I., Maenhaut W., Marsden N., Saarikoski S., Schnelle-Kreis J., Surratt J.D., Szidat S., Szmigielski R. and Wisthaler A., 2015. The molecular identification of organic compounds in the atmosphere: State of the art and challenges. *Chem. Rev.* 115, 3919-3983. DOI: 10.1021/cr5003485
- Orlović-Leko P., Kozarac Z., Čosović B., Strmečki S. and Plavšić M., 2010. Characterization of atmospheric surfactants in the bulk precipitation by electrochemical tools. *J. Atmos. Chem.* 66, 11-26. DOI 10.1007/s10874-011-9189-y
- Pan Y., Wang Y., Xin J., Tang G., Song T., Wang Y., Li X. and Wu F., 2010. Study on dissolved organic carbon in precipitation in Northern China. *Atmos. Environ.* 44, 2350-2357. DOI: 10.1016/j.atmosenv.2010.03.033
- Perrino C., Catrambone M., di Bucchianico A.D.M. and Allegrini I., 2002. Gaseous ammonia in the urban area of Rome, Italy and its relationship with traffic emissions. *Atmos. Environ.* 36, 5385-5394. DOI: 10.1016/S1352-2310(02)00469-7
- Pretsch E., Bhuhlmann P. and Badertscher M., 2009. ¹H NMR Spectroscopy. In: Structure determination of organic compounds. Tables of spectral data (Badertscher M., Bühlmann P. and Pretsch E., Eds.). Springer-Verlag, Berlin, pp. 157-238. DOI: 10.1007/978-3-540-93810-1_4
- Putaud J., Raes F., van Dingenen R., Brüggemann E., Facchini M.C., Decesari S., Fuzzi S., Gehrig R., Hüglin C., Laj P., Lorbeer G., Maenhaut W., Mihalopoulos N., Müller K., Querol X., Rodriguez S., Schneider J., Spindler G., ten Brink H., Tørseth K. and Wiedensohler A., 2004. A European aerosol phenomenology – 2: Chemical characteristics of particulate matter at kerbside, urban, rural and background sites in Europe. *Atmos. Environ.* 38, 2579-2595. DOI: 10.1016/j.atmosenv.2004.01.041
- Saarikoski S., Timonen H., Saarnio K., Aurela M., Järvi L., Keronen P., Kerminen V.-M. and Hillamo R., 2008. Sources of organic carbon in fine particulate matter in northern European urban air. *Atmos. Chem. Phys.* 8, 6281-6295. DOI: 10.5194/acp-8-6281-2008
- Salcedo D., Onasch T.B., Dzepina K., Canagaratna, M.R., Zhang Q., Huffman J.A., DeCarlo P.F., Jayne J.T., Mortimer P., Worsnop D.R., Kolb C.E., Johnson K.S., Zuberi B., Marr L.C., Volkamer R., Molina L.T., Molina M.J., Cárdenas B., Bernabé R.M., Márquez C., Gaffney J.S., Marley N.A., Laskin A., Shutthanandan V., Xie Y., Brune W., Leshner R., Shirley T. and Jiménez J.L., 2006. Characterization of ambient aerosols in Mexico City during the MCMA-2003 campaign with aerosol mass spectrometry: Results from the CENICA supersite. *Atmos. Chem. Phys.* 6, 925-946. DOI: 10.5194/acp-6-925-2006
- Santos P.S., Otero M., Duarte R.M. and Duarte A.C., 2009. Spectroscopic characterization of dissolved organic matter isolated from rainwater. *Chemosphere* 74, 1053-1061. DOI: 10.1016/j.chemosphere.2008.10.061
- Saxena P. and Hildemann L.M., 1996. Water-soluble organics in atmospheric particles: A critical review of the literature and application of thermodynamics to identify candidate compounds. *J. Atmos. Chem.* 24, 57-109. DOI: 10.1007/BF00053823
- Seinfeld J.H. and Pandis S.N., 2006. Atmospheric Chemistry and Physics. From air pollution to climate change. John Wiley and Sons, New Jersey, pp. 761-923.

- Sempere R. and Kawamura K., 1994. Comparative distributions of dicarboxylic acids and related polar compounds in snow, rain and aerosols from urban atmosphere. *Atmos. Environ.* 28, 449-459.
DOI: 10.1016/1352-2310(94)90123-6
- Shukla S.P. and Sharma M., 2010. Neutralization of rain-water acidity at Kanpur, India. *Tellus B* 62, 172-180.
DOI: 10.1111/j.1600-0889.2010.00454.x
- Siudek P., Frankowski M. and Siepak J., 2015. Seasonal variations of dissolved organic carbon in precipitation over urban and forest sites in central Poland. *Environ. Sci. Pollut. Res.* 22, 11087-11096.
DOI: 10.1007/s11356-015-4356-3
- Stein A.F., Draxler R.R., Rolph G.D., Stunder J.B., Cohen M.D. and Ngan F., 2015. NOAA's HYSPLIT atmospheric transport and dispersion modeling system. *Bull. Amer. Meteor. Soc.* 96, 2059-2077.
DOI: 10.1175/BAMS-D-14-00110.1
- Sullivan A.P., Peltier R.E., Brock C.A., de Gouw J.A., Holloway J.S., Warneke C., Wollny A.G. and Weber R.J., 2006. Airborne measurements of carbonaceous aerosol soluble in water over northeastern United States: Method development and an investigation into water-soluble organic carbon sources. *J. Geophys. Res.* 111, D23S46. DOI: 10.1029/2006JD007072
- Sun J. and Ariya P.A., 2006. Atmospheric organic and bio-aerosols as cloud condensation nuclei (CCN): A review. *Atmos. Environ.* 40, 795-820.
DOI: 10.1016/j.atmosenv.2005.05.052
- Sun Y.L., Wang Z.F., Fu P.Q., Yang T., Jiang Q., Dong H.B., Li J., and Jia J.J., 2013. Aerosol composition, sources and processes during wintertime in Beijing, China. *Atmos. Chem. Phys.* 13, 4577-4592.
DOI: 10.5194/acp-13-4577-2013
- Suzuki Y., Kawakami M. and Akasaka K., 2001. ¹H NMR application for characterizing water-soluble organic compounds in urban atmospheric particles. *Environ. Sci. Technol.* 35, 3272-3272. DOI: 10.1021/es001861a
- Thermo Scientific, 2007. HiPerTOC User's Manual. Version 2.0.3. Delft, Netherlands.
- Tie X., Madronich S., Li G.H., Ying Z.M., Zhang R. García, A., Lee-Taylor J. and Liu Y., 2007. Characterizations of chemical oxidants in Mexico City: A regional chemical/dynamical model (WRF-Chem) study. *Atmos. Environ.* 41, 1989-2008.
DOI: 10.1016/j.atmosenv.2006.10.053
- Tsimpidi A.P., Karydis V.A., Zavala M., Lei W., Bei N., Molina L. and Pandis S.N., 2011. Sources and production of organic aerosol in Mexico City: Insights from the combination of a chemical transport model (PM-CAMx-2008) and measurements during MILAGRO. *Atmos. Chem. Phys.* 11, 5153-5168.
DOI: 10.5194/acp-11-5153-2011
- Willey J.D., Kieber R.J., Eyman M.S. and Avery G.B., 2000. Rainwater dissolved organic carbon: Concentrations and global flux. *Glob. Biogeochem. Cycles.* 14, 139-148. DOI: 10.1029/1999GB900036
- Xu T., Song Z., Liu J., Wang C., Wei J. and Chen H., 2008. Organic composition in the dry season rainwater of Guangzhou, China. *Environ. Geochem. Health* 30, 53-65. DOI: 10.1007/s10653-007-9107-z
- Yan G. and Kim G., 2012. Dissolved organic carbon in the precipitation of Seoul, Korea: Implications for global wet depositional flux of fossil-fuel derived organic carbon. *Atmos. Environ.* 59, 117-124.
DOI: 10.1016/j.atmosenv.2012.05.044
- You C., Gao S. and Xu C., 2015. Biomass burning emissions contaminate winter snowfalls in urban Beijing: A case study in 2012. *Atmos. Poll. Res.* 6, 376-381.
DOI: 10.5094/APR.2015.041.
- Zhang Q., Jiménez J.L., Canagaratna M.R., Allan J.D., Coe H., Ulbrich I., Alfarra M.R., Takami A., Middlebrook A.M., Sun Y.L., Dzepina K., Dunlea E., Docherty K., DeCarlo P.F., Salcedo D., Onasch T., Jayne J.T., Miyoshi T., Shimo A., Hatakeyama S., Takegawa N., Kondo Y., Schneider J., Drewnick F., Borrmann S., Weimer S., Demerjian K., Williams P., Bower K., Bahreini R., Cottrell L., Griffin R.J., Rautiainen J., Sun J.Y., Zhang Y.M. and Worsnop D.R., 2007. Ubiquity and dominance of oxygenated species in organic aerosols in anthropogenically-influenced Northern Hemisphere midlatitudes. *Geophys. Res. Lett.* 34, L13801.
DOI: 10.1029/2007GL029979