



Atmósfera

ISSN: 0187-6236

editor@atmosfera.unam.mx

Universidad Nacional Autónoma de México
México

Shahid, Imran; Chishtie, Farrukh; Bulbul, Gufran; Shahid,
Muhammad Zeeshaan; Shafique, Shoaib; Lodhi, Arifa
State of air quality in twin cities of Pakistan: Islamabad and Rawalpindi
Atmósfera, vol. 32, no. 1, 2019, January-March, pp. 71-84
Universidad Nacional Autónoma de México
México

DOI: <https://doi.org/10.20937/ATM.2019.32.01.06>

Available in: <https://www.redalyc.org/articulo.oa?id=56559474008>

- How to cite
- Complete issue
- More information about this article
- Journal's webpage in redalyc.org

UNAM  redalyc.org

Scientific Information System Redalyc
Network of Scientific Journals from Latin America and the Caribbean, Spain and
Portugal

Project academic non-profit, developed under the open access initiative

State of air quality in twin cities of Pakistan: Islamabad and Rawalpindi

Imran SHAHID^{1*}, Farrukh CHISHTIE², Gufran BULBUL¹, Muhammad Zeeshan SHAHID³,
Shoaib SHAFIQUE⁴ and Arifa LODHI⁴

¹ *Institute of Space Technology, Islamabad*

² *Asian Disaster Preparedness Centre, Bangkok, Thailand*

³ *Theoretical Research institute Pakistan Academy of Sciences (TRIPAS), Islamabad*

⁴ *Space and Upper Atmosphere Research Commission (SUPARCO), Pakistan*

*Corresponding author; email: imran.shahid@grel.ist.edu.pk

Received: July 10, 2017; accepted: December 4, 2018

RESUMEN

La contaminación atmosférica se ha perfilado como un desastre en países en desarrollo como Pakistán. El objetivo del presente estudio fue observar y analizar el estado de la calidad del aire en las ciudades paquistaníes gemelas de Rawalpindi e Islamabad, cuya población combinada es de 15 millones de habitantes. Se midieron las concentraciones de los principales contaminantes junto con las de CO₂ en cinco sitios dentro de esas ciudades. El muestreo se realizó durante 24 h en cada sitio, utilizando un laboratorio móvil para medir la calidad del aire. Se encontraron concentraciones totales de SO₂, NO_x y O₃ dentro de los límites permitidos de acuerdo con los estándares de la Agencia de Protección Ambiental de EUA y los Estándares Nacionales de Calidad Ambiental de Pakistán. Las concentraciones más altas de PM₁₀ se observaron en Saddar, Rawalpindi (184 µg m⁻³), en tanto que los valores mínimos se obtuvieron en la Zona Azul de Islamabad (121 µg m⁻³). Durante el periodo de estudio, las concentraciones de CO₂ permanecieron altas en casi todos los sitios, con un valor máximo de 409 ppm. En la Zona Azul y la Universidad Nacional de Ciencia y Tecnología de Islamabad se observaron concentraciones de CO₂ de hasta 385.3 y 246 ppm, respectivamente. También se estudió el comportamiento de los contaminantes con diferentes parámetros meteorológicos. Para identificar las fuentes se utilizaron trayectorias inversas generadas con el modelo HYSPLIT.

ABSTRACT

Atmospheric pollution has emerged as a disaster in developing countries like Pakistan. The aim of this study was to observe and analyze the air quality status of the twin cities of Pakistan, Rawalpindi and Islamabad, with a combined population of 15 million. The concentration of criteria pollutants along with CO₂ was measured at five different locations within these cities. Sampling was done for 24 h at each location using a mobile air quality monitoring lab. The overall concentrations of SO₂, NO_x and O₃ were found to be within the permissible limits of the US-EPA standards and the National Environmental Quality Standards of Pakistan, with higher PM₁₀ and CO₂ concentrations at most locations. The highest concentrations of PM₁₀ were observed at Saddar, Rawalpindi (184 µg m⁻³), while minimum values were observed at the Blue Area of Islamabad (121 µg m⁻³). The concentration of CO₂ during the study period was found to be high in almost all locations, with Saddar, Rawalpindi, indicating the highest value (409 ppm). The concentrations of CO₂ at the Blue Area and the National University of Science and Technology of Islamabad were observed to be as high as 385.3 and 246 ppm, respectively. The behavior of pollutants with different meteorological parameters was also studied. For source identification, backward air trajectories were also generated using the HYSPLIT model.

Keywords: PM₁₀, ozone, twin cities, air quality, urban air pollution.

1. Introduction

Clean air is necessary for healthy living, consequently a decline in the quality of air implies heavy threats to both environment and human health (Ali and Athar, 2008; Desauziers, 2004). Nowadays, major cities around the globe are facing problems with deteriorating air quality (Ali and Athar, 2008), which has serious impacts on human health, ecosystem and economy (Ilyas et al. 2009). Industrialization, unplanned urbanization, and massive volumes of traffic are major contributors to air pollution in both developing and developed nations.

Pollutants mainly responsible for such negative impacts include particulate matter (PM), CO, SO₂, NO_x and O₃ (Hopke et al., 2008; Ali and Athar, 2008). Exposure to air pollution has proven to be the origin of severe human diseases, particularly in highly polluted areas. Many studies have reported these negative effects, which range from minor to major diseases such as asthma, leukemia, lung cancer, stroke, premature births, death, etc. (Laden et al., 2000; Jonsson et al., 2004; Calderón-Garcidueñas et al., 2003; Wilhelm and Ritz, 2003; Preutthipan et al., 2004). Additionally, black carbon or “soot” that results from incomplete combustion of biomass and fossil fuels has been recognized as a substantial contributor to climate change (Seinfeld and Pandis, 1998; Shahid et al., 2014). According to a World Health Organization report (WHO, 2009), air pollution is responsible for more than two million premature deaths every year, and developing countries are more vulnerable to these impacts, since half of this disease burden is reported in their respective populations (Krzyzanowski, 2008).

The Asian region has undergone a huge economic development over the last decade with rapid urbanization, extensive energy use and industrialization. Due to this growth, air pollution has emerged as a public health hazard and has inflicted damage to the environment in the region (Kim et al., 2006; Gurjar et al., 2010; Hopke et al., 2008). Pakistan is among the countries seriously affected by air pollution. It has fast population growth, along with rapid urbanization, industrial pollution and increasing traffic loads, which are reported to be major contributors to air pollution (Pak-EPA, 2005; Ministry of Finance, 2011-2012; Colbeck et al., 2010). According to a World Bank report, more than 22 600 deaths per year are caused by impacts of air pollution at the national level

(Hopke et al., 2008; Tanveer et al., 2015). Dust and smoke in Pakistan are generally twice the world average and typically five times higher than levels in the developed countries. Therefore it is important to study the air quality of different cities on a continuous basis, in order to evaluate their likely impacts on environment and human health, and most importantly, to establish a baseline that will highlight possible sources and identify mitigation and abatement measures accordingly (Krzyzanowski, 2008; Lenschow et al., 2001).

Bulbul et al. (2017) reported the morphology and concentrations of PM₁₀ in Islamabad during the winter fog period. Shahid et al. (2015a, b) reported the sectorial contribution of aerosol in Pakistan and Usman-Alvi et al. (2017) reported traffic and industry related exposure to air quality in an industrialized city. Khokhar et al. (2016) studied the temporal variability of aerosol during winter while Zeb et al. (2018) reported the morphology of PM in urban environments of Pakistan. Awais et al. (2018) studied the optical properties of aerosols using remote sensing over the twin cities of Pakistan. Based on the previous studies and available information, in this study we assessed the status of air quality of Rawalpindi and Islamabad, where five major air pollutants were studied, i.e. PM₁₀, SO₂, NO₂, O₃ and CO₂. The concentrations of these pollutants were measured in different locations within the above-mentioned cities and then compared with the standards of the National Environmental Quality Standards (NEQS) of Pakistan and the United States Environmental Protection Agency (US-EPA) (Table I).

Table I. Meteorological parameters at the study site.

Date	Temperature (°C)	Average wind direction	Wind speed (m s ⁻¹)
4/Mar/2014	14	NNE	8.5
5/Mar/2014	16	NNW	9.5
6/Mar/2014	15.5	NW	9.5
7/Mar/2014	15	NE, NW	6
8/Mar/2014	17	WSW, NE	7
9/Mar/2014	15.5	NE	14

2. Materials and methods

2.1 Site description

Islamabad (33.7294° N, 73.0931° E) is the federal capital of Pakistan, adjacent to Rawalpindi (33.5984° N,

73.0441° E). They are commonly referred as twin cities and are often considered as a single unit. The Rawalpindi-Islamabad metropolitan area is considered the third largest in Pakistan, with a population over 15 million. Both cities are located approximately 500–540 masl. Rawalpindi-Islamabad lay in a semi-arid zone with hot summers followed by monsoons and cold winters. The twin cities have with five seasons: winter (November–February), spring (March–April), summer (May–June), rainy monsoon (July–August) and autumn (September–October).

2.2 Monitoring and sampling

The concentration of major air pollutants was measured at selected locations within the twin cities with sampling conducted by a mobile air quality monitoring equipped with the latest instruments capable of measuring major criteria pollutants within a given location and time, including onboard analyzers for NO_x , SO_x , O_3 , CO_2 and PM (in compliance with the US-EPA standard 40 CFR -53). A total of five locations were selected based on their situation and nearby areas. Two sites in Rawalpindi and two sites in Islamabad were chosen along with the Islamabad international airport (Fig. 1). All parameters were monitored continually for 24 h at every location. Hourly means were also calculated to have a clear picture of pollutant concentrations at different hours within the sampling time.




Fig. 1. Study area with the selected sites.

3. Results and discussion

Concentrations of ambient air pollutants (PM_{10} , SO_2 , NO_2 , CO_2 and O_3) were measured at five locations in Rawalpindi-Islamabad from March 4–9, 2014. Table I shows the meteorological conditions in which the measurements were conducted. The sampling was done for 24 h at each site. The concentrations of four pollutants, namely PM_{10} , SO_2 , NO_2 , and O_3 , were measured in micrograms per cubic meter while CO_2 levels were expressed in parts per million (ppm). The concentrations of measured pollutants were also compared with NEQS and US-EPA standards for ambient air pollutants (Table II). Overall, concentrations of most pollutants were found to be within the permissible range of the above-mentioned standards, except for PM_{10} in some locations.

Table II. Limits of air pollutant concentrations according to PAK- and US-EPA.

Parameter	PAK-EPA ($\mu\text{g m}^{-3}$)	US-EPA ($\mu\text{g m}^{-3}$)	Average time (h)
O_3	130	235	1
CO	5000	10 300	8
SO_2	120	365	24
NO_2	80	110	24
PM_{10}	150	150	24

3.1 Particulate matter (PM_{10})

PM is a mix of liquid droplets and solid particles suspended in the air. In recent years, scientists have been focusing on the identification of PM sources and understanding the atmospheric chemical processes that lead to its transport and removal. PM contains major components that comprise a large proportion of the total particle mass, and trace elements that are usually less than 1 % of the total particle mass. According to previous studies, elemental/black carbon, organic matter, nitrate, sulphate, and ammonium are the main PM contributors.

PM_{10} was monitored at selected locations within the twin cities as shown in Figure 1. The highest values for PM_{10} were observed in Saddar, Rawalpindi ($184 \mu\text{g m}^{-3}$), while the minimum values were observed at the Blue Area of Islamabad ($121 \mu\text{g m}^{-3}$). PM_{10} values observed at the National University of Science and Technology (NUST) and the

Islamabad airport were found to be 142 and $135 \mu\text{g m}^{-3}$, respectively. These results are shown in Figure 2.




Fig. 2. Average PM₁₀ concentrations with standard deviation (Mean + SD) at the NUST, Blue Area, Faizabad, Saddar and Airport sites.

In this study, the sites in Rawalpindi (Faizabad and Saddar) showed the highest values. Both locations are congested areas with high traffic flow throughout the day. The Faizabad site is a junction between the twin cities and is one of their busiest roads. It is an entrance point for Rawalpindi and Islamabad from different cities as well as from nearby rural areas. It also serves as a gateway toward the northern region of the country, as traffic coming from Karachi, Lahore, Faisalabad and other big cities passes through this interchange. Moreover, the dense traffic of this site includes heavy trucks and other vehicles. The emissions from vehicular exhaust triggers higher atmospheric levels of pollutants, especially emissions from vehicles driven on diesel. The industrial area of Islamabad (I-9, I-10) is also located near the Faizabad interchange. Emissions from industrial facilities also drive PM into the ambient air of the area. The adjacent rural area, which includes Sowam and other villages, has undergone rapid urbanization during the last couple of years and is still undergoing expansion. Construction activities have become a source of dust and particulate pollution. These small villages and rural areas do not have the facility of natural gas; in fact, they depend on wood for cooking purposes. Wood stoves and biomass burning in the surrounding villages also contribute to PM pollution.

Location 4, Saddar, showed the highest PM concentrations among all the studied areas. It is one of the busiest areas in the twin cities. The area itself is popular for shopping malls, hotels, restaurants and hospitals. For this reason, roads around the area are always full of traffic, which particularly includes a large quantity of two-stroke vehicles, which account for higher PM pollution rates. Another important driving factor for the elevated PM values is the dust from a construction site for the subway and bus station. Heavy traffic and intense construction activities took place during the study period, which may have been the contributing factor for PM pollution.

Hourly average PM₁₀ concentrations were also calculated in order to analyze hourly variations of the pollutant, as shown in Figure 3.

Overall PM₁₀ concentrations at all locations showed elevated values between 8:00 LT and 10:00 LT, lapse in which many people travel from different regions to their offices and schools. High values of PM were also observed between 15:00-18:00 LT, when people return home from offices and schools. Another factor that contributed to high PM concentrations at this time was heavy traffic arriving to the twin cities, since the city administration and the capital's traffic police allowed heavy traffic into the area around 17:00 LT. Possible sources of PM at this location include emissions from traffic, industries (I-9, I-10), windblown dust and biomass burning. The levels of the PM in the road dust were highly dependent on traffic volume and atmospheric spreading from traffic rotaries. Additional factors that affect PM concentration include vehicle speed, frequency of brake use and industrial emissions levels in the areas. Cemented highways usually had higher pollution levels than asphalt highways.

3.2 Sulphur dioxide

SO₂ concentrations were observed at five locations within the twin cities. The overall concentration of SO₂ was found to be within the limits of NEQS and US-EPA standards. Location 3 (Faizabad) showed higher values as compared to the other sites ($40.4 \mu\text{g m}^{-3}$). SO₂ concentrations at other locations (Saddar, Airport, Blue Area and NUST) were 28.9, 27.1, and 22, $22.4 \mu\text{g m}^{-3}$, respectively, as shown in Figure 4.

Areas within Islamabad comparatively showed lower SO₂ concentrations. This might be due to less




Fig. 3. Hourly trends of PM₁₀ concentrations ($\mu\text{g m}^{-3}$) at Faizabad and Saddar.




Fig. 4. SO₂ concentrations with standard deviation (mean + SD) at NUST, Blue Area, Faizabad, Saddar and Airport sites.

traffic loads and pollution within the city. Hourly average SO₂ values were also calculated, which shows variations at different timings and is presented in Figure 5.

The NO₂ concentrations were found to be within the permissible limits of NEQS and US-EPA. Concentrations within the twin cities were found to be

within the range of 29.6 to 36.3 $\mu\text{g m}^{-3}$ as shown in Figure 6.

The highest concentration was observed at location 5 (Benazir Bhutto International Airport) while the minimum was observed at Faizabad, Islamabad.

The values of NO₂ were also observed on an hourly basis and are shown in Figure 7. Higher values are reported in most locations around 9:00-11:00 LT in the morning and 15:00-18:00 LT in the evening. The influx of heavy traffic was observed on the road at these intervals, in which the higher values indicate that emissions from traffic are the main source for NO₂ pollution.

3.4 Ozone

An O₃ monitor (APOA-370, Horiba) with detection limit of 0.5 ppb and range of 0-1 ppm was used to determine O₃ concentrations in ambient air. This monitor works on the principle of UV photometry. The overall concentration of O₃ was found to be within the permissible limits of NEQs and USEPA.




Fig. 5. Hourly trends of SO₂ concentrations (µg m⁻³) at the Faizabad and Blue Area sites.




Fig. 6. NO₂ concentrations with standard deviation (mean + SD) at the NUST, Blue Area, Faizabad, Saddar and Airport sites.

Among the monitored sites, NUST, Benazir international airport and Faizabad showed peak values as compared to others (26.1, 24.3 and 20.9 µg m⁻³, respectively as shown in Figure 8.

Concentrations of O₃ at other sites were comparatively lower. The O₃ concentration at Saddar was 18.3 µg m⁻³, while the Blue Area of Islamabad had the lowest values among all other sites (13.1 µg m⁻³).

Previous studies in Islamabad and nearby areas also reported that in most parts of the twin cities the O₃ concentrations are within the permissible limits. Only a few research articles highlighted the seasonal variations of O₃, and reported higher values during the summer. The hourly behavior of O₃ has also been studied, showing the same pattern as other pollutants, i.e., higher values between 8:00-11:00 LT and 15:00-18:00 LT, as shown in Figure 9.

3.5 Carbon dioxide

Ambient CO₂ was also monitored at the five different locations within the twin cities. CO₂ is one of the main greenhouse gases (GHGs) responsible for climate change. Hence, it is especially important to monitor its status in the ambient air quality.

The CO₂ concentration during the study period was found to be higher in almost all locations. The Saddar, Rawalpindi area, showed the highest value (409 ppm), as shown in Figure 10. This is due to heavier traffic on congested roads, given that




Fig. 7. Hourly trend of NO_2 concentrations ($\mu\text{g m}^{-3}$) at the Faizabad and Saddar sites.




Fig. 8. Ozone concentrations with standard deviation (mean + SD) at the NUST, Blue Area, Faizabad, Saddar and Airport sites.

vehicular emissions are probably the main source of CO_2 , followed by the Westridge industrial area, which also drives its concentrations. CO_2 concentrations at the Blue Area of Islamabad and NUST were found to be 385.3 and 246.0 ppm, respectively. Elevated values in the Blue Area are attributable to traffic emissions, since many people visit this vicinity in

office hours due to the availability of many local and international offices.

Concentrations of CO_2 at the Faizabad and Airport sites have values of 375.9 and 368.8 ppm, respectively. Overall, CO_2 concentrations seem to be high in all the sites. Hourly average concentrations showed that, throughout the day, CO_2 concentrations remained high, as shown in Figure 11.

3.6 Meteorological conditions

Different meteorological parameters were also observed during the study periods, which include humidity, temperature, average wind direction and wind speed. To find out the probable source of different measured pollutants, backward air trajectories were generated using the Hybrid-Single Particle Lagrangian Integrated Trajectory (HYSPLIT) model, generated by the US National Oceanic and Atmospheric Administration's (NOAA) Air Resources Laboratory (ARL) website. Archives from the Global Data Assimilation System (GDAS) of the National




Fig. 9. Hourly trends of ozone concentrations ($\mu\text{g m}^{-3}$) at the NUST and Airport sites.




Fig. 10. CO_2 concentrations with standard deviation (mean + SD) at the NUST, Blue Area, Faizabad, Saddar and Airport sites.

Center for Environmental Prediction/National Center for Atmospheric Research (NCEP/NCAR) was used to generate the backward trajectories, which were calculated for a different heights, i.e., 500, 1000, and 1500 m above ground level for different time scales. The label interval was set to 6 h to track the path of trajectory.

The back-trajectory analysis using NOAA HY-SPLIT shows that during the study period, the northeast

and northwest were important source areas for trajectories that reached Islamabad (Fig. 12).

Wind roses were also plotted using the NOAA air resource lab data (Fig. 13). These wind roses and air trajectories show a general wind pattern from the NW and NE and infrequently from the WS at Faizabad, an area in which the wind direction was along the NE direction. The air inflow blows from Sowan and nearby areas. Sowan and other villages have undergone a rapid urban growth in the last couple of years and are still expanding. Currently, construction activities are likely sources of dust and particulate pollution. At Saddar area, the wind blows from a nearby area in Westridge, located on the Peshawar road. Also, windblown dust comes from road dust and heavy traffic on the major roads like the Kashmir highway, the IJP road, and the Peshawar road.

Monthly mean wind magnitudes and vectors (m s^{-1}) from the MERRA 2 satellite data at a 10 m height during February and March 2014 were also plotted to regional wind patterns during the study period (Fig. 14).




Fig. 11. Hourly trends for CO₂ concentrations ($\mu\text{g m}^{-3}$) at the Faizabad and Saddar sites.




Fig. 12. Backward trajectories from the NOAA HYSPLIT model.




Fig. 13. Wind rose plot for wind direction and speed during the study period.




Fig. 14. Monthly mean wind magnitude and vectors (m s^{-1}) from the MERRA 2 satellite data at a 10 m height during February and March 2014.

Pollutants are influenced by meteorological conditions such as temperature, wind direction, wind speed, and relative humidity. In this study, the Modern Era Retrospective-Analysis for Research and Applications (MERRA-2) meteorological database was used at the annual, seasonal and synoptic scales. Surface air temperature is shown in Figure 15.

Monthly trends of the observed surface area temperature in the study area from 2011 to 2014 are very similar in the different years of the period (Fig. 15). Surface area temperature is lower in winter and elevates from April until July. The maximum surface area temperature can be seen during June in each year.

Precipitation data from 2011 to 2014 showed higher values during the monsoon season (July-August), while lower values are observed in the post-monsoon season (September-December). In January-March higher precipitation rates are observed in all the years, as shown in Figure 16. The pre-monsoon season (April to June) shows comparatively lower precipitation values. Overall, higher precipitation values are observed throughout 2013, followed by 2011. The years 2012 and 2014 have lower precipitation values compared to 2011 and 2013. Monthly precipitation trends show lower values in the pre-monsoon season and higher values during the month of monsoon. The

annual comparison of precipitation data depicts the same trends in all years.

Yearly data from the Moderate Resolution Imaging Spectroradiometer (MODIS) derived aerosol optical depth (AOD) for 2011-2014 is also plotted in Figure 17. Data for each month is plotted in the graph for each respective year. Higher values of AOD are observed in June-August (summer). MODIS AOD started declining after August, which is due to monsoon rains in this month. Higher AOD values are typically observed in June-August. Here it can be inferred that ground PM concentrations and MODIS AOD are relatively higher in June-August. During 2014, we found elevated levels of AOD from May to July in the study area. In contrast, levels of AOD are comparatively lower in March. Seasonally, the variation in MODIS AOD data shows higher values in summer as compared to winter. Trends in the yearly data remained almost similar from 2011 to 2014 in Islamabad.

From the meteorological data we surmise that higher pollutants levels are observed during summer, mainly in June-August. Higher levels of MODIS AOD are also found in these months. The meteorological trends also suggest that levels of pollutants might be lower during winter. The AOD data




Fig. 15. MERRA-2 surface air temperature (°C) over Islamabad




Fig. 16. MERRA-2 total surface precipitation (mm) over Islamabad.




Fig. 17. MODIS AOD over Islamabad.

validates this observation. Levels of MODIS AOD are low in winter. These trends are highly similar from 2011 to 2014.

4. Conclusions

Most cities in Pakistan are rapidly expanding due to migration from rural to urban areas in search of job and facilities, which leads to environmental degradation and rise in air pollution levels. Worsening air quality levels are also due to the lack of efficient mass transit systems, and endurance of old trucks and two

strokes engines. Many government and international organizations have highlighted the worsening air quality issues in major cities of Pakistan. In this study, our aim was to provide air quality measurements of twin cities Islamabad and Rawalpindi.

The levels of SO_2 at most studied sites were within the permissible limits. Location 3 (Faizabad) showed higher values ($40.4 \mu\text{g m}^{-3}$) as compared to other sites. SO_2 concentrations at other locations, i.e. Saddar, Airport, Blue Area and NUST, were 28.9, 27.1, 22 and $22.4 \mu\text{g m}^{-3}$, respectively.

The overall concentration of NO_2 was found to be within the permissible limits of NEQS and US-EPA. Concentrations within the twin cities were found to be within the range of 29.6 to $36.3 \mu\text{g m}^{-3}$. The highest values were observed at location 5, namely the Benazir Bhutto International Airport, while the minimum were observed at Faizabad, Islamabad.

The overall concentration of O_3 was found to be within the permissible limits of NEQS and USEPA. Among the monitored sites, NUST, Benazir International Airport and Faizabad showed peak values as compared to others (26.1 , 24.3 and $20.9 \mu\text{g m}^{-3}$, respectively). Concentrations of O_3 at other sites were comparative lower. The O_3 concentration at Saddar was $18.3 \mu\text{g m}^{-3}$ while blue area Islamabad has lowest values among all other sites, i.e., $13.1 \mu\text{g m}^{-3}$.

Concentrations of CO₂ during the study period were found to be higher in almost all locations. Sadder, Rawalpindi showed the highest value (409 ppm), which is due to heavier traffic on congested roads. Vehicular emissions are probably the main source for the high levels of CO₂, followed by the Westridge industrial area, which is also driving CO₂ concentrations. Concentrations at the Blue Area and NUST in Islamabad were observed to be 385.3 and 246.0 ppm, respectively. At the Faizabad and Airport sites, CO₂ values were 375.9 and 368.8 ppm, respectively.

The highest PM₁₀ values were reported in Sadder, Rawalpindi (184 ppm) while the minimum were observed at the Blue Area of Islamabad (121 ppm). PM₁₀ values observed at the NUST and Airport sites were found to be 142 and 135 ppm, respectively.

Having in mind the limited data on air pollution from the region, the objective of this study was to monitor the levels of air pollutants in the twin cities, caused by vehicular exhausts, industrial emissions and other sources. Our study revealed that most of the pollutant levels in the twin cities are within the permissible limits of US-EPA and NEQS. The levels of PM₁₀ concentrations were found to be higher in most of the sites. This study indicates that much more comprehensive studies are needed to further understand air quality levels and related issues across Pakistan, in particular, urban centers such as Islamabad and Rawalpindi.

Acknowledgments

The authors acknowledge the support of the National University of Science and Technology (NUST) Islamabad for providing a location for air quality monitoring at its main campus.

References

- Ali, M. and Athar M., 2008. Air pollution due to traffic, air quality monitoring along three sections of national highway N-5, Pakistan. *Environ. Monit. Assess.* 136, 219-226. DOI: 10.1007/s10661-007-9677-3
- Awais M., Shahzad M.I., Nazeer M., Mahmood I., Mehmood S., Iqbal M.F., Yasmin N. and Shahid I., 2018. Assessment of aerosol optical properties using remote sensing over highly urbanized twin cities of Pakistan. *J. Atmos. Sol.-Terr. Phys.* 173, 37-49. DOI: 10.1016/j.jastp.2018.04.008
- Bulbul G., Shahid I., Chishtie F., Shahid M.Z., Hundal R.A., Zahra F. and Shahzad M.I., 2017. PM₁₀ Sampling and AOD trends during 2016 winter fog season in the Islamabad region. *Aerosol Air Qual. Res.* 18, 188-199. DOI: 10.4209/aaqr.2017.01.0014
- Calderón-Garcidueñas L., Mora-Tiscareño A., Fordham L.A., Valencia-Salazar G., Chung C.J., Rodríguez-Alcaraz A., Paredes R., Variakojis D., Villarreal-Calderón A., Flores-Camacho L., Antunez-Solís A., Henríquez-Roldán C. and Hazucha M.J., 2003. Respiratory damage in children exposed to urban pollution. *Pediatr. Pulmonol.* 36, 148-61. DOI: 10.1002/ppul.10338
- Colbeck I., Nasir Z.A., and Ali Z., 2010. The state of indoor air quality in Pakistan—A review. *Environ. Sci. Pollut. R.* 17, 1187-1196. DOI: 10.1007/s11356-010-0293-3
- Desauziers V., 2004. Traceability of pollutant measurements for ambient air monitoring, TrAC-Trend. *Anal. Chem.* 23, 252-260. DOI: 10.1016/S0165-9936(04)00310-3
- Gurjar, B.R., Jain A., Sharma A., Agarwal A., Gupta P., Nagpure A.S. and J. Lelieveld, 2010. Human health risks in megacities due to air pollution. *Atmos. Environ.* 44, 4606-4613. DOI: 10.1016/j.atmosenv.2010.08.011
- Hopke P.K., Cohen D.D., Begum B.A., Biswas S.K., Ni B., Pandit G.G., Santoso M., Chung Y.S., Davy P., Markwitz A., Waheed S., Siddique N., Santos F.L., Pabroa P.C.B., Seneviratne M.C.S., Wimolwattana-pun W., Bunprapob S., Vuong T.B., Duy Hien P. and Markowicz A., 2008. Urban air quality in the Asian region. *Sci. Total Environ.* 404, 103-112. DOI: 10.1016/j.scitotenv.2008.05.039
- Ilyas S.Z., Khattak A.I., Nasir S.M., Tabnak Q., Rehana D., 2009. Air pollution assessment in urban areas and its impact on human health in the city of Quetta, Pakistan. 291-299. *Clean Technol. Envir.* 12, 291-299. DOI: 10.1007/s10098-009-0209-4
- Jonsson P., Bennet C., Eliasson I. and Selin Lindgren E., 2004. Suspended particulate matter and its relations to the urban climate in Dar es Salaam, Tanzania. *Atmos. Environ.* 38, 4175-4181. DOI: 10.1016/j.atmosenv.2004.04.021
- Kim Oanh N.T., Upadhyay N., Zhuang Y.H., Hao Z.P., Murthy D.V.S., Lestari P., Villarin J.T., Chengchua K., Co H.X., Dung N.T. and Lindgren E.S., 2006. Particulate air pollution in six Asian cities: Spatial and temporal distributions, and associated sources. *Atmos. Environ.* 40, 3367-3380. DOI: /10.1016/j.atmosenv.2006.01.050

- Khokhar M.F., Yasmin N., Chishtie F. and Shahid I., 2016. Temporal variability and characterization of aerosols across the Pakistan region during the winter fog periods. *Atmosphere* 7, 67. DOI: 10.3390/atmos7050067
- Krzyzanowski M., 2008. WHO Air Quality Guidelines for Europe. *J. Toxicol. Environ. Health A* 71, 47-50. DOI: 10.1080/15287390701557834
- Laden F., Neas L.M., Dockery D.W. and Schwartz J., 2000. Association of fine particulate matter from different sources with daily mortality in six U.S. cities. *Env. Health Persp.* 108, 941-947. DOI: 10.1289/ehp.00108941
- Lenschow P., Abraham H.J., Kutzner K., Lutz M., Preuss J.D. and Reichenbacher W., 2001. Some ideas about the sources of PM₁₀. *Atmos. Environ.* 35, 25-33. DOI: 10.1016/S1352-2310(01)00122-4
- Ministry of Finance, 2011-2012. Economic Survey of Pakistan. Economic Advisor Wing, Government of Pakistan.
- Pak-EPA, 2005. State of the Environment Report. Pakistan Environment Protection Agency, Ministry of the Environment, Government of Pakistan.
- Preutthipan A., Udomsubpayakul U., Chaisupamongkol-larp T. and Pentamwa P., 2004. Effect of PM₁₀ pollution in Bangkok on children with and without asthma. *Pediatr. Pulmonol.* 37, 187-192. DOI: 10.1002/ppul.10398
- Seinfeld J.H. and Pandis S.N., 1998. Atmospheric chemistry and physics: From air pollution to climate change. 2nd ed. Wiley, 1360 pp.
- Shahid I., Kistler M., Mukhtar A., Ramírez-Santa Cruz C., Bauer H. and Puxbaum H., 2014. Chemical composition of particles from traditional burning of Pakistani wood species. *Atmos. Environ.* 111, 113-126. DOI: 10.1016/j.atmosenv.2015.01.041
- Shahid M.Z., Hong L., Yu-Lu Q. and Shahid I., 2015a. Source sector contributions to aerosol levels in Pakistan. *Atmos. Ocean. Sci. Lett.* 8 (5), 308-313. DOI: 10.3878/AOSL20150049
- Shahid M.Z., Liao H., Li J., Shahid I., Lodhi A. and Man-sha M., 2015b. Seasonal variations of aerosols in Pakistan: Contributions of domestic anthropogenic emissions and transboundary transport. *Aerosol Air Qual. Res.* 15, 1580-1600. DOI: 10.4209/aaqr.2014.12.0332
- Tanveer Z., Ilyas H., Akhtar S.N. and Shaffique K., 2015. Outdoor air pollution: A case study of Gujrat City. *J. Glob. Sci. Issues* 3, 57-65.
- Usman-Alvi M., Chishtie F., Shahid I., Mahmud T. and Hussain R., 2017. Traffic-and industry-related air pollution exposure assessment in an Asian megacity. *Clean-Soil Air Water* 46, 1600773. DOI: 10.1002/clen.201600773
- WHO, 2009. Global health risks: Mortality and burden of disease attributable to selected major risks. World Health Organization, Geneva, 62 pp.
- Wilhelm. M. and Ritz B., 2003. Residential proximity to traffic and adverse birth outcomes in Los Angeles County, California, 1994-1996. *Environ. Health Perspect.* 111, 207-216. DOI: 10.1289/ehp.5688
- Zeb B., Alam K., Sorooshian A., Blaschke T., Ahmad I. and Shahid I., 2018. On the morphology and composition of particulate matter in an urban environment. *Aerosol Air Qual. Res.* 18, 1431-1447. DOI: 10.4209/aaqr.2017.09.0340