



ATMOSFERA

Atmósfera

ISSN: 0187-6236

claudio.amescua@atmosfera.unam.mx

Universidad Nacional Autónoma de México
México

Antonio Reyna, Marco; Schwander, Stephan; López Avitia, Roberto;
Bravo-Zanoguera, Miguel Enrique; Reyna, Myrtha Elvia; Nava,
Martha Lorena; Siqueiros, Miriam; Osornio-Vargas, Álvaro Román
**Particulate matter air pollution effects on pulmonary tuberculosis
activation in a semi-desert city on the US-Mexican border**

Atmósfera, vol. 35, núm. 3, 2022, Mayo, pp. 545-556

Universidad Nacional Autónoma de México

Distrito Federal, México

DOI: <https://doi.org/10.20937/ATM.52852>

Disponible en: <https://www.redalyc.org/articulo.oa?id=56581835007>

- Cómo citar el artículo
- Número completo
- Más información del artículo
- Página de la revista en redalyc.org

redalyc.org

Sistema de Información Científica Redalyc

Red de revistas científicas de Acceso Abierto diamante

Infraestructura abierta no comercial propiedad de la academia

Particulate matter air pollution effects on pulmonary tuberculosis activation in a semi-desert city on the US-Mexican border

Marco Antonio REYNA^{1*}, Stephan SCHWANDER², Roberto LÓPEZ AVITIA¹,
Miguel Enrique BRAVO-ZANOQUERA¹, Myrtha Elvia REYNA³, Martha Lorena NAVA⁴,
Miriam SIQUEIROS¹ and Álvaro Román OSORNIO-VARGAS⁵

¹ *Cuerpo Académico de Bioingeniería y Salud Ambiental de la Universidad Autónoma de Baja California. Blv. Benito Juárez y calle de la Normal S/N, Colonia Insurgentes Este C.P. 21280, Mexicali, B.C., México.*

² *Departments Urban-Global Public Health & Environmental and Occupational Health, Rutgers School of Public Health, Piscataway, NJ 08854, USA.*

³ *Department of Pediatrics, The Hospital for Sick Children, Toronto, Canada.*

⁴ *Departamento de Epidemiología del Instituto Mexicano del Seguro Social, Mexicali, B.C., México.*

⁵ *Department of Pediatrics, University of Alberta, 3-591 ECHA, 11405 87th Ave. Edmonton T6G 1C9, Canada.*

*Corresponding author: mreyna@uabc.edu.mx

Received: March 11, 2020; accepted: January 8, 2021

RESUMEN

En este artículo evaluamos las asociaciones (riesgo relativo, RR) entre la exposición a PM₁₀ y PM_{2.5} (continua y por categorías de exposición baja o alta) en la incidencia de Tuberculosis Pulmonar (PTB) en Mexicali, Baja California, México. Las asociaciones se exploraron en forma semanal con y sin efectos de latencia utilizando la regresión múltiple de Poisson. Observamos un efecto retardado de 10 semanas para PM₁₀ y PM_{2.5} en todos los casos de TBP y en el grupo de hombres. En las mujeres el efecto retardado es de 11 semanas. En todos los casos de TBP, el RR aumentó en 2.4% (IC del 95%: 2.1, 2.6, p<0.10) por cada incremento de 10 µg/m³ de PM₁₀ en la exposición continua a la contaminación y en 3.6% (IC: 3.3, 4.0, p<0.05) en la categoría de exposición a contaminación alta, y en 3.2% (IC: 2.9, 3.4, p<0.05) por cada incremento de 10 µg/m³ de PM_{2.5} en la exposición continua y en 3.9% (IC: 3.6, 4.3, p<0.05) en la categoría de exposición alta. En hombres, el RR aumentó en 2.8% (IC: 2.5, 3.1, p<0.10) por cada incremento de 10 µg/m³ de PM₁₀ en la exposición continua y en 4.6% (IC: 4.2, 5.0, p<0.05) en la categoría de alta exposición, y en 3.4% (IC: 3.1, 3.7, p<0.05) por cada incremento de 10 µg/m³ de PM_{2.5} en la exposición continua y en 4.2% (IC: 3.8, 4.6, p<0.05) en la categoría de alta exposición. En mujeres, el RR aumentó en 5.1% (IC: 4.7, 5.5, p<0.05) por cada incremento de 10 µg/m³ de PM₁₀ en la exposición continua y en 5.3% (IC: 4.7, 5.8, p<0.10) en la categoría de alta exposición, y en 4.3% (IC: 3.8, 4.8, p<0.10) por cada incremento de 10 µg/m³ de PM_{2.5} en la exposición continua y en 5.3% (IC: 4.8, 5.9, p<0.10) en la categoría de alta exposición. La contaminación del aire por PM parece estar asociada con la incidencia de PTB en la población de Mexicali.

ABSTRACT

In this paper we assessed the association (relative risk, RR) between the exposure to PM₁₀ and PM_{2.5} (as a continuous variable and as categories of low or high pollution exposure) on the incidence of pulmonary tuberculosis (PTB) in Mexicali, Baja California, Mexico. We used a weekly, lagged multiple Poisson regression model. We observed a 10-week delayed effect for PM₁₀ and PM_{2.5} in all PTB cases and in male cases with PTB. An 11-week delayed effect occurred in the female PTB cases. For all the PTB cases, the RR rose by 2.4% (95% CI: 2.1, 2.6, p<0.10) for each 10 µg/m³ increase of PM₁₀ in the continuous exposure and by 3.6% (CI: 3.3, 4.0, p<0.05) in the high pollution exposure category, and by 3.2% (CI: 2.9, 3.4, p<0.05) for each 10 µg/m³ increase of PM_{2.5} in the continuous exposure and by 3.9% (CI: 3.6, 4.3, p<0.05) in the high pollution

exposure category. In men, the RR rose by 2.8% (CI: 2.5, 3.1, $p < 0.10$) for each $10 \mu\text{g}/\text{m}^3$ increase of PM_{10} in the continuous exposure and by 4.6% (CI: 4.2, 5.0, $p < 0.05$) in the high pollution exposure category, and by 3.4% (CI: 3.1, 3.7, $p < 0.05$) for each $10 \mu\text{g}/\text{m}^3$ increase of $\text{PM}_{2.5}$ in the continuous exposure and by 4.2% (CI: 3.8, 4.6, $p < 0.05$) in the high pollution exposure category. In women, the RR rose by 5.1% (CI: 4.7, 5.5, $p < 0.05$) for each $10 \mu\text{g}/\text{m}^3$ increase of PM_{10} in the continuous exposure and by 5.3% (CI: 4.7, 5.8, $p < 0.10$) in the high pollution exposure category, and by 4.3% (CI: 3.8, 4.8, $p < 0.10$) for each $10 \mu\text{g}/\text{m}^3$ increase of $\text{PM}_{2.5}$ in the continuous exposure and by 5.3% (CI: 4.8, 5.9, $p < 0.10$) in the high pollution exposure category. PM air pollution appears to associate with the incidence of PTB in the population of Mexicali.

Keywords: Mexicali, Mexico, PM_{10} , $\text{PM}_{2.5}$, Tuberculosis, Incidence.

1. Introduction

Tuberculosis (TB) afflicts humanity since ancient times (Daniel, 2009; Smith et al., 2009; Turco and Byrd, 2001; Narasimhan et al., 2013). *Mycobacterium tuberculosis* (*Mtb*) predominantly causes pulmonary TB (PTB) but has the potential to affect any organ of the human body (Stead, 1965; Anderson et al., 2015).

It was not long ago that discussion began on emerging new risk factors for the activation of TB (Smith et al., 2014). One of those emerging risk factors is exposure to air pollution (indoor and outdoor), which can negatively modulate treatment responses (Kim, 2014). Multiple studies suggest that particulate matter (PM) is the air pollutant component related to an increased risk of TB development. The initial findings of associations between outdoor air pollution and PTB appeared at the end of the first decade of the 2000s (Cohen and Metha, 2007). A study conducted in Los Angeles, California, found that exposure to elevated levels of $\text{PM}_{2.5}$ had a statistically significant association with smear-positive TB (Jassal et al., 2013). In North Carolina, a relationship was demonstrated between $\text{PM}_{2.5}$ and the population's susceptibility to developing PTB (Smith et al., 2014). A study conducted in the city of New Taipei, Taiwan, reported that chronic exposure to PM_{10} delays sputum culture conversion in patients with PTB (Chen et al., 2016). Delayed effects of PM_{10} and $\text{PM}_{2.5}$ exposures on the incidence of TB were found in Shanghai (Wang et al., 2019).

The state of Baja California, Mexico, has the highest PTB incidence rate in the country. As of 2016, the state's average incidence rate was 47.6/100,000 inhabitants, while the national average was 13.8/100,000 (POEBC, 2017; CENAPRECE, 2017). Mexicali, the capital of Baja California, also presents a high incidence of PTB, exceeding 423

new cases per year (an incidence rate of around 42.4/100,000) (SUIVE, 2015). In addition to the PTB problem, the city is characterized by sustained high pollution levels that are predominantly caused by PM_{10} and $\text{PM}_{2.5}$ suspended in the air (Osornio et al., 2011; Reyna et al., 2012; Reyna et al., 2017; Reyna et al., 2018). The city, being at the Mexico-United States of America (USA) border, is exceptionally dynamic from a commercial, industrial and agricultural perspective. It has two border crossing checkpoints with the USA, where long queues of vehicles wait to cross into the neighboring country, emitting significant amounts of pollutants. There are two thermoelectric power plants and one geo-thermoelectric power plant in the vicinity. A large part of Mexicali's outskirts does not have paved roads, and roads are poorly maintained. Vehicular density is one of the highest in the country, and vehicle emissions are not controlled. Large agricultural areas on both sides of the border are pesticides are continuously sprayed, and farming fields are burned seasonally. Many restaurants cook with charcoal or firewood, resulting in uncontrolled emissions from charbroiling or frying meat. In addition to these PM sources, the region's extreme weather and aridity contribute to the high levels of PM pollution in the city (Watson and Chow, 2001; Reyna et al., 2012; ERG, 2019).

In the current study we explored the associated risk of PM exposure with PTB. Although most of the existing scientific studies use monthly or annual observation windows, the evidence suggests the possibility of shorter windows of exposure. Therefore, we examined the associations between PM exposures and PTB using weekly observation windows of continuous and categorical exposure (i.e. high and low pollution categories) up to 12 weeks before the PTB diagnosis.

2. Materials and Methods

2.1 Study Population

The study population consisted of individuals with PTB of all age groups and gender diagnosed between 2003 and 2007 who had resided in Mexicali for at least one year. The diagnosis of PTB (ICD-10: A15) required the presence of the following parameters: Ziehl-Neelsen stain positive sputum smear, clinical and chest x-ray manifestations characteristic for TB, and a suspected epidemiological history.

2.2 Clinical Data

Time-series data from 2003-2007 on PTB were made available by the Epidemiology Department (ED) of the Mexican Institute of Social Security, which records all new PTB cases. Individuals with PTB and HIV coinfection or Diabetes mellitus were excluded from the study. As per WHO guidelines (WHO, 1998), the ED obtains and evaluates three sputum smears and defines a new PTB case as a person with at least two out of three Ziehl-Neelsen stain positive sputum smears. In some instances, sputum smear-negative cases require a sputum culture to confirm PTB. However, those cases were not included in the current study, leaving the possibility of missing false-negative PTB cases; and in very rare cases including false positive cases with *Nocardia* sp. infection (Muricy et al., 2014).

Data in the current study is organized into weekly overall counts and counts by gender. We describe the total distribution by age groups (i.e. 0-14, 15-24, 25-34, 35-44, 45-54, 55-64, > 65 years) and age groups according to gender.

2.3 Meteorological Data

We worked with meteorological data [i.e. temperature (Temp) and relative humidity (Rh)] obtained from the Department of Meteorology at the Engineering Institute of the Autonomous University of Baja California, as published previously (Reyna et al., 2012). The data are organized as time-series (2003-2007) of daily averages and weekly averages.

2.4 PM_{10} and $PM_{2.5}$ data

PM_{10} and $PM_{2.5}$ data were provided by the Environmental Protection Secretary (SPA) of the State Government of Baja California, the National Institute of Ecology and Climate Change (INECC), and

the United States California Air Resources Board (CARB). Time-series of daily averages and weekly averages were calculated as reported previously (Reyna et al., 2012).

2.5 Biostatistics

The associations between PM_{10} or $PM_{2.5}$ and PTB were calculated using Poisson regression models and are expressed as relative risk (RR) and estimated 95% confidence intervals (CI), adjusted for over-dispersion. The effects of PM on PTB were analyzed by gender and for the entire study population.

The baseline models were first estimated and then applied to assess PM_{10} and $PM_{2.5}$ exposures. Baseline models were constructed by separately testing each of the dependent variables (e.g. PTB time-series according to gender) and adjusting for different independent control variables (see below). The models were specified with the Poisson regression algorithm in the NCSS9® software (NCSS, 2013).

We used the following control variables for the estimation of our baseline models: unlagged (i.e. Lag0) and lagged versions up to 12 weeks (i.e. Lag1 – Lag12) of Temp, Rh, the square of Temp and Rh and sinusoidal structural terms for seasonality (Katsouyanni et al., 1996), as well as the long-term linear and squared trends; time indicator variables were created for each year, month and week of the PTB diagnosis, taking as reference the first week, first month and first year; for age, the proportion of 65< year olds was taken as reference; for gender, the proportion of females was taken as reference. The control variables were included one-by-one when adjusting the baseline models. We started with the meteorological variables, then the structural and time variables, and, finally, age and gender. Only those variables contributing to a statistically significant ($p < 0.05$) increase in the deviance were accepted for the baseline model specification.

We assumed having adequate models when the residual autocorrelation coefficients did not exceed the confidence limits ($\alpha = 0.05$).

Once we developed the models, the baseline models were tested separately for PM_{10} and $PM_{2.5}$ at Lag0 and up to 12 weeks (Lag1 – Lag12).

The PM levels were assessed as continuous or as categorical exposure (i.e. low and high pollution). To evaluate categorical exposure, we created two

indicator variables. The “Low Pollution” category took the value of 1 when PM levels were equal to or less than the federal 24-hr average limit of $75 \mu\text{g}/\text{m}^3$ for PM_{10} or $45 \mu\text{g}/\text{m}^3$ for $\text{PM}_{2.5}$ as regulated in the Official Mexican Standards (SSA, 2014). Otherwise, it was assigned the value 0. The “High Pollution” variable took the value of 1 when PM levels were above the federal limits for PM_{10} or $\text{PM}_{2.5}$ and otherwise, took the value of 0.

To obtain the effect of the PM pollutant on PTB in the weeks of high pollution, the indicator variable “Low Pollution” and its interaction with the respective pollutant (i.e., PM_{10} or $\text{PM}_{2.5}$) were included in the model. The coefficient associated with the variable of the PM pollutant without interaction is taken as the effect on PTB in the weeks of high pollution. Thus, the PM variable’s coefficient would show the main effect of the category for a high level of PM pollution.

As with the baseline model estimation, it was assumed that the final models constructed were adequate when the residual autocorrelation did not show statically significant coefficients ($\alpha = 0.05$). Any significant residual autocorrelation coefficient was controlled using the corresponding autoregressive PTB terms.

3. Results

3.1 Population and PTB

During the five year study period (2003–2007), 1,748 new PTB cases were recorded in Mexicali, which had 896,394 inhabitants in 2007 (INEGI, 2020). Of the 1,748 PTB cases studied, 1,253 were male and 495 female, with 61 PTB cases under the age of

15 years; 32 male and 29 female. There were noticeable differences in the PTB incidence rate over the study period, with the highest PTB case rate recorded in 2006 and the lowest in 2004. The incidence of PTB was higher in the period from winter to spring in 2004 and 2006, and the period from spring to summer in 2003, 2005, and 2007. Table I shows the PTB incidence rates estimated for 2003–2007, and by season.

In general, the highest number of PTB cases occurred in men (2.5 males per female). Most of the cases occurred in the 25–34 year age group. The 35–44 year age group had the highest rate of men (4.0 males per female). Table II shows the PTB incidence according to age group and gender.

Table II. Total counts of PTB by age group and gender (2003–2007).

Age group	Men	Women	Total	M:F
0-14	32	29	61	1.1
15-24	224	116	340	1.9
25-34	348	107	455	3.3
35-44	271	68	339	4.0
45-54	182	72	254	2.5
55-64	92	53	145	1.7
65 <	104	50	154	2.1
Total	1253	495	1748	2.5

M – male; and, F – female.

3.2 Meteorology and Pollution

In general, the dry-hottest seasons occurred during spring (March 21–Jun 20) and summer (June 21–September 22), and the wet-coolest seasons occurred

Table I. Total counts and rate $[\text{N}(\%)]/100,000$ -inhabitants of PTB by season (2003–2007).

Season	Year					2003-2007
	2003	2004	2005	2006	2007	
Winter	85(10.49)	92(11.04)	74(8.65)	104(11.88)	83(9.26)	438
Spring	98(12.09)	108(12.96)	106(12.38)	114(13.02)	85(9.48)	511
Summer	98(12.09)	60(7.20)	91(10.63)	86(9.83)	97(10.82)	432
Fall	61(7.52)	46(5.52)	80(9.35)	82(9.37)	98(10.93)	367
Total	342(42.21)	306(36.73)	351(41.00)	386(44.10)	363(40.50)	1748

Population for the years 2003, 2004, 2006 and 2007 were estimated by the author, based on official figures reported by INEGI [INEGI, 2000; INEGI, 2005; INEGI, 2009; INEGI, 2010].

between fall (September 23-December 21) and winter (December 22-March 20). Table III shows the Temp, Rh and PM in Mexicali according to year and season for the entire period studied.

The highest PM₁₀ and PM_{2.5} levels occurred during the wet-coolest seasons, with concentrations exceeding the federal annual average limits of 40 µg/m³ and 12 µg/m³, respectively (SSA, 2014), for every year studied. The federal 24-hr average limit of 75 µg/m³ for PM₁₀ was exceeded on 219 days in winters, 181 days in springs, 201 days in summers, and 268 days in falls, resulting in a total of 869 exceedances during the study period. The federal 24-hr average limit of 45 µg/m³ for PM_{2.5} was exceeded on 302 days in winters, 248 days in springs, 262 days in summers, and 351 days in falls, resulting in a total of 1,163 exceedances during the study period.

Figures 1 and 2 present the distribution of PTB cases and PM levels between 2003 and 2007. Figure 1 presents the number of PTB cases and mean PM concentrations according to the months for the duration

of the entire study. Figure 2 presents the distribution of PTB cases for each month, according to PM₁₀ (Figure 2A) and PM_{2.5} (Figure 2B) concentrations.

3.3 Pollution Exposure

The models showed statistically significant results ($p < 0.05$) in some of the Poisson regression coefficients associated with the PM studied for the full population and for both genders. Table IV shows every most significant lag time regression coefficient associated with the PM and their statistics by continuous exposure and by high pollution exposure category (i.e. pollution levels above and below the 24-hr average limit for PM₁₀ (75 µg/m³) and PM_{2.5} (45 µg/m³)). The Poisson regression coefficients were used to calculate changes in the RRs of PTB if the concentration of the PM₁₀ or PM_{2.5} were increased by 10 µg/m³ of air, also shown in Table IV. Figure 3 graphically depicts the RR percentage increases of PTB by groups, for the 10 µg/m³ increment in PM₁₀ or PM_{2.5}.

Table III. Average levels for the meteorological variables and PM, per year and season, in Mexicali (2003-2007).

Season	Year					
	2003	2004	2005	2006	2007	2003-2007
Temperature	25.38	24.62	25.12	25.49	24.90	25.09
Winter	18.01	16.65	16.87	16.95	17.22	17.14
Spring	26.05	27.86	26.67	27.30	26.58	26.89
Summer	34.92	34.44	34.80	35.57	34.78	34.90
Fall	22.53	19.51	22.15	22.13	21.02	21.47
Relative humidity	41.98	39.34	38.99	33.41	34.00	37.54
Winter	46.68	47.29	56.15	37.60	33.30	44.20
Spring	34.90	30.40	30.39	29.39	31.14	31.25
Summer	41.99	33.68	33.87	34.07	35.37	35.59
Fall	45.35	45.98	35.53	32.57	36.19	39.12
PM ₁₀	100.66	93.48	86.29	85.71	90.16	91.26
Winter	111.87	95.13	75.85	104.94	99.28	97.41
Spring	87.99	77.85	74.23	64.53	80.46	77.01
Summer	92.33	92.42	77.55	69.35	83.11	82.95
Fall	110.45	108.53	117.51	104.01	97.77	107.65
PM _{2.5}	56.61	58.04	66.23	60.49	61.85	60.64
Winter	67.72	62.18	58.64	76.63	71.39	67.31
Spring	43.77	47.60	53.66	44.63	52.29	48.39
Summer	45.62	49.33	57.49	41.96	52.04	49.29
Fall	69.32	73.06	95.11	78.73	71.69	77.58

Temperature in °C; relative humidity in %; and PM₁₀ and PM_{2.5} in µg/m³.




Fig. 1. Time series of the monthly mean PM₁₀ (blue line) and PM_{2.5} (orange line) concentrations and the monthly number of PTB cases (gray bars) in Mexicali between 2003 and 2007. Interestingly, peak particle concentration preceded a peak in the number of cases suggesting seasonality. The models used in the study identified a time related statistically significant association in which existing elevated PM concentrations anteceded a ~3% increased risk for the diagnosis of PTB.

The RR for the development of PTB in males and the totality of cases remained similar and was statistically significant ($p < 0.05$), and borderline significant ($p < 0.10$) for the continuous PM₁₀ exposure. The RRs associated with the PM₁₀ or PM_{2.5} were slightly higher for the male PTB cases. Female PTB cases had an increased RR with broad confidence intervals compared with the RR for all the cases. Observed increased RRs were statistically significant only when considering continuous PM₁₀.

After model adjustments, the analysis showed a 10-week delayed effect for PM₁₀ and PM_{2.5} in all PTB and the male PTB cases ($p < 0.05$), and borderline significance ($p < 0.10$) for continuous PM₁₀ exposure. An 11-week delayed effect occurred in the female PTB case population when considering continuous PM₁₀ exposure ($p < 0.05$). In the full PTB case population, the RR rose by 2.4% (95% CI: 2.1, 2.6) for each 10 µg/m³ increase of PM₁₀ in the continuous exposure and by 3.6% (CI: 3.3, 4.0) in the high pollution exposure category, and by 3.2% (CI: 2.9, 3.4) for each 10 µg/m³ increase of PM_{2.5} in the continuous exposure and by 3.9% (CI: 3.6, 4.3) in the high pollution exposure category. In male PTB cases, the RR rose by 2.8% (CI: 2.5, 3.1) for each 10 µg/m³ increase of PM₁₀ in the continuous exposure and by 4.6% (CI: 4.2, 5.0) in the high pollution exposure category, and by 3.4% (CI: 3.1, 3.7) for each 10 µg/m³ increase of PM_{2.5} in the continuous exposure and by 4.2% (CI:

3.8, 4.6) in the high pollution exposure category. In female PTB cases, the RR for development of PTB rose by 5.1% (CI: 4.7, 5.5) for each 10 µg/m³ increase of PM₁₀ in the continuous exposure and by 5.3% (CI: 4.7, 5.8) in the high pollution exposure category, and by 4.3% (CI: 3.8, 4.8) for each 10 µg/m³ increase of PM_{2.5} in the continuous exposure category and by 5.3% (CI: 4.8, 5.9) in the high pollution exposure category.

As an example, Table V shows the Poisson regression coefficients with their respective confidence intervals and p-values obtained by including the continuous exposure of PM_{2.5} delays in the baseline model for the full population. The most significant effect was in Lag 10.

4. Discussion

We found associations between PM levels and PTB case numbers in the population that was studied. Our findings suggest that PM levels above the Official Mexican Standards (i.e. a 24-hr average of 75 µg/m³ for PM₁₀ and 45 µg/m³ for PM_{2.5}) increase the risk of PTB with a greater statistical significance in males than in females. Our results coincide with those in a recent study by Wang et al. (2019) that used months as the observation units and found that the numbers of TB cases increased by 8% and 4% for each PM_{2.5} and PM₁₀ increment of 10 µg/m³, respectively. In




Fig. 2. Distribution of all PTB cases (colored dots) diagnosed by month (1 = January to 12 = December) according to the PM concentration at the diagnosis time. This graphical representation shows the increased PM concentration (A = PM₁₀, B = PM_{2.5}) values during the cold months and the increase in the number of cases during the warmer months. The short horizontal red line indicates the position of the median for each dispersion cloud and the numbers in parentheses indicate the total monthly count of PTB cases.




Fig. 3. RR increases of PTB by group due to an increment of 10 $\mu\text{g}/\text{m}^3$ in PM. 95% confidence interval. * $p<0.10$. ** $p<0.05$.

the current study the number of PTB cases increased by 3.2% and 2.4% for each 10 $\mu\text{g}/\text{m}^3$ increment of PM_{2.5} and PM₁₀, respectively, and for the continuous exposure (borderline significant ($p<0.10$) for PM₁₀). Increases of 3.9% and 3.6% for each 10 $\mu\text{g}/\text{m}^3$ increment of PM_{2.5} and PM₁₀ we observed for the high pollution exposure category.

The RR found in this study showed a ~10-week lagged PM effect in all the groups analyzed. A recent study using four different monthly windows of exposure to PM_{2.5} before PTB diagnosis found an association after three-months of exposure (RR=1.020 for each increment of 10 $\mu\text{g}/\text{m}^3$) (Liu et al., 2018). The RR decreased strongly by increasing the exposure window (i.e. 6, 9, 12 months).

The published articles cited in our study show differences in the estimated risks (including the risks found in our study) for the development of TB on PM exposures. This is likely due to the differences in the methodologies, algorithms, control variables, types of TB studied, regions and sociodemographic characteristics, or the composition of the PM in the various studies and study sites. When looking at various PM exposure periods (i.e., 1, 3, 6, 9, 12 months) it is apparent that the risk of TB development becomes uniformly critical around the three-month exposure period with decreasing significance at longer exposure periods. The results presented in the current study with shorter exposure periods indeed suggest

the existence of a critical three-month lag exposure, identifying a 10 and 11-week delayed (lag) effect when testing for weekly PM and PTB associations. This time lag relates to an apparent seasonality of the two phenomena (new PTB and PM pollution). The adjustments of our model suggest that temperature and relative humidity are not responsible for the associations found between PM levels and PTB case numbers. Further studies are warranted to explore the likely contribution of behavioral and environmental variables that contribute to the seasonality in PTB incidence rates. For example, more people spend more time in indoor environments during colder seasons of the year where they may be exposed additionally to indoor PM levels from household heating.

Three-month PM exposure periods may be critically influencing the reactivation of PTB in persons with preexisting latent infection with *Mtb*, an important aspect that still requires further exploration. In a recent study in Wuhan, China, associations between PM levels and TB incidence rates were explored using the day as a unit of measurement. That study found statistically significant associations ($p<0.05$) only in lag 7 (days) for 10 $\mu\text{g}/\text{m}^3$ increments in PM₁₀ or PM_{2.5} (Huang et al., 2020). However, the meaning of short lag periods remains to be evaluated, and its biological plausibility explored.

The findings from our time-series analysis are supported by recent studies from our group that suggest that PM₁₀ and PM_{2.5} inhibit human antimycobacterial immune mechanisms. These studies also indicate that PM effects on human host immunity induced by *Mtb* are manifested differently, depending on the season during which ambient PM was collected for *in vitro* studies (Rivas-Santiago et al., 2015). Other studies from our group found that the urban PM_{2.5} suppresses antimycobacterial T cell (Ibironke et al., 2019) and alveolar macrophage functions (Torres et al., 2019). Immunosuppression induced by PM exposures may underlie increased risks of new *Mtb* infection or development of PTB in the population of Mexicali, where TB is considered endemic.

This study has several limitations. Diagnosis of PTB based on sputum smear positivity alone does not have the sensitivity and specificity of sputum cultures. Also, as indicated above, associations between weather seasons and PTB incidence rates may be confounded by exposure to indoor air pollution,

Table IV. Summary statistics of the Poisson regression analysis by PM exposure (i.e. continuous and high pollution exposure category) for the full population and by gender, and rate ratios (RRs) increases of PTB due to each increment of PM₁₀ or PM_{2.5} by 10 µg/m³ of air in Mexicali, Baja California. Only the most significant lag time is presented.

Lag (Weeks)			<i>b</i> (CI 95%)	<i>RR</i> (CI 95%)	<i>P</i>
<i>All population</i>					
PM ₁₀	10	Continuous	0.0023 (−0.0001, 0.0048) ^a	1.0238 (1.0212, 1.0263)	0.062
		≤ 75	Ref		
		> 75	0.0036 (0.0002, 0.0070) ^a	1.0365 (1.0330, 1.0401)	0.040
PM _{2.5}	10	Continuous	0.0031 (0.0004, 0.0058) ^a	1.0316 (1.0288, 1.0344)	0.025
		≤ 45	Ref		
		> 45	0.0039 (0.0007, 0.0071) ^a	1.0394 (1.0361, 1.0428)	0.018
<i>Males</i>					
PM ₁₀	10	Continuous	0.0028 (0.0000, 0.0056) ^b	1.0282 (1.0253, 1.0311)	0.055
		≤ 75	Ref		
		> 75	0.0045 (0.0005, 0.0083) ^b	1.0459 (1.0419, 1.0500)	0.023
PM _{2.5}	10	Continuous	0.0033 (0.0000, 0.0064) ^b	1.0340 (1.0308, 1.0371)	0.031
		≤ 45	Ref		
		> 45	0.0041 (0.0001, 0.0077) ^b	1.0418 (1.0381, 1.0456)	0.024
<i>Females</i>					
PM ₁₀	11	Continuous	0.0050 (0.0010, 0.0090) ^c	1.0510 (1.0468, 1.0551)	0.014
		≤ 75	Ref		
		> 75	0.0051 (0.0002, 0.0104) ^c	1.0526 (1.0471, 1.0582)	0.057
PM _{2.5}	11	Continuous	0.0042 (0.0005, 0.0089) ^c	1.0430 (1.0381, 1.0479)	0.077
		≤ 45	Ref		
		> 45	1.0052 (0.0001, 0.0105) ^c	1.0532 (1.0476, 1.0588)	0.056

PM₁₀ and PM_{2.5} in µg/m³.

^a Adjusted for: temperature Lag12; weeks of diagnosis - 13, 36, 47, 51; sinusoidal terms - sixth order; age group proportion - 15-24 years old; year of diagnosis - 2004; and, PTB Lag18.

^b Adjusted for: month of diagnosis - December; weeks of diagnosis - 13, 21, and 47; PTB male Lag7, 12, and 18.

^c Adjusted for: square of relative humidity Lag0; the month of diagnosis - February, March; age group proportion - 0-14 and 15-24 years old; and, PTB female Lag11.

Table V. Poisson regression coefficients (*b*) for PTB incidence in the full population by lags time of the continuous exposure of PM_{2.5}

Lag (Weeks)	<i>b</i> (CI 95%)	<i>P</i>
0	0.0014 (−0.0016, 0.0045)	0.362
1	0.0029 (0.0001, 0.0058)	0.046
2	0.0012 (0.0017, 0.0040)	0.424
3	0.0005 (0.0034, 0.0024)	0.741
4	0.0005 (0.0022, 0.0033)	0.706
5	0.0005 (0.0022, 0.0032)	0.727
6	0.0012 (0.0014, 0.0039)	0.355
7	0.0016 (0.0011, 0.0043)	0.248
8	0.0013 (0.0014, 0.0039)	0.345
9	0.0009 (0.0018, 0.0037)	0.516
10	0.0031 (0.0004, 0.0058)	0.025
11	0.0007 (0.0022, 0.0036)	0.631
12	−0.0010 (0.0040, 0.0019)	0.490

tobacco smoke and crowding effects in congregated settings. It was also not possible to assess the population at risk for PTB development from *Mtb* infection and PM exposure as the proportion of the Mexicali population with positive tuberculin skin test is very high due to universal BCG vaccination coverage.

5. Conclusion

This study suggests that PM air pollution may contribute to the development of PTB in the population of Mexicali. Diagnosis of PTB showed a lag of 10 or 11 weeks relative to seasonally elevated PM levels. Mexicali, Baja California, has a high incidence of TB and high PM₁₀ and PM_{2.5} levels that consistently exceed the acceptable air quality limits of the Official Mexican Standards. PM exposures may be an essential factor in contributing to the high incidence of PTB in Mexicali's population. These preliminary results warrant further research using different statistical models, longitudinal cohort study designs and the assessment of confounders such as socioeconomic and nutritional status, HIV coinfection, Diabetes mellitus, tobacco smoking, indoor air pollution, and the use of illicit drugs and alcohol.

Acknowledgments

We are grateful to the National Council for Science and Technology (CONACyT) for providing a scholarship to Paulina E. Burgos-Tejeda and collaborating in the development of this project.

References

- Anderson L, Dean A, Falzon D, Floyd K, Garcia I, Gilpin C, Glaziou P, Hamada Y, Hiatt T, Kan-char A, Law I, Lienhardt C, Nguyen L, Siroka A, Sismanidis C, Syed L, Timimi H, Gemert W, Zignol M. 2015. Global tuberculosis report. Geneva, World Health Organization. Available at https://apps.who.int/iris/bitstream/handle/10665/191102/9789241565059_eng.pdf?sequence=1 (accessed 2020 March 02).
- CENAPRECE. 2017. Cifras Oficiales de Tuberculosis. Available at http://www.cenaprece.salud.gob.mx/programas/interior/micobacteriosis/tuberculosis/cifras_oficiales.html (accessed 2020 March 02).
- Chen KY, Chuang KJ, Liu HC, Lee KY, Feng PH, Su CL, Lin CL, Lee CN, Chuang HC. 2016. Particulate matter is associated with sputum culture conversion in patients with culture-positive tuberculosis. *Therapeutics and Clinical Risk Management* 12: 41-46. <https://doi.org/10.2147/TCRM.S92927>
- Cohen A, Mehta S. 2007. Pollution and tuberculosis: Outdoor sources. *PLoS Medicine* 4(3): e142. <https://doi.org/10.1371/journal.pmed.0040142>
- Daniel TM. 2009. The history of tuberculosis: past, present, and challenges for the future. In: Schaaf HS, Zumla A, eds. *Tuberculosis: A Comprehensive Clinical Reference*. Elsevier Saunders, London, UK, 1st edition. 1-8.
- Huang S, Xiang H, Yang W, Zhu Z, Tian L, Deng S, Zhang T, Lu Y, Liu F, Li X, Liu S. 2020. Short-term effect of air pollution on Tuberculosis based on Kriged data: A Time-series analysis. *International Journal of Environmental Research and Public Health* 17(5): 1522. <https://doi.org/10.3390/ijerph17051522>
- Ibironke O, Carranza C, Sarkar S, Torres M, Choi HT, Nwoko J, Black K, Quintana-Belmares R, Osornio-Vargas A, Ohman-Strickland P, Schwander S. 2019. Urban air pollution particulates suppress human T-Cell responses to *Mycobacterium Tuberculosis*. *International Journal of Environmental Research and Public Health*. 16(21): 4112. <https://doi.org/10.3390/ijerph16214112>
- INEGI. 2020. Instituto Nacional de Estadística y Geografía. Available at <https://www.inegi.org.mx/app/buscador/default.html?q=Mexicali#tabMCCollapse-Indicadores> (accessed 2020 March 02).
- INEGI. 2000. XII Censo General de Población y Vivienda 2000. México: Instituto Nacional de Estadística y Geografía. Available at <https://www.inegi.org.mx/programas/ccpv/2000/> (accessed 2020 March 02).
- INEGI. 2005. II Conteo de Población y Vivienda 2005. México: Instituto Nacional de Estadística y Geografía. Available at <https://www.inegi.org.mx/programas/ccpv/2005/> (accessed 2020 March 02).
- INEGI. 2009. Estadísticas históricas de México 2009. México: Instituto Nacional de Estadística y Geografía. Available at http://internet.contenidos.inegi.org.mx/contenidos/productos/prod_serv/contenidos/espanol/bvinegi/productos/integracion/pais/historicas10/Tema1_Poblacion.pdf (accessed 2020 March 02).
- INEGI. 2010. Censo de Población y Vivienda 2010: Instituto Nacional de Estadística y Geografía. Available at <https://www.inegi.org.mx/programas/ccpv/2010/> (accessed 2020 March 02).

- Jassal MS, Bakman I, Jones B. 2013. Correlation of ambient pollution levels and heavily-trafficked roadway proximity on the prevalence of smear-positive tuberculosis. *Public Health* 127(3): 268-274. <https://doi.org/10.1016/j.puhe.2012.12.030>
- Katsouyanni K, Schwartz J, Spix C, Touloumi G, Zmirou D, Zanobetti A, Wojtyniak B, Vonk JM, Tobias A, Pönkä A, Medina S, Bachárová L, Anderson HR. 1996. Short-term effects of air pollution on health: a European approach using epidemiological time-series data: the APHEA protocol. *Journal of Epidemiology & Community Health* 50(Sup I): S12-S18. http://dx.doi.org/10.1136/jech.50.Suppl_1.S12
- Kim J. 2014. Is ambient air pollution another risk factor of tuberculosis? *The Korean Journal of Internal Medicine* 29(2): 170-172. <https://doi.org/10.3904/kjim.2014.29.2.170>
- Liu Y, Cui L, Hou L, Yu C, Tao N, Liu J, Li Y, Zhou C, Yang G, Li H. 2018. Ambient air pollution exposures and newly diagnosed pulmonary tuberculosis in Jinan, China: A time series study. *Scientific Reports*. 8: 1-11. <https://doi.org/10.1038/s41598-018-35411-6>
- ERG. 2019. 2014 Northern Baja California Emissions Inventory Project. Eastern Research Group, Inc. Final Report. Sacramento, California, USA, 36 pp. Available at https://ww3.arb.ca.gov/planning/border/erg_final_report.pdf (accessed 2021 January 05).
- Muricy ECM, Lemes RA, Bombarda S, Ferrazoli L, Chimara E. 2014. Differentiation between *Nocardia* spp. and *Mycobacterium* spp.: Critical aspects for bacteriological diagnosis. *Revista do Instituto de Medicina Tropical de São Paulo*. 56(5): 397-401. <https://doi.org/10.1590/S0036-46652014000500005>
- Narasimhan P, Wood J, MacIntyre CR, Mathai D. 2013. Risk factors for tuberculosis. *Pulmonary Medicine* 1-11. <https://doi.org/10.1155/2013/828939>
- NCSS. 2013. NCSS9 Statistical Software Kaysville, Utah, USA: NCSS, LLC.
- Osornio-Vargas AR, Serrano J, Rojas-Bracho L, Miranda J, García-Cuellar C, Reyna MA, Flores G, Zuk M, Quintero M, Vázquez I, Sánchez-Pérez Y, López T, and Rosas I. 2011. In vitro biological effects of airborne PM_{2.5} and PM₁₀ from a semi-desert city on the Mexico-US border. *Chemosphere* 83(4): 618-626. <https://doi.org/10.1016/j.chemosphere.2010.11.073>
- POEBC. 2017. Actualización del Programa Sectorial de Salud 2015-2019. Periódico Oficial del Estado de Baja California. Available at <https://es.readkong.com/page/actualizaci-n-programa-sectorial-de-salud-2015-2019-8063275> (accessed 2022 February 15)
- Reyna MA, Bravo ME, López R, Nieblas EC, and Nava ML. 2012. Relative risk of death from exposure to air pollutants: a short-term (2003-2007) study in Mexicali, Baja California, Mexico. *International Journal of Environmental Health Research* 22(4): 370-386. <https://doi.org/10.1080/09603123.2011.650153>
- Reyna MA, Nieblas EC, Nava ML, Torillo E. 2017. An estimation of costs and public-health benefits by the PM₁₀ mitigation in Mexicali, Baja California, Mexico. *Revista Internacional de Contaminación Ambiental* 33(1): 117-129. <http://dx.doi.org/10.20937/RICA.2017.33.01.11>
- Reyna MA, Mérida JV, Osornio A, Lerma C, Bravo ME, Avitia RL, and Nieblas EC. 2018. Association between personal PM₁₀ exposure and pulmonary function in healthy volunteers from a semi-arid city on the US-Mexican border. *Revista Internacional de Contaminación Ambiental* 34(4): 583-595. <http://dx.doi.org/10.20937/RICA.2018.34.04.03>
- Rivas-Santiago CE, Sarkar S, Cantarella P, Osornio-Vargas AR, Quintana-Belmares R, Meng Q, Schwander S. 2015. Air Pollution Particulate Matter Alters Antimycobacterial Respiratory Epithelium Innate Immunity. *Infection and Immunity* 83(6): 2507-2517. <http://dx.doi.org/10.1128/IAI.03018-14>
- Smith GS, Schoenbach VJ, Richardson DB, Gammon MD. 2014. Particulate air pollution and susceptibility to the development of pulmonary tuberculosis disease in North Carolina: an ecological study. *International Journal of Environmental Health Research* 24(2): 103-112. <https://doi.org/10.1080/09603123.2013.800959>
- Smith NH, Hewinson RG, Kremer K, Brosch R, Gordon SV. 2009. Myths and misconceptions: the origin and evolution of *Mycobacterium tuberculosis*. *Nature Reviews Microbiology* 7(7): 537-544. <https://doi.org/10.1038/nrmicro2165>
- SSA. 2014. Norma Oficial Mexicana NOM-025SSA1-2014, Salud ambiental. Valores límites permisibles para la concentración de partículas suspendidas PM₁₀ y PM_{2.5} en el aire ambiente y criterios para su evaluación. México: Secretaría de Salud. Diario Oficial de la Federación. 04 de agosto de 2014. Available at http://www.dof.gob.mx/nota_detalle.php?codigo=5357042&fecha=20/08/2014 (accessed 2020 March 10).

- Stead WW. 1965. The pathogenesis of pulmonary tuberculosis among older persons. *American Review of Respiratory Disease* 91(6): 811–822.
- SUIVE. 2015. Plataforma Única de Información TB 2015. Available at http://www.cenaprece.salud.gob.mx/programas/interior/micobacteriosis/descargas/pdf/SituacionActualTB_MEXICO2015.pdf (accessed 2020 March 02).
- Torres M, Carranza C, Sarkar S, Gonzalez Y, Osornio-Vargas A, Black K, Meng Q, Quintana-Belmares R, Hernandez M, Angeles-Garcia JJF, Páramo-Figueroa VH, Iñiguez-Garcia MA, Flores JL, Zhang JJ, Gardner CR, Ohman-Strickland P, Schwander S. 2019. Urban airborne particle exposure impairs human lung and blood *Mycobacterium tuberculosis* immunity. *Thorax* 74(7): 675–683. <http://dx.doi.org/10.1136/thorax-jnl-2018-212529>
- Turco J, Byrd M. 2001. An interdisciplinary perspective: infectious diseases and history. *The American Biology Teacher* 63(5): 325–335. <https://doi.org/10.2307/4451119>
- Wang H, Tian C, Wang W, Luo X. 2019. Temporal cross-correlations between ambient air pollutants and seasonality of tuberculosis: A time-series analysis. *International journal of environmental research and public health* 16(9): 1585. <https://doi.org/10.3390/ijerph16091585>
- Watson JG, Chow JC. 2001. Source characterization of major emission sources in the imperial and Mexicali Valleys along the US/Mexico border. *Science of the Total Environment*. 276(1-3): 33–47. [https://doi.org/10.1016/S0048-9697\(01\)00770-7](https://doi.org/10.1016/S0048-9697(01)00770-7)
- WHO. 1998. Laboratory services in tuberculosis control. Geneva: World Health Organization WHO/TB/98.258. Available at [https://apps.who.int/iris/bitstream/handle/10665/65942/WHO_TB_98.258_\(part1\).pdf](https://apps.who.int/iris/bitstream/handle/10665/65942/WHO_TB_98.258_(part1).pdf) (accessed 2020 March 08).