



Geofísica internacional

ISSN: 0016-7169

Instituto de Geofísica, UNAM

Torres-Rivera, S.; Ramos-Leal, J. A.; Moran-Ramírez, J.; Torres- Hernández, J. R.; Padilla-Sánchez, L.; Carranco-Lozada, S. E.; García-Arreola, M. E.  
Hydrogeochemical characterization and quantification of the groundwater mixture in a hydrogeological basin of the middle zone of San Luis Potosí  
Geofísica internacional, vol. 58, no. 4, 2019, October-December, pp. 247-258  
Instituto de Geofísica, UNAM

DOI: <https://doi.org/10.22201/igeof.00167169p.2019.58.4.2056>

Available in: <https://www.redalyc.org/articulo.oa?id=56872287001>

- How to cite
- Complete issue
- More information about this article
- Journal's webpage in redalyc.org

UNAM  
redalyc.org

Scientific Information System Redalyc

Network of Scientific Journals from Latin America and the Caribbean, Spain and Portugal

Project academic non-profit, developed under the open access initiative

## Hydrogeochemical characterization and quantification of the groundwater mixture in a hydrogeological basin of the middle zone of San Luis Potosí

Torres-Rivera S., Ramos-Leal J. A., Moran-Ramírez J., Torres- Hernández J. R., Padilla-Sánchez L., Carranco-Lozada S. E. and García-Arreola M. E.

Received: November 15, 2017; accepted: september 24, 2019; published on line: October 01, 2019

### Resumen

Los procesos de caracterización hidrogeoquímica e interacción agua-roca en las subcuencas de Santa Catarina y Ocampo-Paraíso en la Zona Media de San Luis Potosí han permitido identificar los tipos de aguas subterráneas. Los resultados del análisis químico y el estudio geológico regional han permitido documentar los posibles mecanismos de interacción del agua con la roca. Los resultados indican que la composición del agua subterránea en cada una de estas cuencas refleja la disolución por el agua de lluvia de los minerales de carbonato y magnesio de las rocas carbonatadas de la formación del Cretácico (Formación El Abra del Cretácico) y del flujo subterráneo a través de un medio granular con fuerte influencia del material de origen volcánico. Las aguas del tipo  $\text{Ca-HCO}_3$ ,  $\text{Ca-Mg-HCO}_3$  se modifican durante el flujo subterráneo mediante la incorporación de iones  $\text{Na}^+$  y  $\text{SO}_4^{2-}$ . En el caso de la subcuenca III de San Nicolás, se ha documentado que su

relleno incluye depósitos fluviales que pueden depositarse en condiciones climáticas de regiones semiáridas. El análisis hidrogeoquímico de las concentraciones de iones mayores, litio y bromo, permitieron identificar que el agua que se extrae de la región es producto de un proceso de mezcla en la que intervienen tres miembros extremos (mezcla ternaria): El primero, Santa Catarina ( $C_1$ ), tiene las concentraciones más bajas de bromo y litio en la recarga ocurre en las partes de mayor altitud en la región, hacia la Sierra de Álvarez. El segundo miembro ( $C_2$ ), tiene valores altos del bromo, bajo en litio y se origina hacia el valle de Ocampo Paraíso. El tercer miembro ( $C_3$ ), tiene bajas concentraciones de bromo, alto en litio y proviene de área de San Nicolás Tolentino. Los porcentajes de mezcla indican que  $C_1$  aporta el 50%,  $C_2$  contribuye con el 31% y  $C_3$  aporta 19 %, del agua al sistema.

Palabras clave: hidrogeoquímica, mezcla ternaria, miembro extremo, elemento conservativo, flujo local, Sierra Madre Oriental.

Torres-Rivera S.<sup>1</sup>  
Torres- Hernández J. R.<sup>1</sup>  
Posgrado en Geología  
Facultad de Ingeniería  
Universidad Autónoma de San Luis Potosí (UASLP),  
Manuel Nava #5, Zona Universitaria  
CP. 78240, San Luis Potosí, S.L.P., México  
*\*Corresponding author: iq\_soniatrivera@yahoo.com.mx*

Ramos-Leal J. A.  
Posgrado en Geociencias Aplicadas  
Instituto Potosino de Investigación Científica y Tecnológica  
A.C. (IPICYT), C. Presa San José No. 2055  
Lomas 4ª. Sección, CP 78216  
San Luis Potosí, S. L. P., México

Moran-Ramírez J.  
<sup>3</sup>Cátedra CONACYT-UNAM, Geofísica  
Circuito de la Investigación Científica s/n  
Ciudad Universitaria, 04150  
Coyoacán, Ciudad de México

Padilla-Sánchez L.  
García-Arreola M. E.  
Area de Ciencias de la Tierra  
Facultad de Ingeniería  
Universidad Autónoma de San Luis Potosí (UASLP)  
Manuel Nava #5, Zona Universitaria  
CP. 78240, San Luis Potosí, S.L.P., México

Carranco-Lozada S. E.  
Instituto Politécnico Nacional, CECyT No. 17  
León Guanajuato, México

## Abstract

The hydrogeochemical characterization and water-rock interaction processes in the sub-basins of Santa Catarina and Ocampo-Paraíso in the Zona Media of San Luis Potosí have allowed identifying the types of groundwater. The results of the chemical analysis and the regional geological study show the possible rock-water interaction mechanisms. These results indicate that the composition of the groundwater in each of these basins reflects the dissolution of the carbonate and magnesium minerals of the carbonate rocks of Cretaceous age (El Abra Formation), by rainwater, and of the groundwater through a granular medium with a strong influence of the material of volcanic origin. The  $\text{Ca-HCO}_3$ ,  $\text{Ca-Mg-HCO}_3$ , water type are modified during the groundwater flow by incorporation of  $\text{Na}^+$  and  $\text{SO}_4^{2-}$  ions. The San Nicolás (sub-basin III) has covered by fluvial deposits in

a climatic conditions of semiarid regions. The hydrogeochemical analysis of major ions, lithium and bromine allowed identifying that the extracted water from the region is a product of mixing process, involving three end-members (ternary mixture). The first, Santa Catarina ( $\text{C}_1$ ), has the lowest concentrations of bromine and lithium and its recharge occur in the highest altitude land in the region, towards the Sierra de Álvarez. The second end member ( $\text{C}_2$ ), displays high values of bromine, low in lithium and originates towards the Ocampo Paraíso Valley. The third end member ( $\text{C}_3$ ), has low bromine concentrations, high in lithium, and comes from San Nicolás Tolentino area. The mixing percentages indicate that  $\text{C}_1$  contributes 50%,  $\text{C}_2$  gives 31% and  $\text{C}_3$  adds 19% water to the system.

**Key words:** hydrogeochemistry, ternary mixture, end-member, conservative element, local flow, Sierra Madre Oriental.

## Introduction

The chemical composition of the groundwater results from the interaction of rainwater with the dissolved rocks in the recharge zone (where it acquires part of its components). After filtration, it continues acquiring ions by its contact with the granular material through which it moves. The carbonate rocks undergo a process of chemical alteration by the action of rainwater that is combined with carbonic acid, this acid water dissolves the carbonate rocks, giving rise to a karstic landscape. In karst areas, the groundwater flow is characterized by high speeds, short residence times and significant water-rock interaction (Roback *et al.*, 2001). When the water infiltrates by small fractures, the solution goes deepening and widening these fissures, creating a network of ducts in the karstic rock. This process increases its secondary permeability and induces the formation of karst (Király *et al.*, 1979; Andreo *et al.*, 2004; Antigüedad *et al.*, 2008). In their evolution, groundwater may interact with other currents or groundwater flows, resulting in mixing of waters of different chemical composition due to natural or anthropogenic processes (Harpaz and Bear, 1963; Ramos *et al.*, 2007). A way of analyzing this phenomenon in generating isovalue curves of the different ions contained in the water, which allows to crossing directions of flow and zones where they can occur encounters of subterranean currents with different chemical compositions. Another

way to organize the contributions of water mixtures in the groundwater flow is to use the conservative elements, frequently considered as tracers. The blends may be mainly binary or ternary; simultaneous mixtures with more than three end-members are uncommon to occur in nature whereas binary mixtures are very common. For example, to quantify the mixture of fresh and marine water have been widely explained in numerous publications (Apello and Postma, 1996; Generaux *et al.*, 2004; Wallick, 1981; Lee and Krothe, 2001; Abu-Jaber, 2001; Skalbeck *et al.*, 2002; Valentino and Stanzione, 2002). However, ternary mix models are less common. Rice and Hornberger (1998) evaluated the maximum flow contribution for groundwater in the US, and Generaux *et al.*, (2004) evaluated groundwater mixtures by basin-to-basin transfer in the Costa Rica. Laaksoharju *et al.*, (1999), have addressed simultaneous mixtures with more than three end-members (EM). However, these authors did not explain how this process is carried out in nature. In addition, Douglas *et al.*, (2000) have reported that the mixing of three types of water with different chemical conditions and documented that this process can promote chemical reactions in an aquifer. In this work, we report that how chemical composition of water circulates through the volcanic rocks, limestone and alluvial fill in three different sub-basins. Therefore, the physicochemical characteristics, groundwater types, ion exchange processes and water-rock interaction

of three different aquifers (San Nicolás Tolentino, Ocampo-Paraíso, Santa Catarina) are studied, which situated in the Zona Media of the state of San Luis Potosí, central Mexico. These findings contribute significantly due to their necessity in supplying drinking water as well as agricultural purposes. In addition, it provides valuable geochemical information for hydrogeology at local as well as a regional level.

### Hydrogeological framework

The San Nicolás Tolentino aquifer (SNTA) located in the southern part of the state of San Luis Potosí (Figure 1) and it covers an exposed area of approximately 1,712.99 Km<sup>2</sup>. In this region, the climate is dry, arid with an average annual rainfall of 300 mm (INEGI, 2002) and an average temperature of 25° C (INEGI, 2002).


The SNTA consists of three sub-basins, drained by the Santa Catarina (sub-basin I), Ocampo Paraíso (sub-basin II) and San Nicolás Tolentino (sub-basin III) River (Figure 1).

The Santa Catarina sub-basin (I), is an exorheic sub-basin with an area of 732.5 km<sup>2</sup>, it is drained by the Santa Catarina River, which

corresponds to 84.5% of the total area of the SNTA. Ocampo-Paraíso sub-basin (II) forms a sub-basin of 345.36 km<sup>2</sup>. San Nicolás sub-basin (III) is an endorheic sub-basin of the area of 369.62 km<sup>2</sup> in which the Las Golondrinas dam is located.

### Geology

The oldest stratigraphic unit that outcrop in the region is known as Guaxcama Formation of Lower Cretaceous (Neocomian-Aptian), which is composed of anhydrite, gypsum, and dolomite (Carrillo, 1971). This formation developed large diapiric structures (Carrillo-Bravo, 1971; Torres-Hernández, 1994), and is covered by the thick more than 1000m calcareous sequence of the Abra Formation. The youngest Cretaceous unit is the Cardenas Formation, which is constituted by a sequence of shales and sandstones. Its thickness is irregular, between 50 and 100m. The central training unit to consider in this work is the Abra Formation, it is construed by lagunar and reef facies. This unit is widely distributed, and carved for the most significant karst development. It thicknesses is up to 1,800 m (Zapata-Zapata and Pérez-Venzor, 1979).



**Figure 1.** Location of the San Nicolás Tolentino Aquifer, regional geology of the study area and sampling points.

The Sierra de Alvarez (west of the studied area), constitutes the highest elevation in the region, and exposed as the slope foot facies (pre-reef deposits with 120 m of thickness), which is known as Tamabra Formation (López-Doncel, 2003).

The Cretaceous rocks are covered discordantly by volcanic rock sequences of andesitic and rhyolitic composition (Casa Blanca andesite, Santa María Ignimbrite, Portezuelo Latite, San Martín Avalanche, San Nicolás Formation, and Jagüey Formation; Torres Hernández *et al.*, 2010). This set of Paleogene (Oligocene) volcanic units is mainly distributed in the south-central part of the study area, with thicknesses greater than 140 m (Labarthe-Hernández *et al.*, 1982). Alluvium sediments cover all the units locally. Figure 2 shows a longitudinal geological section of the Peotillos, San Nicolás Tolentino and Santa Catarina valleys, looking towards the east (Figure 2). It can be observed that the Sierra de San Nicolás Tolentino is a remnant that belongs to a flank of an anticlinal that was eroded in Ocampo Paraiso sub-basin.

#### Hydrogeology

In the study area, aquifers are housed in granular material that partially covers Paleogene volcanic rocks, and carbonated rocks of the Cretaceous. Regionally, the largest aquifers are housed in limestone rocks.


The distribution of granular material is limited in the valleys, where it reaches only a few tens of meters, so its potential is not essential. The fracture medium represented by El Abra Formation that is the most crucial aquifer unit in the region because of its wide distribution and high thickness (Zapata and Pérez, 1979). This unit is exposed overall the high mountain, where it functions as a vital

recharge zone, and in general, the flow goes from NW to SE, W to E, and in the southern part, of S to N (Figure 2). Since the origin of this unit belongs to the marine-platform and cover with reef and lagoon facies that provides it primary porosity whereas tectonic processes responsible for secondary porosity. In this region, El Abra Formation is very fractured, especially in the anticline nuclei where it develops numerous karstic structures (from sinkholes and uvalas to poljes). The processes of dissolution form cavities in the sub-soil, developing karstic caverns such as those of La Puente, El Ángel and La Catedral. Volcanic units with less distribution and thickness than limestones also function as aquifer units (Figure 2).

#### Methodology

##### *Sampling and analysis*

We collected a total of 34 water samples (26 correspond to wells, 8 to dugwells) whose distribution is shown in Figure 1, based on the "Groundwater Sample Protocol" (Armienta *et al.*, 1987; Deutsch, 1997). During sample collection, physical-chemical parameters were measured in an isolation cell, which has limited interaction with the atmosphere, preventing changes in its original composition (temperature, electrical conductivity, pH, dissolved oxygen, oxide reduction potential and alkalinity). All groundwater samples were collected in linear high-density polyethylene (HDPE) bottles for the analysis of more abundant ions. The bottles were previously washed three times, with plenty of deionized water. The sample was acidified right after collection to pH <2 with ultrapure nitric acid. The samples for the determination of anions were not acidified. The major ions were analyzed in the Laboratories of Chemical Analysis, Faculty of Engineering, Autonomous



**Figure 2.** Longitudinal geological section of the Sierra de Peotillos, San Nicolás Tolentino and the Santa Catarina valley, facing east.



University of San Luis Potosi. The alkalinity was performed in situ by the titration method of Gran. For the analysis of  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ ,  $\text{Na}^+$  and  $\text{K}^+$ , the Atomic Flame Absorption Spectrometry (FAA) method was used. The chlorides were analyzed by the Argentometric method, the Sulphates, and Fluor with the Turbidimetric method. All chemical analyses show an ionic balance less than close to 5% allowed (Freese and Cherry, 1979).

#### Calculation of Mixtures

The end-members represent the highest and lowest concentrations of the system. The mixtures types can be binary and ternary; the former has been used to quantify mixtures of freshwater and marine waters (Apello and Postman, 1996; Genereux *et al.*, 2002; Wallick, 1981; Abu-Jaber, 2001, Skalbeck *et al.*, 2002; Valentino and Stanzione 2002). (Laaksoharju *et al.*, 1999; Douglas *et al.*, 2000) showed multidimensional models with more than three water mixtures. As part of the methodology for the qualitative and quantitative analysis of a binary or ternary mixture, three end-members representing the evolution of the aquifer under study are identified.

The percentage calculation for each of the sub-basins was obtained using three equations with three unknowns; it is represented as a balanced equation:

$$C_t = C_1 + C_2 + C_3 \quad (1)$$

Where  $C_1$  is the member associated with the local recharge in Santa Catarina,  $C_2$  is the member associated with Ocampo-Paraiso, and  $C_3$  is associated with the flows of San Nicolás Tolentino.  $C_1$ ,  $C_2$ , and  $C_3$  are unknown values, and  $C_w$  is equal to 1.

These components were determined with the equations of the mass balance of two conservative elements  $\text{Li}^+$  and  $\text{Br}^-$ :

$$C_w \text{Li}_w = C_1 * \text{Li}_1 + C_2 * \text{Li}_2 + C_3 * \text{Li}_3 \quad (2)$$

$$C_w \text{Br}_w = C_1 * \text{Br}_1 + C_2 * \text{Br}_2 + C_3 * \text{Br}_3 \quad (3)$$

(Equation 4) is obtained by substituting in Equation 2 in  $C_3$  of Equation 1:

$$C_3 = \frac{C_w(\text{Li}_w - \text{Li}_1) + (\text{Li}_2 - \text{Li}_1)}{\text{Li}_3 - \text{Li}_1} \quad (4)$$

$C_1$  substitute for Equation 3 in Equation 1 and obtain the Equation 5:

$$C_1 = \frac{C_w(\text{Br}_w - \text{Br}_3) + C_2(\text{Br}_3 - \text{Br}_2)}{\text{Br}_1 - \text{Br}_3} \quad (5)$$

The only possible solution for  $C_2$  is when  $C_w = 1$ . The other two components are calculated by clearing  $C_2$  and substituting  $C_1$  and  $C_3$  in Equation 1 and the Equation is obtained:

$$C_2 = C_T - C_1 - C_3 \quad (6)$$

#### Results and discussion

The statistics of the chemical results of the groundwater samples are presented in Table 1. The results show that sub-basin I has a lower temperature, is rich in  $\text{K}^+$ ,  $\text{Ca}^{2+}$ ,  $\text{HCO}_3^-$ ,  $\text{Cl}^-$ . Sub-basin II has an average temperature, lower pH, DO, Eh,  $\text{Na}^+$ ,  $\text{K}^+$ ,  $\text{Ca}^{2+}$  and higher  $\text{Mg}^{2+}$ . The sub-basin III presents higher T, pH, EC, DO, Eh,  $\text{Na}^+$ ,  $\text{SO}_4^{2-}$  and lower Alkalinity, hardness,  $\text{Cl}^-$ .

#### Hydrogeochemical characterization

The chemical composition of groundwater is the result of continuous processes of interaction between precipitation water that infiltrates the subsoil and circulates through geological materials. Part of the chemical components is acquired in the recharge zone. Others are acquired along the pathway, captured by wells and in the discharge area as springs, resulting in different hydrogeochemical signatures.

#### Types of water


The Piper diagram (Piper, 1994) represents the end-members of each region and the classification of the different types of water (Figure 3). In the left part of the diamond, it displays mainly calcium-bicarbonate water. The upper part indicates dissolution of rock and middle part represents mixed waters where ion exchange can occur. On the left side of the diamond, is possibly the most evolved water, with dominance of the carbonate and sulfate ions.

In general, three families were obtained for the study area: calcium bicarbonate ( $\text{Ca-HCO}_3$ ), sodium bicarbonate ( $\text{Na-HCO}_3$ ), and only one exotic sample that is classified as calcium sulfated ( $\text{Ca-SO}_4$ ) (Figures 3a and 3b).

**Table 1.** Statistics of the chemical results of sub-terrestrial water samples.

Sub-basin	T °C	pH	EC μS/cm	DO %	Eh mV	TDS mg/L	Hard Total mg/L	Major Elements (mg/L)						
								Na <sup>+</sup>	K <sup>+</sup>	Ca <sup>2+</sup>	Mg <sup>2+</sup>	HCO <sub>3</sub> <sup>-</sup>	Cl <sup>-</sup>	SO <sub>4</sub> <sup>2-</sup>
I) SC														
MIN	24	7	309	2	234	200	61	24	6	16	2	134	7	14
MAX	31	9	1848	40	1816	1290	836	149	9	255	48	401	12	725
MEAN	27	8	610	18	516	419	210	51	6	62	13	217	9	118
STD	3	1	557	13	575	391	278	44	1	86	17	93	2	268
II) OP														
MIN	21	8	449	2	81	320	148	36	4	41	8	232	7	29
MAX	34	9	810	24	340	600	336	54	8	83	38	366	30	115
MEAN	24.6	7.8	566	7.4	282	422	221	41	6	58	19	280	12	48
STD	4	0.4	118	7.4	80	96	61	6	1	12	10	46	7	29
III) SNT														
MIN	16	8	1	6	275	400	205	27	4.1	59	8.1	267	11	2
MAX	26	8	808	20	344	600	301	77	25	90.1	22.3	390	35	53
MEAN	21	8	545	10	318	447	247	44	9	77	13	322	18	32
STD	3.1	0.2	208.6	4.4	20.6	66.0	29.3	15.4	5.9	10.1	5.6	35.7	8.3	16.1

I) Santa Catarina sub-basin (SC); II) Ocampo Paraiso sub-basin (OP) and III) San Nicolas Tolentino sub-basin (SNT)



**Figure 3.** Piper diagrams: A) sub-basin I, Santa Catarina. B) sub-basins II and III Ocampo Paraiso and San Nicolas Tolentino for groundwater.

In Santa Catarina sub-basin I, two aquifers have been detected (Torres-Rivera, 2012) that is known as free and confined, respectively. The water recharge area of the free aquifer is influenced by the Santa Catarina and San Martín rivers. The predominant water family is  $\text{Ca-Na-HCO}_3$  and  $\text{Ca-HCO}_3$  (Figure 3A). These water families reflect that the most important source corresponds to the Santa Catarina River. Their main salt content corresponds to the interaction of the water with calcareous  $\text{Ca-HCO}_3$  family and movement of the water in the granular material, which consists predominantly products of the San Martín Avalanche (Torres-Hernández *et al.*, 2010; Torres-Rivera, 2012). The mixing of the  $\text{Ca-HCO}_3$  water type with volcanic rocks interaction generates  $\text{Ca-Na-HCO}_3$  family. A single sample corresponds to Sulfate-Bicarbonate-Calcium-Magnesium composition, and additionally, it has 1290 mg/L of TDS; reflecting above all a much higher salt enrichment than all the other samples analyzed in this valley. This high value is considered as evidence that it reflects a regional flow and corresponds to a confined aquifer (Torres-Rivera, 2012). It also represents a confined aquifer, since the well that captures these waters penetrated beneath a thick ignimbrite layer (Santa María Ignimbrite), which functions as a seal rock of the same.


The waters family in the sub-basin II of Ocampo-Paraíso, is  $\text{Ca-Na-HCO}_3$ , reflecting above all the interaction of the currents with calcareous rocks and the incorporation of the sodium liberated from the volcanic clasts in the granular medium.

In sub-basin III of San Nicolás, the  $\text{Ca-Na-HCO}_3$  and  $\text{Ca-Na-HCO}_3\text{-SO}_4$  families were documented at the south of the Morenos town. This type of water reflects the interaction of the water with limestone rocks, gypsum, and ion exchange in the valleys where there is the presence of clay materials.

#### *Hydrogeochemical processes with Gibbs diagrams*

To identify some hydrogeochemical processes that occur in the groundwater, diagrams are used such as Gibbs diagrams (Gibbs, 1970), which was developed for surface water studies; however, in recent decades it has been used for the study of groundwater. The three main mechanisms that control the chemistry of surface water can be defined as the domain of atmospheric precipitation (rain), the domain of the rock, and domain of the evaporation-crystallization process (Gibbs, 1970). The Gibbs

diagram was elaborated with the chemical data (anions and cations) of the groundwater samples (Figure 4).



**Figure 4.** Gibbs diagram, the main evolutionary process that we identify with this diagram is the interaction rock water; the yellow arrow indicates the direction of the evolution of groundwater.

Based on Figure 4, we can infer that during its evolution, the water was influenced mainly by the water-rock interaction and a little evaporation due to the evaporation seems to have been dominant only in two samples. When the water infiltrates the subsoil, the water-rock interaction begins, the water from the recent infiltration has fewer chemical components; while the greater distance traveled, the residence time, the more significant the physical and chemical components increase; this evolutionary behavior, can be seen in Figure 4. Member I is very close to the vertical axis, which is associated with the water in the recharge zone while member II is away from the vertical axis because of its significant evolution.

#### *Flow Systems*

The increase of chemical components during the evolution of the groundwater can be observed in the Mifflin diagram (Mifflin, 1968) and uses hydrogeochemical indicators ( $\text{Na}^+ + \text{K}^+$  vs.  $\text{Cl}^- + \text{SO}_4^{2-}$ ) to understand the evolution of groundwater and establish the different types of water flows (Figure 5). It is divided into three zones, and the area closest to the origin is due to the low circulation and short residence time of the water, the samples in this area are associated with the recharge (local flow). In the next zone, the concentrations increase and may be associated with a long-distance or longer residence time (intermediate flow).



Finally, the higher concentration is associated with more significant water-rock interaction and higher evolution of the groundwater (regional flow). In the study area, there are two types of water flow, one local and the other intermediate (Figure 5).


#### Hydrogeochemical Section

During its pathway, the groundwater interacts with the different materials, which results in the increase of the chemical components and in some cases, the decrease of these, due to the mixture with water of recent infiltration. To corroborate the water-rock interaction and mixtures a hydrogeological-hydrogeochemical, this section was elaborated.


The hydrogeochemical section is 40.5 km, begins in the Armadillo de los Infante, where volcanic rocks contact with the limestones and known as El Abra Formation, which cross the Ocampo valley of volcanic rocks and finally, cuts a valley where gypsum from the Guaxcamá Formation are found a few meters deep (Figure 6). (This section is only justified for the valleys of San Nicolas Tolentino and Ocampo Paraiso).

For the first sub-basin, the  $\text{Ca-Na-HCO}_3$  waters reflect the primary interaction with calcareous rocks of the El Abra Formation that crops out in the Sierra de Álvarez, which are partially dolomitized and contain varying amounts of magnesium in their dominant composition of  $\text{Ca-HCO}_3$ .  $\text{Ca-HCO}_3$  its transit through the granular material that fills the Graben de Peñillos (López-Doncel *et al.*, 2001) could print its sodium content indicates that this filling was deposited as a lacustrine fluvial material, which is due to having a shallow water tie. The location of the valley in an environment with medium to intense evaporation could include to some extent sodium. Another possible source of this sodium would be the decomposition of sodium plagioclase in the ignimbrite rocks, which interacts with the water currents and mix into the San Nicolas Tolentino river.

To the south of the Morenos town, where the San Nicolás River breaks its trajectory. As mentioned above, the first family waters ( $\text{Ca-HCO}_3$ ) acquire a character of Bicarbonate-Sulphated, perhaps due to the nearby influence of the gypsum that emerges in El Potrero anticline.



**Figure 5.** Mifflin diagram, classifies the water according to its evolution.




**Figure 6.** Hydrogeochemical section of Armadillo de los Infante to Paraiso; the numbers indicate several use; the discontinuous blue line indicates the piezometric level.

The physical parameters of sub-basins II and III, in the Armadillo de los Infante reveal that the electrical conductivity in the waters is approximately 650  $\mu\text{S}/\text{cm}$ , along the hydrogeochemical section. The highest point is located at the center of the section in Paraiso, due to the interaction of the groundwater with the limestones whereas lowest point at the end of the section in Ocampo Paraiso town, because there is a mixture of local recharge water through the volcanic rocks. The concentration of chlorides is high at the beginning of the section, with values of 25 mg/L due to the agricultural activity that takes place in that area. Concentrations tend to decrease toward Ocampo Paraiso where it is possible for groundwater to mix with waters of recent infiltration. The alkalinity starts with values of 320 mg/L in Armadillo de los Infante and increases in the Ocampo zone due to the

interaction of the water with the limestones, decreases to 240 mg/L at the end of the section due to mixing with water from recent infiltration (Figure 7).

The pH is 7.2 at the beginning and maintains little variation until Ocampo. However, it increases up to values of 7.6 and end with values of 7.2. The temperature starts at 20 °C and increases throughout the section until reaching 31 °C towards the end of the section in Paraiso. Sulfates begin with values close to 40 mg/L, increase slightly towards Ocampo town (60 mg/L) and on the Paraiso town increase to reach 100 mg/L; this increase can be related to the interaction of water with gypsums, which generates an exothermic reaction and this is congruent with the temperature anomaly in Paraiso.



**Figure 7.** Scatter diagrams of conservative elements (Br- and Li-) for the study area.

#### *Origin of the end-members and mixtures*

To evaluate the contributions to a hydrogeological system, the mixture models are used, for which it is necessary to identify the end-members representative of each region.

Based on the scatter plot of the conditional elements i.e., Li and Br, which do not react chemically with the environment during the evolution of groundwater are used in the present study. The study represents three end members ( $C_1$ ,  $C_2$ ,  $C_3$ ) and corresponds to a ternary mixing. Two members are the most evolved and corresponds to intermediate flows while third one is associated with local recharge.

In Figure 7, a ternary mixing is identified, which is formed by ( $C_1$ ) associated with sub-basin I (Santa Catarina) for local recharge, ( $C_2$ ) associated with sub-basin II (Ocampo-Paraíso) and ( $C_3$ ) associated with the flow of sub-basin III (San Nicolás Tolentino). This figure display a triangle, in whose vertices the end-members are located. The remaining samples are between the limits of the mixture lines and are considered as a mixture of the fractions of the end member. Any water sample from the system can be generated utilizing the three end-members.

Sub-basin I, is represented by sample 30, which located to the south of the study area (Santa Catarina), towards the Sierra de Álvarez (Figure 1) and associated with local recharge and content low values of  $Cl^-$ ,  $SO_4^{2-}$ ,  $Br^-$ , and  $Li^+$ . In this sample the lowest values were measured for  $Cl^-$  (0.22 meq/L),  $Li^+$  (0.128 meq/L) and 0.012 meq/L of  $Br^-$  (Figure 7); the measured temperature was of 14 °C, with a pH of 6.5. The type of this member is  $Ca-HCO_3$ .

Sub-basin II, sample 1, which corresponds to an intermediate flow, is located at the eastern-center of the study area in the Ocampo-Paraíso basin (Figure 1). The temperature measured in this sample was about 24.4°C and  $Cl^-$  concentrations of 0.84. meq/L; 2.46 meq/L of  $Li^+$ ; and, 0.24 meq/L of  $Br^-$  (Figure 7). The water type of this member is  $Na-K-HCO_3$ .

Sub-basin III, is represented by sample 16. This water sample represents an intermediate flow and is located towards the north of the study area, towards the Jagüey de San Francisco, belonging to the basin of San Nicolás Tolentino (Figure 1). It has high temperatures of 20.5°C and have concentrations of  $Cl^-$  of 0.819 meq/L; 0.312 meq/L of  $Li^+$ ; and high concentrations of  $Br^-$  (1.67 meq/L) (Figure 7). This type of water is  $Na-K-HCO_3$ .

In Table 2,  $C_1$  contributes 50 %,  $C_2$  contributes 31%, and  $C_3$  contributes 19% of water to the system (Table 2).

#### **Conclusions**

- The aquifer receives its recharge water that drains through flat platform rocks (El Abra Formation), gypsum, anhydrite (Guaxcama Formation) and volcanic rocks that delimit the aquifer.
- Three different types calcium bicarbonate ( $Ca-HCO_3$ ), sodium bicarbonate ( $Na-HCO_3$ ), calcium sulfate ( $Ca-SO_4$ ) waters were obtained.
- A local and an intermediate regional flow were identified, where each system has a representative sample used as an extra limb.
- A ternary mixture and water-rock interaction processes were identified in the study area.
- The system is fed by three essential regions.  $C_1$ ,  $C_2$  and  $C_3$  contributes 50%, 31% and 19%, respectively of water to the system.

**Table 2.** Mixing fractions considering three extreme members (samples 30, 1 and 16) for the groundwater of the Zona Media of San Luis Potosí.

Sample	C1	C2	C3	CT
1	0.00	1.00	0.00	1.00
2	0.01	0.20	0.79	1.00
3	0.63	0.24	0.13	1.00
4	0.74	0.25	0.01	1.00
5	0.64	0.17	0.19	1.00
6	0.35	0.31	0.33	1.00
7	0.27	0.64	0.09	1.00
8	0.44	0.54	0.02	1.00
9	0.35	0.55	0.11	1.00
10	0.44	0.52	0.04	1.00
11	0.15	0.71	0.14	1.00
12	0.87	0.03	0.09	1.00
13	*	*	*	*
14	0.57	0.35	0.08	1.00
15	0.54	0.38	0.07	1.00
16	0.00	0.00	1.00	1.00
17	0.48	0.39	0.13	1.00
18	0.58	0.44	0.00	1.00
19	0.44	0.42	0.14	1.00
20	0.72	0.28	0.00	1.00
21	0.72	0.01	0.27	1.00
22	0.75	0.20	0.05	1.00
23	0.75	0.21	0.03	1.00
24	0.63	0.29	0.08	1.00
25	0.81	0.15	0.04	1.00
26	0.40	0.26	0.35	1.00
27	0.00	0.09	0.91	1.00
28	0.74	0.26	0.00	1.00
29	0.77	0.23	0.00	1.00
30	1.00	0.00	0.00	1.00
31	0.94	0.02	0.03	1.00
32	0.73	0.27	0.00	1.00
33	0.86	0.13	0.01	1.00
34	0.71	0.17	0.12	1.00

## References

- Abu Jaber, N., 2001, Geochemical evolution and recharge of the shallow aquifers at Tulul al Ashiqif, NE Jordan: *Environmental Geology*, vol. 41, 372-383pp.
- Antigüedad, I., Morales, T., Uriarte, J.A., 2008, Los acuíferos kársticos. Caso del país Vasco. Grupo de Hidrogeología. Geodinamica Siala. Universidad del país Vasco-Euskal Herriko. 8pp.
- Andreo, B., Vadillo, I., Carrasco, F., Neukum, C., Jiménez, P., Goldscheider, N., Hötzl, H., Vías, J.M., Pérez, I. y Göppert, N., 2004, Precisiones sobre el funcionamiento hidrodinámico y la vulnerabilidad a la contaminación del acuífero kárstico de la Sierra de Líbar (provincias de Málaga y Cádiz, sur de España) a partir de un ensayo de trazadores. *Revista de la Sociedad Geológica de España*, 17 (3-4): 187-197pp.
- Apello, C.A., Postma, D., 1996, *Geochemistry, Groundwater and Pollution*: Rotterdam, Netherlands, A.A. Balkema, 536 pp.
- Armienta, E.K., Zamora y Juárez, F., 1987, Manual para el análisis químico de aguas naturales, en campo y laboratorio. Comunidades técnicas. Serie Docencia y Divulgación, Num. 4. México D.F. Instituto de Geofísica de la UNAM, 86pp.
- Carrillo-Bravo, J., 1971, La Plataforma Valles - San Luis Potosí: *Boletín de la Asociación Mexicana de Geólogos Petroleros*, vol. 23, 1-102pp.
- Deutsch, W.J., 1997, *Groundwater Geochemistry: fundamentals and applications to Contamination*: New York, CRP-Press 232pp.
- Douglas, M., Clark, I.D., Raven, K., Bottomley, D., 2000, Groundwater mixing dynamics at a Canadian Shield mine: *Journal of Hydrology*, 235, 88-103pp.
- Freeze, R.A., and Cherry, J.A., 1979, *Groundwater*: Englewood Cliffs, NJ, Prentice-Hall, 604 pp.
- Genereux, D., 2004, Comparison of naturally-occurring chloride and oxygen-18 as tracers of interbasin groundwater transfer in low land rain forest, Costa Rica: *Journal of Hydrology*, 295(1-4), 17-27pp.
- Gibbs, R.J., 1970, Mechanisms controlling world water chemistry: *American Association for the Advancement of Science, New Series*, vol. 170, No.3962, 1088-1090pp.
- Harpaz, Y. and Bear, J., 1963, Investigations on mixing of waters in subsoil storage operations, *International Association Scientific Hydrology*, vol. 64, 132-153pp.
- Instituto Nacional de Estadística y Geografía (INEGI), 2002, *Estudio Hidrológico del estado de San Luis Potosí*.
- Kiraly, L., y Müller, I., 1979, Heterogeneité de la perméabilité et de l'alimentation dans le

- karst, effect sur la variation du chimisme des sources karstiques. Bull. Center Hydrogéol. Neuchatel, vol. 3, 237-285pp.
- Laaksoharju, M., Skarman, C., Skarman, E., 1999, Multivariate mixing and mass balance (M3) calculations, a new tool for decoding hydrogeochemical information: Applied Geochemistry, vol.14, 861-871pp.
- Labarthe Hernández, G., Tristán González, M., Aranda Gómez, J.J., 1982, Revisión estratigráfica del Cenozoico de la parte central del Estado de San Luis Potosí. Universidad Autónoma de San Luis Potosí, Instituto de Geología y Metalurgia, Folleto Técnico No. 85, 205 pp.
- Lee, E.S., Krothe, N.C., 2001, A four component mixing model for water in a karst terrain in south central Indiana, USA using solute concentration and stable isotopes as tracers: Chemical Geology, 179, 129-143pp.
- López-Doncel, R.A., 2003. Formación Tamabra del Cretácico Medio de la porción central del margen occidental de la Plataforma Valles-San Luis Potosí, Centro-Noreste de México. Revista Mexicana de Ciencias Geológicas Vol. 20, n. 1: 1-19.
- López-Doncel, R. A., Soto-Araiza, R. G., Dircio-Castro, D., 2001, Carta Geológico Minera, Peotillos F14-A75 San Luis Potosí, Servicio Geológico Mexicano
- Mifflin, M.D., 1968, Delineation of groundwater flow systems in Nevada, Report series H-W., Desert research Institute Technical, vol. 4, 111pp.
- Piper, A.M., 1994, A graphic procedure in the geochemical interpretation of water analyses. Am. Geophysical Union Trans. Vol. 25, 914-923p.
- Ramos Leal J.A., Martínez Ruiz V.J., Rangel Méndez J.R. y Alfaro de la Torre M.C., 2007, Hidrogeological and mixing process of waters in aquifers in arid regions: a case study in San Luis Potosí Valley, Mexico. Environ Geol 53:325-337pp.
- Rice, K.C., Hornberger, G.M., 1998, Comparison of hydrochemical tracers to estimate source contributions to peak flow in a small, forested, head water catchment: Water Resources Research, 34(7), 1755-1766pp.
- Roback, R.C., Johnson, T.M., McLing, T.L., Murrell, M.T., Luo, S., Ku, T.L., 2001, Uranium isotopic evidence for groundwater chemical evolution and flow patterns in the eastern Snake River Plain Aquifer, Idaho: Geological Society of America Bulletin, 113 (9), 1133-1141pp.
- Skalbeck, J.D., Shevenell, L., Widmer, M.C., 2002, Mixing of thermal and non thermal waters in the Steamboat Hills area, Nevada, USA: Geothermics, 31, 60-90pp.
- Torres-Hernández, J.R., Rodríguez-Ríos, R., Barboza-Gudiño, J.R., Saucedo-Girón, R., Tristán-Capetillo, L.C., García-Pérez, K., 2010, Cartografía Geológico Minera Santa Catarina F14-A-85 San Luis Potosí, Servicio Geológico Mexicano.
- Torres-Rivera, S., 2012 Caracterización hidrogeoquímica y direcciones de flujo de agua subterránea en el acuífero de San Nicolás Tolentino. Tesis de Maestría. Maestría en Ciencias en Geología Aplicada, Facultad de Ingeniería de la Universidad Autónoma de San Luis Potosí. 105 p.
- Valentino, G.M., Stanzione, D., 2002, Source process of the thermal waters from the Phlegraean Fields (Naples, Italy) by means of selected minor and trace elements distribution: Chemical Geology, 245, 245-274pp.
- Wallick, E.I., 1981, Chemical evolution of groundwater in a drainage basin f Holocene age, east-central Alberta, Canada: Journal of Hydrology, 54, 245-283pp.
- Zapata Zapata, J.L., Pérez Venzor, J.A., 1979, Cartografía Geológica Hoja Peotillos, S.L.P, Universidad Autónoma de San Luis Potosí, Instituto de Geología y Metalurgia, Folletos Técnicos N.63, 30pp.