

Autopsy and Case Reports

ISSN: 2236-1960

Hospital Universitário da Universidade de São Paulo

Daisley Jr, Hubert; Rampersad, Arlene; Daisley, Martina; Ramdin, Amit; Acco, Oneka; Narinesingh, Farhaana; Humphrey, Ornella; James, Errol The vasa vasorum of the large pulmonary vessels are involved in COVID-19 Autopsy and Case Reports, vol. 11, e2021304, 2021 Hospital Universitário da Universidade de São Paulo

DOI: https://doi.org/10.4322/acr.2021.304

Available in: https://www.redalyc.org/articulo.oa?id=576067146101



Complete issue

More information about this article

Journal's webpage in redalyc.org



Scientific Information System Redalyc

Network of Scientific Journals from Latin America and the Caribbean, Spain and Portugal

Project academic non-profit, developed under the open access initiative





The vasa vasorum of the large pulmonary vessels are involved in COVID-19

Hubert Daisley Jr¹ , Arlene Rampersad¹ , Martina Daisley² , Amit Ramdin¹ , Oneka Acco¹ , Farhaana Narinesingh¹ , Ornella Humphrey³ , Errol James⁴

How to cite: Daisley Jr. H, Rampersad A, Daisley M, et al. The vasa vasorum of the large pulmonary vessels are involved in COVID-19. Autops Case Rep [Internet]. 2021;11:e2021304. https://doi.org/10.4322/acr.2021.304

Keywords

Blood Vessels; Coronavirus Infection; Vasa Vasorum

DEAR EDITOR

In our manuscript "COVID-19: a closer look at the pathology in two autopsied cases. Is the pericyte at the center of the pathological process in COVID-19?"; published in Volume 11, 2021 issue of Autopsy Case Reports; Figures 3B, 4A, 4B, and 4D show thrombosis within the walls of large pulmonary vessels. These locations of thrombosis may represent the location of the vasa vasorum of these pulmonary vessels. Figures 3B and 4A may represent thrombus within vasorum externa, whilst figures 4B and 4D represent thrombus within vasorum interna.²

The significance of these sites means that both, the alveolar microcirculation and the vasa vasorum of the large pulmonary vessels are involved in the thrombotic processes in COVID-19. The latter thrombotic site would further implicate COVID-19 as a disease of the microcirculation. Boyle and Haverich³ postulated that "the involvement of the large vessels during coronavirus disease 19 (COVID-19) in both children and adults is likely due to dysfunction of their vasa vasorum and SARS-COV-2 induced microthrombosis

of vasa vasorum would lead to hypoxic conditions in the adventitia."

As the world is grappling with the SARS-COV-2 pandemic, it is essential to understand the pathogenesis of COVID-19 to formulate therapeutic options. It may be that the lung is a secondary partner in COVID-19 and the pathogenesis of this disease begins in the nasopharynx,⁴ where there is an abundance of ACE-2 receptor sites. SARS-COV-2 interaction at the port of entry in the nasopharynx produces chemokines/cytokines,^{3,5} which circulates in the systemic circulation, causing endotheliopathy and subsequent thrombotic microangiopathy⁶ being the ultimate result.

REFERENCES

- 1. Daisley H Jr, Rampersad A, Daisley M, et al. COVID-19: a closer look at the pathology in two autopsied cases. Is the pericyte at the center of the pathological process in COVID-19? Autops Case Rep. 2021;11:e2021262. http://dx.doi.org/10.4322/acr.2021.262. PMid:34307223.
- 2. Gössl M, Rosol M, Malyar NM, et al. Functional anatomy and hemodynamic characteristics of vasa vasorum in the

⁴ Beacon Plastic and Cosmetic Surgery Clinic Woodbrook Port- of Spain, Trinidad and Tobago, West Indies



¹ San Fernando General Hospital, Department of Pathology, San Fernando, Trinidad and Tobago

² Scarborough General Hospital, Department of Accident and Emergency, Tobago, West Indies

³ University of the West Indies, Trinidad, West Indies

- walls of porcine coronary arteries. Anat Rec A Discov Mol Cell Evol Biol. 2003;272(2):526-37. http://dx.doi.org/10.1002/ar.a.10060. PMid:12740947.
- 3. Boyle EC, Haverich A. Microvasculature dysfunction as the common thread between atherosclerosis, Kawasaki disease, and severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) associated multi-system inflammatory syndrome in children. Eur J Cardiothorac Surg. 2020;58(6):1109-10. http://dx.doi.org/10.1093/ejcts/ezaa367. PMid:33247736.
- 4. Gallo O, Locatello LG, Mazzoni A, Novelli L, Annunziato F. The central role of the nasal microenvironment in the transmission, modulation, and clinical progression of SARS-

- CoV-2 infection. Mucosal Immunol. 2021;14(2):305-16. http://dx.doi.org/10.1038/s41385-020-00359-2. PMid:33244161.
- 5. Loo J, Spittle DA, Newnham M. COVID-19, immunothrombosis and venous thromboembolism: biological mechanisms. Thorax. 2021;76(4):412-20. http://dx.doi.org/10.1136/thoraxjnl-2020-216243. PMid:33408195.
- 6. Tiwari NR, Phatak S, Sharma VR, Agarwal SK. COVID-19 and thrombotic microangiopathies. Thromb Res. 2021;202:191-8. http://dx.doi.org/10.1016/j. thromres.2021.04.012. PMid:33894421.

Conflict of interest: The authors have no conflict of interest to declare.

Financial support: The authors declare that no financial support was received.

Submitted on: May 15th, 2021 **Accepted on:** June 1st, 2021

Correspondence

Hubert Daisley Jr.

Department of Pathology, San Fernando General Hospital

Paradise Pasture, Independence Avenue, San Fernando, Trinidad and Tobago, West Indies

Phone +1 (868) 225-4325

profhdjr@aol.com