



Autopsy and Case Reports

ISSN: 2236-1960

Hospital Universitário da Universidade de São Paulo

Lopes, José Manuel

Neuroendocrine neoplasms: a brief overview emphasizing gastroenteropancreatic tumors

Autopsy and Case Reports, vol. 7, no. 1, 2017, January-March, pp. 1-4

Hospital Universitário da Universidade de São Paulo

DOI: <https://doi.org/10.4322/acr.2017.004>

Available in: <https://www.redalyc.org/articulo.oa?id=576068163001>

- How to cite
- Complete issue
- More information about this article
- Journal's webpage in redalyc.org

UABM [redalyc.org](https://www.redalyc.org)

Scientific Information System Redalyc

Network of Scientific Journals from Latin America and the Caribbean, Spain and Portugal

Project academic non-profit, developed under the open access initiative

Neuroendocrine neoplasms: a brief overview emphasizing gastroenteropancreatic tumors

José Manuel Lopes^{a,b,c,d}

Lopes JM. Neuroendocrine neoplasms: a brief overview emphasizing gastroenteropancreatic tumors. [editorial]. Autopsy Case Rep [Internet]. 2017;7(1):1-4. <http://dx.doi.org/10.4322/acr.2017.004>

Neuroendocrine neoplasms are considered to be composed of cells with features similar to those of the normal diffuse neuroendocrine system (DNES) scattered throughout the body (e.g. gastroenteropancreatic [GEP], the enteroendocrine hormonal signaling system; respiratory and urogenital systems; paraganglia; adrenal medulla; skin; thymus; heart; middle ear; and other tissues). Despite being considered rare, neuroendocrine neoplasm incidence has been reported to be increasing, with increased survival duration over time, suggesting that neuroendocrine tumors (NETs) are more prevalent than previously reported.¹ The present overview addresses the main focus on DNES cells and the anatomic pathology features of neuroendocrine neoplasms that occur in GEP.

The recognition of GEP DNES seems to have started in the 19th century. In 1870, Heidenhain² reported the presence of chromaffin cells (EC), a term later introduced by Ciaccio,³ in the gastric mucosa, probably the current gastric counterpart of the intestinal Enterochromaffin Cell: EC-Like (ECL) but it was Kultschitzky⁴ who reported a seminal study describing EC in Lieberkühn crypts of the intestinal mucosa. Then, Feyrter⁵ established the concept of a “diffuse” endocrine system (DNES) in the gastrointestinal mucosa. Neuroendocrine cells have been independently recognized by several

authors and variably termed enterochromaffin cells, argentaffin and argyrophil “clear cells,” enteroendocrine cells, and Kultschitzky cells. Pearse⁶ recognized that neuroendocrine cells of the gut uptake 5-hydroxytryptophan, which is decarboxylated to 5-hydroxytryptamine. Rosai,⁷ in a remarkable essay, presented claims favoring the endodermal origin of the GEP DNES, but pointed out that there was still evidence missing to definitively establish this possibility. The neuroendocrine cells express general markers of allegedly neuroendocrine differentiation (e.g. chromogranin and synaptophysin)—since a common phenotype does not mean a common histogenesis—and hormones according to site (e.g. a more “neural-like” differentiation in the larynx, lung, thymus, and thyroid; a more “epithelial-like” differentiation in the GEP system), and that an interplay seems to exist between these neuroendocrine cells and neural crest-derived nerve endings.

The first autopsy report of a GEP neuroendocrine neoplasm is ascribed to Langhans,⁸ which was followed by reports of Lubarsch⁹ and Ramson.¹⁰ Of note, it seems probable that the first description of a carcinoid tumor was reported in the bronchus by Laennec.¹¹ The term “carcinoid” was established (1907) by Oberndofer,¹² who, in 1929,¹³ confirmed the possibility

^a Department of Pathology - Faculty of Medicine - University of Porto, Porto – Portugal.

^b Department of Pathology - Centro Hospitalar São João, Porto – Portugal.

^c Institute of Pathology and Immunology - University of Porto, Porto – Portugal.

^d Institute of Health Investigation and Innovation - University of Porto, Porto – Portugal.



that “karzinoide,” meaning “carcinoma-like,” might exhibit malignant features and metastasize.

Williams and Sandler¹⁴ reported three distinct groups of GEP neuroendocrine neoplasms: foregut (intrathoracic, gastric, and duodenal), midgut (small intestine, appendix, and proximal colon), and hindgut (distal colon and rectum). In 2000, the WHO (reviewed by Klöppel et al.¹⁵) established the following categories based on the degree of differentiation: well-differentiated endocrine tumors (further classified into benign tumors and low-grade malignant tumors, based on the tumor size, mitotic rate, Ki-67 labeling index, lymphovascular invasion, and symptoms, in association with hormonal oversecretion), and poorly differentiated endocrine (small- and large-cell) carcinomas. The WHO 2010¹⁶ classification established GEP neuroendocrine (well- and poorly-differentiated) neoplasms as malignant tumors, except for gangliocytic paraganglioma and pancreatic neuroendocrine microadenomas, which are considered to be benign tumors; and L-cell-type (glucagon-like peptide and peptide YY-producing) NETs and tubular carcinoids, as uncertain malignancies.

Well-differentiated NETs are reported to occur most frequently in GEP sites up to two-thirds (more in the small intestine, but variably according to different reports), a quarter in the bronchopulmonary system, and the remaining 10% scattered throughout the body. In contrast, neuroendocrine carcinomas occur predominantly in the bronchopulmonary system.¹⁷

There are GEP neuroendocrine neoplasms associated with hereditary syndromes¹⁸ (e.g. multiple neuroendocrine neoplasia type 1 [MEN1], von Hippel-Lindau syndrome, neurofibromatosis 1, and tuberous sclerosis [TSC]), but most are considered sporadic. Well-differentiated sporadic pancreatic neuroendocrine neoplasms were reported to display a high frequency of mutations¹⁹ in *MEN1*, *ATRX* (alpha thalassemia/mental retardation syndrome X-linked), and *DAXX* (death-domain associated protein), which seem to associate with better behavior; and low frequency mutations in genes of the mTOR (mammalian target of rapamycin) pathways, such as *PTEN* (phosphatase and tensin homolog), *TSC2*, and *PIC3CA* (phosphatidylinositol-4,5-bisphosphate 3-kinase catalytic subunit alpha). Interestingly, it seems that telomerase promoter mutations in

pancreatic NETs are rare,²⁰ but mainly found in tumors from patients with hereditary syndromes. Strikingly, most of the molecular mechanisms of neuroendocrine neoplasms remain to be clarified.

The current staging of GEP neuroendocrine neoplasms is based on the proposals of the American Joint Committee on Cancer (AJCC)²¹ and the European Neuroendocrine Tumor Society (ENETS).^{22,23} The AJCC and ENETS staging for almost all gastrointestinal neuroendocrine neoplasms—those of the stomach, duodenum, ampulla, jejunum, ileum, and colon-rectum—are identical; whereas some differences exist in the T classification of the pancreatic and appendiceal neuroendocrine neoplasms between AJCC and ENETS staging.

GEP neuroendocrine neoplasms are graded^{22,23} based on the mitotic count (MC)/10 high power fields (HPFs) and the Ki-67 proliferation index (Ki-67 LI), regardless of tumor size, extent, or location; a minimum of 50 HPFs (1 HPF = 2 mm²) should be carefully evaluated to accurately determine the MC, and 500–2000 cells to determine the percentage of Ki-67 LI. GEP neuroendocrine neoplasms are usually graded as (well-differentiated) G1 (MC < 2/10HPF and/or ≤ 2% Ki-67LI) and G2 (MC 2-20/10 HPF and/or 3-20% Ki-67 LI) NETs, and (poorly-differentiated, small or large cell) G3 neuroendocrine carcinomas (NECs; MC > 20/10HPF and/or > 20% Ki-67 LI). It is worth noting that there are reports²⁴ of “well-differentiated G3 neuroendocrine neoplasms” that seem to display less aggressive behavior than bona fide NECs.

It seems that up to 50% of NETs display synchronous or metachronous metastases after diagnosis (more common in poorly differentiated neoplasms), and appear to occur more frequently when the well-differentiated neuroendocrine primary neoplasm is located in the cecum or pancreas.²⁵ Most metastases of neuroendocrine neoplasms occur in the liver, lungs, and bone, and are much rarer in other sites²⁶ (e.g. brain, ear, ovary, breast, thyroid, pancreas, skin, adrenal glands, kidney, spleen, orbit, retroperitoneum, testis, and pituitary gland).

Conclusively, it seems that, as in the first medical identification of a GEP neuroendocrine neoplasm, autopsy studies are still very much justified²⁷ to clarify the pathogenesis of cancer in general, and of NETs in

particular, especially in unusual cases and/or in cases submitted to the new medical therapies of cancer.

REFERENCES

1. Yao JC, Hassan M, Phan A, et al. One hundred years after "Carcinoid": epidemiology of and prognostic factors for neuroendocrine tumors in 35,825 cases in the United States. *J Clin Oncol*. 2008;26(18):3063-72. PMID:18565894. <http://dx.doi.org/10.1200/JCO.2007.15.4377>.
2. Heidenhain R. Untersuchungen über den Bau der Labdrüsen. *Arch Mikr Anat*. 1870;6(1):368-406. <http://dx.doi.org/10.1007/BF02955986>.
3. Ciaccio MC. Sur une nouvelle espèce cellulaire dans les glandes de Lieberkühn. *CR Seances Soc Biol Fil*. 1906;60:76-7.
4. Kultschitzky N. Zur Frage über den Bau des Darmkanals. *Arch Mikr Anat*. 1897;49(1):7-35. <http://dx.doi.org/10.1007/BF02975387>.
5. Feyrter F. Über diffuse endocrine epitheliale Organe. *Zentralbl Innere Med*. 1938;545:31-41.
6. Pearse AG. 5-hydroxytryptophan uptake by dog thyroid "C" cells, and its possible significance in polypeptide hormone production. *Nature*. 1966;211(5049):598-600. PMID:5298184. <http://dx.doi.org/10.1038/211598a0>.
7. Rosai J. The origin of neuroendocrine tumors and the neural crest saga. *Mod Pathol*. 2011;24(Suppl 2):S53-7. PMID:21455201. <http://dx.doi.org/10.1038/modpathol.2010.166>.
8. Langhans T. Ueber einen Drüsenpolyp im ileum. *Virchows Arch Pathol Anat*. 1867;38:550-60.
9. Lubarsch O. Ueber dem primären Krebs des Ileum nebst Bemerkungen über das gleichzeitige Vorkommen von Krebs und Tuberculose. *Virchows Arch Pathol Anat*. 1888;111(2):280-317. <http://dx.doi.org/10.1007/BF01966242>.
10. Ramson W. A case of primary carcinoma of the ileum. *Lancet*. 1890;2:1020-3.
11. Laennec R. *Traité de l'auscultation médiate e des maladies des poumons e du couer*. Paris; 1831.
12. Oberndofer S. Karzinoide tumoren des dünndarms. *Frankf Z Pathol*. 1907;1:426-43.
13. Oberndofer S. *Karzinoide*. Berlin: Verlag von Julius Springer; 1929.
14. Williams ED, Sandler M. The classification of carcinoid tumors. *Lancet*. 1963;1(7275):238-9. PMID:14000847. [http://dx.doi.org/10.1016/S0140-6736\(63\)90951-6](http://dx.doi.org/10.1016/S0140-6736(63)90951-6).
15. Klöppel G, Perren A, Heitz PU. The gastroenteropancreatic neuroendocrine cell system and its tumors: the WHO classification. *Ann N Y Acad Sci*. 2004;1014(1):13-27. PMID:15153416. <http://dx.doi.org/10.1196/annals.1294.002>.
16. Bosman FT, Carneiro F, Hruban RH, Theise ND, editors. *WHO Classification of Tumours of the Digestive System*. 4th ed. Lyon: IARC Press; 2010.
17. Hauso O, Gustafsson BI, Kidd M, et al. Neuroendocrine tumor epidemiology: contrasting Norway and North America. *Cancer*. 2008;113(10):2655-64. PMID:18853416. <http://dx.doi.org/10.1002/cncr.23883>.
18. Jensen RT, Berna MJ, Bingham DB, Norton JA. Inherited pancreatic endocrine tumor syndromes: advances in molecular pathogenesis, diagnosis, management and controversies. *Cancer*. 2008;113(7, Suppl):1807-43. PMID:18798544. <http://dx.doi.org/10.1002/cncr.23648>.
19. Jiao Y, Shi C, Edil BH, et al. DAXX/ATRX, MEN1, and mTOR pathway genes are frequently altered in pancreatic neuroendocrine tumors. *Science*. 2011;331(6021):1199-203. PMID:21252315. <http://dx.doi.org/10.1126/science.1200609>.
20. Vinagre J, Nabais J, Pinheiro J, et al. TERT promoter mutations in pancreatic endocrine tumours are rare and mainly found in tumours from patients with hereditary syndromes. *Sci Rep*. 2016;6:29714. PMID:27411289. <http://dx.doi.org/10.1038/srep29714>.
21. Edge SB, Compton CC. The American Joint Committee on Cancer: the 7th edition of the AJCC cancer staging manual and the future of TNM. *Ann Surg Oncol*. 2010;17(6):1471-4. PMID:20180029. <http://dx.doi.org/10.1245/s10434-010-0985-4>.
22. Rindi G, Klöppel G, Alhman H, et al. TNM staging of foregut (neuro)endocrine tumors: a consensus proposal including a grading system. *Virchows Arch*. 2006;449(4):395-401. PMID:16967267. <http://dx.doi.org/10.1007/s00428-006-0250-1>.
23. Rindi G, Klöppel G, Couvelard A, et al. TNM staging of midgut and hindgut (neuro)endocrine tumors: a consensus proposal including a grading system. *Virchows Arch*. 2007;451(4):757-62. PMID:17674042. <http://dx.doi.org/10.1007/s00428-007-0452-1>.
24. Basturk O, Yang Z, Tang LH, et al. The high-grade (WHO G3) pancreatic neuroendocrine tumor category is morphologically and biologically heterogeneous and includes both well differentiated and poorly differentiated neoplasms. *Am J Surg Pathol*. 2015;39(5):683-90. PMID:25723112. <http://dx.doi.org/10.1097/PAS.0000000000000408>.
25. Yao JC, Hassan M, Phan A, et al. One hundred years after "Carcinoid": epidemiology of and prognostic factors for neuroendocrine tumors in 35,825 cases in

- the United States. *J Clin Oncol*. 2008;26(18):3063-72. PMID:18565894. <http://dx.doi.org/10.1200/JCO.2007.15.4377>.
26. Pavel M, Grossman A, Arnold R, et al. ENETS Consensus Guidelines for the Management of Brain, Cardiac and Ovarian Metastases from Neuroendocrine Tumors. *Neuroendocrinology*. 2010;91(4):326-32. PMID:20453466. <http://dx.doi.org/10.1159/000287277>.
27. Budczies J, von Winterfeld M, Klauschen F, et al. The landscape of metastatic progression patterns across major human cancers. *Oncotarget*. 2015;6(1):570-83. PMID:25402435.

Correspondence

José Manuel Lopes
Department of Pathology - Centro Hospitalar São João
Alameda Prof. Hernâni Monteiro 4200-319 – Porto – Portugal
Phone: +351 (225) 512-100
jmlopes@ipatimup.pt