

Studies on mutagenic sensitivity of seeds of pummelo (*Citrus maxima* Merr.)

Sankaran, M; Kalaivanan, D; Sunil Gowda, D.C

Studies on mutagenic sensitivity of seeds of pummelo (*Citrus maxima* Merr.)

Journal of Horticultural Sciences, vol. 16, núm. 1, 2021

Society for Promotion of Horticulture, India

Disponible en: <https://www.redalyc.org/articulo.oa?id=577074023013>

DOI: <https://doi.org/10.24154/JHS.2021.v16i01.013>



Esta obra está bajo una Licencia Creative Commons Atribución-NoComercial-CompartirIgual 4.0 Internacional.

Studies on mutagenic sensitivity of seeds of pummelo (*Citrus maxima* Merr.)

M Sankaran kmsankaran@gmail.com

Institute of Horticultural Research Hesaraghatta Lake Post, India

D Kalaivanan

ICAR-Indian Institute of Horticultural Research Hesaraghatta Lake Post, India

D.C Sunil Gowda

ICAR-Indian Institute of Horticultural Research Hesaraghatta Lake Post, India

Journal of Horticultural Sciences, vol. 16, núm. 1, 2021

Society for Promotion of Horticulture, India

Recepción: 28 Octubre 2020
Aprobación: 15 Abril 2021

DOI: <https://doi.org/10.24154/JHS.2021.v16i01.013>

Redalyc: <https://www.redalyc.org/articulo.oa?id=577074023013>

Abstract: Mutation breeding is a key method of generating large number of heritable variations. Effective dose (LD50) needs to be standardized for inducing sufficient variation in a crop. In the present study, seeds were irradiated with different doses of Gamma rays and found that 66.94 Gy could suppress germination close to 50 per cent (LD50) in pummelo. This 60 Gy gamma dose can effectively be used for raising the mutant populations to identify a desirable mutation in pummelo.

Keywords: Gamma Irradiation, Germination, Gray (Gy), LD⁵⁰, Mutation and Pummelo.

Induced mutation plays a significant role in the crop improvement of horticultural crops. It is an important tool for induction of variation in quantitative and qualitative characters. It can be a supplement to conventional breeding methods when it is desired to improve one or two characters in a well-adapted variety. Induced mutation improving crops cultivars, enhancing biodiversity and Gamma irradiations are safe for human and environment and can be used widely to develop new varieties in fruit crops. In the recent past, mutation using gamma rays is regularly attempted in the banana breeding program (Smith *et al.*, 2006 and Mishra *et al.*, 2007). Among the different strategies to enhance crop improvement programs, induced mutagenesis has contributed immensely by creating mutant varieties with improved and desirable genetic changes in agronomically important traits in crops. Much progress has been made in generating superior genotype with favourable attributes through induced mutations in fruit crops. Pummelo is the largest citrus fruit and known in the western world mainly as the principal ancestor of the grapefruit. Pummelo fruit, like all fruits of the citrus family has several health benefits because of its super – rich Vitamin C and Vitamin B content. It also contains Vitamin A, Vitamin B1, B2 and C, bioflavonoid, healthy fats, protein, fibre, antioxidants and enzymes. It bears medium sized fruits (1-1.5kg) with good TSS (8-10 OB) value. The pulp texture of pummelo fruit is fleshy and pulp colour is pink.

Excessive bitter taste in citrus juice is a major problem in citrus industry worldwide because it reduces the quality and commercial value of the product (Mongkolkul *et al.*, 2006). The bitterness in citrus fruit is affected by limonin and naringin, which are generally recognized as the major two bitter compounds. Limonin is the bitter limonoid found in major citrus cultivars such as grapefruit, the Navel orange, and the Shamouti orange (Guadagni *et al.*, 1973). Naringin is not transported after being synthesized in the fruit or leaves. However, till date, there is no information about availability of sweet pummelo varieties or bitter free pummelo varieties across the globe. Hence, the present investigation is aimed to determine the optimal dosage of gamma irradiation (LD50) for mutation induction in seeds of pummelo genotype “Kallar Selection” (Deep pink pulp with high bitterness) for developing a mutant with desirable horticultural traits.

The present investigation was carried out at ICAR- Indian Institute Horticultural Research, Bengaluru- 560089. The seeds of Kallar Selection (Accession-6) pummelo were irradiated with different doses of gamma rays. The fresh and physically pure quality seeds of pummelo fruit were used for the irradiation purpose. In first step of experiment, a preliminary study was conducted to know the sensitivity of pummelo dry seeds to gamma radiation in which, pummelo seeds irradiated with gamma rays at the doses of 100 Gy, 200 Gy, 300 Gy, 400 Gy and 500 Gy. In second step of the experiment, based on preliminary results the pummelo fresh seeds were irradiated with gamma doses of 25Gy, 50Gy, 60Gy, 75Gy and 100Gy to determine the exact LD50. The irradiated seeds along with non-irradiated seeds (control) were sown in protrays filled with cocopeat and each protray was marked with given gamma dose. Thereafter, water was sprinkled over the protray to provide enough moisture for seed germination and it was kept in controlled condition. Different parameters (seed germination per cent, number of seed germinated and number of seed deformities during germination) related to determining LD50 was recorded in 60 days after sowing. The data were subjected to Chi-Square analysis and Chi- Square table was used for the calculation error degrees of freedom.

For any induced mutagenesis programme, it is necessary to fix the LD50 value based on which larger population can be raised to isolate the desirable mutant progeny/progenies. The LD50 value varies according to crop species, varieties, seeds or other planting materials, nature of treatment, method of raising, climate, cultural practices and other parameters (Singh, 1994). In Citrus, several attempts to induce variability with some traits of seedless, thorn less, color changed fruits and juices (Maluszynski *et al.*, 2000). The radio-sensitivity (LD50) of acute citrus exposure ranges from 40 to 100 Gy (Sanada and Amano, 1998; Sparrow *et al.*, 1968), depending on the species and variety. The bud wood of pummelo Nambangan was irradiated with the dosage of 20, 40, and 60 Gy. The buds then grafted to Japanese Citron rootstocks. After three years of selection based on performance and fruit evaluation on MV2 generation, a mutant plant derived from 20 Gy irradiation treatments

were obtained with improved character on the number of seeds. Pummelo Nambangan has more than 40 seeds/ fruit and the mutant has less than 10 seeds/fruit on average and it shows higher volume of juice compared to that in pummelo Nambangan. However, the fruit appearance between mutant plant and parent in term of the fruit shape had no difference, showing a combination of spheroid and pyriform shape. The difference between them was shown prominently when the fruit was cut, indicating the less seed contained and more intense red color of flesh of the mutant fruit compared to that of parent (Mariana *et al.*, 2018.) In the initial study, pummelo seeds were treated with 100, 200, 300, 400 and 500 Gy). The effect of different doses of gamma rays ranging from 0 to 100 Gy on seed germination are shown in Table 1 and shown in Fig.1. The percentage of seed germination ranged from 18.31 per cent to 95.07percent with different doses of gamma rays in pummelo (Acc-06) as compared to 97.96 per cent in control. Gradual reduction/ decrease in pummelo seed germination was observed with increase in gamma irradiation dose. The inhibitory effect on seed germination was directly proportional to the dose of gamma radiation. Similar results were reported by Dhatt et al. (2000) and Latado et al. (2001) with gamma radiation in citrus. Decrease in percent seed germination by gamma irradiation might be due to its effect on genetical and cytological processes coupled with the changes induced in metabolic processes. The decrease in seed germination was mainly due to the interference of mutagens with metabolic activities of the seeds (Sjodin, 1962). Sinha and Godward (1972) opined that the reduction in percentage of seed germination was due to the disturbances caused at the physiological level coupled with chromosomal damage. Disturbance in the formation of enzymes involved in the germination process may be one of the physiological effects caused by mutagenic treatments (Kulkarni,2011). Gamma radiation is well known for their action causing point mutations, enzyme inhibitions and chromosomal aberrations. The observed reduction in seed germination in pummelo as a result of gamma radiation might be due to point mutations or the injuries caused to the genetic material. This may eventually lead to decrease the rate of respiration and energy production, which finally caused decrease in seed germination. Days taken to germination and number of seeds with deformity were increased with increasing dose of gamma radiation. Based on probit analysis, 66.94Gy dose was found to be effective as

LD50 for irradiation of pummelo seeds (Table 2) to induce sufficient variation. This finding may assist as reference dose for large scale gamma irradiation of pummelo genotypes to induce genetic variation. However, considering the practical difficulty, its better fix 60 Gy as LD50 to induce sufficient mutation to select the desirable one. This result was in line with other studies on the effect of gamma rays in citrus. On lemon, Gulsen et al. (2007) obtained most seedless fruit from 50 Gy treatment while Spiegel-Roy et al. (2007) successfully obtained seedless fruit with 13.3 Gy. On mandarin, Kafa et al. (2015) obtained most mutant plants with seedless

fruit with 30 Gy while Montanola et al. (2015) obtained seedless fruit with 40-50 Gy.

Table 1
Effect of gamma irradiation in pummelo seed germination

Treatment	No of seeds sown	No of seeds germinated	No of under developed seedlings	Days to germination	Germination percentage
Control	98	96	0	10	97.96
Dosage 25 Gy	98	93	0	11	94.90
Dosage 50 Gy	98	88	0	11	89.90
Dosage 60 Gy	98	46	21	14	46.94
Dosage 75 Gy	98	24	16	20	24.49
Dosage 100 Gy	98	18	14	20	18.37

Table 2
Lethal dose calculation

Pummelo	na	LD50 LCL-UCL (95% confidence limit)	LD90 LCL-UCL (95% confidence limit)	χ^2	df
Gamma irradiation	98	66.94 (60.06–74.61)	164.27 (127.03–212.42)	9.4	4

LD50 hal dose that kill 50% of the population; na= population number
LCL lower confidence limit
UCL upper confidence limit




Fig.1
Mutagenic sensitivity of pummelo seedlings to different doses of gamma radiation

Acknowledgments

Authors are thankful to the Indian Council of Agricultural Research, Ministry of Agriculture and Farmer's Welfare, GOI, for funding the project.

Referencias

- Dhatt, A S., Rattanpal, H. S., Khokhar, R. and Singh, S. N. 2000. Standardising LD50 dose of gamma rays and EMS in kinnow seeds. *Journal of Research Punjab Agricultural University*, **37** (3-4): 194–7.
- Guadagni, D.G., Maier, V.P and Turnbaugh, J.G, 1973. Effects of some citrus constituents on taste thresholds for limonin and naringin bitterness. *J. Sci. Food Agric.*, **24**: 1199-1205.
- Gulsen, O., Uzun, A., Pala, H., Canihos, E. and Kafa, G. 2007. Development of seedless and mal secco tolerant mutant lemons through bud wood irradiation. *Scientia Horticulturae*, **112**: 184- 190.
- Kafa, G., Seday, U., Uysa, I. O. and Polatoz, S. 2015. Effect of budwood irradiation on seed number of 'Clementine', 'Nova' and 'Robinson' mandarin mutants. Proc. XII Intl. Citrus Congress. Acta Horticulturae No. 1065, International Society for Horticultural Science, Leuven, Belgium.
- Kulkarni, G. B. 2011. Effect of mutagen on pollen fertility and other parameters in horse gram (*Macrotyloma unguiculatum* (Lam.) Verdc). *Bio. Sci. Discovery*, **1**(1): 146-150.
- Mishra, P. J., Ganapathi, T. R., Suprasanna, P., Bapat, V. A. 2007. Effect of Single and recurrent gamma irradiation on *in vitro* shoot cultures of banana. *Int. J. Fruit Sci.*, **7**: 45-57.
- Latado, R. R., Tulmann Neto, A., Ando A Iemma, A., F, Pompeu Junior, J., Figueiredo, J. O., Pio, R. M., Machado, M. A., Namekata, T., Ceravolo, L. and Rossi, A. C. 2001. Sweet orange 'Pera' mutants with low number of seeds obtained through mutation induction. *Revista Brasileira de Fruticultura*, **23** (2): 339–44.
- Maluszynski, M., Nichterlein, K., Vanzanten, L. and Ahloowalia, B. S. 2000. Officially released mutant varieties – the FAO/IAEA database. *Mutation Breeding Review*, **12**: 1–84.
- Mariana, B., Arisah, H., Yenni and Selvawajayanti. M. 2018. Seedless fruit pummelo induced by Gamma Ray irradiation: Fruit morphological characters and stability evaluation. *Biodiversitas*. **19**(2): 706-711.
- Mongkolkul, P., Rodart, P., Pipatthitikon, T., Meksut, L. and Sanguandeekul, R. 2006 Debittering of Tangerine (*Citrus reticulata* Blanco.) Juice by β -Cyclodextrin polymer. *J Incl Phenom Macrocycl Chem.*, **56**: 167– 170.
- Montanola, M. J., Galaz, A., Gambardella, M. and Mártiz, J. 2015. New low seeded mandarin (*Citrus reticulata*) and lemon (*C. limon*) selections obtained by Gamma irradiation. Proc. XII Intl. Citrus Congress. Acta Horticulturae No. 1065, International Society for Horticultural Science, Leuven, Belgium.
- Sanada, T. and Amano, E. 1998. Induced mutation in fruit trees. In Jain, S.M., Brar, D. S. and Ahloowalia, B. S. Soma clonal variation and induced

- mutations in crop improvement. Kluwer Academic Publishers. England. p. 401- 419.
- Singh, B.D. 1994. Mutations in crop improvement In Plant Breeding Kalyani Publ., Ludhiana p 434- 450
- Sinha, S. S and Godward, N. 1972. Radiation studies in *Lens culanaris*. Induction of mutations and type of mutants. *J. Cytol. Genet.*, : 131- 136.
- Sjodin, J. 1962. Some observations in X1 and X2 of *Vicia faba*(L.) after treatment with different mutagens. *Hereditas*, **48**: 565-586.
- Smith, M. K., Hamill, S. D., Langdon, P. W., Giles, J. E., Doogan, V. J. and Pegg, K. G. 2006. Towards the development of a Cavendish banana resistant to race 4 of Fusarium wilt: gamma irradiation of micro propagated Dwarf Parfitt (*Musa* spp., AAA group, Cavendish subgroup). *Aus. J. Exp. Agric.*, **46**(1):107-113.
- Sparrow, A. K., Rogers, A. F. and Schwenmer, S. S. 1968. Radio sensitivity studies which woody plants. I. Acute gamma irradiation survival data for 28 species and predictions for 190 species. *Radiation Botany*, :149-186.
- Spiegel-Roy, P., Vardi, A., Yaniv, Y., Fanberstein, L., Elhanati, A. and Carmi, N. 2007. 'Ayelet' and 'Galya': New seedless lemon cultivars. *Hort science*, **42**(7): 1723-1724.