



INNOTEC

ISSN: 1688-3691

ISSN: 1688-6593

innotec@latu.org.uy

Laboratorio Tecnológico del Uruguay
Uruguay

Ferrari, Graciela

**Water flow and temperature as main factors that regulate
phytoplankton and cyanobacterial blooms in a large subtropical river**

INNOTEC, no. 20, 2020, July-, pp. 30-66

Laboratorio Tecnológico del Uruguay
Uruguay

DOI: <https://doi.org/10.12461/20.07>

Available in: <https://www.redalyc.org/articulo.oa?id=606163454002>

- How to cite
- Complete issue
- More information about this article
- Journal's webpage in redalyc.org

UDEM [redalyc.org](https://www.redalyc.org)

Scientific Information System Redalyc

Network of Scientific Journals from Latin America and the Caribbean, Spain and
Portugal

Project academic non-profit, developed under the open access initiative

Water flow and temperature as main factors that regulate phytoplankton and cyanobacterial blooms in a large subtropical river

El caudal y la temperatura del agua son los principales factores que regulan el fitoplancton y las floraciones de cianobacterias en un gran río subtropical

O caudal e a temperatura da água são os principais fatores que regulam o fitoplâncton e as florações de cianobactérias em um grande rio subtropical

RECIBIDO: 20/4/2020 → APROBADO: 17/6/2020 ✉ gferrari@latu.org.uy



FERRARI, GRACIELA (1)

(1) DEPARTAMENTO DE CALIDAD DE AGUA Y EVALUACIÓN AMBIENTAL DEL LABORATORIO TECNOLÓGICO DEL URUGUAY, LATU. MONTEVIDEO, URUGUAY.

ABSTRACT

Seasonal variations and interannual changes (2006-2019) of the phytoplankton community, related to variations in water quality and river discharge, in the lower part of the Uruguay River in South America are presented. In total, 422 microalgae and cyanobacterial taxa were identified. Cryptophytes, cyanobacteria, and diatoms were the predominant groups. Although nanoflagellates and cryptophytes were always present in plankton, statistical differences were found in seasonal assemblages of phytoplankton. According to the similarity analysis SIMPER, the typical species for each season were identified: *Aulacoseira* spp. in winter, *Dolichospermum uruguayense*, *Durinskia baltica* and *Ankistrodesmus arcuatus* in spring, *Microcystis aeruginosa*, *M. wesenbergii*, *Dolichospermum uruguayense* and *D. circinale* in summer and *Ceratium* cf. *furcoides*, *A. arcuatus*, *Raphidiopsis* and *Jaaginema* sp. in autumn. Under lentic river conditions ($<2000 \text{ m}^3 \text{ s}^{-1}$) and above 22°C , high biomass was recorded due to the blooms of *Microcystis* and *Dolichospermum*. In 2010, *M. panniformis* reached $10.4 \mu\text{g l}^{-1}$ microcystin-LR. *Raphidiopsis mediterranea* and *R. raciborskii* began to be registered in spring (2009 and 2014 respectively) reaching $1041 \text{ cell ml}^{-1}$ in spring 2017. This paper shows that flow, transparency and temperature are the main factors that regulate phytoplankton biomass and cyanobacterial blooms in the Uruguay River. In the summer, when the Uruguay

River is in low flow conditions, cyanobacteria remain in the river. When flow increases, cyanobacteria are transported downstream and could reach the estuary of the Río de la Plata, impacting its beaches.

KEY WORDS: harmful algal blooms, large rivers, Río de la Plata, diversity.

RESUMEN

Se presentan las variaciones estacionales y los cambios interanuales (2006-2019) de la comunidad de fitoplancton, relacionados con las variaciones en la calidad del agua y la descarga del río, en la parte inferior del río Uruguay, en América del Sur. En total, se identificaron 422 taxones de microalgas y cianobacterias; cryptomonas, cianobacterias y diatomeas fueron los grupos predominantes. Aunque los nanoflagelados y las cryptomonas estuvieron siempre presentes en el plancton, se encontraron diferencias estadísticas en los ensambles estacionales del fitoplancton. Según el análisis de similitud SIMPER, fueron identificadas las especies típicas para cada estación. Para el invierno, *Aulacoseira* spp.; para la primavera, *Dolichospermum uruguayense*, *Durinskia baltica* y *Ankistrodesmus arcuatus*; para el verano, *Microcystis aeruginosa*, *M. wesenbergii*, *Dolichospermum uruguayense* y *D. circinale*; y para el otoño, *Ceratium* cf. *furcoides*, *A. arcuatus*, *Raphidiopsis* y *Jaaginema* sp. En condiciones lénticas del río ($<2000 \text{ m}^3 \text{ s}^{-1}$) y por encima de los 22°C se registraron altas biomásas debido a las floraciones de *Microcystis* y *Dolichospermum*. En 2010, *M. panniformis* alcanzó los $10.4 \mu\text{g l}^{-1}$ de microcistina-LR. *Raphidiopsis mediterranea* y *R. raciborskii* comenzaron a registrarse en la primavera de 2009 y de 2014, respectivamente, alcanzando las 1041 cel ml^{-1} en la primavera de 2017. Los principales factores que regulan la biomasa del fitoplancton son el caudal, la transparencia y la temperatura. En verano, cuando el río Uruguay se encuentra en bajas condiciones de caudal, las cianobacterias permanecen en el río. Si el caudal aumenta, son transportadas aguas abajo, pudiendo alcanzar el estuario del Río de la Plata, afectando sus playas.

PALABRAS CLAVE: floraciones de algas nocivas, grandes ríos, Río de la Plata, diversidad.

RESUMO

São apresentadas as variações estacionais e as mudanças interanuais (2006-2019) da comunidade de fitoplâncton, relacionadas com as variações na qualidade da água e a descarga do rio, na parte inferior do Rio Uruguay, na América do Sul. No total, foram identificados 422 táxons de microalgas e cianobactérias. Cryptomonas, cianobactérias e diatoméias foram os grupos predominantes. Embora os nanoflagelados e cymomonas estiveram sempre presentes no plâncton, foram achadas diferenças estatísticas nas assembleias sazonais do fitoplâncton. Segundo a análise de similitude SIMPER, as espécies típicas para cada estação do ano foram identificadas. Para o inverno, *Aulacoseira* spp.; para a primavera, *Dolichospermum uruguayense*, *Durinskia baltica* e *Ankistrodesmus arcuatus*; para o verão, cianobactérias *Microcystis aeruginosa*, *M. wesenbergii*, *Dolichospermum*

uruguayense e *D. Circinale*; e para o outono, *Ceratium* cf. *furcoides*, *A. arcuatus*, *Raphidiopsis* sp. e *Jaaginema* sp. Nas condições lênticas do rio (<2000 m³ s⁻¹) e por cima dos 22 °C, foram registradas altas biomassas devido às florações de *Microcystis* e *Dolichospermum*. Em 2010, *M. panniformis* alcançou os 10,4 µg l⁻¹ de microcistina-LR. *Raphidiopsis mediterranea* e *R. raciborskii* começaram a ser registradas na primavera de 2009 e 2014, respectivamente, chegando às 1041 cel ml⁻¹ na primavera de 2017. Os principais fatores que regulam a biomassa de fitoplâncton são o caudal, a transparência e a temperatura. No verão, quando o Rio Uruguai tem caudal baixo, as cianobactérias permanecem no rio. Se o caudal do rio aumenta, as cianobactérias são transportadas água abaixo, podendo chegar até o estuário do Rio da Prata, afetando as praias.

PALAVRAS CHAVE: florações de algas nocivas, grandes rios, Rio da Prata, diversidade.

INTRODUCTION

In large rivers, significant autotrophic primary production of phytoplankton is expected and is limited to middle river sections, or to lowland areas of higher-order rivers presuming favorable conditions for phytoplankton growth (Reynolds and Descy, 1996). In subtropical rivers, the development of phytoplankton biomass is strongly correlated with the concentration of nutrients, turbidity or light availability (Basu and Pick, 1995; Dodds, 2006). According to Reynolds (2000), hydrological factors such as discharge and dead zones (areas of high resilience) are the most important factors in the development of the rivers' phytoplankton. In the subtropical region, seasonal changes in the phytoplankton community are related to seasonal changes in temperature (Cloern, et al., 2014).

The Uruguay River is one of the largest rivers in South America (>1,800 km); it belongs to the La Plata Basin, forming the triple border of Brazil, Argentina and Uruguay. One of the main factors affecting the river's water quality was the construction of more than ten hydroelectric dams (i.e. Itá, Machadinho, Itaipiranga, Roncador, Garabí, San Pedro, Salto Grande) to meet the growing energy demands which accompanied the region's economic growth, (National Water Agency, ANA, 2005). In Uruguay, the Salto Grande Dam, located 300 km upstream from the mouth of the river, has an important regulatory effect over the discharges above and within the Río de la Plata estuary (EcoMetrix, 2006). Based on historical values (1980-2015), the average annual discharge of the river is 5527 m³ s⁻¹, but in 2017, it reached 7598 m³ s⁻¹ (Salto Grande Mixed Technical Commission, [s.d.]).

Agriculture is an important activity in the Lower Uruguay River Basin with a significant production of citrus fruit and other crops (O'Farrell and Izaguirre, 2014). The most important Uruguayan city in the area studied is Fray Bentos, which is also an active port. This lower section of the Uruguay river gained importance after the installation of a pulp mill in 2006, and the subsequent start of its operations in November 2007 (Uruguayan Technological Laboratory, LATU, 2019). Periodical reports on navigation, cyanobacterial blooms and fisheries surveys are published locally by the Salto Grande Mixed Technical Commission

and the Uruguay River Administrative Commission (CARU, [s.d.]). In addition to this, binational monitoring in the pulp mill's area of influence has been carried out since 2012 and along the binational Argentina-Uruguay section since 2018. CARU surveys the state of the river through water, sediment and biota samples monitoring on both sides of the river and in the center of the channel showing the acceptable limits. Thirteen water quality parameters measured showed concentration values above the limits established in CARU's Standard for 179 variables. Parameters that systematically exceed CARU Digest levels are Iron, Oils and Fats, and Total Phenolic Substances (CARU, [s.d.]).

Although several authors have conducted phytoplankton studies in the Uruguay river basin (de Díaz, 1961; et al, 1986; O'Farrell, 1994; O'Farrell and Izaguirre, 1994; Bonilla, 1997; Pérez, et al., 1999; Pérez, 2002), most of the existing studies on the Uruguay River were focused on the Salto Grande dam (Quirós and Luchini, 1982; De León and Chalar, 2003; Chalar, 2006, 2009; O'Farrell, et al., 2012; Chalar, et al., 2002, 2017;) since 1979, when the lake reservoir was formed. Little research has been done on the topic of phytoplankton composition and ecology in its lower part. O'Farrell and Izaguirre, (1994), showed a gradient along the Argentinean banks and the Uruguay River's tributaries during 1986-1987 where the potamoplankton was dominated by nanoflagellates, cryptophytes and *Aulacoseira* genus. They found that the seasonal succession of phytoplankton diversity was mainly regulated by temperature, transparency and water discharge, with turbidity showing up as a limiting factor for algae development during summer's low water periods, but a correspondence analysis was not performed. Many other reports focused on cyanobacterial blooms. Ferrari and others (2011) reports two possible scenarios which could account for the dynamics of cyanobacterial blooms in the Lower Uruguay River. The first one is that in high flow conditions, the bloom is concentrated in the dam and reaches the open waters of the Río de la Plata estuary as a result of rainfall, or because of the management of the Salto Grande dam. During the summer, when low flow conditions (lentic conditions) are present, the bloom is concentrated locally by a stable high water column. A marked gradient was observed in the meteorological, physical, and chemical parameters of the plankton assemblages from Salto Grande to Punta del Este, in a comprehensive monitoring during 2013-2014, along the Uruguay River's banks and the Río de la Plata coastline (Kruk, et al., 2015). In the sampling from both summers (2013 and 2014), a *Microcystis* bloom was registered in the Salto Grande dam. An exceptional bloom observed in the summer of 2019 reached Uruguay's Atlantic ocean coast. This event occurred under extreme precipitation conditions that induced high water flows, which transported cyanobacterial blooms downstream (Kruk, et al., 2019). Recently, a synoptic study showed the relevance of temperature and precipitation on the distribution of cyanobacteria, comparing lentic and lotic freshwaters in the country (Haakonsson, et al., 2017). The study found that the filamentous *Dolichospermum* was the most widely distributed and the most frequent genus in the region surveyed.

The aim of this study was to identify the main factors that produce seasonal changes in the phytoplankton community of the lower Uruguay River, as well as its relationship with water quality. This work is based on the data compiled over a fourteen-year period (2006 to 2019). We reported on the composition, abundance, carbon biomass of microalgae and cyanobacteria and water quality variables.

MATERIAL AND METHODS

Study area

Uruguay belongs to a subtropical climate zone (Kottek, et al., 2006) characterized by hot and humid summers and generally mild winters with the following features: mild moderate rain (1000-1600mm annual average between 1961-1991), humid subtropical climate (*Cfa* type) (INUMET, [s.d.]) and annual average temperature of 17.5 °C with temperatures above 22 °C in the warmest months. During the study period (2006-2019) the average values of air temperature were 17.9 °C with precipitation values of 1207 mm, (data provided by UPM weather station).

The study area comprises three sites along the lower part of the Uruguay River: an upstream site located in Nuevo Berlín (NB), a central site in Fray Bentos (FB) and a downstream site at Las Cañas (LC). A long-term (2006-2019) seasonal sampling was carried out at the three sites. In each site, three samples were taken covering littoral, center and channel zones of the Uruguayan section of the river (Figure 1). The NB site is located between natural channels and islands. The central site, FB, is located close to the international bridge and the city of Fray Bentos, which is also an active port. In the downstream site at LC, the river becomes wider, forming an extensive plain with sandy beaches that attract a significant number of tourists during summertime.




Figure 1. A) Study area of the Uruguay River and sampling sites from upstream to downstream: Nuevo Berlín (NB), Fray Bentos (FB) and Las Cañas (LC) (black-white circles). B) Sampling design.

Samplings and analysis

Fourteen seasonal sampling surveys (summer: February, autumn: May, winter: August and spring: November) on three sites with a transect with three points (littoral, center and channel) were carried out during July 2006 to November 2019, with a total of 486 sampled points (Figure 1). Conductivity, temperature, oxygen and pH were measured *in situ* with a YSI 6600 V2 sensor and the transparency was estimated with a Secchi disc. Water samples were taken directly for nutrients and chlorophyll analyses, using standard methods as described in detail by Ferrari and others (2011), Tana and others (2013), Uruguayan Technological Laboratory, LATU (2016, 2017, 2018, 2019). Discharge at the Salto Grande dam displays considerable variability related to regional and global processes, with high discharges during El Niño events in phase with El Niño Southern Oscillation (ENSO) due to heavy rainfall (Colombo, et al., 2015). The opening and closing of the floodgates is the main influence on the flow of the river, since in this lower section there is no significant contribution from tributaries that could modify its flow rate. The daily average of the Salto Grande dam discharge was obtained for the period of study from the river's binational monitoring entity (Salto Grande Mixed Technical Commission, [s.d.]). The total nitrogen and total phosphorus were used to calculate N:P atomic ratios. The monthly geometric mean total phosphorus concentration following IET (PT) = $10^{(6 - ((0,42 - 0,36 * (\ln PT)) / \ln 2))} - 20$ was used to determine the Uruguay River's trophic status (Lamparelli, 2004). Subsurface phytoplankton samples were collected with bottles and fixed *in situ* with Lugol's solution for quantification, and net samples were collected and fixed with formaldehyde (4%) for taxonomical studies (Sournia, 1978). When cyanobacterial blooms occurred, cyanobacteria samples were gathered for microcystin-LR analysis were taken using HPLC, according to the ISO 20179 (International Organization for Standardization, 2005). During the sampling period, three extra samplings were carried out in the summer of 2009 and the spring of 2010, as a result of a cyanobacterial bloom warning in the area near the Fray Bentos site.

The sedimentation method based on Utermöhl (1958), was used to identify and quantify phytoplankton. Half or the entire chamber was screened under low magnification (100 or 200x) for organisms >50 µm, while smaller organisms were counted under 400x magnification according to cell density in each sample. Phytoplankton organisms were identified at the lowest possible taxonomic level and divided into groups or Classes. For cyanobacteria, the most recent taxonomical classification presented by Komárek and others (2014) was used. Determinations were made following Komárek and Anagnostidis (2008a, 2008b) and Komárek (2013). To identify some diatoms, the oxidation technique was followed (Battarbee, 1986). Counts included at least 100 cells of the most abundant species, in order to achieve a confidence interval of 95% (Lund, et al., 1958). In the case of *Microcystis* spp. blooms, the methodology followed Box (1981), counting the number of cells per ml in a gridded Sedgewick-Rafter counting chamber. Biovolume was estimated from mean cell dimensions and shape based on Hillebrand and others (1999). The cell biovolume was transformed to biomass as cell carbon (pgC cel⁻¹), according to Menden-Deuer and Lessard (2000). Biovolume was multiplied by the density (cell

ml⁻¹) to estimate phytoplankton carbon biomass (ngC ml⁻¹). Picoplankton cells (<2 µm) were not counted because this requires a special methodology.

Exploratory analyses were performed to identify major patterns and define final tests. In order to evaluate spatial (site and zone) differences among samplings, analyses of similarities ANOSIM were performed with phytoplankton and environmental matrixes. The carbon biomass of the main phytoplankton groups (Cyanobacteria, Chlorophyta, Bacillariophyta, Cryptophyta and “other”) were logarithmically transformed and similarity matrices were constructed with the Bray-Curtis index. ANOSIM results showed significant temporal differences between sampling surveys regarding biomass matrix (RGlobal = 0.544), meanwhile no significant spatial differences were found in either sites (RGlobal = 0.011, $p > 0.001$) or in zones, (RGlobal = 0.016, $p > 0.001$). The same results were found with environmental matrices, (RGlobal temporal = 0.876, $p > 0.001$) with no differences either per sites (RGlobal sites = -0.016, $p > 0.001$) or per zones (RGlobal zones = 0.017, $p = 0.001$). In addition, the analysis of variance ANOVA showed no significant differences either between the three sites (NB, FB and LC) or between zones (littoral, center and channel) for phytoplankton density $F_{(4, 477)} = 0.79$, $p = 0.52$, biomass $F_{(4, 477)} = 0.92$, $p = 0.45$, richness $F_{(4, 477)} = 0.16$, $p = 0.95$ and Shannon diversity index $F_{(4, 477)} = 1.21$, $p = 0.30$. Those analyses enable the use of monthly averages. For those analyses, environmental variables were used: (nitrates (NO₃ mgL⁻¹), total nitrogen (TN mgL⁻¹), nitrate (NO₃ mgL⁻¹), total phosphorus (TP µgL⁻¹), soluble reactive phosphorus (PO₄ µgL⁻¹), chlorophyll *a* (Chl *a* µgL⁻¹), Secchi disk (SD cm), conductivity (mS cm⁻¹K), temperature (T °C), pH and discharge (Qm³s⁻¹), were normalized and Euclidean indexes were used for the similarity matrix. Flow data of the sampling dates were used to perform the analysis.

To evaluate the role of each species and its contribution to the season, the coefficients of similarity between samples and the average similarity / dissimilarity between all the pairs of samples in each group was calculated, applying the similarity percentages breakdown (SIMPER) using the PRIMER routine (Clarke and Warwick, 2001). This analysis identifies the species that are mostly responsible for the observed patterns. Species that contributed the most to the similarity within a season were considered typifying. The species that contributed the most to the dissimilarity within a group, were considered discriminates between groups (Clarke, 1993, Clarke and Gorley, 2006). For the SIMPER analysis, the biomass matrix was used. The small Cryptophytes heterotrophic organisms were not considered for this analysis. A one way ANOSIM was performed to analyze differences between seasons.

For the environmental matrix, Spearman's correlation coefficient was used. Bubble biplots were used to highlight the relationship between flow and temperature in phytoplankton biomass. For XY graph, Cryptophytes were not considered. To determine the set of environmental variables which contribute to explain the variability of phytoplankton biomass, we performed a Canonical Correspondence Analysis (CCA) using the unimodal method. This analysis is valid to apply for any size of the environmental gradient (Legendre and Legendre, 2012). A biomass matrix for species with >25% in the total Log10 (x +1) sample was used. The

environmental matrix was normalized and the significance of the contributions from environmental variables with the ordination of species was tested through Monte Carlo permutations test ($p < 0.05$). The significance of canonical correlation coefficients of each environmental variable for axes 1 and 2 was evaluated by means of an approximation of t-Student statistic (Ter Braak, 1986). A Multivariate Statistical Package (MVSP software) was used.

RESULTS

Environmental variables

Surface water temperature varied seasonally between 11.5 °C in winter and 28.8 °C in summer. Conductivity showed low variation (32-279 $\mu\text{S cm}^{-1}$) throughout the period. pH did not show a wide range of variation (6.1-9.5); the Secchi disc values ranged between 0.20 and 1.60 m and the turbidity reached 78 NTU (Table 1).

Table 1. Seasonal average and maximum-minimum during 2006-2019. Average and standard deviation of nutrients and *in situ* parameters for each season. nd: Not determined.

SEASON	Total Nitrogen (mg l ⁻¹)	Total Phosphorus (µg l ⁻¹)	Nitrate (mg l ⁻¹)	Soluble Phosphorus (µg l ⁻¹)	Chlorophyll (µg l ⁻¹)	Turbidity (NTU)	Secchi Disc (cm)	Temperature (°C)	Conductivity (µS/cm)	pH	Dissolved oxygen (mg l ⁻¹)	River flow (m ³ s ⁻¹)
SUMMER Average	0.9	61.4	0.5	42.0	5.3	19.6	60.9	26.4	67.9	7.4	7.5	4525
stdev	0.2	45.1	0.2	30.9	18.9	7.4	19.2	1.1	11.7	0.4	0.7	2893
AUTOMN Average	1.1	73.3	0.6	34.0	1.4	25.4	61.2	18.4	69.2	7.2	8.5	11698
stdev	0.7	42.1	0.3	23.1	4.4	13.1	32.6	1.5	17.5	0.4	1.4	14073
WINTER Average	1.3	89.2	0.8	24.8	0.6	27.0	45.7	14.5	64.4	7.3	9.5	7296
stdev	0.2	95.5	0.3	19.1	0.8	11.0	13.4	1.5	13.4	0.4	0.9	4461
SPRING Average	1.1	83.9	0.7	23.6	1.6	29.9	48.3	23.3	73.2	7.2	7.9	11365
stdev	0.3	123.8	0.2	18.5	2.8	11.6	12.1	2.8	48.9	0.7	1.6	8325
Annual average	1.1	77.2	0.6	30.8	2.2	25.4	53.8	20.6	68.7	7.3	8.4	8744
minimum	0.1	nd	nd	nd	nd	9.0	20.0	11.6	32.4	6.1	2.3	522
maximum	6.8	1340.0	1.7	163.0	185.0	78.0	160.0	28.8	279.0	9.5	13.0	56070

Nutrient fluctuation at the three sites was similar, with an increase in total nitrogen at NB (in winter), while soluble phosphorus showed three peaks in autumn 2007, 2011 (near 100 $\mu\text{g l}^{-1}$). Fluctuations of total phosphorus remained within historical ranges, with the exception of two peaks, one in August 2011 at the three sites, with the highest concentration at FB reaching values of 887 $\mu\text{g l}^{-1}$, and in November 2019 with 1340 $\mu\text{g l}^{-1}$ at FB (Table 1). The general trophic state of the Uruguay River was mesotrophic (52-59), presenting some eutrophic (60-63) sampling in Aug-11, Feb-17, May-19, Nov-19 due to peak of total phosphorus (Figure 2).




Figure 2. Monthly Redfield ratio N:P (solid line) and monthly trophic status index IET (dotted line) during 2006-2019. Bold solid lines show the limit of 59 for IET. Above, eutrophic state is indicated. Orange squares: annual average and orange line: tendency line.

The atomic Redfield ratio showed no nitrogen limitation condition for phytoplankton growth. Most of the time, the ratio was values above 20, but in Nov-06, Aug-11, Feb-17 and Nov-19, this ratio was below 16 at times when PT concentration was above 100 $\mu\text{g l}^{-1}$ (Figure 2).

Phytoplankton diversity biomass and chlorophyll *a*

During the study period, from July 2006 to November 2019, 422 taxa of algae and cyanobacteria were recorded at three selected sites (Annex 1). The phytoplankton groups were: Cyanobacteria, Bacillariophyta, Chlorophyta *s.l.*, Zygnemaphyta, Cryptophyta, Chrysophyta, Dinophyta, Euglenophyta, Raphidophyta, Xanthophyta, and undetermined flagellates. For the purpose of graphics, an additional group called “Others” composed of Chrysophyta, Dinophyta, Euglenophyta, Raphidophyta, Xanthophyta and undetermined flagellates were grouped together.

The diatoms presented the highest number of taxa (158), followed by the Chlorophyta *s.l.* with 95 taxa, including 55 green coccal algae, 56 Euglenoids (Annex 1). Cyanobacteria were represented by 49 taxa. From a total of 158 diatoms recorded, 134 taxa belonged to pennates and 24 taxa to centric diatoms. During high flow periods, most of the diatom taxa were tychoplankton. Dinoflagellates were represented with 18 taxa and Cryptophyta and Chrysophyta with 9 taxa each. Raphidophyta was represented with the species *Gonyostomum semen* (Annex 1).

The lowest number of taxa per sample (4) were found in winter 2008 (NB), winter 2009 (LC), spring 2008 (FB and LC) and in autumn 2010 (NB and LC). The highest taxa number per sample (69) was found in summer 2016 and 2013: 52 taxa both NB, followed by 43 taxa in LC and 42 in FB. The genera *Nitzschia* and *Trachelomonas* showed the highest number of species (21 and 17). The diatom *Hydrosera whampoensis* and the cyanobacteria *Jaaginema* sp. and *Microcrocis* sp. were recorded for the first time in Uruguay. The most frequent species, found in more than 30% of the samples were: *Plagioselmis lacustris* (92,6%), *Cryptomonas* cf. *marssonii* (86,6%), *Aulacoseira granulata* (71,6%), *Ankistrodesmus arcuatus* (57,8%), *Aulacoseira granulata* var. *angustissima* (54,5%), *Plagioselmis nannoplantica* (53,5%), *Aulacoseira* cf. *muzzanensis*

(45%), *Cryptomonas* cf. *ovata*, *Cryptomonas* cf. *reflexa* (40%), *Trachelomonas rugulosa* and the centric sp. 1 (30%) that constitute the potamoplankton.

Figure 3 shows phytoplankton carbon biomass variations of the main four groups in each sampling site during the study period. The cryptophytes, cyanobacteria and diatoms were the most representative groups and to a lesser extent, the chlorophytes. The maximum total biomass was 1270 ngC ml⁻¹ in NB (Nov-10). *Dolichospermum uruguayense* was the most abundant cyanobacteria with 6903 cell ml⁻¹ in low flow conditions (1275 m³ s⁻¹) and high temperature (24.4 °C). In the same sampling survey in FB and LC, this species, *D. uruguayense*, was recorded with values of 1414 and 1597 cell ml⁻¹, respectively. In the sampling of spring 2010, high values of the three parameters were recorded in FB (8073 cell ml⁻¹, 126.8 ngC ml⁻¹, 9.2 µg l⁻¹) and in LC (5734 cell ml⁻¹, 132 ngC ml⁻¹, 7.0 µg l⁻¹), although the most abundant species was the cyanobacteria *Pseudanabaena mucicola*. Another high peak of phytoplankton abundance occurred in summer 2008 in NB (9497 cell ml⁻¹, 347 ngC ml⁻¹, 6.8 µg l⁻¹) when the flow was 2861 m³ s⁻¹ and the temperature reached the highest value (28 °C). In this opportunity, the dominant Cyanobacteria was *Microcystis aeruginosa* with 6705 cell ml⁻¹ (Figure 3).

Only three phytoplankton biomass peaks of other groups occurred, almost 130 ngC ml⁻¹ (Aug-08 and 09, Feb-13 in NB) with 99% of Cryptophyta and 126 ngC ml⁻¹ (Feb-15, NB) of Chlorophyta (*Eudorina elegans*). In the summer of 2017 and 2019, the biomass of “other groups” increased through the presence of the dinoflagellates and euglenoids species. Big size species, like *Ceratium* and *Trachelomonas*, contributed to the total biomass (Figure 3).

Extra monitoring samples in the summer due to a cyanobacterial bloom were taken at the Fray Bentos site. In 2006, the highest microcystin-LR was recorded with 2010 µg l⁻¹ in a *M. aeruginosa* bloom. In 2009, the values were 6.2×10⁶ cell ml⁻¹. *M. aeruginosa* represented 90% of the total abundance followed by *D. uruguayense*, *M. wesenbergii* and *D. circinale*. The other case was in spring 2010, when a high concentration of microcystin-LR was recorded at the FB site with 10.4 µg l⁻¹. In this bloom, *M. panniformis* reached 1.66×10⁶ cell ml⁻¹ and *D. uruguayense* was observed with values of 2×10⁵ cell ml⁻¹ at the NB site. Recently, in 2012 in FB Remeros marina, *M. aeruginosa* reached 8.5×10⁶ cell ml⁻¹.

Other potentially toxic species began to be registered in 2009: *Raphidiopsis mediterranea* and *R. raciborskii* in spring of 2014, reaching 1041 cell ml⁻¹ in spring 2017. *Cuspidothrix* sp., *Microcrocis* sp. and *Aphanizomenon gracile* appeared in the summer 2015.




Figure 3. Monthly average carbon biomass (ngC ml⁻¹), for each group: Cyanobacteria (CYANO), Chlorophyta (CHLORO), Bacillariophyta (DIATO), Cryptophyta (CRYPTO) and OTHERS for each sampling between July 2006 and November 2019 on sites: Nuevo Berlín (NB), Fray Bentos (FB) and Las Cañas (LC).

Spearman's correlation coefficient showed high correlation between abundance and carbon biomass ($r = 0.52$) while relatively low with secchi disc ($cc = 0.25$), temperature ($cc = 0.26$) and flow (-0.16). Bubble biplots are used to highlight the relationship between flow and temperature in phytoplankton biomass. The diagram shows that the highest biomass was recorded at the highest temperatures and with flow rates of less than $2000 \text{ m}^3 \text{ s}^{-1}$ (Figure 4).




Figure 4. Increasing bubble diagram showing total biomass (ng C ml⁻¹) with regards to flow (m³ s⁻¹) and temperature (°C) during July 2006 - November 2019.

Tables 2 and 3 show the SIMPER analysis carried out with carbon biomass values. *A. granulata* was present in all seasons throughout the sampling period. The typifying species for each season were: *Aulacoseira* spp. in winter, *Dolichospermum uruguayense*, *Durinskia baltica* and *Ankistrodesmus arcuatus* for spring, the cyanobacteria *Microcystis aeruginosa*, *M. wesenbergi* and *Dolichospermum uruguayense* and *D. circinale* in summer, and *Ceratium* cf. *furcoides*, *A. arcuatus*, *Raphidiopsis* and *Jaginema* sp. in autumn. (Table 2). The discriminant species for winter-spring were *Dolichospermum uruguayense* and *Ceratium* cf. *furcoides*, for summer-winter, *M. aeruginosa* and *Ceratium* cf. *furcoides* and for summer-autumn, *D. uruguayense* and *C. furcoides* (Table 3). There were significant differences between seasons (ANOSIM, Global R: 0.085; p = 0.001) (Table 4).

Table 2. Summary of SIMPER results. Species contributions in each season.
Average similarity (Av. Sim.) of typifying species in each season, their contribution (%) to the within-group similarity, and cumulative total (%) of contributions.

Species WINTER Average similarity: 13.13	Av.Sim.	Contribution%	Cumulative total%
<i>Aulacoseira granulata</i>	7.0	53.0	53.0
<i>A. granulata</i> var. <i>angustissima</i>	2.0	15.2	68.2
<i>Aulacoseira</i> cf. <i>muzzanensis</i>	0.7	5.3	73.5
<i>Ceratium furcoides</i>	0.5	4.0	77.5
<i>Ankistrodesmus arcuatus</i>	0.4	3.0	80.5
<i>Trachelomonas rugulosa</i>	0.4	2.8	83.3
Species SPRING Average similarity: 15.19	Av.Sim	Contribution%	Cumulative total%
<i>Aulacoseira granulata</i>	7.8	51.3	51.3
<i>A. granulata</i> var. <i>angustissima</i>	1.5	9.8	61.1
<i>Dolichospermum uruguayense</i>	1.3	8.5	69.7
<i>Durinskia baltica</i>	1.1	7.0	76.7
<i>Ankistrodesmus arcuatus</i>	0.6	3.8	80.4
<i>Aulacoseira italica</i>	0.5	3.6	84.0
<i>Aulacoseira muzzanensis</i>	0.4	2.9	86.9
Species SUMMER Average similarity: 20.45	Av.Sim	Contribution%	Cumulative total%
<i>Aulacoseira granulata</i>	7.3	35.8	35.8
<i>Aulacoseira granulata</i> var. <i>angustissima</i>	5.2	25.4	61.1
<i>Microcystis aeruginosa</i>	1.1	5.2	66.4
<i>Dolichospermum uruguayense</i>	0.8	4.1	70.5
<i>Microcystis wesenbergii</i>	0.8	4.0	74.5
<i>Dolichospermum circinale</i>	0.8	4.0	78.4
Species AUTUMN Average similarity: 10.17	Av.Sim	Contribution%	Cumulative total%
<i>Aulacoseira granulata</i>	4.5	43.8	43.8
<i>Ceratium</i> cf. <i>furcoides</i>	1.3	12.3	56.1
<i>Aulacoseira granulata</i> var. <i>angustissima</i>	1.1	11.1	67.2
<i>Ankistrodesmus arcuatus</i>	0.7	6.6	73.7
<i>Aulacoseira</i> cf. <i>muzzanensis</i>	0.5	4.8	78.5
<i>Trachelomonas rugulosa</i>	0.4	3.7	82.2
<i>Raphidiopsis</i> sp.	0.4	3.5	85.8
<i>Jaaginema</i> sp.	0.3	3.1	88.8

Table 3. Average Dissimilarity percentages (Av. Diss) and species contributions (%) in each season. Discriminant species by group determined by SIMPER analysis with biomass matrix.

Groups WINTER & SPRING	Av. Diss	Contributions%
Average dissimilarity = 87.53		
<i>Aulacoseira granulata</i>	10.1	11.6
<i>A. granulata</i> var. <i>angustissima</i>	5.2	17.5
<i>Dolichospermum uruguayense</i>	4.5	22.6
<i>Ceratium</i> cf. <i>furcoides</i>	3.8	27.0
<i>Peridinium</i> sp. 1	3.7	31.2
Groups WINTER & SUMMER	Av. Diss	Contributions%
Average dissimilarity = 87.05		
<i>Aulacoseira granulata</i>	7.7	8.9
<i>A. granulata</i> var. <i>angustissima</i>	6.9	16.9
<i>Microcystis aeruginosa</i>	3.5	20.9
<i>Ceratium</i> cf. <i>furcoides</i>	3.2	24.5
<i>Dolichospermum uruguayense</i>	3.0	28.0
Groups SPRING & SUMMER	Av. Diss	Contributions%
Average dissimilarity = 85.16		
<i>Aulacoseira granulata</i>	7.1	8.3
<i>A. granulata</i> var. <i>angustissima</i>	6.0	15.3
<i>Dolichospermum uruguayense</i>	5.0	21.2
<i>Microcystis aeruginosa</i>	3.3	25.0
<i>Durinskia baltica</i>	3.1	28.6
Groups WINTER & AUTUMN	Av. Diss	Contributions%
Average dissimilarity = 89.43		
<i>Aulacoseira granulata</i>	12.0	13.4
<i>A. granulata</i> var. <i>angustissima</i>	6.7	20.9
<i>Ceratium</i> cf. <i>furcoides</i>	6.2	27.8
<i>Jaaginema</i> sp.	2.8	30.9
<i>Aulacoseira</i> cf. <i>muzzanensis</i>	2.8	34.0
<i>Trachelomonas rugulosa</i>	2.4	36.7
Groups SPRING & AUTUMN	Av. Diss	Contributions%
Average dissimilarity = 89.73		
<i>Aulacoseira granulata</i>	11.5	12.8
<i>A. granulata</i> var. <i>angustissima</i>	5.0	18.4
<i>Dolichospermum uruguayense</i>	5.0	24.0
<i>Ceratium</i> cf. <i>furcoides</i>	4.8	29.3
<i>Durinskia baltica</i>	4.3	34.2
Groups SUMMER & AUTUMN	Av. Diss	Contributions%
Average dissimilarity = 88.67		
<i>Aulacoseira granulata</i>	8.9	10.1
<i>A. granulata</i> var. <i>angustissima</i>	7.7	18.8
<i>Microcystis aeruginosa</i>	4.2	23.5
<i>Ceratium</i> cf. <i>furcoides</i>	3.7	27.6
<i>Dolichospermum circinale</i>	3.5	31.5
<i>Dolichospermum uruguayense</i>	3.3	35.2
<i>Microcystis wesenbergii</i>	2.9	38.4

Table 4. Analysis of Similarity (ANOSIM) results of biomass matrix for each season.
R statistic (Global R) and significance level of sample statistic: $p < 0.001$

Seasons	R	p
Winter – Spring	0.100	0.001
Winter - Summer	0.129	0.001
Winter – Autumn	0.034	0.001
Spring - Summer	0.099	0.001
Spring - Autumn	0.077	0.001
Summer - Autumn	0.151	0.001

Correlation between environmental variables and carbon biomass

Transparency, flow and temperature were the most important variables explaining phytoplankton structure. According to the CCA analysis, 57.7% of the total variance was explained in the first two axes. The variables that contributed to the first axis were flow ($cc = 0.71$), Secchi disc depth ($SD\ cc = -0.64$) and temperature ($cc = -0.63$) (Figure 5 y Annex 2).

There was a positive correlation between the euglenoids *Lepocinclis caudata*, *Trachelomonas planctonica*, the diatoms *Ulnaria ulna*, the dinoflagellate *Ceratium furcoides* and the cyanobacteria *Jaaginema* and *Raphidiopsis mediterranea* with flow and total phosphorus and nitrogen, while cyanobacteria (*Dolichospermum* cf. *planctonicum*, *D. uruguayense* and *Microcystis* spp.) correlated with temperature, pH, Chl *a* and transparency (SD) (Figure 5).




Figure 5. Biplot diagram Canonical Correspondence Analysis for species with biomass over 25%. Cond: conductivity, T: temperature, pH, TN: total nitrogen, TP: total phosphorus, SD: Secchi disc, Chl *a*: chlorophyll *a*, ODO: oxygen and flow. Code of species are presented.

The river flow fluctuated from 2006 to 2019 a minimum of 522 m³.s⁻¹ (February 2008) to a maximum of 56070 m³.s⁻¹ (May 2017) showing several peaks. When flow rate decreased, transparency increased, favoring the development of phytoplankton, increasing the biomass during the summer period (Figure 6).




Figure 6. Monthly flow average for the three zones and the three sites ($\text{m}^3 \text{s}^{-1}$) (grey area), phytoplankton biomass groups (ngC ml^{-1}) (bars) and temperature (solid black line), from 2006 to 2019. CYANO: cyanobacteria (green bar), CHLORO: Chlorophyta (dark green bar) DIATO: Bacillariophyta (red bar) CRYPTO: Cryptophyta (black bar) and OTHERS: Chrysophyta, Dinophyta, Euglenophyta, Raphidophyta, Xanthophyta: Blue bar)

DISCUSSION

After fourteen years of monitoring in the lower Uruguay River, phytoplankton biomass showed a seasonal pattern in which the flow, transparency and temperature were the structuring factors of the phytoplankton community. Potamoplankton is subjected to unidirectional transport, and it is affected by the physical, hydrological and geographical characteristics of the watershed. In the lower Uruguay River, like in any other lotic ecosystem (Giblin, et al., 2010, Zalocar De Domitrovic, et al., 2007, Sabater, et al., 2008), turbulence, associated with flow, has a direct effect on transparency, attenuating the penetration of light. In general, when flow increases, not only do turbulence and suspended solids increase due to erosion, but so do phosphorus and nitrogen levels. In high discharge conditions, the basin bottom is stirred, and resuspended sediments decrease transparency and increase the light extinction coefficient, resulting in decreasing values of phytoplankton biomass (Kirk, 1994). Paraná River, a similar regional river, showed the same behavior with regards to biogeochemical processes and phytoplankton ecology (Bonetto, et al., 1983, O’Farrell, et al., 1996, Zalocar de Domitrovic, et al., 2007, Colombo, et al., 2015, De Cabo, et al., 2003). High densities of phytoplankton at the lower Paraná River during low flow were also recorded, similar to our results for the Uruguay River and to Ferrari and others (2011), O’Farrell and Izaguirre (2014) and Haakonsson and others (2017), have shown that water temperature and rainfall regime may play a key role in the proliferation of cyanobacteria, with precipitation providing indirect meteorological data of the river’s hydrological condition.

Temperature defines the typical seasonal patterns of rivers in the region (O’Farrell and Izaguirre, 2014); Zalocar de Domitrovic, et al., 2007, Soares, et al., 2007). Seasonal changes in phytoplankton community are related to seasonal

changes in temperature, as it happens in other regions (Salmaso, N. and Braioni, 2008, Desortová, B. and Punčochář, 2011). This pattern of seasonal phytoplankton development is generally observed in temperate European rivers, such as in the middle stretch of the Elbe River in Germany, as well as in several rivers in England (Desortová, B. and Punčochář, 2011).

This is the first study that shows this seasonality with different phytoplankton community assemblages. Wehr and Descy (1998) believed that the most successful algal groups in large rivers were Bacillariophyta and Chlorophyta. Prevalence of diatoms and coccal green algae in phytoplankton composition is common for the large temperate rivers in Europe (Desortová, B. and Punčochář, 2011, Sabater et al., 2009, Reynolds and Descy 1996). Dokulil (2014) states that the phytoplankton species composition has always been dominated by diatoms, particularly centric taxa in the Danube River and others, like the Loire River (Descy, et al., 2012) and the lower Ebro River (Pérez, et al., 2009). Montesanto and others (2000), recorded a total of 122 phytoplankton taxa in the lower branches of the Aliakmon river (Greece), where the diatoms were one of the two more important groups. Garnier and others (1995) also reported similar patterns in the River Seine that contained a mixed group comprised of approximately 200 taxa, of which 90 were diatoms and 60 were Chlorophyceae, similar to findings for regional studies in the Paraguay-Paraná Rivers (Zalocar De Domitrovic, 2005), Negro River (Pérez, 2002), and Salado River (O'Farrell, 1993). However, in the Uruguay River, cryptophytes, cyanobacteria and diatoms were the most representative groups, followed by chlorophytes, dinoflagellates and euglenoids. Descy and others (2012) have proved that the presence of a few dominant species accompanied by a large number of sporadic species is the main feature of phytoplankton community structures in large river ecosystems, and similar results have also been reported in many European rivers. The different phytoplankton studies conducted in the Uruguay River have coincided in showing a relatively high species richness (around 200–400 taxa). The phytoplankton community composition and species richness (422 taxa) turned out to be lower than in previous studies in this region of the river, such as during 1986-1987, when 730 taxa were observed (O'Farrell, 1994). This could be due to the fact that most sampling points were on tributaries, not on the Uruguay River itself. The species richness was similar to other regulated rivers in the region, such as the Negro River, with 218 taxa (Pérez, 2002), the Salado River with 291 taxa (O'Farrell, 1993), the Paraná River (García de Emiliani, 1990) and to other tributaries of the Paraná River (García de Emiliani and de Manavella, 1981). Nanoflagellates, *Cryptomonas* spp. and *Aulacoseira* spp. were the most frequent genus throughout the whole period, since they are organisms well adapted to systems with high suspended material content. This was also described by O'Farrell and Izaguirre, (2014), for the potamoplankton of the Uruguay River. *Cryptomonas* are capable of moving through the water column and respond well to conditions of high nutrient content. They can also convert to mixotrophic nutrition during light-limiting conditions. These organisms are very palatable for zooplankton, and their high density makes the system very rich for the next trophic level (Klaveness, 1988).

Total phosphorus concentration in Uruguay's aquatic systems has been shown to be the most critical variable for water quality. The Uruguay River is not a limited nutrient system—it has appeared as mesotrophic with some peaks in total phosphorus, above the current limit set at 25 µg/l, leaning towards eutrophic. Consequently, nutrients would not be a limiting factor for phytoplankton development. According to previous studies upstream, in the Salto Grande reservoir, the main access route of nutrients into the system is also the flow of the Uruguay River (Berón, 1990; Conde, et al., 1996; Chalar, et al., 2017; EcoMetrix, 2006). The lower stretches of the river are a natural receptor for nutrient discharge.

This study showed the presence of 49 Cyanobacteria taxa. Different species of the genera *Dolichospermum* and *Microcystis* were dominant among the bloom-forming species, in agreement with what was reported at the Argentinean banks (O'Farrell, et al., 2014). A new species of *Dolichospermum* was also described: *Dolichospermum uruguayense*, referring to the Uruguay River (Kozlíková-Zapomělová, et al., 2016). Cyanobacterial blooms commonly occur during summer in the lower Uruguay River and they have been reported since 1974 (Quirós and Luchini, 1982; Di Persia, et al., 1986; Berón, 1990; Conde, et al., 1996; Chalar, et al., 2017). *M. aeruginosa* had already been reported in this study area, reaching high concentrations from September 2006 to November 2008 (O'Farrell and Izaguirre, 2014) and its toxin, microcystin, was detected in the Uruguay River on several occasions (Vidal y Britos, 2012; UNESCO, 2009; Bonilla, et al., 2015; Kruk, et al., 2015). Those blooms, together with the Río Negro dam blooms, could be an important source of the Río de la Plata estuary cyanobacterial blooms, affecting the balneability of the coastal zone and its beaches (Sienra and Ferrari, 2006; Kruk, et al., 2019). This work showed a small upward tendency in the IET index over fourteen years. This implies a tendency towards increasing eutrophication and, consequently, more cyanobacterial blooms could be expected in the River. This work also shows that flow and temperature are the main factors that regulate cyanobacterial blooms in the Uruguay River: under low flow conditions, the cyanobacteria remain in the Uruguay River, whereas when flow increases during the summer months, they reach the Río de la Plata estuary, impacting its beaches. According to Haakonsson and others (2017), *Raphidiopsis* and other Nostocales are spreading out their distribution as competitive genera among bloom-forming cyanobacteria in freshwater ecosystems. In a future scenario, complicated by global warming, with predicted increases in temperature and rainfall, this saxitoxin-producing species could have a much higher biomass.

ANNEXES

Annex 1. Species composition showing the current name and occurrence frequency of phytoplankton for each season during 2006-2019. Numbers reflect the frequency of occurrence in each season through the years.

	WINTER	SPRING	SUMMER	AUTUMN
CYANOBACTERIA				
<i>Anabaena</i> sp.	4	2	2	1
<i>Anathece minutissima</i> Komárek et al. 2011	0	0	1	0
<i>Aphanizomenon gracile</i> Lemm	1	0	0	0
<i>Aphanizomenon</i> sp.	0	1	4	0
<i>Aphanotece</i> sp.	0	1	0	0
<i>Coclosphaerium</i> cf. <i>dubium</i> Grun. in Rabenh.	0	0	0	1
<i>Cuspidothrix</i> sp.	4	4	4	1
<i>Dolichospermum circinale</i> (Rabenh. ex Bornet et Flahault) Wacklin et al.	2	12	44	10
<i>Dolichospermum crassum</i> (Lemm.) Wacklin et al.	0	5	5	3
<i>Dolichospermum planctonicum</i> (Brunnthal) Wacklin et al.	1	0	4	2
<i>Dolichospermum spiroides</i> (Kleb.) Wacklin et al.	0	1	0	0
<i>Dolichospermum uruguayense</i> Kozlíková-Zapomělová et al.	1	35	34	3
<i>Dolichospermum viguieri</i> (Denis & Frémy) Wacklin et al.	7	7	15	1
<i>Geitlerinema</i> sp.	0	2	1	0
<i>Geitlerinema splendidum</i> (Grev. ex Gomont) Anagn.	0	0	3	0
<i>Jaaginema</i> sp.	23	10	18	25
<i>Johanseninema constrictum</i> (Szafer) Hasler et al.	0	0	0	1
<i>Komvophoron crassum</i> (Vozzen) Anagn. & Kom.	0	2	1	5
<i>Komvophoron</i> sp.	2	0	3	6
<i>Leptolyngbya</i> sp.	0	0	3	0
<i>Limnothrix</i> sp.	1	0	0	4
<i>Lyngbya</i> sp.	2	2	0	0
<i>Merismopedia glauca</i> (Ehr.) Kütz.	0	1	4	2
<i>Merismopedia</i> sp.	0	0	1	3
<i>Merismopedia tenuissima</i> Lemm.	0	1	1	0
<i>Microcrocis</i> sp.	0	0	1	0
<i>Microcystis aeruginosa</i> (Kütz.) Kütz.	4	10	41	15
<i>Microcystis flos-aquae</i> (Witt.) Kirchn.	0	0	2	0
<i>Microcystis novacekii</i> (Kom.) Comp.	0	2	7	0
<i>Microcystis panniformis</i> Kom. et al.	0	0	1	0
<i>Microcystis</i> sp.	0	6	9	3
<i>Microcystis wesenbergii</i> (Kom.) Kom.	0	0	39	1
<i>Oscillatoria</i> cf. <i>ornata</i> Kütz. ex Gom.	0	1	0	0
<i>Phormidium</i> sp.	3	0	7	2
<i>Planktolyngbya</i> cf. <i>holsatica</i> (Lemm.) Anagn. & Kom.	0	0	0	1
<i>Planktolyngbya limnetica</i> (Lemm.) Komark. & Cronberg	1	0	4	1
<i>Planktothrix isothrix</i> (Skuja) Kom. & Komark.	1	0	3	3
<i>Planktothrix</i> sp.	0	3	4	0
<i>Planktothrix</i> sp. 2	0	1	0	0
<i>Pseudanabaena catenata</i> Lauter.	12	1	10	8

<i>Pseudanabaena moniliformis</i> Kom. & Kling	0	1	0	0
<i>Pseudanabaena mucicola</i> (Naum. & Hub.-Pest.) Schwabe	2	8	19	4
<i>Pseudanabaena</i> sp.	3	6	12	2
<i>Raphidiopsis raciborskii</i> (Wolos.) Aguil. et al.	0	1	9	2
<i>Raphidiopsis curvata</i> Fritsch & Rich	2	1	5	0
<i>Raphidiopsis mediterranea</i> Skuja 1937	10	21	26	29
<i>Snowella lacustris</i> (Chod.) Kom. & Hind.	0	0	1	0
Undetermined cyanobacteria	7	1	0	2
Undetermined oscillatoriales	13	7	6	2
CHLOROPHYCEAE s.l.				
<i>Actinastrum gracillimum</i> G.M. Smith	0	0	0	1
<i>Actinastrum hantzschii</i> Lagerh.	9	11	41	8
<i>Ankistrodesmus arcuatus</i> Kors.	67	79	74	61
<i>Ankistrodesmus fusiformis</i> Corda	0	1	0	0
<i>Binuclearia lauterbornii</i> (Schm.) Pros.-Lav.	4	1	5	1
<i>Chlamydomonas</i> cf. <i>duplex</i> Skuja	1	1	7	0
<i>Chlamydomonas</i> sp. 1	5	5	8	7
<i>Chlorella</i> sp.	7	3	0	1
<i>Chlorella vulgaris</i> Biej.	2	2	13	7
Undetermined unicellular chlorococcales	1	1	2	0
Undetermined chlorophytes	2	2	0	0
<i>Closteriopsis longissima</i> (Lemm.) Lemm.	9	1	5	12
<i>Coclastrum microporum</i> Näg.	0	3	11	2
<i>Coclastrum pseudomicroporum</i> Kors.	1	0	2	0
<i>Coclastrum</i> sp.	3	1	4	0
<i>Cocnochloris fottii</i> (Hindák) Tsarenko	1	0	9	0
<i>Comasiella arcuata</i> var. <i>platydisca</i> (G.M.Smith) E.Heg. & M.Wolf	0	0	0	1
<i>Crucigenia tetrapedia</i> (Kirchn.) Kunt.	2	2	2	0
<i>Crucigeniella apiculata</i> (Lemm.) Kom.	0	0	5	0
<i>Crucigeniella</i> sp.	1	0	2	0
<i>Desmodesmus arthrodesmiformis</i> (Schr.) An et al.	3	1	6	0
<i>Desmodesmus bicaudatus</i> (Dedus.) Tsarenko	0	0	3	0
<i>Desmodesmus</i> cf. <i>abundans</i> (Kirchn.) Hegew.	0	0	1	0
<i>Desmodesmus</i> cf. <i>maximus</i> (West & West) Heg.	0	1	0	0
<i>Desmodesmus communis</i> (Hegew.) Hegew.	10	15	41	9
<i>Desmodesmus denticulatus</i> (Lagerh.) Hegew.	0	0	3	1
<i>Desmodesmus intermedius</i> (Chod.) Hegew.	1	1	1	0
<i>Desmodesmus opoliensis</i> (Richt.) An et al.	0	2	3	0
<i>Desmodesmus protuberans</i> (Frit. & Rich) Hegew.	0	3	1	0
<i>Desmodesmus serratus</i> (Corda) An et al.	0	0	0	2
<i>Desmodesmus</i> sp.	0	1	0	3
<i>Dictyosphaerium chrenbergianum</i> Nägeli	0	0	3	0
<i>Dictyosphaerium</i> sp.	0	0	4	4
<i>Eudorina</i> cf. <i>cyindrica</i> Thomason	0	0	0	3
<i>Eudorina elegans</i> Ehr.	5	5	21	3
<i>Eudorina</i> sp.	0	3	8	1
<i>Genicularia</i> sp.	2	0	0	0

<i>Golenkinia</i> sp.	0	0	0	1
<i>Gonium pectorale</i> Müller	0	0	2	0
<i>Hariotina reticulata</i> Dang.	0	0	0	4
<i>Hindakia tetrachotoma</i> (Printz) Bock et al.	1	0	17	0
<i>Kirchneriella lunaris</i> (Kichn.) Schmidle	0	1	1	0
<i>Messastrum gracile</i> (Rein.) Gar.	3	7	4	0
<i>Micractinium bornhemiense</i> (Conr.) Korshikov	0	0	3	0
<i>Micractinium pusillum</i> Fres.	1	1	8	1
<i>Monactinus simplex</i> (Meyen) Corda	2	1	12	7
<i>Monoraphidium</i> cf. <i>flexuosum</i> Kom.	1	9	3	7
<i>Monoraphidium</i> cf. <i>irregulare</i> G.M. Smith	6	21	22	4
<i>Monoraphidium</i> cf. <i>tortile</i> (West & West) Komark.	0	0	1	1
<i>Monoraphidium contortum</i> (Thur.) Komark.	16	5	20	2
<i>Monoraphidium griffithii</i> (Berkeley) Komark.	23	23	14	16
<i>Monoraphidium komarkovae</i> Nyg.	13	4	12	0
<i>Monoraphidium minutum</i> (Näg.) Komark.	3	3	1	0
<i>Oocystis elliptica</i> West	0	1	0	0
<i>Oocystis parva</i> West & West	0	0	5	1
<i>Oocystis</i> sp.	0	0	2	0
<i>Pandorina morum</i> (Müller) Bory	12	11	40	8
<i>Pandorina morum</i> var. <i>major</i>	0	0	1	0
<i>Paradoxia multiseta</i> Svirenko	0	0	4	0
<i>Pediastrum duplex</i> Meyen	1	1	11	0
<i>Pseudoschroederia antillarum</i> (Kom.) Hegew. & Schnepf	2	2	9	0
<i>Pyramimonas</i> sp.	0	2	1	0
<i>Raphidocelis danubiana</i> (Hind.) Marv. et al	0	0	0	1
<i>Scenedesmus ecornis</i> (Ehr.) Chod.	2	3	12	0
<i>Scenedesmus obtusus</i> (Chodat) Meyen	0	0	1	0
<i>Scenedesmus</i> sp.	0	6	3	3
<i>Schroederia setigera</i> (Schroeder) Lemm.	1	1	8	0
<i>Selenastrum bibrainum</i> Reinsch.	0	0	3	0
<i>Spermatozopsis</i> cf. <i>exsultans</i> Korsh.	12	8	17	1
<i>Sphaerocystis schroeteri</i> Chod.	1	9	12	4
<i>Stauridium tetras</i> (Ehrenb.) Corda	0	2	2	0
<i>Tetradesmus incrassatulus</i> (Bohlin) Wyn.	2	0	5	0
<i>Tetradesmus lagerheimii</i> Wyn. & Guiry	4	6	7	3
<i>Tetradesmus obliquus</i> (Tur.) Wyn.	2	0	0	0
<i>Tetraedron minutum</i> (Braun) Hans.	1	1	8	3
<i>Tetrachlorella alternans</i> (Smith) Korsh.	1	0	2	0
<i>Tetraselmis cordiformis</i> (Carter) Stein	19	15	44	22
<i>Tetrastrum</i> sp.	5	0	3	2
Ulothrichal undetermined	0	0	0	2
<i>Ulothrix</i> sp.	2	0	0	1
<i>Volvocal</i> sp. 1	1	7	7	6
<i>Volvocal</i> sp. 2	3	1	6	4
<i>Volvox aureus</i> Ehr.	0	0	0	1
<i>Volvox</i> cf. <i>spermatosphaera</i> Powers	0	0	4	5
<i>Volvox</i> sp.	0	0	0	0

<i>Willca apiculata</i> (Lemm.) John et al.	0	0	0	0
<i>Willca crucifera</i> (Wolle) John et al.	1	0	3	0
Chlorococcal sp.1	2	0	0	0
Chlorococcal sp.2	4	4	1	0
Chlorococcal sp.3	8	2	1	1
Undetermined Chlorococcal	2	4	0	1
Undetermined colonia 10µ	1	0	4	0
Undetermined colonia 3µ	1	6	8	0
Undetermined colonia 5µ	0	1	5	0
Undetermined Chlorophyceae	0	1	0	2
ZYGNEMAPHYCEAE				
<i>Closterium aciculare</i> West	8	1	2	1
<i>Closterium acutum</i> Bréb.	21	45	50	18
<i>Closterium ceratium</i> Perty	0	1	0	2
<i>Closterium</i> cf. <i>calosporum</i> Witt	0	1	3	0
<i>Closterium</i> cf. <i>cornu</i> Ehr. ex Ralfs	0	2	1	1
<i>Closterium</i> cf. <i>kuetzingii</i> Bréb.	0	0	0	1
<i>Closterium</i> cf. <i>moniliferum</i> Ehr. ex Ralfs	1	1	2	1
<i>Closterium gracile</i> Bréb. ex Ralfs	0	5	0	0
<i>Closterium</i> sp. 1	2	3	1	0
<i>Closterium</i> sp. 2	2	0	2	0
<i>Closterium</i> sp. 3	3	0	4	0
<i>Closterium</i> sp. 4	0	0	2	0
<i>Closterium</i> sp. 5	0	3	0	3
<i>Cosmarium</i> sp.	0	1	0	1
<i>Cosmarium granatum</i> Bréb. ex Ralfs	0	0	0	0
<i>Cosmarium punctulatum</i> Bréb.	0	0	0	0
<i>Mougeotia</i> sp.	1	0	0	0
<i>Sphaerosma aubertianum</i> f. <i>archeri</i> (Gutw.) Petl.	0	0	0	2
<i>Staurastrum</i> cf. <i>anatinum</i> Cooke & Wills	1	0	0	0
<i>Staurastrum leptocladum</i> Nordstedt	1	1	0	0
<i>Staurastrum mutabile</i> Turner	0	0	1	0
<i>Staurastrum</i> sp. 1	0	0	3	1
BACILLARIOPHYCEAE				
<i>Achnanthes</i> cf. <i>inflata</i> grandis Metzeltin et al.	1	2	1	0
<i>Achnanthes</i> sp.	6	0	1	2
<i>Achnanthidium minutissimum</i> (Kütz.) Czarn.	6	2	6	0
<i>Actinocyclus normanii</i> (Greg.) Hust.	4	1	0	0
<i>Actinocyclus normanii</i> f. <i>subsalsus</i> (Juhlin-Dannfelt) Hustedt	3	0	0	0
<i>Amphipleura pellucida</i> (Kütz.) Kütz.	2	3	2	4
<i>Amphora</i> cf. <i>copulata</i> (Kütz.) Sch.& Arch.	3	0	0	1
<i>Amphora</i> sp.	12	6	3	7
<i>Amphora</i> sp. 2	0	2	0	5
<i>Asterionella formosa</i> Hassall	1	0	0	0
<i>Aulacoseira ambigua</i> (Grun.) Sim.	3	9	6	4
<i>Aulacoseira ambigua</i> f. <i>spiralis</i> (Skuja) Ludwig	13	18	39	7
<i>Aulacoseira</i> cf. <i>distans</i> (Ehr.) Sim.	5	2	37	2
<i>Aulacoseira</i> cf. <i>italica</i> (Ehr.) Sim.	11	26	8	1

<i>Aulacoseira</i> cf. <i>muzzanensis</i> (Meis.) Kram.	57	53	62	47
<i>Aulacoseira</i> <i>granulata</i> (Ehr.) Sim.	84	92	103	69
<i>Aulacoseira</i> <i>granulata</i> var. <i>angustissima</i> (Mull.) Sim.	61	64	90	50
<i>Aulacoseira</i> <i>herzogii</i> (Lemm.) Sim.	3	11	20	1
<i>Aulacoseira</i> <i>islandica</i>	0	0	6	5
<i>Biremis</i> <i>bicontracta</i> (Østrup) Cantonati & Lange-Bertalot	1	0	0	0
Centric sp. 1	60	47	45	18
Centric sp. 2	8	12	28	1
Centric sp. 3	12	0	15	1
Centric sp. 4	1	0	2	0
<i>Centronella</i> <i>reicheltii</i> Voigt	2	0	0	1
<i>Cocconeis</i> <i>placentula</i> Ehr.	7	3	5	2
<i>Cocconeis</i> sp.	1	0	0	0
<i>Craticula</i> sp.	0	3	1	2
<i>Cyclotella</i> <i>meneghiniana</i> Kütz.	8	8	1	11
<i>Cyclotella</i> sp.	1	1	6	0
<i>Cymbella</i> cf. <i>charrua</i> Metz. et al.	0	1	0	0
<i>Cymbella</i> cf. <i>cistula</i> (Ehr.) Kirch.	2	0	1	0
<i>Cymbella</i> cf. <i>cymbelloides</i> (Grun.) Gus.	0	0	0	1
<i>Cymbella</i> cf. <i>dorsenotata</i> Østrup	0	0	0	1
<i>Cymbella</i> sp. 1	4	8	3	8
<i>Cymbella</i> sp. 2	0	0	0	0
<i>Denticula</i> <i>kuetzingii</i> Grun.	2	0	0	4
<i>Diatoma</i> sp.	0	1	0	0
<i>Diatoma</i> <i>vulgare</i> Bory	0	0	0	1
<i>Diploneis</i> sp.	0	0	1	0
<i>Encyonema</i> cf. <i>minutum</i> (Hilse) Mann	0	2	0	0
<i>Encyonema</i> <i>jemtlandicum</i> Krammer	7	3	3	3
<i>Encyonema</i> sp. 1	1	5	2	0
<i>Encyonema</i> <i>vulgare</i> Krammer	0	0	0	1
<i>Encyonopsis</i> <i>microcephala</i> (Grun.) Krammer	2	0	0	0
<i>Entomoncis</i> sp.	0	2	0	0
<i>Epithemia</i> sp.	1	0	1	1
<i>Eunotia</i> <i>bilunaris</i> (Ehr.) Scha.	1	4	0	2
<i>Eunotia</i> cf. <i>naegeli</i> Migula	0	1	0	1
<i>Eunotia</i> sp.	7	4	4	3
<i>Fragilaria</i> cf. <i>capucina</i> Desm.	4	0	0	0
<i>Fragilaria</i> <i>crotonensis</i> Kitton	0	4	3	0
<i>Fragilaria</i> <i>goulardii</i> (Bréb. ex Grun.) Lange-Bertalot	5	0	9	1
<i>Fragilaria</i> sp.	5	3	1	0
<i>Fragilaria</i> sp. 2	2	2	7	0
<i>Fragilaria</i> sp. 3	2	0	0	0
<i>Frustulia</i> sp.	1	2	0	0
<i>Geissleria</i> sp.	5	4	2	3
<i>Gomphonema</i> <i>capitatum</i> Ehr.	1	0	0	2
<i>Gomphonema</i> cf. <i>acuminatum</i> Ehr.	1	2	2	1
<i>Gomphonema</i> cf. <i>affine</i> Kütz.	0	0	0	1
<i>Gomphonema</i> cf. <i>anglicum</i> Ehr.	1	0	0	0

<i>Gomphonema cf. brasiliense</i> Grun.	0	2	0	0	
<i>Gomphonema</i> cf. <i>salae</i> Lange-Bertalot & Reich.	0	4	0	0	
<i>Gomphonema</i> cf. <i>turris</i> Ehr.	2	1	0	1	
<i>Gomphonema parvulum</i> (Kütz.) Kütz.	1	2	2	4	
<i>Gomphonema respectabile</i> Metz. et al.	1	2	0	0	
<i>Gomphonema</i> / <i>Gomphoneis</i> sp. 1	14	6	2	2	
<i>Gomphonema</i> sp. 2	2	0	1	0	
<i>Gomphonema</i> sp. 3	0	1	0	0	
<i>Gomphonema</i> sp. 4	1	0	0	1	
<i>Gomphonema</i> sp. 5	0	1	0	0	
<i>Gyrosigma acuminatum</i> (Küt.) Rabenhorst	0	1	1	0	
<i>Gyrosigma</i> cf. <i>kuetzingii</i> (Grunow) Cleve	5	5	1	2	
<i>Gyrosigma</i> cf. <i>obtusatum</i> (Sulliv. & Worm.) Boyer	0	0	0	0	
<i>Gyrosigma</i> sp. 1	9	6	8	2	
<i>Halamphora montana</i> (Krass.) Lev.	2	6	0	4	
<i>Hantzschia abundans</i> Lange-Bertalot	0	0	0	1	
<i>Hantzschia amphiozys</i> (Ehr.) Grun.	0	4	0	6	
<i>Hantzschia</i> sp.	1	1	1	1	
<i>Hydrosera whampoensis</i> (Schw.) Deby	1	3	0	0	
<i>Iconella guatimalensis</i> (Ehr.) Ruck & Nakov	9	2	10	3	
<i>Lemnicola erigua</i> (Grun.) Kulik. et al.	0	0	1	1	
<i>Melosira varians</i> Agardh	27	15	13	5	
<i>Navicula capitatoradiata</i> Germain ex Gasse	2	1	1	2	
<i>Navicula</i> cf. <i>gregaria</i> Donkin	6	2	0	5	
<i>Navicula</i> cf. <i>neomundana</i> (Lange-Bertalot & Runrich) Lange-Bertalot et al.	1	0	0	1	
<i>Navicula kuseliana</i> Lange-Bertalot & Runrich	24	16	25	4	
<i>Navicula</i> sp. 1	11	6	5	7	
<i>Navicula</i> sp. 2	14	4	4	2	
<i>Navicula</i> sp. 3	14	4	1	14	
<i>Navicula</i> sp. 4	3	1	1	0	
<i>Navicula</i> sp. 5	0	1	6	0	
<i>Navicula</i> sp. 6	0	1	1	0	
<i>Navicula</i> sp. 7	0	2	0	0	
<i>Nitzschia acicularis</i> (Kütz.) Smith	12	3	12	2	17
<i>Nitzschia amphibia</i> Grun.	5	0	1	1	
<i>Nitzschia</i> cf. <i>filiformis</i> (Smith) Van Heurck	3	3	8	1	
<i>Nitzschia</i> cf. <i>incurva</i> var. <i>lorenziana</i> Ross	2	1	1	0	
<i>Nitzschia</i> cf. <i>scapelliformis</i> Grun.	0	1	0	0	
<i>Nitzschia</i> cf. <i>sigma</i> (Kütz.) Smith	2	1	0	0	
<i>Nitzschia</i> cf. <i>sigmoidea</i> (Nitz.) Smith	3	1	1	0	
<i>Nitzschia frustulum</i> (Kütz.) Grun.	0	0	0	5	
<i>Nitzschia fruticosa</i> Hustedt	0	7	11	0	
<i>Nitzschia linearis</i> Smith	1	2	4	1	
<i>Nitzschia palea</i> (Kütz.) Smith	25	13	28	18	
<i>Nitzschia reversa</i> Smith	0	0	3	0	
<i>Nitzschia</i> sp. 1	3	20	12	11	
<i>Nitzschia</i> sp. 2	2	3	0	1	

<i>Nitzschia</i> sp. 3	1	0	0	0
<i>Nitzschia</i> sp. 4	5	2	7	0
<i>Nitzschia</i> sp. 5	0	1	0	0
<i>Pinnularia acrosphaeria</i> Smith	1	0	0	0
<i>Pinnularia</i> cf. <i>minoricapitata</i> Kram.	0	1	0	2
<i>Pinnularia</i> cf. <i>similis</i> Hust.	1	0	0	0
<i>Pinnularia dubitabilis</i> Hust.	1	0	0	0
<i>Pinnularia fistuciformis</i> Metz. et al.	0	0	0	1
<i>Pinnularia microstauron</i> (Ehr.) Cleve	2	2	0	2
<i>Pinnularia</i> sp. 1	5	0	5	5
<i>Pinnularia</i> sp. 2	6	2	0	0
<i>Pinnularia</i> sp. 3	0	2	0	0
<i>Pinnularia</i> sp. 4	0	1	0	0
<i>Pinnularia</i> sp. 5	1	0	0	0
<i>Placoneis</i> cf. <i>elegans</i> Metz. et al.	0	1	0	0
<i>Placoneis</i> cf. <i>gracilis</i> Metz. et al.	1	1	2	0
<i>Placoneis clementis</i> (Grunow) E.J.Cox	5	5	6	4
<i>Placoneis serena</i> (Freng.) Metz.	5	0	3	4
<i>Placoneis</i> sp.	4	0	1	2
<i>Planotidium</i> sp.	1	0	0	0
<i>Pleurosira laevis</i> (Ehr.) Comp.	0	1	1	0
<i>Pseudostaurosira parasitica</i> (Smith) Morales	1	0	0	0
<i>Rhoicosphenia abbreviata</i> (Agardh) Lange-Bertalot	0	1	0	0
<i>Sellaphora parapupula</i> Lange-Bertalot	1	2	0	0
<i>Sellaphora</i> sp.	6	0	2	1
<i>Skeletonema potamos</i> (Weber) Hasle	26	20	27	12
<i>Staurosira</i> / <i>Staurostrella</i> sp.	0	0	1	0
<i>Surirella angusta</i> Kütz.	0	1	0	1
<i>Surirella apiculata</i> Smith	2	3	7	0
<i>Surirella guatemalensis</i>	1	0	4	0
<i>Surirella librile</i> (Ehrenberg) Ehrenberg	1	0	0	0
<i>Surirella ovalis</i> Bréb.	0	0	0	0
<i>Surirella</i> sp. 1	0	0	0	0
<i>Surirella splendida</i> (Ehr.) Kütz.	2	1	2	1
<i>Tabularia</i> sp.	4	1	1	4
<i>Terpsinoe musica</i> Ehr.	1	1	1	0
<i>Thalassiosira</i> sp.	14	0	5	4
<i>Tryblionella</i> cf. <i>hungarica</i> (Grun.) Freng.	0	0	1	0
<i>Tryblionella levidensis</i> Smith	0	0	1	1
<i>Ulnaria acus</i> (Kütz.) Aboal	23	8	3	3
<i>Ulnaria ulna</i> (Nitzsch.) Comp.	25	18	8	15
<i>Pennada</i> sp. 1	9	12	6	2
<i>Pennada</i> sp. 2	2	10	20	1
<i>Pennada</i> sp. 3	2	3	2	1
<i>Pennada</i> sp.4	27	26	16	9
<i>Pennada</i> sp. 5	0	1	1	0
<i>Pennada</i> sp. 6	0	0	0	0
<i>Pennada</i> sp. 7	0	0	3	0

<i>Pennada</i> sp. 8	1	0	3	0
CRYPTOPHYCEAE				
<i>Chroomonas</i> cf. <i>nordstedtii</i> Hansgirg	0	1	0	6
<i>Chroomonas</i> sp.	5	15	16	1
<i>Cryptomonas</i> cf. <i>marssonii</i> Skuja	109	106	111	95
<i>Cryptomonas</i> cf. <i>ovata</i> Ehr.	62	56	56	26
<i>Cryptomonas</i> cf. <i>pyrenoidifera</i> Geitler	5	0	0	0
<i>Cryptomonas</i> cf. <i>refleza</i> Skuja	52	58	51	27
<i>Cryptomonas</i> <i>curvata</i> Ehr.	11	23	9	13
<i>Plagioselmis</i> cf. <i>lacustris</i> (Pasch. & Rutt.) Javorn.	126	105	110	109
<i>Plagioselmis</i> <i>nannoplantica</i> (Skuja) Novarino et al.	83	52	68	57
CRYSOPHYCEAE				
<i>Dinobryon</i> <i>divergens</i> Imhof	1	0	2	0
<i>Goniochloris</i> cf. <i>fallax</i>	0	2	4	0
<i>Kephyrion</i> sp.	2	5	2	3
<i>Mallomonas</i> cf. <i>caudata</i> Iwanoff	0	2	0	0
<i>Mallomonas</i> cf. <i>tonsurata</i> Teiling	5	4	11	3
<i>Mallomonas</i> sp. 1	1	2	0	0
<i>Mallomonas</i> sp. 2	0	1	0	0
<i>Stylococcus</i> sp.	0	0	6	0
<i>Synura</i> cf. <i>uvella</i> Ehr.	13	2	1	0
DINOPHYCEAE				
<i>Ceratium</i> cf. <i>furcoides</i> (Levander) Langhans	20	17	22	19
<i>Ceratium</i> cf. <i>rhomvodes</i> Hickel	2	9	1	0
<i>Ceratium</i> <i>hirundinella</i> (Müller) Dujardin	3	4	1	0
Undetermined athecate dinoflagellate	3	1	7	2
Thecate dinoflagellate 15µ	3	4	4	3
<i>Durinskia</i> <i>dybowskii</i> (Wolos.) Carty (= <i>Durinskia</i> <i>baltica</i> (K.M.Levander) Carty & Cox)	3	16	17	2
<i>Fusiperidinium</i> cf. <i>wisconsinense</i> (Eddy) McCarthy et. al	0	3	2	0
<i>Gonyaulax</i> cf. <i>apiculata</i>	0	0	1	0
<i>Gymnodinium</i> sp.	0	4	1	2
<i>Peridinium</i> cf. <i>willei</i> Huitfeldt-Kaas	0	1	13	11
<i>Peridinium</i> sp. 1	9	38	27	14
<i>Peridinium</i> sp. 2	3	1	12	4
<i>Peridinium</i> sp. 3	1	4	5	3
Peridinales sp. 4	0	0	3	1
Peridinales sp. 5	2	3	4	0
Peridinales sp. 6	1	1	2	1
Gymnodinales	0	6	1	2
Cyst of dinoflagellate	0	0	14	1
EUGLENOPHYCEAE				
<i>Euglena</i> cf. <i>gaumei</i> Allorge & Lefèvre	17	17	36	14
<i>Euglena</i> cf. <i>sanguinea</i> Ehr.	4	9	11	1
<i>Euglena</i> <i>ehrenbergii</i> Klebs	0	1	2	1
<i>Euglena</i> sp. 1	0	6	6	3
<i>Euglena</i> sp. 2	4	6	6	1
<i>Euglena</i> sp. 3	0	0	1	0

Undetermined euglenoids	5	0	0	0
<i>Lepocinclis acus</i> (Müller) Marin & Melk.	4	8	10	6
<i>Lepocinclis caudata</i> (da Cunha) Pascher	6	5	3	1
<i>Lepocinclis</i> cf. <i>fusiformis</i> (Carter) Lemm.	0	0	2	0
<i>Lepocinclis</i> cf. <i>playfairiana</i> (Def.) Def.	1	2	0	1
<i>Lepocinclis</i> cf. <i>salina</i> Fritsch.	0	0	1	0
<i>Lepocinclis ovum</i> (Ehr.) Lemm.	1	2	9	3
<i>Lepocinclis oxyuris</i> (Schm.) Marin & Melk.	3	1	11	0
<i>Lepocinclis</i> sp.1 L=22 a=10	0	2	0	0
<i>Lepocinclis</i> sp.2 L=22,5 a=15	0	2	0	0
<i>Phacus acuminatus</i> var. <i>acuticauda</i> (Roll) Huber-Pest.	0	0	0	1
<i>Phacus</i> cf. <i>orbicularis</i> Hübner	0	0	1	0
<i>Phacus longicauda</i> (Ehr.) Duj.	0	2	1	1
<i>Phacus onyx</i> Poch.	0	3	2	0
<i>Phacus</i> cf. <i>succicus</i>	0	0	1	0
<i>Phacus tortus</i>	0	0	1	0
<i>Phacus</i> sp. 1	5	2	7	1
<i>Strombomonas</i> cf. <i>fluviatilis</i> (Lemm.) Defl.	3	3	15	3
<i>Strombomonas</i> cf. <i>longicauda</i> (Swir.) Defl.	0	0	1	0
<i>Strombomonas</i> cf. <i>ovalis</i> (Playf.) Defl.	1	1	7	1
<i>Strombomonas</i> cf. <i>treubii</i> (Wolosz.) Defl.	0	0	3	0
<i>Strombomonas</i> cf. <i>urceolata</i> (Stokes) Defl.	0	0	1	0
<i>Strombomonas</i> cf. <i>verrucosa</i> (Daday) Defl.	0	0	3	0
<i>Strombomonas deflandrei</i> (Roll) Defl.	0	0	0	0
<i>Strombomonas girardiana</i> (Playf.) Defl.	4	4	12	0
<i>Strombomonas scabra</i> (Playf.) Tell & Conforti	19	20	33	9
<i>Strombomonas triquetra</i> (Playf.) Defl.	0	1	1	0
<i>Strombomonas</i> cf. <i>gibberosa</i>	3	0	1	0
<i>Strombomonas</i> sp.	0	0	1	0
<i>Trachelomonas armata</i> (Ehr.) Stein	0	2	1	0
<i>Trachelomonas bacillifera</i> Playfair	1	2	2	1
<i>Trachelomonas</i> cf. <i>hispida</i> (Perty) Stein.	2	6	7	0
<i>Trachelomonas</i> cf. <i>oblonga</i> Lemm.	12	14	24	7
<i>Trachelomonas</i> cf. <i>raciborskii</i> Wolosz.	3	3	2	0
<i>Trachelomonas</i> cf. <i>superba</i> Svir.	2	0	1	0
<i>Trachelomonas</i> cf. <i>tuberculata</i> Midd.	0	0	1	0
<i>Trachelomonas</i> cf. <i>verrucosa</i> Stokes	8	10	14	2
<i>Trachelomonas hispida</i> var. <i>punctata</i> Lemm.	1	0	0	1
<i>Trachelomonas lacustris</i> Drez.	0	1	0	1
<i>Trachelomonas planctonica</i> Swirenko	21	20	23	6
<i>Trachelomonas rugulosa</i> f. <i>paralella</i> Tell & Domit.	2	1	0	0
<i>Trachelomonas rugulosa</i> Stein	42	30	39	34
<i>Trachelomonas sculpta</i> Balech	4	11	10	5
<i>Trachelomonas similis</i> var. <i>spinosa</i> Hub.-Pestalozzi	0	0	0	0
<i>Trachelomonas</i> sp. 1	1	6	1	3
<i>Trachelomonas</i> sp. 2	0	2	3	0
<i>Trachelomonas</i> sp. 3	2	1	1	0
<i>Trachelomonas</i> sp. 4	0	5	1	1

<i>Trachelomonas volvocina</i> (Ehr.) Ehr.	10	12	21	13
<i>Trachelomonas volvocinopsis</i> Swirenko	16	10	15	1
XANTOPHYCEAE				
<i>Tetraëdriella regularis</i> (Kütz.) Fott	1	0	0	0
<i>Goniochloris</i> cf. <i>fallax</i> Fott	0	0	0	0
<i>Ophiocytium capitatum</i> Wolle	0	1	0	0
RAPHIDOPHYCEAE				
<i>Gonyostomum semen</i> (Ehr.) Dies.	4	9	8	0
OTHERS				
Undetermined flagellates	3	3	6	1
Undetermined organisms	5	0	2	0

Annex 2. Canonical correspondence analysis (CCA), principal axes and abiotic variables contribution made to the main species biomass matrix

Eigenvalues	Axis 1	Axis 2
Eigenvalues	0.32	0.11
Percentage	3.37	1.13
Cumulative, Percentage	3.37	4.50
Cumulative constraint	43.17	57.70
Species-environmental correlations	0.78	0.55
Correlations between env. variables & CCA axes	Envl. Axis 1	Envl. Axis 2
Total Nitrogen	0.33	0.19
Chlorophyll <i>a</i>	-0.31	0.19
Temperature	-0.63	0.51
Conductivity	-0.13	-0.14
pH	-0.61	0.21
Oxygen	0.30	0.15
Flow	0.71	0.55
Transparency (Secchi disc depth)	-0.64	-0.17
Total Phosphorus	0.28	-0.16

Acknowledgments

The authors would like to thank Profs. Jiří Komárek, Nora Maidana and Augusto Comas for their valuable comments and discussions about taxonomic issues. I am grateful to M. Dabezies, A. Kröger, L. Boccardi, G. Useta, Macarena Silva, Joe Reboledo, Marcelo Bado, J. Clemente and technicians of LATU-Fray Bentos and Montevideo for their assistance with samplings. To José and his “Coraje” vessel who are always ready to set sail. To Carlos Saizar, Elina Ordoqui and UPM pulp mill for their support. A special thanks to Maricarmen Pérez, who analyzed part of the phytoplankton samples, for her input and invaluable help with taxonomy, and for her friendship during this long and arduous monitoring study. I am very grateful to the anonymous reviewers for their comments, which helped me improve the original version of the manuscript.

REFERENCES

- Basu, B. and Pick, F.R., 1995. Longitudinal and seasonal development of planktonic chlorophyll a in the Rideau River, Ontario. In: *Canadian Journal of Fisheries and Aquatic Sciences*, 52(4), pp.804–815. DOI:10.1139/f95-080
- Battarbee, R. W., 1986. Diatom analysis. In: Berglund, B.E. And Ralska-Jasiewiczowa M., eds. *Handbook of Holocene palaeoecology and palaeohydrology*. Chichester: Wiley, pp. 527–570.
- Berón, L.E., 1990. Features of the limnological behavior of Salto Grande's reservoir Argentina-Uruguay. In: *Ecological Modelling*, 52, pp.87–102.
- Bonetto, C.A., Zalocar De Domitrovic, Y. and Vallejos, E.R., 1983. Fitoplancton y producción primaria del río Alto Paraná Argentina. In: *Physis*, 41(101), pp.81–93.
- Bonilla, S., Haakonsson, S., Somma, A., Gravier, A., Britos, A., Vidal, L., De León, L., Brena, B., Pérez, M., Piccini, C., Martínez de la Escalera, G., Chalar, G., González-Piana, M., Martigani, F. and Aubriot, L., 2015. Cianobacterias y cianotoxinas en ecosistemas límnicos de Uruguay. En: *Innotec*, 10, pp. 9-22.
- Bonilla, S., 1997. Composición y abundancia fitoplanctónica de tres embalses en cadena sobre el Río Negro, Uruguay. In: *Iheringia, Ser. Bot*, 49, pp.47–61.
- Box, J.D., 1981. Enumeration of cell concentrations in suspensions of colonial freshwater microalgae, with particular reference to *Microcystis aeruginosa*. In: *British Phycological Journal*, 16(2), pp.153–164.
- CARU, [s.d.]. *Archivo de Plan de concientización sobre floraciones algales* [On line]. Paysandú, Colón: CARU. [Accessed: May 21, 2020]. Available at: <http://www.caru.org.uy/web/category/plan-de-concientizacion-sobre-floraciones-algales/>
- Clarke, K. and Gorley, R. 2006. *Primer v6, user manual/tutorial, plymouth routine in multivariate ecological research*. Auckland: Primer-e.
- Clarke, K.R. and Warwick, R.M., 2001. *Change in marine communities: an approach to statistical analysis and interpretation*. 2nd ed. [s.l.]: Primer-E Ltd.
- Clarke, K.R., 1993. Non-parametric multivariate analyses of changes in community structure. In: *Australian Journal of Ecology*, 18(1), pp.117–143. <https://doi.org/10.1111/j.1442-9993.1993.tb00438.x>
- Cloern, J.E., Foster, S.Q. and Kleckner, A.E., 2014. Phytoplankton primary production in the world's estuarine-coastal ecosystems. In: *Biogeosciences*, 11, pp.2477-2501. doi:10.5194/bg-11-2477-2014

- Colombo, J.C., Carlos, N., Skorupka, C., Bilos, L., Tatone, N., Cappelletti, M., Migoya, C., Astoviza, M. and Speranza, E., 2015. Seasonal and inter-annual variability of water quality in the Uruguay River, Argentina. In: *Hydrological Sciences Journal*, 60(6), pp.1155-1163. DOI: 10.1080/02626667.2014.905690
- Conde, D. Pintos, W., Gorga, J., De Leon, R., Chalar, G. and Sommaruga, R., 1996. The main factors inducing chemical spatial heterogeneity in the Salto Grande, a reservoir on the Uruguay River. In: *Large Rivers*, 10 (1-4), pp.571-578.
- Chalar, G., 2009. The use of phytoplankton patterns of diversity for algal bloom management. In: *Limnologia*, 39(3), pp.200-208. <https://doi.org/10.1016/j.limno.2008.04.001>
- Chalar, G., 2006. Dinámica de la eutrofización a diferentes escalas temporales: embalse Salto Grande Argentina-Uruguay. In: Tundisi, J.G., Matsumura-Tundisi, T. and Sidagis, C., eds. *Eutrophication in South America: causes, consequences and technologies for management and control*. São Carlos: International Institute of Ecology, pp.87-101.
- Chalar, G., De Leon, L., Brugnoli, E., Clemente, J., and Paradiso, M., 2002. Antecedentes y nuevos aportes al conocimiento de la estructura y dinámica del Embalse Salto Grande. In: Fernández Cirelli, A. and Chalar, G. eds. *El agua en Iberoamérica: de la limnología a la gestión en Sudamérica*. Buenos Aires: CYTED, pp.123-142.
- Chalar, G., García-Pesenti, P., Silva-Pablo, M., Perdomo, C., Olivero, V. and Arocena, R., 2017. Weighting the impacts to stream water quality in small basins devoted to forage crops, dairy and beef cow production. In: *Limnologia, Ecology and Management of Inland Waters*, 65, pp.76-84.
- De Cabo, L., Arreghini, S., Olgún, H.F., Seoane, R. and Obertello, I., 2003. Physicochemical variables and plankton from the Lower Delta of the Paraná River Argentina. in relation to flow. In: *Hydrological Processes*, 17(7), pp.1279-1290.
- de Díaz, E.N., 1961. Algas de aguas continentales de la Argentina: Fitoplancton de Ibicuy Entre Ríos. In: *Darwiniana*, 122, pp.191-202.
- De León, L. and Chalar, G., 2003. Abundancia y diversidad del fitoplancton en el Embalse de Salto Grande Argentina-Uruguay. Ciclo estacional y distribución espacial. In: *Limnetica*, 22(1), pp.103-113.
- Descy, J.P., Leitão, M., Everbecq, E. and Deliège, J.F., 2012. Phytoplankton of the River Loire, France: a biodiversity and modelling study. In: *Journal of Plankton Research*, 34(2), pp.120-135.

- Desortová, B. and Punčochář, P., 2011. Variability of phytoplankton biomass in a lowland river: response to climate conditions. In: *Limnologica*, 41(3), pp.160-166.
- Di Persia, D.H., Neiff, J.J. and Olazarri, J., 1986. The Uruguay river system. In: Davies, B.R. and Walker K.F., eds. *The ecology of river systems*. Dordrecht: Springer. (Monographiae Biologicae, vol. 60). pp.599-629. https://doi.org/10.1007/978-94-017-3290-1_12
- Dodds, W.K., 2006. Eutrophication and trophic state in rivers and streams. In: *Limnology and Oceanography*, 51(1) part 2, pp.671–680.
- Dokulil, M.T., 2014. Phytoplankton of the River Danube: composition, seasonality and long-term dynamics. In: Liska, I., ed. *The Danube River basin*. Berlin: Springer. (The Handbook of Environmental Chemistry, vol 39).
- EcoMetrix, 2006. *Cumulative impact study-Uruguay pulp mills*. [s.l.]: Internantional Finance Corporation
- Ferrari, G., Pérez, M., Dabezies, M., Míguez, D. and Saizar, C., 2011. Planktic cyanobacteria in the lower Uruguay river, South America. In: *Fottea*, 111 (1), pp. 225–234.
- García de Emiliani, M.O. and de Manavella, M.I.A., 1981. Fitoplancton de los principales cauces y tributarios del valle aluvial del río Paraná: tramo Goya-Diamante. In: *Rev. Asoc. Cienc. Nat. Litoral*, 12, pp.112–125.
- García de Emiliani, M.O., 1990. Phytoplankton ecology of the middle Paraná River. In: *Acta Limnol. Brasil*, 3, pp.391–417.
- Garnier, J., Billen, G. and Coste, M., 1995. Seasonal succession of diatoms and Chlorophyceae in the drainage network of the Seine River: observations and modeling. In: *Limnol. Oceanogr.*, 40(4), pp.750–765.
- Giblin, S., Kraig, H., Jim, F. and Terry, D., 2010. *Evaluation of light penetration on Navigation Pools 8 and 13 of the Upper Mississippi River*: Virginia: U.S. Geological Survey. (Technical Report 2010–T001).
- Haakonsson, S., Rodríguez-Gallego, L., Somma, A. and Bonilla, S., 2017. Temperature and precipitation shape the distribution of harmful cyanobacteria in subtropical lotic and lentic ecosystems. In: *Science of the Total Environment*, 609, pp.1132–1139.
- Hillebrand, H., Dürselen, C., Kirschtel, D. Pollinger, U. and Zohary, T., 1999. Biovolume calculation for pelagic and benthic microalgae. In: *Journal of Phycology*, 35(2), pp.403–424.

- International Organization for Standardization, 2005. ISO 20179: *Water quality — Determination of microcystins — Method using solid phase extraction (SPE) and high performance liquid chromatography (HPLC) with ultraviolet (UV) detection*. Ginebra: ISO.
- INUMET, [s.d]. *Instituto Uruguayo de Meteorología* [On line]. Montevideo: INUMET. [Accessed: January 9, 2019]. Available at: <http://www.meteorologia.com.uy>
- Kirk, J.T.O., 1994. *Light and photosynthesis in aquatic ecosystems*. Cambridge: Cambridge University Press.
- Klaveness, D., 1988. Ecology of the Cryptomonadida: a first review. In: Sandgren, Craig D., ed. *Growth and reproductive strategies of freshwater phytoplankton*. Cambridge: Cambridge University Press. pp.105–133.
- Komárek, J., Kaštovský, J., Mareš, J. and Johansen, J.R., 2014. Taxonomic classification of cyanoprokaryotes cyanobacterial genera 2014, using a polyphasic approach. In: *Preslia*, 864., pp.295–335.
- Komárek, J., 2013. *Süßwasserflora von Mitteleuropa, Bd. 19/3: Cyanoprokaryota 3. Teil / 3rd part: Heterocytous Genera*. Berlín: Springer ISBN: 978-3-8274-0932-4
- Komárek, J. and Anagnostidis, K., 2008a. *Cyanoprokaryota. Teil 1 / Part 1: Chroococcales*. Wiesbaden: Springer Spektrum
- Komárek, J. and Anagnostidis, K., 2008b. *Cyanoprokaryota. Teil 2 / Part 2: Oscillatoriales, Süßwasserflora von Mitteleuropa*. Wiesbaden: Springer Spektrum
- Kottek, M., Grieser, J., Beck, C., Rudolf, B. and Rubel, F., 2006. World map of the Köppen-Geiger climate classification updated. In: *Meteorologische Zeitschrift*, 15(3), pp.259–263. DOI: 10.1127/0941-2948/2006/0130
- Kozlíková-Zapomělová, E., Ferrari, G. and del Carmen Pérez, M. 2016. *Dolichospermum uruguayense* sp. nov., a planktic nostocacean cyanobacterium from the Lower Uruguay River, South America. In: *Fottea*, 16(2), pp.189–200.
- Kruk, C., Martinez, A., Martínez de la Escalera Siri, G., Trinchin, R., Manta G., Segura, A., Piccini, C., Berna, B., Fabiano, G., Pirez, M., Gabito, L., Alcántara, I. and Yannicelli B., 2019. Floración excepcional de cianobacterias tóxicas en la costa de Uruguay, verano 2019. In: *Innotec*, 18, pp.36-68.
- Kruk, C., Piccini, C., Segura, A., Nogueira, L., Carballo, C., Martínez de la Escalera Siri, G., Calliari, D., Ferrari, G., Simoens, M., Cea, J., Alcántara, I., Vico, P. and Míguez, D., 2015. Herramientas para el monitoreo y sistema de alerta de floraciones de cianobacterias nocivas: Río Uruguay y Río de la Plata. In: *Innotec*, 10, pp.23–39.

- Lamparelli, M., 2004. *Graus de trofia em corpos d'água do estado de São Paulo: avaliação dos métodos de monitoramento* [On line]. San Pablo: Universidad de San Pablo. (Tesis de Doctorado). [Accessed: May 27, 2020]. Available at: <https://www.teses.usp.br/teses/disponiveis/41/41134/tde-20032006-075813/publico/TeseLamparelli2004.pdf>
- Legendre, P. y Legendre, L., 2012. *Numerical ecology*. 3ra ed. Amsterdam: Elsevier. 1006p.
- Lund, J.W.G., Kipling, C. and Le Cren, E.D., 1958. The inverted microscope method of estimating algal numbers and the statistical basis of estimations by counting. In: *Hydrobiologia*, 11, pp.143–170.
- Menden-Deuer, S. and Lessard, E.J., 2000. Carbon to volume relationships for dinoflagellates, diatoms, and other protist plankton. In: *Limnology and Oceanography*, 45(3), pp.569–579.
- Montesanto, B., Ziller, S., Danielidis, D. and Economou-Amilli, A., 2000. Phytoplankton community structure in the lower reaches of a Mediterranean river Aliakmon, Greece. In: *Arch. Hydrobiol.*, 147(2), pp.171–191. DOI: 10.1127/archiv-hydrobiol/147/1999/171
- National Water Agency, ANA, 2005. *Aproveitamento do potencial hidráulico para geração de energia* [On line]. Brasília: ANA. (Caderno de Recursos Hídricos). [Accessed: May 21, 2020]. Available at: <http://arquivos.ana.gov.br/planejamento/planos/pnrh/VF%20Gera%C3%A7aoEnergia.pdf>
- O'Farrell, I., 1993. Phytoplankton ecology and limnology of the Salado river Buenos Aires, Argentina. In: *Hydrobiologia*, 271, pp.169–178.
- O'Farrell, I., 1994. Comparative analysis of the phytoplankton of fifteen lowland fluvial systems of the River Plate Basin Argentina. In: *Hydrobiologia*, 289 (1), pp.109–117.
- O'Farrell, I. and Izaguirre, I. 1994. Phytoplankton ecology and limnology of the River Uruguay Lower Basin Argentina. In: *Arch Hydrobiol Suppl*, 99, pp.155–179.
- O'Farrell, I. and Izaguirre, I., 2014. Phytoplankton of the middle and lower stretches of the Uruguay River. In: *Advances in Limnology*, 65, pp.113–126. DOI: 10.1127/1612-166X/2014/0065-0037
- O'Farrell, I., Bordet, F. and Chaparro, G., 2012. Bloom forming cyanobacterial complexes co-occurring in a subtropical large reservoir: validation of dominant eco-strategies. In: *Hydrobiologia*, 698, pp.175–190.

- O'Farrell, I., Izaguirre, I. and Vinocur, A., 1996. Phytoplankton ecology of the lower Paraná River Argentina. In: *Arch. Hydrobiol. Suppl.*, 11(1), pp.75–89.
- Pérez, M. del C., 2002. Fitoplancton del río Negro, Uruguay. In: *Limnetica*, 21(1), pp.81–92.
- Pérez, M. del C., Bonilla, S. and Martínez, G., 1999. Phytoplankton community of a polymictic reservoir, La Plata River basin, Uruguay. In: *Revista Brasileira de Biologia*, 594, pp.535–541.
- Pérez, M. del C., Maidana, N.I. and Comas, A., 2009. Phytoplankton composition of the Ebro River estuary, Spain. In: *Acta Botanica Croatica*, 681, pp.11–27.
- Quirós, R. and Luchini, L., 1982. Características limnológicas del embalse de Salto Grande, III: fitoplancton y su relación con parámetros ambientales. In: *Revista de la Asociación de Ciencias Naturales del Litoral*, 131, pp.49–66.
- Reynolds, C.S., 2000. Hydroecology of river plankton: the role of variability in channel flow. In: *Hydrological Processes*, 14(16-17), pp.3119–3132. [https://doi.org/10.1002/1099-1085\(200011/12\)14:16/17<3119::AID-HYP137>3.0.CO;2-6](https://doi.org/10.1002/1099-1085(200011/12)14:16/17<3119::AID-HYP137>3.0.CO;2-6)
- Reynolds, C.S. and Descy, J.P., 1996. The production, biomass and structure of phytoplankton in large rivers. In: *Large Rivers*, 10, pp.161–187.
- Sabater, S., Artigas, J., Duran, C., Pardos, M., Roman, A. M., Tornes, E. and Ylla, I., 2008. Longitudinal development of chlorophyll and phytoplankton assemblages in a regulated large river (the Ebro River). In: *Science of the total environment*, 404, pp.196-206.
- Sabater, S., Muñoz, I., João Feio, M., Romani, A.M. and Graça, Manuel, 2009. The Iberian rivers. In: Tockner, K., Robinson, C., and Uehlinger, U., eds. *Rivers of Europe*. Amsterdam: Elsevier, pp.113–150.
- Salmaso, N. and Braioni, M.G., 2008. Factors controlling the seasonal development and distribution of the phytoplankton community in the lowland course of a large river in Northern Italy (River Adige). In: *Aquatic Ecology*, 42, pp.533-545.
- Salto Grande Mixed Technical Commission, 2017. *Memorias de gestión* [On line]. *Concordia, Salto: CTMSG*. [Accessed: May 21, 2020]. Available at: https://www.saltogrande.org/memoria_gestion/Memoria%20Gestion%202017.pdf.
- Sienra, D. and Ferrari, G., 2006. Monitoreo de cianobacterias en la costa de Montevideo Uruguay. In: Menafrá, R., Conde, D., Scarabino, F., Menafrá, R. and Rodríguez-Gallego, L., eds. *Bases para la conservación y el manejo de la costa uruguaya*. Montevideo: Vida Silvestre, pp. 413–420.

- Soares, M. C. S., Huszar, V. and Roland, F., 2007. Phytoplankton Dynamics in two tropical rivers with different degrees of human impact (Southeast Brazil). In: *River Research and Applications*, 23, pp.698-714.
- Sournia, A., ed., 1978. *Phytoplankton manual*. Paris: UNESCO. (Monographs on oceanographic methodology 6). doi:10.1002/iroh.19800650312
- Tana, J., Ferrari, G., Dabezies, M., Boccardi, L., Clemente, J., Teixeira de Mello, F. and Gonzalez, I., 2013. *Effects of Pulp Mill effluents in Río Uruguay: a review of monitoring studies in the receiving waters of UPM PULP Mill during 2008-2011*. Informe Inédito. (Informe de Asesoramiento N° 1361295). 60p.
- Ter Braak, C.J.F., 1986. Canonical correspondence analysis: a new eigenvector technique for multivariate direct gradient analysis. In: *Ecology*, 67(5), pp.1167–1179.
- UNESCO y Bonilla, S., ed., 2009. *Cianobacterias Planctónicas del Uruguay. Manual para la identificación y medidas de gestión*. Montevideo: UNESCO. (Documento Técnico PHI-LAC, 16). ISBN 978-92-9089-138-3
- Utermöhl, H., 1958. Zur vervollkommnung der quantitativen phytoplankton-methodik: mit 1 Tabelle und 15 abbildungen im Text und auf 1 Tafel. In: *SIL Communications*, 9(1), pp.1–38. <https://doi.org/10.1080/05384680.1958.11904091>
- Uruguayan Technological Laboratory, LATU, 2019. *Informe sobre caracterización biológica en el tramo inferior del Río Uruguay, febrero - noviembre 2018 y evaluación 2006-2018*. Informe inédito. (Informe de Ensayo N° 1751258). 82 p.
- Uruguayan Technological Laboratory, LATU, 2018. *Informe sobre caracterización biológica en el tramo inferior del Río Uruguay, febrero - noviembre 2017 y evaluación 2006-2017*. Informe inédito. (Informe de Ensayo N° 1669101). 84 p.
- Uruguayan Technological Laboratory, LATU, 2017. *Informe sobre caracterización biológica en el tramo inferior del Río Uruguay, febrero-noviembre 2016*. Informe inédito. (Informe de Ensayo N° 1597867). 84 p.
- Uruguayan Technological Laboratory, LATU, 2016. *Informe sobre caracterización biológica en el tramo inferior del Río Uruguay, febrero-noviembre 2015*. Informe inédito. (Informe de Ensayo N° 1508147). 81 p.
- Vidal, L., y Britos, A., 2012. Uruguay: occurrence, toxicity and regulation of Cyanobacteria. In: Chorus, I., ed. *Current approaches to cyanotoxin risk assessment, risk management and regulations in different countries*. Dessau: Umweltbundesamt. pp.130-136. ISSN 1862-4804

Wehr, J.D. and Descy, J.P., 1998. Use of phytoplankton in large river management.
In: *Journal of Phycology*, 34, pp.741-749. doi:10.1046/j.1529-8817.1998.340741.x

Zalocar De Domitrovic, Y., 2005. Biodiversidad del fitoplancton en el eje fluvial
Paraguay-Paraná. In: *Miscelánea*, 14, p. 230.

Zalocar De Domitrovic, Y., Devercelli, M. and García de Emiliani, M.O., 2007.
Phytoplankton. In: Iriondo, M. H., Paggi, J.C., and Parma, M.J., eds. *The
middle Parana River, limnology of a subtropical wetland*. Berlín: Springer-
Verlag. pp.175-203.