



DEVELOPMENT AND USE OF VITAFERT IN THE AGRICULTURAL SECTOR DESARROLLO Y UTILIZACIÓN DEL VITAFERT EN EL SECTOR AGROPECUARIO

Elaine C. Valiño Cabrera^{1*}, Yaneisy García Hernández¹,
Daymara Bustamante García¹, A. Beruvides Rodríguez²

¹Instituto de Ciencia Animal, C. Central, km 47 ½, San José de las Lajas, Mayabeque, Cuba

²Universidad de Matanzas Camilo Cienfuegos, Vía Blanca, km 3 ½, Matanzas, Cuba

*Email: elainevalino@gmail.com

At Instituto de Ciencia Animal (Cuba), an additive called Vitafert was obtained, with probiotic activity, for animal production. This review mainly deals with the aspects related to the process for the development of this zootechnical additive, which includes microbial inoculants, percentages of raw matters used, the production process, technological variants and its use in the agricultural sector. Information about the progress made with this zootechnical additive is also collected, with positive effects on physiological, immunological, health and production indicators of different species of economic interest. Due to its importance, it is proposed to continue the researchers carried out with Vitafert, with the aim of improving its production and optimization process, as well as the mechanisms of action that generate beneficial responses in animals, based on advanced technologies in health and nutrition.

Key words: animals (Source CAB), fermentation, microbial additive

En el Instituto de Ciencia Animal (Cuba) se obtuvo un aditivo denominado Vitafert, con actividad probiótica, destinado a la producción animal. Esta reseña aborda, fundamentalmente, los aspectos relacionados con el proceso para el desarrollo de este aditivo zootécnico, que incluye inóculos microbianos, porcentajes de las materias primas utilizadas, proceso de obtención, variantes tecnológicas y su utilización en el sector agropecuario. Se recopila, además, información acerca de los avances alcanzados con este aditivo zootécnico, con efectos positivos en indicadores fisiológicos, inmunológicos, sanitarios y productivos de diferentes especies de interés económico. Por su importancia, se propone continuar las investigaciones realizadas con Vitafert, con el propósito de mejorar su proceso de obtención y optimización, así como los mecanismos de acción que generan las respuestas benéficas en los animales, basadas en las tecnologías de avanzada en la salud y la nutrición.

Palabras clave: aditivo microbiano, animales (Fuente CAB), fermentación

Introduction

The International Organization for Epizootics, the World Organization of the United Nations for Feeding and Agriculture and the World Health Organization are working to introduce new products and technologies into animal production systems to obtain healthy foods, which will allow for high productions with adequate economic sustainability and biological guarantee, in order to protect animals and humans. Among these products, in the category of zootechnical additives, microbial additives highlighted,

which represent a potentially significant and safe alternative (Barba-Vidal *et al.* 2019 and Pandey *et al.* 2019).

Microbial additives are products that do not cause problems of microbial resistance or the residual effects produced by antimicrobials when they are used as animal growth promoters, indiscriminately and for a long time. These additives are mainly made from lactic bacteria, *Bacillus*, fungi and yeasts, which contribute to maintaining a favorable ecological balance in the gastrointestinal tract and the proper functioning of the immune system (Quach *et al.* 2021, Konieczka *et al.* 2023 and Rasaei *et al.* 2023).

Received: January 15, 2024

Accepted: April 30, 2024

Conflict of interest: The authors declare that there is no conflict of interest with the results of the research and the publication of this paper.

CRedit authorship contribution statement: Elaine Valiño Cabrera: Investigation, Formal analysis, Writing - original draft. Yaneisy García Hernández: Investigation, Formal analysis, Writing - original draft. Daymara Bustamante García: Investigation, Formal analysis, Writing - original draft. A. Beruvides Rodríguez: Investigation, Formal analysis, Writing - original draft.



This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial (CC BY-NC 4.0). <https://creativecommons.org/licenses/by-nc/4.0/>



Also, they can exert specific activity or multifunctional effects, which through different mechanisms manage to improve the productive and health indicators of animals (Al-Shawi et al. 2020, Lambo et al. 2021, Sun et al. 2021 and Crespo-Piazuelo et al. 2022).

In Cuba, at the end of the 1990s, the Instituto de Ciencia Animal (ICA) developed a microbial preparation called Vitafert, composed of lactic acid bacteria, yeasts, organic acids and other metabolites. A large number of researchers describe its use as a zootechnical additive and show its ability to control the development of enterobacteria, reduce the incidence of diarrhea, increase live weight gain and increase energy and nitrogen retention in different species of productive animals (García et al. 2008, Bustamante et al. 2016, Savón et al. 2020 and Beruvides et al. 2021). In addition, technological variants were used which were differed in the form of presentation, the inoculum and raw materials used in their preparation (Bustamante et al. 2016 and Beruvides et al. 2021). This review mainly deals with the aspects related to the process for the development of this additive, which includes microbial inoculants, percentages of raw matters used, the production process, technological variants and its use in the agricultural sector.

Obtaining and characterization of Vitafert and its technological variants

In the available scientific literature, different formulations are described for obtaining the additive, adapted to the availability of raw matters and existing conditions in each region or locality of Cuba or in other countries where it was applied (López 2010, 2012 and Ruiz 2014). The production process is simple and does not require specialized infrastructure to produce small volumes, which offers economic advantages and facilitates its preparation under rustic and sustainable production conditions in its two forms of presentation, liquid and solid (Bustamante et al. 2016 and Beruvides et al. 2021).

In general, it is obtained through a discontinuous fermentation process, for 48 hours and at room temperature (28 ± 2 °C), of a mixture of easily fermented carbohydrates (sugar cane molasses, sugar cane juice, whey and raw sugar), energy meals (corn meal and wheat meal), a source of vegetable protein (soybean cake meal), a source of non-protein nitrogen, such as urea and mineral salts. This additive involves the addition of an inoculum of fresh excrement, ruminal content or lactic bacteria from yogurt.

Qualitative and quantitative composition

Vitafert liquid additive: It is made up of raw matters (inert ingredients) corn meal (4 %), soybean meal (4 %), urea (0.5 %), magnesium sulfate (0.25 %), mineral premixture (0.5 %). The composition of the premix per kg of product is described as follows: vit. A 10 000 IU, vit. D3 2 000 IU,

vit. E 10 mg, vit. K 2 mg, thiamine 1 mg, riboflavin 5 mg, pyridoxine 2 mg, vit. B12 15.4 µg, nicotinic acid 125 mg, calcium pantothenate 10 mg, folic acid 0.25 mg, biotin 0.02 mg, selenium 0.1 mg, iron 40 mg, copper 12 mg, zinc 120 mg, magnesium 100 mg, iodine 2.5 mg, cobalt 0.75 mg) and final sugar cane molasses (10 %) or raw sugar (15 %). Microbial and lactic acid bacteria (LAB) sources are inoculated at 1 % (active ingredients). They are stable for 90 days or more.

The fermentation process lasts 48 hours at room temperature (26 ± 2 °C). Under production conditions, it can be extended up to 96 hours. The liquid presentation is acidic (pH, 3.5-4.0), reddish brown in color and has a pleasant smell.

Dry Fermented Vitafert Additive: The dry fermented Vitafert additive is made from the mentioned product (it contains the active and inert ingredients). Corn meal (new inert ingredient) is added to liquid VITAFERT in a 1:1 (w/v) ratio. This starchy source used as an absorbent support has the following chemical composition: dry matter (DM) 88.28 % and crude protein (CP) 8.4 %. Drying is carried out by direct exposure to the sun for 72 h. The average maximum temperature and relative humidity values are 27.3 °C and 76.0 %, respectively. Dry Vitafert has a particle size of 1 mm. The presentation of the solid Vitafert additive is golden meal, sweet smell, according to the formulation used in the production of the liquid VITAFER additive, as part of the mixture with corn meal.

The safe and harmless effect of this product was proven in monogastric animals, ruminants and small ruminants.

Use of Vitafert in the agricultural sector

The first publication about Vitafert was made by Díaz et al. (1996) based on scientific reports from Instituto de Ciencia Animal, which stated that it is a solid product, made from aerobic fermentation in an open-air dish of a mixture of ground sugar cane and fresh chicken manure, with the addition of urea and a premixture of minerals in a process similar to the obtaining of Saccharina (Elías et al. 1990). In addition, at the laboratory level, it was reported that the results with this product showed the presence of a high content of antibodies, vitamins and other nutrients of importance for the pre-fattening category of pigs. The protein content of the product makes it comparable in composition to corn meal, although its level of crude fiber is much higher, as well as that of ash and non-protein nitrogen. The work highlighted that during the fermentation process colibacilli disappear and lactobacilli, streptococci and yeasts predominate. This product, as a substitute for cereals in the diet of pre-fattening pigs, starting at 10 kg of live weight, had no effect on daily gain, final weight and total conversion. However, its inclusion at 10 and 20 % improved the conversion of traditional products with a favorable economic impact.

González and Enríquez (1997) reported the effect of a new food, called liquid Vitafert (Elías 1995, unpublished) on the nutritional value of a ration for ruminants, based on star grass hay (*Cynodon nlemfuensis*). This new food was obtained by liquid fermentation from the final sugar cane molasses. Valdivié *et al.* (1997) compiled researchers on new variants of Saccharinas, from combinations of sugar cane with other substrates. Among these variants, “Sacchasoyamaíz” inoculated was presented, a product made from a primary substrate of ground corn (30 %), ground soybean cake (5 %), ground cane (63 %), urea (1.5 %) and mineral salts (0.5 %), to which after mixing 10 % of Vitafert MN was added (primary name due to the addition of a mixture of minerals) as an inoculants, components that were dried and fermented, according to the rustic technology of Saccharina. Also, Elías *et al.* (2001 and 2009), Valdivié and Elías (2006) and Díaz (2010) referred to Vitafert MN as an inoculum to generate other variants of Saccharina.

After the year 2000, a large number of researchers were carried out with Vitafert and its variants, all with the principle or line to produce and apply efficient biologically activated microorganisms (MEBA) (Elías and Herrera 2015).

García *et al.* (2008) carried out the first studies on the dynamics of fermentation of excreta from broilers and laying hens, the main microbial inoculants for obtaining Vitafert, the first variant used. These researchers mainly focused on the proposal to improve the quality of the additive and the benefit of the products derived from mixed fermentation, such as lactic acid and other organic acids. These fermentation processes potentially eliminated pathogenic microorganisms. Regardless of the obtained

results, poultry manure was no longer used in Cuba as inoculum for reasons of biological safety.

In their research, Díaz *et al.* (2014) isolated and characterized strains of lactic acid bacteria from native microbial preparations, made from agro-industrial wastes (whey, bovine manure and ruminal content), designed as biological inoculants for post-harvest organic waste silage processes. These authors proposed that other crop wastes could be studied with these inoculants.

Table 1 summarizes the most significant elements of different studies on Vitafert, in which chicken manure, bovine excrement, whey, rumen content were used as inoculum and final molasses as an energy source. Their effects on biological activity are also described.

The introduction of natural yogurt to obtain Vitafert gave the zootechnical additive a high concentration of metabolites during fermentation, due to its content of lactic acid bacteria. This formulation was obtained as a safer alternative to previous biological inoculants. Table 2 shows the different researchers carried out with Vitafert, made with yogurt as inoculum, in ruminant and monogastric animals, obtained not only under laboratory conditions but also under production conditions.

The production of Vitafert using sugar cane molasses as an energy source in the fermentation process showed difficulties due to its limited availability and the quantities needed to be used under production conditions. Hence, its replacement in the formulation with raw sugar not suitable for human consumption was evaluated. The new variant was produced under production conditions and its effect as an additive was proven to make it accessible to medium and small pig producers in Cuba, mainly (table 3).

Table 1. Researchers carried out with Vitafert made with final sugar cane molasses and different microbial inoculants

Microbial inoculum	Purpose of the research	Results	References
Fresh poultry manure	Evaluation of the inoculum applied by spraying on coffee husk beds and poultry manure. Supplementation of grazing Pelibuey sheep with beds inoculated with Vitafert.	- Reduction of enteropathogenic organisms in beds and poultry excrement. - Increase of the productive indicators in the cycle of laying hens and their replacements. - Improvement of the productive indicators of growing and fattening sheep, by applying 20 g of inoculated bed/kg of live weight of the animal supplemented with final sugar cane molasses to guarantee energy requirements.	Calderón <i>et al.</i> (2005)
Bovine manure, fresh whey or ruminal content	Obtaining microbial consortia with lactic acid activity for silages.	- Fresh whey was the best inoculum for microbial preparations. - A low-cost, high biological value silage product was obtained, with the capacity to stimulate production and improve the quality of milk in cows.	Díaz <i>et al.</i> (2014)
Ruminal content	Supplementation of broilers with a dry Vitafert.	It improved the health and histological indicators of the animals that intake it at 3 %.	Savón <i>et al.</i> (2020)

Table 2. Results of researchers carried out with Vitafert, made with yogurt as inoculum in ruminant and monogastric animals

Purpose of the research	Results	References
- Evaluation of Vitafert on intake, feeding behaviour, ruminal fermentation and <i>in situ</i> degradation of DM and NDF in goats fed with <i>Brachiaria brizantha</i> hay. - Evaluation of the best dose in a stable system under production conditions, its effect on DM intake, feed efficiency, production and chemical composition of milk.	- With the addition of 6 mL/kg of live weight (best dose), an increase was achieved in the voluntary intake of DM and total NDF and in its positive relation with live and metabolic weight under controlled and production conditions. - Increase of ingestion and rumination. - Increased fermentation capacity of the goat rumen. - Reduction of the colonization time of the neutral detergent fiber and increase of the degradation rate and effective degradability of the DM and NDF - Productive and economic advantages in relation to gross profitability per goat and increase of milk quality.	Gutiérrez et al. (2012)
Inclusion of the microbial additive Vitafert in the <i>in vitro</i> ruminal fermentation of a diet for goats.	- The inclusion of Vitafert did not affect gas production or the kinetic parameters of diet fermentation. - The degradability of DM and neutral detergent fiber was increased with 4.5 mL of the additive/kg of animal live weight. - Microbial biomass production and synthesis efficiency was improved with 8.5 mL/kg of live weight.	Rodríguez et al. (2013)
Evaluation of the MUSS-LACTIBIOL supplement (Vitafert variant) in beef production.	- Average daily gain increased (higher than 1000 g in bull fattening). - Vitafert variant dose of 1.37 kg/d per animal.	Castellón et al. (2014)
Evaluation of the inclusion of Vitafert as a microbial additive in the behavior and health of growing pigs	- Live weight, average daily gain, feed conversion and viability of animals supplemented with Vitafert (10 mL/kg of live weight) were increased. - Morbidity, mortality and lethality due to diarrheal episodes were reduced - Lower risk of death in pigs that intake it, which represents lower economic loss due to death.	Lazo-Pérez et al. (2017)
Effect of beneficial microorganisms activated in the finishing of bulls in silvopastoral system with leucaena, supplemented with sugar cane.	The use of the biological product Vitafert® (vtf) generated a contrasting productive effect on average daily gain, weight increase and yield per hectare. The intake of sugar cane was also encouraged.	Iraola et al. (2017)
Biological response in broiler that intake 2 % of Vitafert + corn meal (1:1), sun-dried and a diet with <i>Moringa oleifera</i> foliage meal, Supergenius variety (10 %)	The apparent ileal digestibility of amino acids was increased and the productive performance and meat yield of chickens were improved.	Bustamante et al. (2021)

In the previous researchers has shown that Vitafert positively influences on the physiology and productive performance of animals. Effects that were attributed, mainly, to the presence of beneficial microorganisms with probiotic activity, proven in birds, pigs, calves and to its content of organic acids with antimicrobial action and modulation of the immune system. Likewise, other studies showed that Vitafert stimulated the fermentation process and increased the nutritional value of grasses, foliages, forages, fruits, legumes and protein plants, crop residues or mixtures of these sources, as a starter culture (table 4).

Final considerations

In most of the researchers, where Vitafert was applied in its different formulations and technological variants, improvements were obtained in the indicators evaluated in the animals that intake it or in the quality of the plant materials where it was applied. The microorganisms present in Vitafert and its metabolites are considered to have been

able to promote the absorption of nutrients, strengthen the barrier function of the intestinal mucosa, improve the balance of the intestinal microbiota, promote the growth of beneficial microorganisms, inhibit the growth of pathogens and modulate immunity. However, the effectiveness of the zootechnical additive depended on its chemical and microbiological composition, the dose, mode and frequency of application, as well as the species or category of animal that intake it.

Due to its importance, it is proposed to continue the researchers carried out with the Vitafert additive to improve the process of obtaining and optimizing it, in addition to its mechanism of action that generates beneficial responses in animals, based on advanced technologies in health and nutrition.

Acknowledgments

Special thanks to the laboratory technician Félix Herrera for his collaboration in the data collection.

Table 3. Researchers carried out with Vitafert, made with raw sugar and yogurt, in breeding sows, pre-fattening and piglets

Purpose of the researcher	Results	References
Evaluation of different levels of Vitafert in growing and fattening pigs.	With the use of Vitafert in the feeding of growing-fattening pigs, with inclusion levels of 5, 10, 15 mL/kg of live weight, improvements in productive performance and health were obtained.	Vitaluña (2014)
Evaluation of the zootechnical additive Vitafert in the productive performance and health of pre-fattening piglets.	The weaning-estrus interval was shortened and the weight loss of the breeder during lactation was reduced	Beruvídes <i>et al.</i> (2018)
Effect of the zootechnical additive Vitafert on the biological response of piglets.	Improvements in the digestive physiology of breeding and pre-fattening pig categories were achieved under production conditions	Beruvídes <i>et al.</i> (2020)
Evaluation of productive and health response in pigs using the Vitafert additive. Chemical and microbiological characterization.	A positive effect on the performance of microbiological, hematological, immunological, and productive and health indicators was obtained, under intensive production conditions.	Beruvídes <i>et al.</i> (2021)
Productive and health performance in suckling piglets, supplemented with sugar, fermented with yogurt	Growth rate, feed efficiency and reduction of intestinal infections were favored	Beruvídes <i>et al.</i> (2023)

Table 4. Researchers carried out with Vitafert as an inoculum in fermentation processes of agricultural products and wastes

Purpose of the research	Results	References
Obtaining a food, fermented in a solid state from the returned bagasse, polished rice and inoculants.	- The inclusion of rice polishing + Vitafert at 5 % increased the TP to 11.1 %. - The NDF decreased to 57.4 % and pH values ranged from 3.85 to 4.67. - Lactic acid production was high and VFA performance ranged from 2.8 to 6 % and 9.6 to 33 mg 100g of DM⁻¹. - The results suggest that returned bagasse can be used as a nutritious food for cattle.	Cárdenas <i>et al.</i> (2008)
Inclusion of Vitafert as an inoculum in the solid fermentation of breadfruit meal for diets of pre-fattening pigs.	The chemical composition of the fermented food was improved with the inclusion of 5 % Vitafert.	Brea <i>et al.</i> (2014)
Evaluation of the chemical composition and ruminal degradability of DM from silage mixtures of <i>Tithonia diversifolia</i> : <i>Pennisetum purpureum</i> cv. Cuba CT-169, inoculated with Vitafert.	- Crude protein and ash content were increased and NDF was reduced in all inoculated silages. - The best results were obtained with the inclusion of 4.5 and 6.0 % of Vitafert. - The 20:80 combination of <i>Tithonia</i>:<i>Pennisetum</i> with 4.5 % of the additive achieved the highest degradability in the shortest time, highest degradation rate and effective degradability.	Gutiérrez <i>et al.</i> (2014)
Inclusion of Vitafert in the nutritional value of <i>Tithonia diversifolia</i> and <i>Pennisetum purpureum</i> silages.	- The chemical composition of silages was improved with 4.5 and 6.0 % of Vitafert. - The best indicators of <i>in vitro</i> fermentation of mixed silages were obtained with 60 % <i>T. diversifolia</i> and 8.0 % Vitafert.	Morales <i>et al.</i> (2016)
Obtaining an inoculant that enhances solid fermentation from postharvest wastes of <i>Solanum tuberosum</i>	The microbial preparation with lactic acid bacteria favored the fermentation process of potato harvest wastes.	Borrás <i>et al.</i> (2017, 2020 a, b; 2021)
Evaluation of the nutritional value of mixed silages of <i>Cenchrus purpureus</i> cv. Cuba CT-169 and <i>Moringa oleifera</i> with the inclusion of 1 % Vitafert and <i>Ipomoea batata</i> (0, 5, 10 and 15 %)	Both additives improved the fermentation characteristics of silages by reducing butyric acid levels and increasing propionic acid levels.	Rodríguez <i>et al.</i> (2019)

References

- Al-Shawi, S.G., Dang, D.S., Yousif, A.Y., Al-Younis, Z.K., Najm, T.A. & Matarnah, S.K. 2020. The potential use of probiotics to improve animal health, efficiency, and meat quality: a review. *Agriculture*, 10(10): 452-463, ISSN: 2077-0472. <https://doi.org/10.3390/agriculture10100452>.
- Barba-Vidal, E., Martín-Orúe, S.M. & Castillejos, L. 2019. Practical aspects of the use of probiotics in pig production: a review. *Livestock Science*, 223: 84-96, ISSN: 1878-0490. <https://doi.org/10.1016/j.livsci.2019.02.017>.
- Beruides, A., Elías, A., Valiño, E.C., Milián, G., Lezcano, Y., Moliner, J.L., Rodríguez, M. & Zamora, H. 2018. Evaluation of the zootechnical additive Vitafert in the productive performance and health of pre-fattening piglets. *Cuban Journal of Agricultural Science*, 52(1): 49-56, ISSN: 2079-3480. <https://www.cjascience.com/index.php/CJAS/article/view/780>.
- Beruides, A., Elías, A., Valiño, E.C., Milián, G., Rodríguez, M. & González, R. 2023. Comportamiento productivo y de salud en lechones lactantes suplementadas con azúcar fermentado con yogurt. *Livestock Research for Rural Development*, 30, Article #72, ISSN: 0121-3784. <http://www.lrrd.org/lrrd30/4/agust30072.html>.
- Beruides, A., Elías, A., Valiño, E.C., Milián, G., Rondón, A.J., Rodríguez, M. & Milián, J. 2021. Chemical and microbiological characterization of the zootechnical additive Vitafert, under small-scale production conditions. *Cuban Journal of Agricultural Science*, 55(1): ISSN: 2079-3480. <https://cjascience.com/index.php/CJAS/article/view/998>.
- Beruides, A., Elías, A., Valiño, E.C., Milián, G., Rondón, A., Rodríguez, M., Lezcano, Ya., Brea, O., Milián, J., Laurencio, M., Valdivia, A., Sánchez, L., Sarduy, L., Herrera, M., Cobo, R., Caro, Y., Fuentes, L., Matos, M., Bustamante, D., Vera, R., Betancourt, C. & Mendoza, M. 2020. Efecto del aditivo zootécnico Vitafert en la respuesta biológica de crías porcinas. Libro: Editorial académica española. Is trademark of international Book Market Service Ltd., KS OmniScriptum Publishing group, p. 1-100, ISBN: 978-620-0-41183-9.
- Borrás, L.M., Valiño, E.C. & Rodríguez, C. 2017. Preparado microbiano con actividad ácido láctica como acelerante biológico en los procesos de fermentación para alimento animal. *Ciencia y agricultura*, 14(1): 7-13, ISSN: 0122-8420. <https://www.redalyc.org/journal/5600/560062845001/560062845001>.
- Borrás, L.M., Valiño, E.C., Elías, A., Martínez, J.J., Sanabria, A.M. & Becerra, M.L. 2020a. Solid-state fermentation of post-harvest wastes of *Solanum tuberosum* and a microbial preparation. *Cuban Journal of Agricultural Science*, 54(4): 525-533, ISSN: 2079-3480. <https://cjascience.com/index.php/CJAS/article/view/985>.
- Borrás, L.M., Valiño, E.C., Elías, A., Martínez, J.J., Sanabria, A.M. & Becerra, M.L. 2020b. Efecto del carbonato de calcio CaCO₃ en la fermentación en estado sólido de residuos de poscosecha de *Solanum tuberosum* con un preparado microbiano. *Cuban Journal of Agricultural Science*, 54(4): 535-545, ISSN: 2079-3480. http://scielo.sld.cu/scielo.php?script=sci_arttext&pid=S2079-34802020000400535&nrm=iso.
- Borrás, L.M., Valiño, E.C., Elías, A., Martínez, J.J., Sanabria, A.M. & Becerra, M.L. 2021. Efecto de la inclusión de materiales fibrosos en la fermentación en estado sólido de residuos poscosecha de *S. tuberosum* inoculado con un preparado microbiano. *Cuban Journal of Agricultural Science*, 55(1): 31-42, ISSN: 2079-3480. http://scielo.sld.cu/scielo.php?pid=S2079-34802021000100004&script=sci_arttext&tlng=en.
- Brea, O., Ortiz, A., Elías, A., Herrera, F. & Motta, W. 2014. Solid-state fermentation of the fruit meal from the breadtree (*Artocarpus altilis*) in diets for pre-fattening pigs. *Cuban Journal of Agricultural Science*, 48(4): 391-398, ISSN: 2079-3480. <https://cjascience.com/index.php/CJAS/article/view/569>.
- Bustamante, D., Savón, L., Elías, A., Caro, Y., Sierra, F. & Almeida, M. 2016. Effect of Vitafert on the digestive use of nitrogen in roosters that consume *Moringa oleifera* forage meal. Technical note. *Cuban Journal of Agricultural Science*, 50(2): 273-278, ISSN: 2079-3480. <https://www.redalyc.org/pdf/6537/653768179003>.
- Bustamante, D.G., Savón, L., Elías, A. I., Caro, Y.R., Valiño, E.C., Rojas, M.V., Martín, C.N. & Mireles, S. 2021. Chemical and microbiological characterization of a technological variant of Vitafert intended for animal production. Technical note. *Cuban Journal of Agricultural Science*, 55(2): 61-73, ISSN: 2079-3480. <http://cjascience.com/index.php/CJAS/article/view/940>.
- Calderón JO., Elías, A. & Valdivié, N.M. 2005. Dynamics of the fermentation in solid state of the husk beds of coffee in beginning of ponedoras inoculated with Vitafert. *Revista Electrónica de Veterinaria REDVET®*, VI (05), ISSN: 1695-7504. <http://www.veterinaria.org/revistas/redvet/n050505.html>.
- Cárdenas, J.R., Aranda, E.M., Hernández, D., Lagunes, L.C., Ramos, J.A. & Salgado, S. 2008. Obtención de un alimento fermentado en estado sólido a partir del bagacillo de retorno, pulido de arroz e inóculos. Su utilización en la alimentación animal. *Cuban Journal of Agricultural Science*, 42(2): 173-176, ISSN: 0034-7485. <https://www.redalyc.org/articulo.oa?id=193015494009>.
- Castellón, M., Elías, A. & Jordán, H. 2014. Evaluation of peanut hay (*Arachis hypogaea*) with grass silage and a protein-energy supplement for fattening bulls with different starting live weights. *Cuban Journal of Agricultural Science*, 48(3): 235-239, ISSN: 2079-3480. <https://www.cjascience.com/index.php/CJAS/article/view/577>.

- Crespo-Piazuelo D, Gardiner, G.E., Ranjitkar, S., Bouwhuis, M.A., Ham, R., Phelan, J.P., Marsh, A. & Lawlor, P.G. 2022. Maternal supplementation with *Bacillus altitudinis* spores improves porcine offspring growth performance and carcass weight. *British Journal of Nutrition*, 127(3): 403-420, ISSN: 0007-1145. <https://doi.org/10.1017/S0007114521001203>.
- Díaz, B., Elías, A. & Valiño E. 2014. Consorcios microbianos con actividad ácido-láctica promisorios aislados desde inoculantes bacterianos nativos para ensilajes. *Revista Ciencia y Agricultura*, 11(1): 17-25, ISSN: 2539-0899. <https://dialnet.unirioja.es/servlet/articulo?codigo=5039258>.
- Díaz, J., Elías, A. & Achan, J. 1996. Nota sobre el estudio de niveles de Vitafert (caña de azúcar + gallinaza) en precebas porcinas. *Cuban Journal of Agricultural Science*, 30(3): 309-311, ISSN: 2079-3480. <https://www.cabidigitallibrary.org/doi/full/10.5555/19971>.
- Díaz, Q.V. 2010. Efecto del vitafer en el comportamiento de ovinos en finalización en pastoreo suplementados con Sacchapulido. Tesis en opción al grado Master en Producción agropecuaria en el trópico. (1-51). Tabasco, México. Colegio de Postgraduados. <http://193.122.196.39:8080/xmlui/handle/10521/342>.
- Elías, A., Aguilera, L., Rodríguez, Y. & Herrera, F.R. 2009. Inclusión de niveles de harina de granos de *Canavalia ensiformis* en la fermentación de la caña de azúcar en estado sólido (Sacchacanavalia). *Cuban Journal of Agricultural Science*, 43(1): 51-54, ISSN: 0034-7485. <https://www.redalyc.org/pdf/1930/193015398009>.
- Elías, A. & Herrera, F.R. 2015. Producción de alimento para animales a través de procesos biotecnológicos sencillos con el empleo de Microorganismos Benéficos Activados (MEBA). Vitafert. En: Memorias del V Congreso de Producción Animal Tropical 2015. ISBN: 978-959-7171-70-6 Instituto de Ciencia Animal, San José de las Lajas, Mayabeque, Cuba, 8-13.
- Elías, A., Lezcano, O. & Herrera, F.R. 2001. Algunos indicadores bromatológicos y productos finales de la fermentación para la obtención de cuatro tipos de Saccharina inoculados con Vitafert. *Cuban Journal of Agricultural Science*, 35(2): 153-158, ISSN: 0034-7485. <https://www.redalyc.org/pdf/1930/193018220010>.
- Elías, A., Lezcano, O., Lezcano, P., Cordero, J. & Quintana, L. 1990. Reseña descriptiva sobre el desarrollo de una tecnología de enriquecimiento proteico en la caña de azúcar mediante fermentación en estado sólido (Saccharina). *Cuban Journal of Agricultural Science*, 24(1): 1-12, ISSN: 0034-7485. <https://www.cabidigitallibrary.org/doi/full/10.5555/19906773998>.
- García, Y., Elías, A., Albelo, N., Herrera, F.R., Núñez, O. & Dieppa, O. 2008. Growth of acid-lactic bacteria and yeasts during the liquid fermentation of broilers feces for the obtainment of probiotics. *Cuban Journal of Agricultural Science*, 42(2): 195-197. ISSN: 0034-7485. <https://cjas.science.com/index.php/CJAS/article/view/531>.
- González, R. & Enríquez, A.V. 1997. Nota sobre el efecto del Vitafert líquido (producido por la fermentación de la miel final y aditivos) en el valor nutritivo de una ración basada en heno de pasto estrella (*Cynodon nlemfuensis*). *Cuban Journal of Agricultural Science*, 31(1): 57-60, ISSN: 0034-7485. <https://agris.fao.org/search/en/providers/122595/records/647229>.
- Gutiérrez, D., Elías, A., García, R., Herrera, F., Jordán, H. & Sarduy, L. 2012. Effect of the microbial additive VITAFERT on the intake of dry matter and neutral detergent fiber in Saanen goats fed *Brachiaria brizantha* hay. *Cuban Journal of Agricultural Science*, 46(3): 267-271, ISSN: 2079-3480. <https://cjas.science.com/index.php/CJAS/article/view/42>.
- Gutiérrez, D., Morales A. Elías, A. García, R. & Sarduy L. 2014. Chemical composition and in situ ruminal degradability of dry matter in mixed silages of *Tithonia diversifolia*: *Pennisetum purpureum* cv. Cuba CT-169, inoculated with Vitafert. *Cuban Journal of Agricultural Science*, 48(4): 379-385, ISSN: 2079-3480. <https://cjas.science.com/index.php/CJAS/article/view/567>.
- Iraola, J., Elías, A., Gutiérrez, D., García, Y., Fraga, L.M., Vázquez, A., Barros-Rodríguez, M., Hernández, J.L. & Herrera, F. 2017. Efecto de microorganismos beneficiosos activados en la finalización de toros en silvopastoreo con leucaena, complementados con caña de azúcar. *Revista Científica FCV-LUZ*, XXVII 27 (6): 403-410, ISSN: 2521-9715. <https://www.redalyc.org/journal/959/95953773009>.
- Konieczka, P., Ferenc, K., Jørgensen, J.N., Hansen, L.H.B., Zabielski, R., Olszewski, J., Gajewski, Z., Mazur-Kusnerek, M., Szkopek, D., Szyrnska, N. & Lipinski, K. 2023. Feeding *Bacillus*-based probiotics to gestating and lactating sows is an efficient method for improving immunity, gut functional status and biofilm formation by probiotic bacteria in piglets at weaning. *Animal Nutrition*, 13: 361-372, ISSN: 2405-6545. <https://www.sciencedirect.com/science/article/pii/S2405654523000331>.
- Lambo, M.T., Chang, X. & Liu, D. 2021. The recent trend in the use of multistrain probiotics in livestock production: An Overview. *Animals*, 11(10): 2805-2814, ISSN: 2076-2615. <https://doi.org/10.3390/ani11102805>.
- Lazo-Pérez, L., Ruiz, Q.D., Elías, A., Herrera, F.R. & Zamora I. 2017. Efecto de un aditivo microbiano VITAFERT en algunos indicadores bioproductivos y de salud en cerdos en crecimiento. *Cuban Journal of Agricultural Science*, 51(3): 321-328, ISSN: 2079-3480. http://scielo.sld.cu/scielo.php?pid=S2079-34802017000300005&script=sci_arttext.

- López, C.A. 2010. Efecto de los niveles de Vitafert y melaza en la pollinaza fermentada aeróbica. Tesis en opción al grado de Master en Producción Agroalimentaria Trópico (1-40). Institución de enseñanza e investigación en ciencias, Agrícolas, Campus Tabasco, México <http://193.122.196.39:8080/xmlui/handle/10521/287>.
- López, C.A. 2012. Producción de un alimento fermentado en estado sólido a partir de la pollinaza y Vitafert. Tesis en opción al grado de Master en Producción Agroalimentaria Trópico (1-66). Institución de enseñanza e investigación en ciencias, Agrícolas, Campus Tabasco, México. <http://colposdigital.colpos.mx:8080/xmlui/handle/10521/1830>.
- Morales, A., Rodríguez, R., Gutiérrez, D., Elías, A., Gómez, S. & Sarduy L. 2016. Evaluación de la inclusión de VITAFERT en el valor nutritivo de ensilajes de *Tithonia diversifolia* y *Pennisetum purpureum*. *Cuban Journal of Agricultural Science*, 50(4): 619-630, ISSN: 2079-3480. https://scielo.sld.cu/scielo.php?pid=S2079-34802016000400011&script=sci_arttext.
- Pandey, A.K., Kumar, P. & Saxena, M.J. 2019. Feed additives in animal health. In: Gupta R, Srivastava A, Lall R, editors. *Nutraceuticals in veterinary medicine*. Cham: Springer; 345-462, ISBN: Print 978-3-030-04623-1, Online: 978-3-030-04624-8. https://doi.org/10.1007/978-3-030-04624-8_23.
- Quach, N.T., Vu, T.H.N., Nguyen, N.A., Nguyen, V.T., Bui, T.L. & Ky, S.C. 2021. Phenotypic features and analysis of genes supporting probiotic action unravel underlying perspectives of *Bacillus velezensis* VTX9 as a potential feed additive for swine. *Annals of Microbiology*, 71(1): 1-14. ISSN: 1869-2044. <https://doi.org/10.1186/s13213-021-01646-4>.
- Rasaei, D., Alemeh, S.H., Asasi, K., Shahram, S.S. & Khodakaram, A.T. 2023. The beneficial effects of spraying of probiotic *Bacillus* and *Lactobacillus* bacteria on broiler chickens experimentally infected with avian influenza virus H9N2. *Poultry Science*, 102(7): 1-12, ISSN: 1525-3171. <https://doi.org/10.1016/j.psj.2023.102669>.
- Rodríguez, R., Herrera, F., Gómez, S., González, N., Alonso G.J., Elías, A., Moreira, O., Sarduy, L. & Medina, Y. 2019. Effects of including sweet potato (*Ipomoea batatas*) and Vitafert as additives on the nutritional value of *Cenchrus purpureus* cv. CUBA CT-169 and *Moringa oleifera* silages. *Cuban Journal of Agricultural Science*, 53(2): 119-133. ISSN: 2079-3480. <https://www.cjascience.com/index.php/CJAS/article/view/877>.
- Rodríguez, R., Lores J., Gutiérrez, D., Ramirez A., Gómez S., Elías, A., Aldana A.I., Moreira O., Sarduy L. & Jay O. 2013. Inclusión del aditivo microbiano Vitafert en la fermentación ruminal *in vitro* de una dieta para cabras. *Cuban Journal of Agricultural Science*, 47(2): 171-178, ISSN: 0034-7485. <https://www.redalyc.org/pdf/1930/193028751011.pdf>.
- Ruiz, P.A. 2014. Efecto de la adición a la dieta de un producto biológicamente activo en el crecimiento y calidad de la carne de corderos Suffolk-Hampshire. Ganadería. Tesis en opción al grado de Master en Zootecnia. Montecillo (1-67), Texcoco, México. <http://colposdigital.colpos.mx:8080/xmlui/handle/10521/2465>.
- Savón, L., Sánchez, B., Elías, A., Ortega, H.J., Gutiérrez, M., Scull, I. & Herrera, M. 2020. Effect of a dry fermented product on morphological, immunological, histological and health indicators of broilers. *Cuban Journal of Agricultural Science*, 54(1): 85-94, ISSN: 2079-3480. <https://www.cjascience.com/index.php/CJAS/article/view/947>.
- Sun, Y., Duarte, M.E. & Kim, S.W. 2021. Dietary inclusion of multispecies probiotics to reduce the severity of post-weaning diarrhea caused by *Escherichia coli* F18 (p) in pigs. *Animal Nutrition*, 7(2): 326-333, ISSN: 2405-6383. <https://doi.org/10.1016/j.aninu.2020.08.012>.
- Valdivié, M. & Elías, A. 2006. Posibilidades del grano de *Canavalia ensiformis* fermentado con caña (*Sacchacana*) en pollos de ceba. *Cuban Journal of Agricultural Science*, 40(4): 459-464, ISSN: 2079-3480. <https://www.redalyc.org/pdf/1930/193017672010>.
- Valdivié, M., González, L.M. & Elías, A. 1997. Nuevos tipos de Saccharina para aves. *Cuban Journal of Agricultural Science*, 31(3): 231-236, ISSN: 0034-7485. <https://www.cabidigitallibrary.org/doi/full/10.5555/19971401347>.
- Vitaluña, O.V.M. 2014. Evaluación de diferentes niveles de VITAFERT en crecimiento – engorde de cerdos. Trabajo de Titulación para Ing. Zootecnista. Riobamba, Ecuador. p. 48. <http://dspace.espace.edu.Ec>.



Available in:

<https://www.redalyc.org/articulo.oa?id=653777768017>

How to cite

Complete issue

More information about this article

Journal's webpage in redalyc.org

Scientific Information System Redalyc
Diamond Open Access scientific journal network
Non-commercial open infrastructure owned by academia

Elaine C. Valiño Cabrera, Yaneisy García Hernández,
Daymara Bustamante García, A. Beruvides Rodríguez

Development and use of Vitafert in the agricultural sector
Desarrollo y utilización del Vitafert en el sector
agropecuario

Cuban Journal of Agricultural Science

vol. 58, e17, 2024

Ediciones ICA,

ISSN: 0864-0408

ISSN-E: 2079-3480