



Siembra

ISSN: 1390-8928

ISSN: 2477-8850

siembra.fag@uce.edu.ec

Universidad Central del Ecuador

Ecuador

Campos, Edison  
Recovery of hardpan «cangahua» soils: experience in Chimborazo and new conservation perspectives  
Siembra, vol. 13, no. 3, Esp., e9603, 2026  
Universidad Central del Ecuador  
Quito, Ecuador

DOI: [https://doi.org/10.29166/siembra.v13i3\(Especial\).9603](https://doi.org/10.29166/siembra.v13i3(Especial).9603)

Available in: <https://www.redalyc.org/articulo.oa?id=653883476008>

- ▶ How to cite
- ▶ Complete issue
- ▶ More information about this article
- ▶ Journal's webpage in redalyc.org

redalyc.org

Scientific Information System Redalyc

Diamond Open Access scientific journal network

Non-commercial open infrastructure owned by academia

## Recovery of hardpan «cangahua» soils: experience in Chimborazo and new conservation perspectives

Edison Campos<sup>1</sup>

*Siembra* 13 (3) (2026): Special Edition  
PROCEEDINGS OF FIRST INTERNACIONAL  
SYMPOSIUM ON SOIL HEALTH

DOI: [10.29166/siembra.v13i3\(Especial\).9603](https://doi.org/10.29166/siembra.v13i3(Especial).9603)



<sup>1</sup> Universidad Nacional Agraria La Molina, Facultad de Agronomía. Lima, Perú\*

Correspondence: [fmontesdeoca@uce.edu.ec](mailto:fmontesdeoca@uce.edu.ec)

### Abstract

### Abstract

Chimborazo is one of the poorest provinces in Ecuador, 6.30% of its territory is hardpan «cangahua» soils, which corresponds to 82,778.05 ha. The cangahuas are sterile volcanic tuffs, their rehabilitation and management present several environmental, social and economic challenges that considerably slow down agricultural development. The recovery of the cangahuas opens the possibility of revitalizing depressed agriculture. The experience of restoring eroded and degraded soils through physical and biological systems carried out in 3 cantons and 10 communities of Chimborazo allowed the recovery of 79.15 ha for agricultural production. Through soil plowing, incorporation of green manures, delivery of native plants and ongoing technical assistance and training. This allowed the soil organic matter to increase from 1.7% to 1.91% in recovered soil, also increased the economic income of 438 producers by an average of 10.12%.

**Keywords:** Degraded soils, Soil organic matter, Green manures

SIEMBRA  
<https://revistadigital.uce.edu.ec/index.php/SIEMBRA>  
ISSN-e: 2477-8850  
Frequency: half-yearly  
vol. 13, issue 3 Special, 2026  
[siembra.fag@uce.edu.ec](mailto:siembra.fag@uce.edu.ec)



© The Authors 2026

This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License