



Vojnotehnicki glasnik/Military Technical Courier
ISSN: 0042-8469
ISSN: 2217-4753
vojnotehnicki.glasnik@mod.gov.rs
University of Defence
Serbia

Some new observations on fixed point results in rectangular metric spaces with applications to chemical sciences

Younis, Mudasir; Fabiano, Nicola; Fadail, Zaid M.; Mitrović, Zoran D.; Radenović, Stojan N.

Some new observations on fixed point results in rectangular metric spaces with applications to chemical sciences

Vojnotehnicki glasnik/Military Technical Courier, vol. 69, no. 1, 2021

University of Defence, Serbia

Available in: <https://www.redalyc.org/articulo.oa?id=661771133002>

DOI: <https://doi.org/10.5937/vojtehg69-29517>

<http://www.vtg.mod.gov.rs/copyright-notice-and-self-archiving-policy.html>



This work is licensed under Creative Commons Attribution 4.0 International.

Some new observations on fixed point results in rectangular metric spaces with applications to chemical sciences

Примечания к некоторым результатам в области неподвижных точек в прямоугольных метрических пространствах, с их применением в химии

Напомене о неким резултатима о непокретним тачкама управоугаоним метричким просторима са применама у хемији

Mudasir Younis
University Institute of Technology - RGPV, India
mudasiryouniscuk@gmail.com
 <https://orcid.org/0000-0001-5499-4272>

DOI: <https://doi.org/10.5937/vojtehg69-29517>
Redalyc: <https://www.redalyc.org/articulo.oa?id=661771133002>

Nicola Fabiano
Independent researcher, Italy
nicola.fabiano@gmail.com
 <https://orcid.org/0000-0003-1645-2071>

Zaid M. Fadail
Thamar University, Yemen
zaid.fadail@tu.edu.ye
 <https://orcid.org/0000-0003-0899-0056>

Zoran D. Mitrović
University of Banja Luka, Bosnia and Herzegovina
zoran.mitrovic@etf.unibl.org
 <https://orcid.org/0000-0001-9993-9082>

Stojan N. Radenović
University of Belgrade, Serbia
radens@beotel.rs
 <https://orcid.org/0000-0001-8254-6688>

Received: 21 November 2020

Revised document received: 21 December 2020

Accepted: 23 December 2020

ABSTRACT:

Introduction/purpose: This paper considers, generalizes and improves recent results on fixed points in rectangular metric spaces. The aim of this paper is to provide much simpler and shorter proofs of some new results in rectangular metric spaces.

Methods: Some standard methods from the fixed point theory in generalized metric spaces are used.

Results: The obtained results improve the well-known results in the literature. The new approach has proved that the Picard sequence is Cauchy in rectangular metric spaces. The obtained results are used to prove the existence of solutions to some nonlinear problems related to chemical sciences. Finally, an open question is given for generalized contractile mappings in rectangular metric spaces.

AUTHOR NOTES

zoran.mitrovic@etf.unibl.org

Conclusions: New results are given for fixed points in rectangular metric spaces with application to some problems in chemical sciences.

KEYWORDS: fixed point, rectangular metric space, contractive map, Green function.

Р е з ю м е :

Введение/цель: В данной статье обсуждаются, суммируются и улучшаются недавние результаты о неподвижных точках в прямоугольных метрических пространствах. Целью данной статьи является представление гораздо более простых и коротких доказательств некоторых новых результатов в области прямоугольных метрических пространств.

Методы: В статье применены стандартные методы теории неподвижной точки в обобщенных метрических пространствах.

Результаты: Полученные результаты данного исследования улучшают известные результаты существующей литературы. Благодаря новому подходу доказана последовательность Коши-Пикара в прямоугольных метрических пространствах. Полученные результаты также используются для доказательства экзистенциальных решений некоторых нелинейных задач, относящихся к химическим наукам. В конце статьи задается открытый вопрос в связи с обобщенными сжатыми отображениями в прямоугольных метрических пространствах.

Выводы: В статье приведены новые результаты, касающиеся теории неподвижных точек в прямоугольных метрических пространствах, примененные в решении некоторых проблем в области химических наук.

К л ю ч е в ы е с л о в а : неподвижная точка, прямоугольное метрическое пространство, сжатое отображение, функция Грина.

ABSTRACT:

Увод/ціль: У овом раду се разматрају, уопштавају и побољшавају недавни резултати о непокретним тачкама у оквиру правоугаоних метричких простора. Циљ овог рада је да пружи много једноставније и краће доказе о неким новим резултатима у правоугаоним метричким просторима.

Метод: Користе се стандардне методе из теорије непокретне тачке у генерализованим метричким просторима.

Резултати: Добијени резултати побољшавају добро познате резултате у литератури. Користећи нови приступ доказује се да је Пикаров низ Кошијев у оквиру правоугаоних метричких простора. Добијени резултати користе се за доказ егзистенције решења неких нелинеарних проблема који се примењују у хемијским наукама. На крају се даје једно отворено питање за генерализована контрактивна пресликавања у правоугаоним метричким просторима.

Закључак: Дати су нови резултати за непокретне тачке у правоугаоним метричким просторима са применом на неке проблеме у хемијским наукама.

KEYWORDS: непокретна тачка, правоугаони метрички простор, контрактивно пресликавање, Гринава функција.

INTRODUCTION AND PRELIMINARIES

It is well known that the Banach contraction principle (Banach, 1922) is one of the most important and attractive results in nonlinear analysis and mathematical analysis in general. The whole fixed point theory is a significant subject in different fields: geometry, differential equations, informatics, physics, economics, engineering, and many others. After solutions are guaranteed, numerical methodology is established to obtain the approximated solution. The fixed point of functions depends heavily on considered spaces defined using intuitive axioms. In particular, variants of generalized metric spaces are proposed, e.g. partial metric space, b-metric, partial b-metric, extended b-metric, rectangular metric, rectangular b-metric, Gmetric, G_b -metric, S-metric, S_b -metric, cone metric, cone b-metric, fuzzy metric, fuzzy b-metric, probabilistic metric, etc. For more details on all variants of generalized metric spaces, see (Budhia et al, 2017), (Collaco & Silva, 1997).

In this paper, we will discuss some results recently established in (Alsulami et al, 2015) and (Budhia et al, 2017). Firstly, we give the basic notion of a rectangular metric space (g.m.s or RMS by some authors).

Definition 1. Let X be a nonempty set and let $dr : X \times X \rightarrow [0, +\infty)$ satisfy the following conditions: for all $x, y \in X$ and all distinct $u, v \in X$ each of them different from x and y .

(i) $dr(x, y) = 0$ if and only if $x = y$,

(ii) $dr(x, y) = dr(y, x)$,

(iii) $dr(x, y) \leq dr(x, u) + dr(u, v) + dr(v, y)$ (quadrilateral inequality).

Then the function d_r is called a rectangular metric and the pair (X, d_r) is called a rectangular metric space (RMS for short).

Notice that the definitions of convergence and Cauchyness of the sequences in rectangular metric spaces are the same as the ones found in the standard metric spaces. Also, a rectangular metric space (X, d_r) is complete if each Cauchy sequence in it is convergent. Samet et al. (Samet et al, 2012) introduced the concept of α - ψ -contractive mappings and proved the fixed point theorems for such mappings. In (Karapinar, 2014), Karapinar gave contractive conditions to obtain the existence and uniqueness of a fixed point of α - ψ contraction mappings in rectangular metric spaces. Salimi et al. (Salimi et al, 2013) introduced modified α - ψ contractive mappings and obtained some fixed point theorems in a complete metric space. Alsulami et al. (Alsulami et al, 2015) established some fixed point theorems for α - ψ -rational type contractive mappings in a rectangular metric space.

Let Ψ be the family of all functions $\psi: [0, +\infty) \rightarrow [0, +\infty)$ such that ψ is nondecreasing and $\sum_{n=1}^{\infty} \psi^n(t) < +\infty$ for each $t > 0$. Obviously, if $\psi \in \Psi$, then $\psi(t) < t$ for each $t > 0$.

Definition 2. (Salimi et al, 2013) Let T be a self mapping on a metric space (X, d_r) and let $\alpha, \eta: X \times X \rightarrow [0, +\infty)$ be two functions. It is called an α -admissible mapping with respect to η if $\alpha(x, y) \geq \eta(x, y)$ implies that $\alpha(Tx, Ty) \geq \eta(Tx, Ty)$ for all $x, y \in X$.

If $\eta(x, y) = 1$ for all $x, y \in X$, then T is called an α -admissible mapping.

It is called a triangular α -admissible mapping if for all $x, y, z \in X$ holds: $(\alpha(x, y) \geq 1 \text{ and } \alpha(y, z) \geq 1)$ implies $\alpha(x, z) \geq 1$.

Otherwise, a rectangular metric space (X, d_r) is α -regular with respect to η if for any sequence in X such that $\alpha(x_n, x_{n+1}) \geq \eta(x_n, x_{n+1})$ for all $n \in \mathbb{N}$ and $x_n \rightarrow x$ as $n \rightarrow +\infty$, then $\alpha(x_n, x) \geq \eta(x_n, x)$.

For more details on a triangular α -admissible mapping, see (Karapinar et al, 2013), pages 1 and 2. In this paper, we will use the following result:

Lemma 1. (Karapinar et al, 2013), Lemma 7. Let T be a triangular α -admissible mapping. Assume that there exists $x_0 \in X$ such that $\alpha(x_0, Tx_0) \geq 1$. Define the sequence $\{x_n\}$ by $x_n = T^n x_0$. Then

$$\alpha(x_m, x_n) \geq 1 \text{ for all } m, n \in \mathbb{N} \cup \{0\} \text{ with } m < n.$$

In (Budhia et al, 2017), the authors proved the following result:

Theorem 1. Let (X, d_r) be a Hausdorff and complete rectangular metric space, and let $T: X \rightarrow X$ be an α -admissible mapping with respect to η . Assume that there exists a continuous function $\psi \in \Psi$ such that

$$x, y \in X, \alpha(x, y) \geq \eta(x, y) \text{ implies } d_r(Tx, Ty) \leq \psi(M(x, y)),$$

Theorem 1. Let (X, d_r) be a Hausdorff and complete rectangular metric space, and let $T: X \rightarrow X$ be an α -admissible mapping with respect to η . Assume that there exists a continuous function $\psi \in \Psi$ such that where

$$M(x, y) = \max \left\{ d_r(x, y), d_r(x, Tx), d_r(y, Ty), \frac{d_r(x, Tx) d_r(y, Ty)}{1 + d_r(x, y)}, \frac{d_r(x, Tx) d_r(y, Ty)}{1 + d_r(Tx, Ty)} \right\}.$$

Also, suppose that the following assertions are hold:

1. there exists $x_0 \in X$ such that $\alpha(x_0, Tx_0) \geq \eta(x_0, Tx_0)$,
2. for all $x, y, z \in X$, $(\alpha(x, y) \geq \eta(x, y) \text{ and } \alpha(y, z) \geq \eta(y, z))$ implies $\alpha(x, z) \geq \eta(x, z)$,

3. either T is continuous or X is α -regular with respect to η .

Then T has a periodic point $a \in X$ and if $\alpha(a, Ta) \geq \eta(a, Ta)$ holds for each periodic point, then T has a fixed point. Moreover, if for all $x, y \in F(T)$, we have $\alpha(x, y) \geq \eta(x, y)$, then the fixed point is unique.

Taking $\eta(x, y) = 1$ for $x, y \in X$, the authors obtained the following corollary:

Corollary 1. Let (X, d_r) be a Hausdorff and complete rectangular metric space, and let $T : X \rightarrow X$ be an α -admissible mapping. Assume that there exists a continuous function $\psi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ such that

$$x, y \in X, \alpha(x, y) \geq 1 \text{ implies } d_r(Tx, Ty) \leq \psi(M(x, y))$$

where

$$M(x, y) = \max \left\{ d_r(x, y), d_r(x, Tx), d_r(y, Ty), \frac{d_r(x, Tx) d_r(y, Ty)}{1 + d_r(x, y)}, \frac{d_r(x, Tx) d_r(y, Ty)}{1 + d_r(Tx, Ty)} \right\}.$$

Also, suppose that the following assertions are hold:

1. there exists $x_0 \in X$ such that $\alpha(x_0, Tx_0) \geq 1$,
2. for all $x, y, z \in X$ ($\alpha(x, y) \geq 1$ and $\alpha(y, z) \geq 1$) implies $\alpha(x, z) \geq 1$,
3. either T is continuous or (X, d_r) is α -regular.

Then T has a periodic point $a \in X$ and if $\alpha(a, Ta) \geq 1$ holds T has a fixed point. Moreover, if for all $x, y \in F(T)$, we have $\alpha(x, y) \geq 1$, then the fixed point is unique.

Further, taking $\alpha(x, y) = 1$ for $x, y \in X$ authors obtained the following corollary:

Corollary 2. Let (X, d_r) be a Hausdorff and complete rectangular metric space, and let $T : X \rightarrow X$ be an α -admissible mapping. Assume that there exists a continuous function $\psi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ such that

$$x, y \in X, 1 \geq \eta(x, y) \text{ implies } d_r(Tx, Ty) \leq \psi(M(x, y))$$

where

$$M(x, y) = \max \left\{ d_r(x, y), d_r(x, Tx), d_r(y, Ty), \frac{d_r(x, Tx) d_r(y, Ty)}{1 + d_r(x, y)}, \frac{d_r(x, Tx) d_r(y, Ty)}{1 + d_r(Tx, Ty)} \right\}.$$

Also, suppose that the following assertions are hold:

1. there exists $x_0 \in X$ such that $1 \geq \eta(x_0, Tx_0)$,
2. for all $x, y, z \in X$ ($1 \geq \eta(x, y)$ and $1 \geq \eta(y, z)$) implies $1 \geq \eta(x, z)$,

For $\psi(t) = kt$, $0 < k < 1$ then the authors obtained

Corollary 3. Let (X, d_r) be a Hausdorff and complete rectangular metric space, and let $T : X \rightarrow X$ be an α -admissible mapping with respect to η . Assume that

$$x, y \in X, \alpha(x, y) \geq \eta(x, y) \text{ implies } d_r(Tx, Ty) \leq kM(x, y),$$

where

$$M(x, y) = \max \left\{ d_r(x, y), d_r(x, Tx), d_r(y, Ty), \frac{d_r(x, Tx) d_r(y, Ty)}{1 + d_r(x, y)}, \right. \\ \left. \frac{d_r(x, Tx) d_r(y, Ty)}{1 + d_r(Tx, Ty)} \right\}.$$

Also, suppose that the following assertions are hold:

1. there exists $x_0 \in X$ such that $\alpha(x_0, Tx_0) \geq \eta(x_0, Tx_0)$,
2. for all $x, y, z \in X$ ($\alpha(x, y) \geq \eta(x, y)$ and $\alpha(y, z) \geq \eta(y, z)$) implies $\alpha(x, z) \geq \eta(x, z)$,
3. either T is continuous or (X, d_r) is α -regular.

Then T has a periodic point $a \in X$ and if $\alpha(a, Ta) \geq \eta(a, Ta)$ holds, T has a fixed point. Moreover, if for all $x, y \in F(T)$, we have $\alpha(x, y) \geq \eta(x, y)$, then the fixed point is unique.

The following two lemmas are a rectangular metric space modification of a result which is well known in the metric space, see, e.g. (Radenović et al, 2012), Lemma 2.1. Many known proofs of fixed point results in rectangular metric spaces become much more straightforward and shorter using both lemmas. Also, in the proofs of the main results in this paper, we will use both lemmas:

Lemma 2. (Kadelburg & Radenović, 2014a), (Kadelburg & Radenović, 2014b) *Let (X, d_r) be a rectangular metric space and let $\{x_n\}$ be a sequence in it with distinct elements ($x_n \neq x_m$ for $n \neq m$). Suppose that $d_r(x_n, x_{n+1})$ and $d_r(x_n, x_{n+2})$ tend to 0 as $n \rightarrow +\infty$ and that $\{x_n\}$ is not a Cauchy sequence. Then there exists $\varepsilon > 0$ and two sequences $\{m(k)\}$ and $\{n(k)\}$ of positive integers such that $n(k) > m(k) > k$ and the following sequences tend to ε as $k \rightarrow +\infty$:*

$$\{d_r(x_{n(k)}, x_{m(k)})\}, \{d_r(x_{n(k)+1}, x_{m(k)})\}, \{d_r(x_{n(k)}, x_{m(k)-1})\}, \\ \{d_r(x_{n(k)+1}, x_{m(k)-1})\}, \{d_r(x_{n(k)+1}, x_{m(k)+1})\}.$$

Lemma 3. *Let $\{x_{n+1}\}_{n \in \mathbb{N} \cup \{0\}} = \{Tx_n\}_{n \in \mathbb{N} \cup \{0\}} = \{T^n x_0\}_{n \in \mathbb{N} \cup \{0\}}$, $T^0 x_0 = x_0$ be a Picard sequence in a rectangular metric space (X, d_r) induced by the mapping $T: X \rightarrow X$ and the initial point $x_0 \in X$. If $d_r(x_n, x_{n+1}) < d_r(x_{n-1}, x_n)$ for all $n \in \mathbb{N}$ then $x_n \neq x_m$ whenever $n \neq m$.*

Proof. Let $x_n = x_m$ for some $n, m \in \mathbb{N}$ with $n < m$. Then $x_{n+1} = Tx_n = Tx_m = x_{m+1}$. Further, we get

$$d_r(x_n, x_{n+1}) = d_r(x_m, x_{m+1}) < d_r(x_{m-1}, x_m) < \dots < d_r(x_n, x_{n+1}),$$

which is a contradiction.

In some proofs, we will also use and the following interesting as well as significant result in the context of rectangular metric spaces:

Proposition 1. (Kirk & Shahzad, 2014), *Proposition 3. Suppose that $\{q_n\}$ is a Cauchy sequence in a rectangular metric space (X, d_r) and suppose $\lim_{n \rightarrow +\infty} d_r(q_n, q) = 0$. Then $\lim_{n \rightarrow +\infty} d_r(q_n, p) = d_r(q, p)$ for all $p \in X$. In particular, $\{q_n\}$ does not converge to p if $p \neq q$.*

MAIN RESULTS

In this section, we generalize and improve Theorem 2 and all its corollaries. The obtained generalizations extend the result in several directions. Namely, we will use only one function $\alpha : X \times X \rightarrow [0, +\infty)$ instead of two α and η as in (Budhia et al, 2017), Definition 2.3. and Definition 3.1. This is possible according to the (Mohammadi & Rezapour, 2013), Page 2, after Theorem 1.2. Note that we assume neither that the rectangular metric space is Hausdorff, nor that the mapping d_r is continuous.

The authors (Alsulami et al, 2015), page 6, line 6+, say that the sequence $\{x_n\}$ in a rectangular metric space (X, d_r) is a Cauchy if $\lim_{n \rightarrow +\infty} d_r(x_n, x_{n+k}) = 0$, for all $k \in \mathbb{N}$. However, it is well know that this claim is dubious. Therefore, we also improve the proof that the sequence $\{x_n\}$ is Cauchy

Our first new result in this paper is the following:

Theorem 2. *Let (X, d_r) be a complete rectangular metric space and let $T : X \rightarrow X$ be a triangular α -admissible mapping. Assume that there exists continuous function $\psi \# \Psi$ such that*

$$x, y \in X, \alpha(x, y) \geq 1 \text{ implies } d_r(Tx, Ty) \leq \psi(M(x, y)), \quad (1)$$

where

$$M(x, y) = \max \left\{ d_r(x, y), d_r(x, Tx), d_r(y, Ty), \frac{d_r(x, Tx) d_r(y, Ty)}{1 + d_r(x, y)}, \right. \\ \left. \frac{d_r(x, Tx) d_r(y, Ty)}{1 + d_r(Tx, Ty)} \right\}.$$

Also, suppose that the following assertions are hold:

1. there exists $x_0 \in X$ such that $\alpha(x_0, Tx_0) \geq 1$
2. either T is continuous or (X, d) α -regular

Then T has a fixed point.

Morover, if for all $x, y \in F(T)$ implies $\alpha(x, y) \geq 1$ then the fixed point is unique.

Proof. Given $x_0 \in X$ such that

$$\alpha(x_0, Tx_0) \geq 1. \quad (2)$$

Define a sequence $\{x_n\}$ in X by $x_n = Tx_{n-1} = T^n x_0$ for all $n \in \mathbb{N}$. If $x_{k+1} = x_k$ for some $k \in \mathbb{N}$, then $Tx_k = x_k$, i.e., x_k is a fixed point of T and the proof is finished. From now on, suppose that $x_n \neq x_{n+1}$ for all $n \in \mathbb{N} \cup \{0\}$. Using (2) and the fact that T is an α -admissible mapping, we have

$$\alpha(x_1, x_2) = \alpha(Tx_0, Tx_1) \geq 1.$$

By induction, we get

$$\alpha(x_n, x_{n+1}) \geq 1 \text{ for all } n \in \mathbb{N} \cup \{0\}.$$

In the first step, we will show that the sequence $\{d_r(x_n, x_{n+1})\}$ is nonincreasing and $d_r(x_n, x_{n+1}) \rightarrow 0$ as $n \rightarrow +\infty$. From (1), recall that

$$d_r(x_n, x_{n+1}) = d_r(Tx_{n-1}, Tx_n) \leq \psi(M(x_{n-1}, x_n)), \quad (3)$$

where

$$\begin{aligned} M(x_{n-1}, x_n) &= \max \{d_r(x_{n-1}, x_n), d_r(x_{n-1}, x_n), d_r(x_n, x_{n+1}), \\ &\quad \frac{d_r(x_{n-1}, x_n) d_r(x_n, x_{n+1})}{1 + d_r(x_{n-1}, x_n)}, \frac{d_r(x_{n-1}, x_n) d_r(x_n, x_{n+1})}{1 + d_r(x_n, x_{n+1})}\} \\ &\leq \max \{d_r(x_{n-1}, x_n), d_r(x_n, x_{n+1})\}. \end{aligned}$$

Now from (3) follows

$$d_r(x_n, x_{n+1}) \leq \psi(\max \{d_r(x_{n-1}, x_n), d_r(x_n, x_{n+1})\}). \quad (4)$$

If $\max \{d_r(x_{n-1}, x_n), d_r(x_n, x_{n+1})\} = d_r(x_n, x_{n+1})$, we get a contradiction. Indeed, (4) implies

$$d_r(x_n, x_{n+1}) \leq \psi(d_r(x_n, x_{n+1})) < d_r(x_n, x_{n+1}).$$

Therefore, we get that $d_r(x_n, x_{n+1}) < d_r(x_{n-1}, x_n)$. This means that there exists $\lim_{n \rightarrow +\infty} d_r(x_n, x_{n+1}) = d_r^* \geq 0$. If $d_r^* > 0$, then from (3) follows

$$\begin{aligned} d_r^* &\leq \psi \left(\max \left\{ d_r^*, d_r^*, d_r^*, \frac{d_r^{*2}}{1 + d_r^*}, \frac{d_r^{*2}}{1 + d_r^*} \right\} \right) \\ &\leq \psi(\max \{d_r^*, d_r^*\}) < d_r^*, \end{aligned}$$

which is a contradiction. Hence $\lim_{n \rightarrow +\infty} d_r(x_n, x_{n+1}) = 0$.

Further, we will also show that $\lim_{n \rightarrow +\infty} d_r(x_n, x_{n+2}) = 0$. Firstly, we have that $\alpha(x_{n-1}, x_n) \geq 1$, i.e., $\alpha(x_{n-1}, x_{n+1}) \geq 1$, because T is a triangular α -admissible mapping. Therefore,

$$d_r(x_n, x_{n+2}) = d_r(Tx_{n-1}, Tx_{n+1}) \leq \psi(M(x_{n-1}, x_{n+1})),$$

where

$$M(x_{n-1}, x_{n+1}) = \max \{d_r(x_{n-1}, x_{n+1}), d_r(x_{n-1}, x_n), d_r(x_{n+1}, x_{n+2}), \\ \frac{d_r(x_{n-1}, x_n) d_r(x_{n+1}, x_{n+2})}{1 + d_r(x_{n-1}, x_{n+1})}, \frac{d_r(x_{n-1}, x_n) d_r(x_{n+1}, x_{n+2})}{1 + d_r(x_n, x_{n+2})}\}.$$

Since $\frac{d_r(x_{n-1}, x_n) d_r(x_{n+1}, x_{n+2})}{1 + d_r(x_{n-1}, x_{n+1})} \leq d_r(x_{n-1}, x_n) d_r(x_{n+1}, x_{n+2})$ and $\frac{d_r(x_{n-1}, x_n) d_r(x_{n+1}, x_{n+2})}{1 + d_r(x_n, x_{n+2})} \leq d_r(x_{n-1}, x_n) d_r(x_{n+1}, x_{n+2})$ we get that

$$M(x_{n-1}, x_{n+1}) \leq \max \{d_r(x_{n-1}, x_{n+1}), d_r(x_{n-1}, x_n), \\ d_r(x_{n+1}, x_{n+2}), d_r(x_{n-1}, x_n) d_r(x_{n+1}, x_{n+2})\}$$

that is,

$$M(x_{n-1}, x_{n+1}) \leq \max \{d_r(x_{n-1}, x_{n+1}), d_r(x_{n-1}, x_n), d_r^2(x_{n-1}, x_n)\} \\ \leq \max \{d_r(x_{n-1}, x_{n+1}), d_r(x_{n-1}, x_n)\}.$$

The last relation follows from the fact that $d_r(x_{n-1}, x_n) \rightarrow 0$ as $n \rightarrow +\infty$. Hence, for some $n_1 \in \mathbb{N}$, we have that

$$d_r(x_n, x_{n+2}) \leq \max \{d_r(x_{n-1}, x_{n+1}), d_r(x_{n-1}, x_n)\},$$

whenever $n \geq n_1$. Since, $d_r(x_{n-1}, x_n) \rightarrow 0$ as $n \rightarrow +\infty$ it is not hard to check that also $d_r(x_n, x_{n+2}) \rightarrow 0$ as $n \rightarrow +\infty$.

In order to prove that the sequence $\{x_n\}$ is a Cauchy one, we use Lemma 6. Namely, since according to Lemma 1, $\alpha(x_{n(k)}, x_{m(k)}) \geq 1$ if $m(k) < n(k)$, then, by putting in (1) $x = x_{n(k)}$, $y = x_{m(k)}$, we obtain

$$d_r(x_{n(k)+1}, x_{m(k)+1}) \leq \psi(M(x_{n(k)}, x_{m(k)})), \quad (5)$$

where

$$M(x_{n(k)}, x_{m(k)}) = \\ = \max \{d_r(x_{n(k)}, x_{m(k)}), d_r(x_{n(k)}, x_{n(k)+1}), d_r(x_{m(k)}, x_{m(k)+1}), \\ \frac{d_r(x_{n(k)}, x_{n(k)+1}) d_r(x_{m(k)}, x_{m(k)+1})}{1 + d_r(x_{n(k)}, x_{m(k)})}, \frac{d_r(x_{n(k)}, x_{n(k)+1}) d_r(x_{m(k)}, x_{m(k)+1})}{1 + d_r(x_{n(k)+1}, x_{m(k)+1})}\} \\ \xrightarrow{k \rightarrow +\infty} M(x_{n(k)}, x_{m(k)}) = \max \left\{ \varepsilon, 0, 0, \frac{0 \cdot 0}{1 + \varepsilon}, \frac{0 \cdot 0}{1 + \varepsilon} \right\} = \varepsilon.$$

Now, taking in (5) the limit as

$$\varepsilon \leq \psi(\varepsilon) < \varepsilon,$$

which is a contradiction. The sequence $\{x_n\}$ is hence a Cauchy one. Since (X, d_r) is a complete rectangular metric space, there exists a point $x^* \in X$ such that $x_n \rightarrow x^*$ as $n \rightarrow +\infty$. If T is continuous, we get that $x_{n+1} = Tx_n \rightarrow Tx^*$ as $n \rightarrow +\infty$. Let $Tx^* \neq x^*$. Since $d_r(x_n, x_{n+1}) < d_r(x_{n-1}, x_n)$ for all $n \in \mathbb{N} \cup \{0\}$, then, according to

Lemma 7, we have that all x_n are distinct. Therefore, there exists $n_2 \in \mathbb{N}$ such that $x^*, Tx^* \notin \{X_n\}_{n \geq n_2}$. Further, by (iii) follows:

$$d_r(x^*, Tx^*) \leq d_r(x^*, x_n) + d_r(x_n, x_{n+1}) + d_r(x_{n+1}, Tx^*),$$

whenever $n \geq n_2$, taking the limit, we obtain $d_r(x^*, Tx^*) = 0$, i.e. $x^* = Tx^*$, which is a contradiction.

In the case that (X, d_r) is α -regular, we get the following: Since $\alpha(x_n, x^*) \geq 1$ for all $n \in \mathbb{N}$, then from (1) follows

$$d_r(Tx_n, Tx^*) \leq \psi(M(x_n, x^*)), \quad (6)$$

where

$$M(x_n, x^*) = \max\{d_r(x_n, x^*), d_r(x_n, x_{n+1}), d_r(x^*, Tx^*), \\ \frac{d_r(x_n, x_{n+1})d_r(x^*, Tx^*)}{1 + d_r(x_n, x^*)}, \frac{d_r(x_n, x_{n+1})d_r(x^*, Tx^*)}{1 + d_r(x_{n+1}, Tx^*)}\} \xrightarrow{n \rightarrow +\infty} d_r(x^*, Tx^*).$$

By taking in (6) the limit as $n \rightarrow +\infty$ and by using Proposition 8 and the continuity of the function ψ , we get $d_r(x^*, Tx^*) \leq \psi(d_r(x^*, Tx^*)) < d_r(x^*, Tx^*)$ if $x^* \neq Tx^*$, which is a contradiction. Hence, x^* is a fixed point of T .

Now, we show that the fixed point is unique if $\alpha(x, y) \geq 1$ whenever $x, y \in F(T)$. Indeed, in this case, by contractive condition (1), for such possible fixed points x, y we have

$$d_r(x, y) = d_r(Tx, Ty) \leq \psi(M(x, y)), \quad (7)$$

where

$$M(x, y) = \max\left\{d_r(x, y), d_r(x, Tx), d_r(y, Ty), \frac{d_r(x, Tx)d_r(y, Ty)}{1 + d_r(x, y)}, \right. \\ \left. \frac{d_r(x, Tx)d_r(y, Ty)}{1 + d_r(Tx, Ty)}\right\} \\ = \max\left\{d_r(x, y), 0, 0, \frac{0 \cdot 0}{1 + 0}, \frac{0 \cdot 0}{1 + 0}\right\} = d_r(x, y).$$

Hence, (7) becomes

which is a contradiction. The proof of Theorem 9 is complete.

Remark 1. In the proof of the case 2 on Page 96, the authors used the fact that the rectangular metric d_r (see the condition (3.12)) is continuous, which is not given in the formulation of (Budhia et al, 2017), Theorem 3.2.

By putting in (1) instead of $M(x, y)$, one of the following sets

$$\{d_r(x, y)\}, \max\{d_r(x, y), d_r(x, Tx), d_r(y, Ty)\}, \\ \max\left\{\frac{d_r(x, Tx)d_r(y, Ty)}{1+d_r(x, y)}, \frac{d_r(x, Tx)d_r(y, Ty)}{1+d_r(Tx, Ty)}\right\},$$

immediately follows as a consequences of Theorem 9.

Corollary 4. Let (X, d_r) be a complete rectangular metric space and let $T: X \rightarrow X$ be a triangular α -admissible mapping. Assume that there exists a continuous function $\psi \# \Psi$ such that

$$x, y \in X, \alpha(x, y) \geq 1 \text{ implies } d_r(Tx, Ty) \leq \psi(d_r(x, y)). \quad (8)$$

Also, suppose that the following assertions hold:

1. there exists $x_0 \in X$ such that $\alpha(x_0, Tx_0) \geq 1$,
2. either T is continuous or (X, d_r) is α -regular.

Then T has a fixed point. Moreover, if for all $x, y \in F(T)$, we have $\alpha(x, y) \geq 1$, then the fixed point is unique.

Corollary 5. Let (X, d_r) be a complete rectangular metric space and let $T: X \rightarrow X$ be a triangular α -admissible mapping. Assume that there exists a continuous function $\psi \# \Psi$ such that for $x, y \# X$,

$$\alpha(x, y) \geq 1 \text{ yields } d_r(Tx, Ty) \leq \psi(\max\{d_r(x, y), d_r(x, Tx), d_r(y, Ty)\}). \quad (9)$$

Also, suppose that the following assertions hold:

1. there exists $x_0 \in X$ such that $\alpha(x_0, Tx_0) \geq 1$,
2. either T is continuous or (X, d_r) is α -regular

Then T has a fixed point. Moreover, if for all $x, y \in F(T)$, we have $\alpha(x, y) \geq 1$, then the fixed point is unique.

Corollary 6. Let (X, d_r) be a complete rectangular metric space and let $T: X \rightarrow X$ be a triangular α -admissible mapping. Assume that there exists a continuous function $\psi \# \Psi$ such that for $x, y \# X$, $\alpha(x, y) \geq 1$

$$\text{yields } d_r(Tx, Ty) \leq \psi\left(\max\left\{\frac{d_r(x, Tx)d_r(y, Ty)}{1+d_r(x, y)}, \frac{d_r(x, Tx)d_r(y, Ty)}{1+d_r(Tx, Ty)}\right\}\right). \quad (10)$$

Also, suppose that the following assertions hold:

1. there exists $x_0 \in X$ such that $\alpha(x_0, Tx_0) \geq 1$
2. either T is continuous or (X, d_r) is α -regular

Then T has a fixed point. Moreover, if for all $x, y \in F(T)$, we have $\alpha(x, y) \geq 1$, then the fixed point is unique.

In the book (Ćirić, 2003), Ćirić collected various contractive mappings in the usual metric spaces, see also (Rhoades, 1977) and (Collaco & Silva, 1997). The next three contractive conditions are well known in the existing literature:

- Ćirić 1: Ćirić's generalized contraction of first order: there exists $k_1 \in [0, 1)$ such that for all $x, y \in X$ holds:

$$d(Tx, Ty) \leq k_1 \max \left\{ d(x, y), \frac{d(x, Tx) + d(y, Ty)}{2}, \frac{d(x, Ty) + d(y, Tx)}{2} \right\}.$$

- Ćirić 2: Ćirić's generalized contraction of second order: there exists $k_2 \in [0, 1)$ such that for all $x, y \in X$ holds:

$$d(Tx, Ty) \leq k_2 \max \left\{ d(x, y), d(x, Tx), d(y, Ty), \frac{d(x, Ty) + d(y, Tx)}{2} \right\}. \quad (12)$$

In both cases, (X, d) is a metric space, $T : X \rightarrow X$ is a given selfmapping of the set X .

In (Ćirić, 2003), Ćirić introduced one of the most generalized contractive conditions (so-called quasicontraction) in the context of a metric space as follows:

- Ćirić 3: The self-mapping $T : X \rightarrow X$ on a metric space (X, d) is called a quasicontraction (in the sense of Ćirić) if there exists $k_3 \in [0, 1)$ such that for all $x, y \in X$ holds:

$$d(Tx, Ty) \leq k_3 \max \{d(x, y), d(x, Tx), d(y, Ty), d(x, Ty), d(y, Tx)\}. \quad (13)$$

Since,

$$\frac{d(x, Tx) + d(y, Ty)}{2} \leq \max \{d(x, Tx), d(y, Ty)\}$$

and

$$\frac{d(x, Ty) + d(y, Tx)}{2} \leq \max \{d(x, Ty), d(y, Tx)\}$$

it follows that (11) implies (12) and (12) implies (13). In (Ćirić, 2003), Ćirić proved the following result:
In (Ćirić, 2003), Ćirić proved the following result:

Theorem 3. *Each quasicontraction T on a complete metric space (X, d) has a unique fixed point (say) z . Moreover, for all $x \in X$, the sequence $\{T^n x\}_{n=0}^{+\infty}$, $T^0 x = x$ converges to the fixed point z as $n \rightarrow +\infty$.*

Now we can formulate the following notion and one open question:

Definition 3. *Let (X, d_r) be a rectangular metric space and let $\alpha : X \times X \rightarrow [0, +\infty)$ be a mapping. The mapping $T : X \rightarrow X$ is said to be a modified triangular α -admissible mapping if there exists a continuous function $\psi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ such that*

$$x, y \in X, \alpha(x, y) \geq 1 \text{ implies } d_r(Tx, Ty) \leq \psi(M(x, y)), \quad (14)$$

where $M(x, y)$ is one of the sets:

$$\begin{aligned} & \max \left\{ d_r(x, y), \frac{d_r(x, Tx) + d_r(y, Ty)}{2}, \frac{d_r(x, Ty) + d_r(y, Tx)}{2} \right\} \\ & \max \left\{ d_r(x, y), d_r(x, Tx), d_r(y, Ty), \frac{d_r(x, Ty) + d_r(y, Tx)}{2} \right\} \\ & \max \{ d_r(x, y), d_r(x, Tx), d_r(y, Ty), d_r(x, Ty), d_r(y, Tx) \}. \end{aligned}$$

AN OPEN PROBLEM

A suggestion for further research - it is logical natural to ask the following question:

Problem 0.1. *Let T be a modified triangular α -admissible mapping defined on a complete rectangular metric space (X, d_r) such that T is continuous or (X, d_r) is α -regular. Show that T has a fixed point.*

APPLICATIONS

In this section, we will focus on the applicability of the acquired obtained results.

An application to chemical sciences

Consider a diffusing substance placed in an absorbing medium between parallel walls such that δ_1, δ_2 are the stipulated concentrations at walls. Moreover, let $\Omega(r)$ be the given source density and $\Xi(r)$ be the known absorption coefficient. Then the concentration $\varkappa(r)$ of the substance under the aforementioned hypothesis governs the following boundary value problem

$$\begin{cases} -\varkappa'' + \Xi(r)\varkappa = \Omega(r) & ; r \in [0, 1] = I \\ \varkappa(0) = \delta_1, \varkappa(1) = \delta_2, \end{cases} \quad (1)$$

Problem (1) is equivalent to the succeeding integral equation

$$\varkappa(r) = \delta_1 + (\delta_2 - \delta_1)r + \int_0^1 \Theta(r, \varpi) (\Omega(\varpi) - \Xi(\varpi)\varkappa(\varpi)) d\varpi, \quad r \in [0, 1], \quad (2)$$

where $\Theta(r, \varpi) : [0, 1] \times \mathbb{R} \rightarrow \mathbb{R}$ is the Green's function which is continuous and is given by

$$\Theta(r, \varpi) = \begin{cases} r(1 - \varpi) & 0 \leq r \leq \varpi \leq 1, \\ \varpi(1 - r) & 0 \leq \varpi \leq r \leq 1. \end{cases} \quad (3)$$

Suppose that $C(I, \mathbb{R}) = X$ is the space of all real valued continuous functions defined on I and let X be endowed with the rectangular b-metric d_r defined by

$$d_r(\kappa, \kappa^*) = \|(\kappa - \kappa^*)\|,$$

where $\|\kappa\| = \sup\{|\kappa(r)| : r \in I\}$. Obviously (X, d_r) is a complete rectangular metric space.

Let the operator $\Xi : X \rightarrow X$ be defined by

$$\Xi \kappa(r) = \kappa(r) = \delta_1 + (\delta_2 - \delta_1)r + \int_0^1 \Theta(r, \varpi) (\Omega(\varpi) - \Xi(\varpi)\kappa(\varpi)) d\varpi.$$

Then κ^* is a unique solution of (2) if and only if it is a fixed point of Ξ . The subsequent Theorem is furnished for the assertion of the existence of a fixed point of Ξ .

Theorem 4. Consider problem (2) and suppose that there exists $\# > 0$ and a continuous function $\Xi(\varpi) : I \rightarrow \mathbb{R}$ such that the following assertion holds:

$$\alpha(\kappa(\varpi), \kappa^*(\varpi)) \geq 1 \implies 0 \leq |\Xi(\varpi)\kappa(\varpi) - \Xi(\varpi)\kappa^*(\varpi)| \leq \kappa^*(\varpi) - \kappa(\varpi).$$

Then the integral equation (2) and, consequently, the boundary value problem (1) governing the concentration of the diffusing substance has a unique solution in X .

Proof. Clearly, for $\kappa \in X$ and $r \in I$, the mapping $\Xi : X \rightarrow X$ is well defined. Also Ξ is triangular α -admissible.

$$\begin{aligned} & |\Xi \kappa(r) - \Xi \kappa^*(r)| \\ &= \left| \int_0^1 \Theta(r, \varpi) (\Omega(\varpi) - \Xi(\varpi)\kappa(\varpi)) d\varpi - h(r) \right. \\ &\quad \left. - \int_0^1 \Theta(r, \varpi) (\Omega(\varpi) - \Xi(\varpi)\kappa^*(\varpi)) d\varpi \right| \\ &\leq \int_0^1 \Theta(r, \varpi) |(\Omega(\varpi) - \Xi(\varpi)\kappa(\varpi)) - (\Omega(\varpi) - \Xi(\varpi)\kappa^*(\varpi))| d\varpi \\ &= \int_0^1 \Theta(r, \varpi) |\Xi(\varpi)\kappa(\varpi) - \Xi(\varpi)\kappa^*(\varpi)| d\varpi \\ &\leq \int_0^1 \Theta(r, \varpi) |\kappa(\varpi) - \kappa^*(\varpi)| d\varpi \\ &\leq \int_0^1 \Theta(r, \varpi) \|(\kappa(\varpi) - \kappa^*(\varpi))\| d\varpi \\ &\leq \|(\kappa - \kappa^*)\| \sup_{r \in [0,1]} \int_0^1 \Theta(r, \varpi) d\varpi. \end{aligned}$$

Since $\int_0^1 \Theta(r, \varpi) d\varpi = \frac{r-r^2}{2}$ and so $\sup_{r \in [0,1]} \int_0^1 \Theta(r, \varpi) d\varpi = \frac{1}{8}$.

Hence for all $\kappa, \kappa^* \in X$, we obtain

$$d_r(\Xi \kappa, \Xi \kappa^*) \leq \frac{d_r(\kappa, \kappa^*)}{8} \leq \frac{M(\kappa, \kappa^*)}{8},$$

where

$$M(\varkappa, \varkappa^*) = \max \{d_r(\varkappa, \varkappa^*), d_r(\varkappa, T\varkappa), d_r(\varkappa^*, T\varkappa^*), \\ \frac{d_r(\varkappa, T\varkappa) d_r(\varkappa^*, T\varkappa^*)}{1 + d_r(\varkappa, \varkappa^*)}, \frac{d_r(\varkappa, T\varkappa) d_r(\varkappa^*, T\varkappa^*)}{1 + d_r(T\varkappa, T\varkappa^*)}\}$$

Taking $\psi(M(\varkappa, \varkappa^*)) = \frac{1}{8}$, we obtain

$$d_r(T\varkappa, T\varkappa^*) \leq \psi(M(\varkappa, \varkappa^*))$$

Hence, all the hypotheses of Theorem 2 are contented. We conclude that Ξ has a unique fixed point $\varkappa$ in X , which guarantees that the integral equation (2) has a unique solution and, consequently, the boundary value problem (1) has a unique solution.

Application to a class of integral equations for an unknown function

We present the application of the existence of a fixed point for a generalized contraction to the following class of integral equations for an unknown function u :

$$u(t) = g(t) + \int_a^b \chi(t, z) f(z, u(z)) dz, \quad t \in [a, b], \quad (4)$$

where $f : [a, b] \times \mathbb{R} \rightarrow \mathbb{R}$, $K : [a, b] \times [a, b] \rightarrow [0, \infty)$, $g : [a, b] \rightarrow \mathbb{R}$ are the given continuous functions.

Let X be the set $C[a, b]$ of real continuous functions defined on $[a, b]$ and let $d_r : X \times X \rightarrow [0, \infty)$ be equipped with the metric defined by

$$d_r(u, v) = \sup_{a \leq t \leq b} |u(t) - v(t)|. \quad (5)$$

One can easily verify that (X, d_r) is a complete rectangular metric space. Let the self map $T : X \rightarrow X$ be defined by

$$Tu(t) = g(t) + \int_a^b \chi(t, z) f(z, u(z)) dz, \quad t \in [a, b], \quad (6)$$

then u is a fixed point of T if and only if it is a solution of (4). Also, we can easily check that T is triangular α -admissible. Now, we formulate the following subsequent theorem to show the existence of a solution of the underlying integral equation.

Theorem 5. Assume that the following assumptions hold:

(1) $\sup_{a \leq t \leq b} \int_a^b |\chi(t, z)| dz \leq \frac{1}{b-a}$;

2) Suppose that for all $x, y \in$,

$$\alpha(x(t), y(t)) \geq 1 \implies |f(z, x) - f(z, y)| \leq \frac{1}{2}|x(t) - y(t)|.$$

Then integral equation (4) has a solution.

Proof. Employing conditions (1) – (2) along with inequality (4), we have

$$\begin{aligned} d_r(Tu_1, Tu_2) &= \sup_{a \leq t \leq b} |Tu_1(t) - Tu_2(t)| \\ &= \sup_{a \leq t \leq b} \left| g(t) + \int_a^b \chi(t, z) f(z, u_1(z)) dz - \left(g(t) + \int_a^b \chi(t, z) f(z, u_2(z)) dz \right) \right| \\ &= \sup_{a \leq t \leq b} \left\{ \left| \int_a^b (\chi(t, z) f(z, u_1(z)) - \chi(t, z) f(z, u_2(z))) dz \right| \right\} \\ &\leq \sup_{a \leq t \leq b} \left\{ \int_a^b |\chi(t, z)| dz \cdot \int_a^b |f(z, u_1(z)) - f(z, u_2(z))| dz \right\} \\ &= \left\{ \sup_{a \leq t \leq b} \int_a^b |\chi(t, z)| dz \right\} \cdot \left\{ \int_a^b |f(z, u_1(z)) - f(z, u_2(z))| dz \right\} \\ &= \left\{ \sup_{a \leq t \leq b} \int_a^b |\chi(t, z)| dz \right\} \cdot \left\{ \int_a^b |f(z, u_1(z)) - f(z, u_2(z))| dz \right\} \\ &\leq \left\{ \frac{1}{b-a} \right\} \cdot \left\{ \frac{1}{2} \int_a^b |u_1(z) - u_2(z)| dz \right\} \\ &\leq \frac{1}{2(b-a)} \int_a^b \sup_{a \leq t \leq b} |u_1(t) - u_2(t)| dz \\ &= \frac{1}{2} \sup_{a \leq t \leq b} |u_1(t) - u_2(t)| \end{aligned}$$

i.e. $d_r(Tu_1, Tu_2) = \frac{1}{2} (d_r(u_1, u_2)) \leq \frac{M(u_1, u_2)}{2}$. Which amounts to say that

$$d_r(Tu_1, Tu_2) \leq \frac{M(u_1, u_2)}{2},$$

where

$$M(u_1, u_2) = \max \{ d_r(u_1, u_2), d_r(u_1, Tu_1), d_r(u_2, Tu_2),$$

$$\frac{d_r(u_1, Tu_1) d_r(u_2, Tu_2)}{1 + d_r(u_1, u_2)}, \frac{d_r(u_1, Tu_1) d_r(u_2, Tu_2)}{1 + d_r(Tu_1, Tu_2)} \}.$$

Taking $\psi(M(u_1, u_2)) = 1/2$, the above inequality turns into

$$d_r(Tu_1, Tu_2) \leq \psi(M(u_1, u_2))$$

Thus, all the hypotheses Theorem 2 are satisfied and we conclude that T has a unique fixed point x^* in X , which amounts to say that integral equation (4) has a unique solution which belongs to $X = C[a, b]$.

REFERENCES

- Alsulami, H.H., Chandok, S., Taoudi, M-A. & Erhan, I.M. 2015. Some fixed point theorems for (α, ψ) -rational type contractive mappings. *Fixed Point Theory and Applications*, 2015(art.number:97). Available at: <https://doi.org/10.1186/s13663-015-0332-3>.
- Banach, S. 1922. Sur les opérations dans les ensembles abstraits et leur applications aux équations intégrales. *Fundamenta Mathematicae*, 3, pp.133-181 (in French). Available at: <https://doi.org/10.4064/fm-3-1-133-181>.
- Budhia, L., Kir, M., Gopal, D. & Kiziltunc, H. 2017. New fixed point results in rectangular metric space and application to fractional calculus. *Tbilisi Mathematical Journal*, 10(1), pp.91-104. Available at: <https://doi.org/10.1515/tmj-2017-0006>.
- Collaco, P. & Silva, J.C.E. 1997. A complete comparison of 25 contraction conditions. *Nonlinear Analysis: Theory, Methods & Applications*, 30(1), pp.471-476. Available at: [https://doi.org/10.1016/S0362-546X\(97\)00353-2](https://doi.org/10.1016/S0362-546X(97)00353-2).
- Ćirić, Lj. 2003. *Some Recent Results in Metrical Fixed Point Theory*. Belgrade: University of Belgrade.
- Kadelburg, Z. & Radenović, S. 2014a. Fixed point results in generalized metric spaces without Hausdorff property. *Mathematical Sciences*, 8(art.number:125). Available at: <https://doi.org/10.1007/s40096-014-0125-6>.
- Kadelburg, Z. & Radenović, S. 2014b. On generalized metric spaces: A survey. *TWMS Journal of Pure and Applied Mathematics*, 5(1), pp.3-13 [online]. Available at: <http://www.twmsj.az/Files/Contents%20V.5,%20%20N.1,%20%202014/pp.3-13.pdf> [Accessed: 15 November 2020].
- Karapinar, E. 2014. Discussion on α - ψ Contractions on Generalized Metric Spaces. *Abstract and Applied Analysis*, 2014(art.ID:962784). Available at: <https://doi.org/10.1155/2014/962784>.
- Karapinar, E., Kumam, P. & Salimi, P. 2013. On α - ψ -Meir-Keeler contractive mappings. *Fixed Point Theory and Applications*, art.number:94. Available at: <https://doi.org/10.1186/1687-1812-2013-94>.
- Kirk, W.A. & Shahzad, N. 2014. *Fixed Point Theory in Distance Spaces*. Springer International Publishing Switzerland. ISBN: 978-3-319-10927-5.
- Mohammadi, B. & Rezapour, Sh. 2013. On modified α - ψ -contractions. *Journal of Advanced Mathematical Studies*, 6(2), pp.162-166 [online].
- Radenović, S., Kadelburg, Z., Jandrlić, D. & Jandrlić, A. 2012. Some results on weakly contractive maps. *Bulletin of the Iranian Mathematical Society*, 38(3), pp.625-645 [online]. Available at: http://bims.iranjournals.ir/article_229.html [Accessed: 15 March 2018].
- Rhoades, B.E. 1977. A comparison of various definitions of contractive mappings. *Trans. Amer. Math. Soc.*, 226, pp.257-290. Available at: <https://doi.org/10.1090/S0002-9947-1977-0433430-4>.
- Samet, B., Vetro, C. & Vetro, P. 2012. Fixed point theorem for α - ψ -contractive type mappings. *Nonlinear Analysis: Theory, Methods & Applications*, 75(4), pp.2154-2165. Available at: <https://doi.org/10.1016/j.na.2011.10.014>.
- Salimi, P., Latif, A. & Hussain, N. 2013. Modified α - ψ -contractive mappings with applications. *Fixed Point Theory and Applications*, art.number:151. Available at: <https://doi.org/10.1186/1687-1812-2013-151>.

ADDITIONAL INFORMATION

FIELD: Mathematics

ARTICLE TYPE: Original scientific paper

ALTERNATIVE LINK

<https://scindeks.ceon.rs/article.aspx?artid=0042-84692101008Y> (html)

<https://scindeks-clanci.ceon.rs/data/pdf/0042-8469/2021/0042-84692101008Y.pdf> (pdf)

<https://www.elibrary.ru/item.asp?id=44584147> (pdf)

<http://www.vtg.mod.gov.rs/archive/2021/military-technical-courier-1-2021.pdf> (pdf)