



Darwiniana, nueva serie

ISSN: 0011-6793

ISSN: 1850-1702

Instituto de Botánica Darwinion & Museo Botánico de Córdoba

González, Favio; Valenzuela, Luis; Pabón-Mora, Natalia
ARISTOLOCHIA BRACHYLIMBA (ARISTOLOCHIACEAE; PIPERALES), A NEW SPECIES
FROM THE PERUVIAN AMAZONIA WITH AN UNUSUALLY SHORT PERIANTH LIMB

Darwiniana, nueva serie, vol. 11, no. 1, 2023, pp. 347-356
Instituto de Botánica Darwinion & Museo Botánico de Córdoba

DOI: <https://doi.org/10.14522/darwiniana.2023.111.1139>

Available in: <https://www.redalyc.org/articulo.oa?id=66976167015>

- ▶ How to cite
- ▶ Complete issue
- ▶ More information about this article
- ▶ Journal's webpage in redalyc.org

redalyc.org

Scientific Information System Redalyc

Network of Scientific Journals from Latin America and the Caribbean, Spain and Portugal

Project academic non-profit, developed under the open access initiative

ARISTOLOCHIA BRACHYLIMBA (ARISTOLOCHIACEAE; PIPERALES), A NEW SPECIES FROM THE PERUVIAN AMAZONIA WITH AN UNUSUALLY SHORT PERIANTH LIMB

Favio González¹ , Luis Valenzuela²  & Natalia Pabón-Mora³ 

¹ Instituto de Ciencias Naturales, Facultad de Ciencias, Universidad Nacional de Colombia, Sede Bogotá, AA 7495, Bogotá, Colombia.

² Jardín Botánico de Missouri - Perú - Herbario Selva Central Oxapampa (HOXA), 19231 Oxapampa, Perú.

³ Instituto de Biología, Facultad de Ciencias Exactas y Naturales, Universidad de Antioquia, Medellín, ZIP Code 050010, Colombia; lucia.pabon@udea.edu.co (author for correspondence).

Abstract. González, F.; L. Valenzuela & N. Pabón-Mora. 2023. *Aristolochia brachylimba* (Aristolochiaceae; Piperales), a new species from the Peruvian Amazonia with an unusually short perianth limb. *Darwiniana*, nueva serie 11(1): 347-356.

Aristolochia brachylimba, a new species from Amazonian forests of Peru, is described, illustrated, and discussed with respect to its most similar species, *A. stomachoidis* Hoehne, from northern, central and eastern Brazil (Goiás, Maranhão, Mato Grosso, Mato Grosso do Sul, Pará and Tocantins), and northeastern Argentina (Misiones). The new species differs from its relatives mainly by the size of the utricle, which is considerably larger than the short perianth limb, the conical shape of the tube (i.e. gradually narrowing down towards its distal end), and the winged, flattened seeds. So far, the new species has been found in Amazonian forests of two disjunct localities in Ataz (Oxapampa, Pasco) and the Cordillera Azul National Park (Ucayali, Loreto), at elevations between 340 and 479 m a.s.l.

Keywords. Hexandrous neotropical *Aristolochia*; Pasco flora; Peruvian endemics; Piperales; Ucayali flora.

Resumen. González, F.; L. Valenzuela & N. Pabón-Mora. 2023. *Aristolochia brachylimba* (Aristolochiaceae; Piperales), una nueva especie de la Amazonia peruana con el limbo del perianto inusualmente corto. *Darwiniana*, nueva serie 11(1): 347-356.

Aristolochia brachylimba, especie nueva de bosques amazónicos del Perú, es descrita e ilustrada, y son discutidas sus afinidades con respecto a su especie más similar, *A. stomachoidis* Hoehne, del norte, centro y este de Brasil (Goiás, Maranhão, Mato Grosso, Mato Grosso do Sul, Pará y Tocantins) y el noreste de Argentina (Misiones). La nueva especie difiere principalmente por el tamaño del utrículo, que es considerablemente más grande que el limbo del perianto, la forma cónica del tubo (es decir, que se estrecha gradualmente hacia su extremo distal), y las semillas aladas y aplanadas. Hasta ahora, la nueva especie ha sido recolectada en bosques amazónicos de dos localidades disjuntas en Ataz (Oxapampa, Pasco) y el Parque Nacional Cordillera Azul (Ucayali, Loreto), entre 340 y 479 m de altura sobre el nivel del mar.

Palabras clave. *Aristolochia* hexandras neotropicales; especies endémicas del Perú; Flora de Pasco; Flora de Ucayali; Piperales.

INTRODUCTION

Aristolochia L., the largest genus of the family Aristolochiaceae, comprises nearly 560 species mostly diversified in the tropics and subtropics

of the Americas (González & Pabón-Mora, 2017; Freitas et al., 2020a). All members of the genus possess alternate, distichous leaves, a perianth formed by the early fusion of three sepals, and an inferior ovary forming capsular fruits.

The highly elaborate petaloid perianth is often differentiated into a proximal, inflated portion called the utricle, a narrow middle portion called the tube, and an expanded, laminar portion called the limb (Pfeifer, 1966; González, 1994). In most species of *Aristolochia* the utricle is considerably shorter than the limb, and the tube is funnel-shaped as it gradually widens towards its distal end. The perianth morphology, epidermal elaboration, and vascular patterning provide key traits to reconstruct the phylogenetic relationships within the genus (González & Stevenson, 2000a), as well as to discover new species (e.g. Freitas et al., 2020b).

All South American species of *Aristolochia* belong to the subgenus *Aristolochia* section *Gymnolobus* Duch. subseries *Hexandrae*, as the gynostemium is formed by six stamens fused to six stigmatic lobes that lack a transverse roof-like process above the stamens (Duchartre, 1854a, 1864; González, 1991a; González & Stevenson, 2000b). With nearly 140 species, *Aristolochia* subseries *Hexandrae* is one of the the most species-rich infrageneric groups worldwide. The series has been divided in two informal groups, namely 'Exstipulosae' and 'Pseudostipulosae', with ca. 70 species each (Hoehne, 1942), based on the presence or absence of pseudostipules, respectively. The *Aristolochia* pseudostipules correspond to reduced prophylls of axillary branches that resemble stipules (for details see Duchartre, 1854b; Hoehne, 1942; González, 1990). While working on the monograph of *Aristolochia* for Flora Neotropica, three specimens collected in Pasco and Ucayali (Peru) called our attention as it does not match any of the previously described species of the genus in Peru or neighboring countries (cf. Masters, 1875; Malme, 1904; Ule, 1905; Moore, 1915; Hoehne, 1927, 1942; Schmidt, 1927; Macbride, 1937; Ahumada, 1977, 2010; González, 1990, 1991b, 1994, 1998, 2000; Brako & Zarucchi, 1993; León & La Torre, 2006; González et al., 2015; Freitas et al., 2020b). Notably, the utricle in this specimen is considerably longer than the extremely short limb, the tube diameter gradually narrows down towards its distal portion resulting in a conical contour of the tube, and the seeds are winged and flattened.

The relative size of the utricle *versus* the limb and the atypical tube contour were consistently found at various stages from early preanthesis to anthesis (Fig. 1A). The presence of all these uncommon traits simultaneously strongly indicates that it corresponds to an undescribed taxon. In addition to the description of the new species, remarks on the conservation status and the geographical distribution are provided, along with a key to identify all the *Exstipulosae* species of *Aristolochia* subser. *Hexandrae*.

MATERIALS AND METHODS

The type specimens of the new species are deposited at HOXA (Herbario Selva Central Oxapampa, Pasco, Peru), MO (Missouri Botanical Garden herbarium, USA), and USM (Herbario Universidad Nacional Mayor de San Marcos, Lima, Peru). Comparison with similar species was based on extensive herbarium search at B, BM, COL, CONN, CTES, F, GH, K, MO, NY, and US (acronyms follow Thiers, 2023). Additionally, the Jstor Global Plants database (JSTOR, 2021) was consulted, and the available barcode for original material was included. Measurements of leaves, flowers, fruits and seeds were taken from fully formed organs. Terminology used to describe *Aristolochia* flowers follows Pfeifer (1966), and González (1990, 1994).

The traits of *Aristolochia stomachoidis* Hoehne, the most similar species to the taxon here described (Table 1), were taken from the descriptions provided by Hoehne (1910, 1927, 1942), Keller (2010) and González et al. (2015), and from direct examination of the following specimens: ARGENTINA. **Misiones.** Depto. San Ignacio, Club del Río, 27-VII-2008, *H. A. Keller 5675* (CTES). BRAZIL. **Goiás:** Queixada, Yatai, 11-XI-1948 (fl), *A. Macedo 1470* (BM, MO); Aragarças, 15°51'S, 52°15'W, km 51 of Aragarças, 14-I-1968 (fl), *D. Philcox & E. Ferreira 4050* (K). **Maranhão:** Barra do Corda, Cocal Grande, 34 km NE of Barra do Corda along the rio Mearim, 5°24'S, 45°06'W, 7-III-1983 (fl), *G. E. Schatz et al. 889* (COL, NY). **Mato Grosso:** Corrego do Porco, rio Suiazinho, c. 24 km along the Suiá-miçu road, 24-X-1968 (fl),




Fig. 1. *Aristolochia brachylimba* F. González, Pabón-Mora & L. Valenz., sp. nov. (from Valenzuela et al. 12765, HOXA, holotype). **A**, habit, leaves and flowers. **B**, capsule at dehiscence. Scale bars: 1 cm.

Table 1. Differential traits between *Aristolochia brachylimba* and its closely related species *A. stomachoidis*.

	<i>A. brachylimba</i>	<i>A. stomachoidis</i>
Petiole length	1.8-6 cm	5-7 cm
Leaf blade size	6-12 × 3.5-6.4 cm	10-17 × 6-9 cm
Leaf blade base	lobes not overlapping, forming sinuses 1-1.8 cm deep	lobes frequently overlapping, forming sinuses 1-3.5 cm deep
Leaf blade apex	acute to shortly acuminate	obtuse to acute, rarely acuminate
Peduncle plus ovary length	3.5-6 cm	6-12 cm
Utricle plus tube shape	conspicuously distinct; utricle widely obovoid, 1.4-2 cm long	indistinct, together forming a widely obovoid ventricose structure ('U' shaped in lateral view), 3.3-6 cm long
Limb shape and size	broadly ovate, 1-1.9 × 0.6-1.2 cm, forming an angle of 100-120° with respect to the tube, base cordate, apex acuminate	ovate to narrowly ovate, 1.5-4 × 0.8-1.4 cm, forming an angle of 180° with respect to the tube, base constricted and emarginate, apex acute
Capsule length	3-5 cm long, ending in a rostrum 3-5 mm long	5-6 cm long, ending in a rostrum 2-3 mm long
Geographical distribution	Peru (Oxapampa and Ucayali)	Brazil (Goiás, Maranhão, Mato Grosso, Mato Grosso do Sul, Pará and Tocantins), and Argentina (Misiones).

R. M. Harley et al. 10834 (K); ao longo dos rios Taruman e Sepotuba, perto de Tapirapoan, Mar 1909 (fl), *F. C. Hoehne 1443* (lectotype SP[000413]!); vicinity of Barra do Garças, roadside thicket ca. 50 km N of Barra do Garças on road to Xavantina, 15°53'S, 52°15'W, 300-400 m, 14-X-1964 (fl), *H. S. Irwin & T. R. Soderstrom 6830* (MO, NY); Barra do Garças-Xavantina, (fl), *E. Onishi & S. G. Fonseca 332/1101* (K); Serra do Roncador, road Chavantina to Barra do Garças, 55 km N of Barra do Garças, 15-X-1964 (fl), *G. T. Prance & N. T. Silva 59413* (COL, CONN, NY); ½ km N of the rio Suiá-miçu Ferry, appr. 290 km N of Xavantina, 11-V-1968 (fl, fr), *J. A. Ratter et al. 1393* (K); rio Suaizinha, appr. 280 km N of Xavantina, 11-V-1968 (fl), *J. A. Ratter et al. 1395* (K). **Mato Grosso do Sul:** Bataguassu, Porto XV, 14-II-1970 (fl), *G. Hatschbach 23524* (CTES). **Pará:** Conceição do Araguaia, Alacilândia, 42 km W of Conceição do Araguaia along highway PA-287 at rio Arraias do Araguaia, ca. 8°13'S, 49°36'W, 300 m, 23-II-1980 (fl), *T. Plowman et al. 9029* (COL); Sete Varas airstrip on Rio Curuá, 54°92'W, 0°95'S, 5-VIII-1981 (fl), *J. J. Strudwick & G. L. Sobel 4194* (COL, NY). **Tocantins:** Rod. Transamazonica, Araguatins, 27-III-1976 (fl), *G. Hatschbach 38421* (B, MO, NY).

RESULTS

Aristolochia brachylimba F. González, Pabón-Mora & L. Valenz., **sp. nov.** TYPE: Perú, Pasco, Oxapampa, Palcazú, Ataz, Yanesha Reserva Camunal, 10°10'12''S, 75°18'08''W, 340 m, 21-V-2009 (fl, fr), *L. Valenzuela, A. Peña & J. L. Mateo 12765* (holotype HOXA!; isotypes, MO!). Figs. 1-3.

Diagnosis. *Aristolochia brachylimba* resembles *A. stomachoidis*, from which it differs mainly by the petiole 1.8-6 cm long, the leaf blade 6-12 × 3.5-6.4 cm, the peduncle plus ovary 3.5-6 cm long, the utricle 1.4-2 cm long, distinct from the tube, and the limb broadly ovate, 1-1.9 × 0.6-1.2 cm, forming an angle of 100-120° with respect to the tube, with cordate base and acuminate apex (*versus* the petiole 5-7 cm long, the leaf blade 10-17 × 6-9 cm, the peduncle plus ovary 6-12 cm, the utricle indistinct from the tube forming an obovoid ventricose structure ('U' shaped in lateral view), 3.3-6 cm long, and the limb ovate to narrowly ovate, 1.5-4 × 0.8-1.4 cm, at an angle of 180° with respect to the tube, with constricted and emarginate base and acute apex in *A. stomachoidis*). The two species have allopatric distribution.




Fig. 2. *Aristolochia brachylimba*, photographs of type specimens (A–J, *L. Valenzuela et al.* 39374, paratype; K, *L. Valenzuela et al.* 39378, paratype; L, M, *L. Valenzuela et al.* 12765, holotype). **A**, leaf. **B**, flower at anthesis. **C**, perianth, lateral view; note the narrowing of the tube distal portion (arrowhead). **D**, **E**, limb, frontal view. **F**, detail of inner limb surface. **G**, detail of perianth throat. **H**, dissected utricle and gynostemium. **I**, detail of syrinx. **J**, gynostemium, sagittal section. **K**, twig with capsule prior dehiscence. **L**, **M**, deids, adaxial (L) and abaxial (M) view. Abbreviations: **l**, limb; **o**, ovary; **t**, tube; **u**, utricle. Scale bars: 2 cm in A; 5 mm in B–E; 1 mm in F–J, L, M; 1 cm in K. Color version at <https://www.ojs.darwin.edu.ar/index.php/darwiniana/article/view/1139/1305>

Vines. Stems glabrescent, cylindrical, slender, with internodes 6-11 cm long. Petiole 1.8-6 cm long, glabrescent; blade oblong to narrowly ovate, 6-12 × 3.5-6.4 cm, membranous, adaxial surface glabrous, abaxial surface glabrescent, base cordate, not peltate, lobes parallel to convergent leaving narrow sinuses 1-1.8 cm deep, apex acute to shortly acuminate, basal primary veins 3(5). Pseudostipules absent. Flowers axillary, solitary, ebracteolate. Peduncle plus ovary 3.5-6 cm long, glabrous, with a short (to 0.5 mm) rostrum at the dorsal portion of the ovary tip. Perianth glabrous on its outer surface, reflexed, vinaceous to purple; utricle widely obovoid, 1.4-2 × 0.4-1 cm, syrinx present, aequilateral, ca. 1 mm broad; tube reflexed forming an angle of 80-90° with respect to the utricle, 0.8-1.4 cm long, 5.5-6.5 mm proximal diameter, 3.5-4.5 mm distal diameter, gradually narrowed down towards its distal portion resulting in a conical contour; limb broadly ovate, 1-1.9 cm long (including an acumen 1.5-3 mm), 0.6-1.2 cm wide, forming an angle of 100-120° with respect to the tube, warty to shortly fimbriate on its inner surface, base cordate, apex acuminate; gynostemium 6-lobed, sessile, 4-5 mm long; anthers six, equidistant, oblong, 2.5-3 mm long. Capsule broadly cylindrical, 3-5 cm long including an apical rostrum 3-5 mm long, 1.5-1.8 cm in diameter at its mid-level, basipetally dehiscent, glabrous and smooth externally, midvein of each carpel prominent, ca. 1 mm, septae entire. Seeds broadly ovoid, 8-9 × 6.5-7.5 mm including a membranous, peripheral wing, flattened, warty on both surfaces, seed proper cordiform, 4.0-4.3 × 3.9-4.1, brown, darker than the wing, raphe prominulous.

Distribution, phenology and habitat. *Aristolochia brachylimba* has been collected in two disjunct localities in Ataz (Oxapampa, Pasco region), and the Cordillera Azul National Park (Ucayali province, Loreto region)(Fig. 3), at elevations between 340 and 479 m a.s.l. The new species sets flowers and fruits in May.

This new species belongs to the Exstipulosae group (sensu Hoehne, 1942), and possesses all diagnostic traits of *Aristolochia* subser. Hexandrae (sensu González, 1990, 1991a), including the lack of an abscission zone at the base of the petiole, and the solitary, axillary, ebracteolate flowers.




Fig. 3. Map of distribution of *Aristolochia brachylimba* (diamonds) and its closely related species *A. stomachoidis* (circles).

Five additional Exstipulosae species of *Aristolochia* subser. Hexandrae occur in Peru, namely, *A. barbouri* Barringer (Fig. 4A), endemic to the Department of Madre de Dios; *A. chachapoyensis* Ahumada (Fig. 4B), from the departments of Amazonas, Cajamarca and San Martín; *A. pilosa* Kunth (Fig. 4C), widespread from Central America to Bolivia; *A. trilabiata* Glaziou (Fig. 4D, formerly *A. didyma* S. Moore; see González & Pabón-Mora, 2018), widely distributed in Panama, Colombia, Ecuador, Venezuela, Peru, Guyanas, Brazil and Bolivia; and *A. xerophytica* R.E. Schult. (Fig. 4E), from the departments of Cajamarca, La Libertad and Piura. All these species possess a perianth with the tube funnel-shaped, and the utricle shorter than the limb, sharply contrasting to the perianth with the obconic tube and the utricle longer than the limb in the new species. A key to identify these species is provided below.

Etymology. The epithet refers to the unusually short perianth limb that characterizes the new species.

Provisional conservation status. According to the IUCN Red List criteria (IUCN, 2014, 2022), *Aristolochia brachylimba* should be provisionally considered as Endangered (EN B1ab(iii) + 2ab(iii), D), due to population reduction based




Fig. 4. Outline drawings of leaf and perianth (side view) of the species of *Aristolochia* subser. Hexandrae from Peru (A-E), and its close relative *A. stomachoidis* (F), from Brazil. **A.**, *A. barbouri* (from *Barbour 5405*, holotype F[F0042307F!]). **B.**, *A. chachapoyensis* (from *Hutchison & Wright 5447*, holotype UC[UC1335070!]). **C.**, *A. pilosa* (from *Humboldt & Bonpland s.n.*, holotype P [P00669996!]). **D.**, *A. trilabiata* (from *Glaziou 10033*, lectotype P [P01657954!]). **E.**, *A. xerophytica* (from *Evinger 540*, holotype US[US00610968!]). **F.**, *A. stomachoidis* (from *Hoehne 1443*, lectotype SP [SP000413!]). Note in A-E the funnel-shaped tube and the shorter utricle with respect to the limb. Abbreviations: l, limb; t, tube; u, utricle. Scale bars: 1 cm.

on the number of mature individuals known (<1000), small extent of occurrence, small area of occupancy with just one station of occurrence, and habitat reduction in the area.

Notes. The narrowly ovate to oblong leaves, the perianth utricle larger than the limb, the tube gradually narrowing down towards its distal portion, and the capsules and seeds of *Aristolochia brachylimba* resemble those of *A. stomachoidis* (Figs. 3, 4F), from northern, central and southeastern Brazil (Goiás, Maranhão, Mato Grosso, Mato Grosso do Sul, Pará and Tocantins),

and northeastern Argentina (Misiones). Apart from the allopatric distribution of these two species, the new species differs in the petiole 1.8-6 cm long; the leaf blade 6-12 × 3.5-6.4 cm, with two non-overlapping lobes forming sinuses 1-1.8 cm deep at its base, and an acute to shortly acuminate apex; the peduncle plus ovary 3.5-6 cm long; the utricle widely obovoid, 1.4-2 cm long, distinct from the tube; the limb broadly ovate, 1-1.9 × 0.6-1.2 cm, forming an angle of 100-120° with the tube, cordate at its base and acuminate at its apex; and the capsule 3-5 cm long, ending in a rostrum 3-5 mm long in *A. brachylimba* (Figs. 1, 2).

Rather, *A. stomachoides* has longer (5-7 cm) petioles, larger leaf blades (10-17 × 6-9 cm) with two lobes overlapping forming sinuses 1-3.5 cm deep at its base, and an obtuse to acute (rarely shortly acuminate) apex; peduncle plus ovary 6-12 cm; utricle indistinct from the tube, together forming a widely obovoid ventricose structure ('U' shaped in lateral view), 3.3-6 cm long; limb ovate to narrowly ovate, 1.5-4 × 0.8-1.4 cm, forming an angle of 180° with respect to the tube, basally constricted and emarginate and apically acute; and capsules 5-6 cm long, ending in a rostrum 2-3 mm long (Fig. 4F, Table 1).

Additional specimens studied (Paratypes)

PERU. **Loreto Region.** Ucayali Province, Pampa Hermosa District, Cordillera Azul National Park, Shanshuico Surveillance Post, Amazon forest, silt-clay soil, 07°21'22.2''S, 076°00'27.6''W, 479 m, 11-V-11, 2021 (fl), *L. Valenzuela A. García, R. García & J. Salazar 39374* (HOXA, MO, USM); Ucayali Province, Pampa Hermosa District, Cordillera Azul National Park, Shanshuico Surveillance Post, primary Amazon Forest, silt-clay soil, 07°21'22.2''S, 076°00'27.6''W, 479 m, 11-V-2021 (fr), *L. Valenzuela, A. García, R. García & J. Salazar 39378* (HOXA, MO, USM).

Key to the Exstipulosae species of *Aristolochia* subser. Hexandrae from Peru

1. Perianth tube conical (i.e. gradually narrowing down towards its distal end), and limb shorter than the utricle. Seeds flattened, winged *A. brachylimba*
1. Perianth tube funnel-shaped, and limb 2 to 5 times longer than the utricle (if shorter, then the limb trilobed in *A. trilabiata*). Seeds concave-convex, not winged 2
- 2(1). Stems, leaves, flowers and fruits densely hirsute-pilose, trichomes septate *A. pilosa* Kunth
2. Stems, leaves, flowers and fruits glabrous to puberulous but never hirsute-pilose nor with septate trichomes 3
- 3(2). Petiole 5-30 cm long, glabrous; leaf blade ovate to broadly ovate, 9-29 × 10-28 cm, almost as long as wide. Floral peduncle and ovary 11-21 cm long, glabrous. Perianth strongly reflexed between the utricle and the tube, these forming an angle of ca. 30°; utricle 46-75 × 22-40 mm; tube 2.5-8 cm long; limb trilobed, the upper two lips formed at the flanks of the dorsal midvein, each 3.5-6.8 × 2-3.5 cm, separated by a sinus 1-3.5 cm depth, with marginal fimbriae to 6 mm long, the lower lip median, widely ovate, 1-2 × 1.4-2.5 cm; gynostemium 10-15 × 6-10 mm. Capsule very narrowly cylindrical, 10-25 × 1.5-2.5 cm; seeds with a sticky aril around the raphe. Widespread in rainforests of Panamá, Colombia, Ecuador, Venezuela, Perú, Guyanas, Brazil, and Bolivia *A. trilabiata* Glaziou
3. Petiole 0.8-3.5 cm long; leaf blade narrowly to very narrowly ovate (rarely ovate), 3-16 × 1.4-7 cm, 2 to 7 times longer than wide. Floral peduncle and ovary 1.8-5 cm long, puberulous. Perianth very slightly arched between the utricle and the tube, these forming angles ≥ 110°; utricle 5-14 × 4-8 mm; tube 8-26 mm long; limb unilobed, narrowly ovate, ovate or spatulate, 2-6.5 × 0.4-1.6 cm, efimbriate or, in *A. xerophytica*, with fimbriae less than 3 mm long; gynostemium 2.5-5 × 3.5-4.5 mm. Capsule cylindrical, 1.5-4 × 1-1.8 cm; seeds concave-convex, exarillate. Endemic to Peru 4
- 4(3). Leaf blade very narrowly ovate, 10-16 × 1.4-2 cm, 5 to 7 times longer than wide, not constricted at its mid-portion, with basal lobes parallel to convergent; primary basal veins three. Gynostemium 4-5 mm long. Capsule 3.5-4 cm long, including a rostrum 6-8 mm long. Endemic to Madre de Dios (Peru), in Amazonian rainforests at elevations below 300 m a.s.l. *A. barbouri* Barringer
4. Leaf blade narrowly ovate, 3.5-11.4 × 2-7 cm, 2 to 4 times longer than wide, often constricted at its mid-portion, with basal lobes divergent, rarely parallel; primary basal veins five. Gynostemium 2.5-4 mm long. Capsule 2.5-3.2 cm long, including a rostrum 1-5 mm long. Dry thickets of Amazonas, Cajamarca, La Libertad, Piura and San Martín (Peru), at elevations between 700 and 1500 m a.s.l. 5
- 5(4). Stems, petiole, and floral peduncle glabrescent. Limb of the perianth narrowly to very narrowly ovate, not spatulate, 4-6.5 cm long, slightly cuculate, forming a wide angle of 110-120° with respect to the tube, efimbriate, apex acuminate *A. chachapoyensis* Ahumada
5. Stems, petiole, and floral peduncle puberulous. Limb of the perianth ovate to slightly spatulate, 2-4 cm long, not cuculate, forming a sharp angle of 90-120° with respect to the tube, fimbriate on its margins and its inner surface, apex acute to obtuse, sometimes mucronulate *A. xerophytica* R.E. Schult.

ACKNOWLEDGEMENTS

We thank Dr. Kanchi Gandhi (Harvard University) for helping with the etymology of the new species, and all the staff of the Harvard University Herbaria and Library for logistic support during a short term visit of two of the authors (FG and NLPM) to this Institution. We also thank the curators of the herbaria B, BM, CONN, CTES, F, GH, HOXA, K, MO, NY, US, and USM, for making available the specimens examined. The first author also thanks the Faculty of Sciences, Universidad Nacional de Colombia, for granting a leave of absence to advance the monograph of the family Aristolochiaceae for Flora Neotropica. The last author thanks the Fulbright 'Beca Investigador Visitante Colombiano 2022' for a short term scholarship at Harvard University. The authors also want to thank the Central Selva HOXA Herbarium and the Missouri Botanical Garden program in Peru, for allowing us to review the type collections.

BIBLIOGRAPHY

- Ahumada, L. Z. 1977. Novedades sistemáticas en el género *Aristolochia* (Aristolochiaceae) em Sudamérica. I. *Darwiniana* 21: 65-80.
- Ahumada, L. Z. 2010. Aristolochiaceae. In: L. Ramella & R. Perret (eds.), Flora of Paraguay, vol. 41. Conservatoire et Jardin Botanique de la Ville de Genève, Switzerland.
- Brako, L. & J. L. Zarucchi. 1993. Catalogue of the Flowering Plants and Gymnosperms of Perú. *Monographs in Systematic Botany from the Missouri Botanical Garden* 5: i-xl, 1-1286.
- Duchartre, P. 1854a. Tentamen methodicae divisionis generis *Aristolochia*. *Annales des Sciences Naturelles, series 4*, 2: 29-76.
- Duchartre, P. 1854b. Sur les prétendues stipules des Aristoloches. *Bulletin de la Société Botanique de France* 1: 56-60.
- Duchartre, P. 1864. Aristolochiaceae. In: A. P. De Candolle (ed.), *Prodromus systematis naturalis regni vegetabilis* 15: 421-498.
- Freitas, J.; E. J. Lirio, F. Barros & F. González. 2020a. Aristolochiaceae. In: Flora do Brasil 2020. Jardim Botânico do Rio de Janeiro. Available at: <https://floradobrasil.jbrj.gov.br/reflora/floradobrasil/FB54>
- Freitas, J.; F. González, O. Poncy, C. Feuillet & A. Alves-Araújo. 2020b. Floral geometric morphometrics unveils a new cauliflorous species of *Aristolochia* (Aristolochiaceae) from the Guiana Shield. *Phytotaxa* 474: 1-14.
- González, F. 1990. Flora de Colombia. Aristolochiaceae. Monografía no. 12. Universidad Nacional de Colombia, Instituto de Ciencias Naturales.
- González, F. 1991a. Notes on the systematics of *Aristolochia* subsect. *Hexandrae*. *Annals of the Missouri Botanical Garden* 78: 497-503.
- González, F. 1991b. Acerca de la identidad de *Aristolochia xerophytica* R. E. Schultes (Aristolochiaceae). *Caldasia* 16: 295-300.
- González, F. 1994. Aristolochiaceae. In: G. Harling & L. Andersson (eds.), Flora of Ecuador. Monograph No. 51, pp. 1-42. Council for Nordic Publications in Botany.
- González, F. 1998. Two new species of *Aristolochia* (Aristolochiaceae) from Brazil and Peru. *Brittonia* 50: 5-10.
- González, F. 2000. Notes on the Central Andean species of *Aristolochia* (Aristolochiaceae) with the description of a new species from Bolivia. *Kew Bulletin* 55: 905-916.
- González, F.; J. C. Ospina & C. Zanotti. 2015. Sinopsis y novedades taxonómicas de la familia Aristolochiaceae para la Argentina. *Darwiniana*, nueva serie 3(1): 38-64. DOI: <https://doi.org/10.14522/darwiniana.2015.31.644>
- González, F. & N. L. Pabón-Mora. 2017. *Aristolochia keratuma* (Aristolochiaceae), nueva especie de la serie *Thyrsicae* del Chocó (Colombia) y clave de identificación para sus especies. *Caldasia* 39: 50-58.
- González, F. & N. L. Pabón-Mora. 2018. Sinopsis actualizada de *Aristolochia* (Aristolochiaceae, Piperales) en Panamá. *Acta Botanica Mexicana* 122: 109-140.
- González, F. & D. W. Stevenson. 2000a. Perianth development and systematics of *Aristolochia*. *Flora* 195: 370-391.
- González, F. & D. W. Stevenson. 2000b. Gynostemium development in *Aristolochia*. *Botanische Jahrbücher für Systematik, Pflanzengeschichte und Pflanzengeographie* 122: 249-291.
- Hoehne, F. C. 1910. Aristolochiaceae. *Comissão de Linhas Telegráficas e Estratégicas de Mato Grosso ao Amazonas 5, Bot. pt. 1*: 65, pl. 60.
- Hoehne, F. C. 1927. Monografía Ilustrada das Aristolochiaceae Brasileiras. *Memórias do Instituto Oswaldo Cruz* 20: 67-175, tabs. 16-103.
- Hoehne, F. C. 1942. Aristolochiaceae. In: Flora Brasílica 15: 1-141, t. 1-123.
- IUCN. 2014. Guidelines for using the IUCN Red List Categories and Criteria. Version 11. Prepared by the Standards and Petitions Subcommittee. Available at: <https://www.iucnredlist.org/documents/RedListGuidelines.pdf>
- IUCN Standards and Petitions Committee. 2022. Guidelines for using the IUCN Red List categories and criteria, Version 15.1. Prepared by the IUCN Standards and Petitions Committee. Available at: <https://www.iucnredlist.org/documents/RedListGuidelines.pdf>

- JSTOR. 2021. JSTOR Global Plants. <https://plants.jstor.org/collection/TYPSPE> (accessed April 2023).
- Keller, H. A. 2010. *Aristolochia stomachioides* (Aristolochiaceae), nueva cita para la Flora Argentina. *Bonplandia* 19: 121-125.
- León, B. & M. I. La Torre. 2006. Aristolochiaceae endémicas del Perú. *Revista Peruana de Biología* 13: 56s-57s.
- Macbride, J. F. 1937. Aristolochiaceae, Flora of Peru. *Field Museum of Natural History Botanical Series* 13: 431-443.
- Malme, G. O. A. 1904. Beiträge zur Kenntnis der südamerikanischen Aristolochiaceen. *Arkiv för Botanik* 1: 521-548.
- Masters, M. T. 1875. Aristolochiaceae. In: C. F. P. Martius et al. (eds.), *Flora Brasiliensis* 4: 77-114.
- Moore, S. 1915. The Aristolochias of Pará (Brazil). *Journal of Botany* 53: 1-8.
- Pfeifer, H. W. 1966. Revision of the North and Central American hexandrous species of *Aristolochia* (Aristolochiaceae). *Annals of the Missouri Botanical Garden* 53: 115-196.
- Schmidt, O. C. 1927. Beiträge zur Kenntnis der Aristolochiaceen. I. *Repertorium Specierum Novarum Regni Vegetabilis* 23: 282-299.
- Thiers, B. [2023, permanently updated, accessed April 2023]. Index Herbariorum: A global directory of public herbaria and associated staff. New York Botanical Garden's virtual herbarium. <https://sweetgum.nybg.org/ih>
- Ule, E. H. G. 1905. Aristolochiaceae. *Verhandlungen des Botanischen Vereins für die Provinz Brandenburg* 47: 118-124.