



Biotecnia

ISSN: 1665-1456

Universidad de Sonora, División de Ciencias Biológicas y de la Salud

Soto-García, Marcela; Rosales-Castro, Martha; García-Sánchez, José Rubén; Rivas-Arreola, María José  
In vitro cytotoxic activity of bark extracts from *Pinus durangensis* Martínez and *Quercus sideroxyla* Bonpl  
Biotecnia, vol. 24, no. 1, 2022, January-April, pp. 55-61  
Universidad de Sonora, División de Ciencias Biológicas y de la Salud

DOI: <https://doi.org/10.18633/biotecnia.v24i1.1520>

Available in: <https://www.redalyc.org/articulo.oa?id=672974928001>

- How to cite
- Complete issue
- More information about this article
- Journal's webpage in redalyc.org

redalyc.org

Scientific Information System Redalyc  
Network of Scientific Journals from Latin America and the Caribbean, Spain and Portugal

Project academic non-profit, developed under the open access initiative

# In vitro cytotoxic activity of bark extracts from *Pinus durangensis* Martinez and *Quercus sideroxyla* Bonpl.

Actividad citotóxica in vitro de los extractos de *Pinus durangensis* Martinez y *Quercus sideroxyla* Bonpl.

Marcela Soto-García<sup>1</sup>, Martha Rosales-Castro<sup>2</sup>, José Rubén García-Sánchez<sup>3</sup>, María José Rivas- Arreola<sup>\*4</sup>

<sup>1</sup> Universidad Autónoma de Ciudad Juárez, Instituto de Ciencias Biomédicas-DMCU, C.P.32310. Ciudad Juárez, Chih., México.

<sup>2</sup> Instituto Politécnico Nacional, CIIDIR- Unidad Durango, Sigma 119 Fraccionamiento 20 de noviembre, C.P. 34220, Durango, Dgo. México.

<sup>3</sup> Instituto Politécnico Nacional, Sección de Estudios de Posgrado e Investigación. Escuela Superior de Medicina del, Plan de San Luis y Díaz Mirón s/n, Col. Casto de Santo Tomas, Delegación Miguel Hidalgo, C.P. 113340, Ciudad de México, México.

<sup>4</sup> Universidad Iberoamericana Puebla Blvd. del Niño Poblano 2901 Col. Reserva Territorial Atlxícáyolt San Andrés Cholula, Puebla, México C.P. 72820.

## ABSTRACT

Nowadays, plants bioactive compounds represent a great potential for discovering novel drugs that could act in cancer treatment. Previous word reports that different pine and oak species possess cytotoxic activities against various cancer cell lines. The present study evaluated the phenolic profile and cytotoxic activities of crude and organic extracts from *P. durangensis* and *Q. sideroxyla* bark. Among ESI-MS identified compounds in extracts from both species, are taxifolin and procyanidin dimers. The cytotoxic activity was performed on MDA-MB-231 (breast cancer), HeLa (cervix cancer), MCF-10A (breast non-tumorous cell), and HSF-1184 (human skin fibroblast cells) by MTT assay. All extracts showed a dose-dependent cytotoxic effect against cancer cell lines and small or no activity against the non-tumorous cells. The results provide important information about the cytotoxic activity of bark extracts from *P. durangensis* and *Q. sideroxyla*. This strong cytotoxic effect represents an opportunity for the valorization of a by-product of the wood industry.

**Keywords:** Bark extracts, cytotoxicity, phenolic profile.

## RESUMEN

En la actualidad, los compuestos bioactivos de las plantas representan un gran potencial para descubrir un fármaco novedoso que pueda actuar en el tratamiento del cáncer. Se ha informado que diferentes especies de pino y roble poseen actividades citotóxicas contra varias líneas celulares cancerosas. En el presente estudio, se evaluó el perfil fenólico y las actividades citotóxicas de extractos crudos y orgánicos de la corteza de *P. durangensis* y *Q. sideroxyla*. Algunos compuestos, como la taxifolina y dímeros de procianidina, fueron identificados en extractos de ambas especies mediante ESI-MS. La actividad citotóxica se realizó en MDA-MB-231 (cáncer de mama), HeLa (cáncer de cuello uterino), MCF-10A (células no tumorales de mama) y HSF-1184 (células de fibroblastos de piel humana) mediante ensayo MTT. Todos los extractos mostraron un efecto citotóxico dependiente de la dosis contra las líneas celulares cancerosas y poca o ninguna actividad contra las no tumorales. Los resultados proporcionan infor-

mación importante sobre la actividad citotóxica de extractos de corteza de *P. durangensis* y *Q. sideroxyla*. Este fuerte efecto citotóxico representa una oportunidad para la valorización de un subproducto de la industria de la madera.

**Palabras clave:** Extractos de corteza, citotoxicidad, perfil fenólico

## INTRODUCTION

Breast and cervix cancer are the most common cancer-related death cause among females worldwide. Thus, it is crucial to search for newer therapies that can help prevent and treat this illness, besides lessening the side effects of available therapies (Khosropanah *et al.*, 2016; Saenglee *et al.*, 2016). The use of medicinal plants in the treatment of diseases is increasing due to the benefits against chronic diseases like cancer, since plants have bioactive compounds, playing a vital role in the development of pharmaceuticals (Abdulla *et al.*, 2014).

The therapeutic potential of medicinal plants generally associates to the antioxidant activity of phytochemicals, mainly phenols and flavonoids, closely linked to their ability to suppress cancer cells' growth through reduced oxidative stress (Marvibaigi *et al.*, 2016). However, the cellular mechanisms by which the phenols elicit these anticancer effects are multifaceted (Sorice *et al.*, 2016).

*Q. sideroxyla* and *P. durangensis* are forest timber species in Mexico. Their primary non-timber uses are for edible harvesting seeds and even as medicine in the treatment of disorders like ulcers and inflammatory problems (Bermejo and Pontones 1999; Luna *et al.*, 2003). Previous studies reported antibacterial effects, high phenolic content, and antioxidant properties in *P. durangensis* extracts (Rosales-Castro and González-Laredo, 2003; Rosales-Castro *et al.*, 2006). Extracts from *Q. sideroxyla* have shown hypoglycemic and genotoxic effects, amelioration of oxidative stress evaluated in a murine model, high antioxidant capacity, anti-inflammatory effects, and anticarcinogenic activity in rat colon (Moreno-Jimenez *et al.*, 2015; Soto-García *et al.*, 2016). The main bioactive compound of these species identified is flavan-3-ols in *Q. sideroxyla* (Rosales *et al.*, 2012), and flavonoids like taxifolin

\*Autor para correspondencia: María José Rivas Arreola  
Correo electrónico: mariajose.rivas@iberopuebla.mx

Recibido: 14 de junio de 2021

Aceptado: 26 de octubre de 2021

and quercetin in *P. durangensis* (Soto-García and Rosales-Castro, 2016).

Many investigations reported catechin, taxifolin, and quercetin as compounds with cytotoxic activities (Moreira et al., 2013; Evacuasiyany et al., 2014; Moreno-Jimenez et al., 2015; Alzaharna et al., 2017; Rameshthangam and Chitra, 2017). Thus, this research aimed to determine the cytotoxic activities of polyphenolic extracts from *P. durangensis* and *Q. sideroxyla* bark against MDA MB-231, MCF 10A HeLa, and HSF-1184 cell lines.

## METHODS

### Plant material

The bark collection was in January 2014 in Pueblo Nuevo, Durango, México. María Del Socorro Gonzalez Elizondo (taxonomist) identified the samples, and botanical specimens deposited at the Herbarium in CIIDIR-IPN, Durango, with vouchers number 42842, 428443, 42844 for *Q. sideroxyla* and 93, 97, 99, 100, 101 for *P. durangensis*. The bark preparation consisted in a mixture to make a unique sample, drying at room temperature, milled (mesh 40), and finally stored at 4 °C in paper bags until further use.

### Extracts preparation and purification

The bark powder (10 g) was twice soaked with 50% ethanol (ethanol/water 50:50 v/v) (2 x 300 mL) with stirring at room temperature for 24 h, then filtered through a Whatman filter paper no. 1. The extracts were combined, filtered, and then vacuum evaporated at 40 °C until ethanol was removed. A portion of the remaining aqueous extract was dried and identified as crude extract (CE); the other portion went through a liquid partition process with ethyl acetate (3 x 100 mL). The organic phase was vacuum evaporated at 40 °C and identified as the organic extract (OE).

### ESI-MS analysis

To obtain the *P. durangensis* and *Q. sideroxyla* OE spectra, we used a microTOF-QII mass spectrometer (Bruker Daltonics, Bremen, Germany) equipped with an electrospray ionization source (ESI). The parameters were set as capillary 2700 V, nebulizer pressure 1.2 bar, dry gas flow 11 L/min, and dry gas temperature 200 °C and the sample ran in the negative ion mode. The scan range was from 50 to 3000 m/z.

### Cytotoxic evaluation

#### Cell lines and culture conditions

Human breast cancer (MDA-MB-231), human cervical cancer (HeLa), and non-cancer (MCF-10A and H-1184) cell lines were from the American Tissue Culture Collection (ATCC, USA). The cultures of HeLa, MCF-7, MDA-MB-231, and H-1184 cells were in DMEM high glucose (Gibco Lab, Grand Island, NY) in 10% (MDA-MB-231 was 5%) bovine fetal serum and antibiotic 1X (streptomycin/penicillin 10000 U/mL, Gibco LabGrand Island, NY). The culture of MCF-10A was in DMEM/F12 medium with 10% bovine fetal serum, incubated at 37 °C in a humid atmosphere of 5% CO<sub>2</sub>.

### Cytotoxic assay

The cytotoxic assay was measured using MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) assay. Each cell line was plated in 96-well plates at a density of  $1 \times 10^4$  cells/well and incubated for 24 h (37 °C, 5% CO<sub>2</sub> air humidified) to allow them to adhere to the plate. Different concentrations of the CE and OE from *P. durangensis* and *Q. sideroxyla* were prepared (from the stock solution) and added to the culture medium to treat the cells, and cells incubated for another 72 h under the same conditions, using cells without treatment as a negative control. At the end of the experimental period, 100 µL of MTT dissolved in medium (5 mg/mL) were added to each well and incubated for another 2.5 h. Then, the medium was removed, the formazan crystals dissolved in 150 µL DMSO, and the absorbance measured at 550 nm in a spectrophotometer (Sinergy Biotek Instruments, Winooski, VT). In order to determine viability and the concentration leading to 50% inhibition of the viability, the determination of IC<sub>50</sub> was by regression analysis using the following equation:

$$\% \text{ Viability} = \frac{\text{Abs treated cells}}{\text{Abs control cells}} \times 100$$


**Statistical analysis**

The results reported are as the mean and the standard deviation (SD) of at least two independent experiments. The data analysis was by One-way analysis of variance (ANOVA), followed by post-hoc Turkey test, using a significance level of  $\alpha \leq 0.5$  to determine statistical significance. The statistical analysis software used for these analyses was Statistica 7.

## RESULTS AND DISCUSSION

The characterization of organic extracts (OE) from *P. durangensis* and *Q. sideroxyla* was by the analysis of the major peaks identified based on elution order, and that of all compounds was by interpreting their mass spectra obtained by the HPLC-ESI-MS, considering previously reported data. Figure 1 (A and B) shows the total ion chromatogram (TIC) of extracts from species studied. Several investigations reported cytotoxic effects of some Pine and oaks species on different cell lines that were related to compounds with the biological activities (Şöhretoğlu et al., 2012; Moradi et al., 2016; Li et al., 2016; Yang et al., 2016; Amessis-Ouchemoukh et al., 2017; Sarmeili et al., 2017).

Table 1 shows the main constituents of pine bark, where the compounds of the peaks 11, 12, 15, 16, 25, 36, 40, 41, and 58 coincide with compounds recently reported by Rosales (2017) in *P. durangensis* bark, demonstrating the presence of flavonoids like procyanidins and the abundance of taxifolin. Other TIC peak constituents were compared with the relevant literature in order to obtain a tentative identification (most studies on compounds of *Pinus* and *Quercus* bark) finding: kaempferol and eriodictyol ([M-H]<sup>-</sup> 285 and 287 respectively; De la Luz Cádiz-Gurrea et al., 2014), taxifolin-o-hexoside ([M-H]<sup>-</sup> 465; Cretu et al., 2013), β-hydroxypropiovanillone glucoside ([M-H]<sup>-</sup> 357; Karonen et al., 2004), HHDP-galoyl-glucose ([M-H]<sup>-</sup> 357; Fernández et al., 2009),



**Fig. 1.** Total ion chromatogram. (A) OE from *P. durangensis*. 11= Procyanidin dimer, 16= (+)-Catechin/(-)-Epicatechin, 25= Dihydroquercetin (Taxifolin), 40= Myricetin, 70= Kaempferol; (B) OE from *Q. sideroxyla*. 6= Procyanidin dimer, 10= Procyanidin dimer B1, 12= (+)-Catechin/(-)-Epicatechin; Procyanidin dimer, 13= (+)-Catechin/(-)-Epicatechin.

**Fig. 1.** Cromatografía de iones totales. (A) OE de *P. durangensis*. 11= Dímero de procianidina, 16= (+)-Catequina/(-)-Epicatequina, 25= Dihidroquercetina (Taxifolina), 40= Miricetina, 70= Kaempferol; (B) OE de *Q. sideroxyla*. 6= Dímero procianidina, 10= Dímero procianidina B1, 12= (+)-Catequina/(-)-Epicatequina; dímero Procianidina, 13= (+)-Catequina/(-)-Epicatequina.

trigalloyl glucose ([M-H]<sup>-</sup>635; Mucilli *et al.*, 2017), (Epi)catechin-3-O-glucoside-gallate ([M-H]<sup>-</sup>603; Jiménez-Sánchez *et al.*, 2015), t-caftaric acid ([M-H]<sup>-</sup>311; Pardo-García *et al.*, 2014). The polyphenolic profiles of *Q. sideroxyla* shown in Table 2 includes procyanidin dimer of catechin/epicatechin compounds (peaks 6, 10, 12, and 13) which is, by far, the dominant polyphenols in bark extract from this species, also reported in previous studies (Rosales *et al.*, 2012). Other compounds present in the extract also compared with previously reported literature, resulted in the tentative identification of constituents such as Quercetin 3-O-glucoside and t-caftaric acid ([M-H]<sup>-</sup>463 and 311 respectively; Pardo-García *et al.*, 2014), Ellagic acid-rhamnoside ([M-H]<sup>-</sup>447; Santos *et al.*, 2013) gallic acid hexoside ([M-H]<sup>-</sup>331; Lorenz *et al.*, 2016), taxifolin ([M-H]<sup>-</sup>303; Rosales *et al.*, 2017). Other unknown compounds reported in barks detected in both extracts are also included ([M-H]<sup>-</sup>293; Mämmelä *et al.*, 2000).

In previous studies, we have identified some of the major phenolic compounds of crude *P. durangensis* and *Q. sideroxyla* bark extracts by HPLC-DAD (Soto-García and

**Table 1.** Compounds identified by ESI-MS in the *P. durangensis* bark organic extract.

**Tabla 1.** Identificación de compuestos por ESI-MS en extractos orgánicos de corteza de *P. durangensis*.

| # Peak | RT (min) | Mass [m/z]- | Suggested compound                  |
|--------|----------|-------------|-------------------------------------|
| 11     | 2.6      | 577.1320    | Procyanidin dimer                   |
| 12     | 2.7      | 289.0699    | (+)-Catechin/(-)-Epicatechin        |
| 15     | 3.3      | 577.1321    | Procyanidin dimer                   |
| 15     | 3.3      | 865.1916    | Procyanidin trimer                  |
| 16     | 3.5      | 289.0706    | (+)-Catechin/(-)-Epicatechin        |
| 16     | 3.5      | 579.1490    | Catechin dimer                      |
| 19     | 3.9      | 465.1041    | Taxifolin-O-hexoside                |
| 25     | 4.6      | 303.0506    | Dihydroquercetin (Taxifolin)        |
| 34     | 5.9      | 633.1210    | HHDP-galoyl-glucose                 |
| 36     | 6.1      | 317.0616    | Myricetin                           |
| 36     | 6.1      | 635.1369    | Trigalloyl glucose                  |
| 39     | 6.5      | 287.0539    | Eriodictyol                         |
| 40     | 6.9      | 317.0639    | Myricetin                           |
| 40     | 6.9      | 635.1369    | Trigalloyl glucose                  |
| 41     | 7.1      | 331.0448    | Pinomyricetin (myricetin-6Me)       |
| 50     | 8.7      | 357.1330    | b-hydroxypropiovanillone glucoside  |
| 58     | 9.9      | 301.0679    | Quercetin                           |
| 58     | 9.9      | 603.1440    | (Epi)catechin-3-O-glucoside-gallate |
| 70     | 11.6     | 285.0759    | Kaempferol                          |
| 77     | 12.5     | 311.1854    | t-caftaric acid                     |
| 79     | 12.9     | 293.1740    | Unknown                             |

Rosales-Castro, 2016); these results, together with those obtained in this research, confirm the reproducibility of their bioactive compounds in bark extracts, like catechin in both species.

The *P. durangensis* and *Q. sideroxyla* CE and OE bark cytotoxic effects were also evaluated on a cell line from human breast carcinoma, estrogen receptor-negative (ER-) MDA-MB-231, and cervix carcinoma cell line HeLa. Non-tumorous MCF-10A and HSF-1184 cells, used as controls, allowed comparing the effects produced by the extracts in healthy and tumor cells.

Results showed that crude extract and organic extract of both species have cytotoxic activity on cancer cell lines growth. A concentration-dependent manner (Fig. 2 A, B, C), becomes statistically different ( $p \leq 0.05$ ) between treated and untreated cells (MDA-MB-231 and HeLa cell lines), nevertheless they were not cytotoxic to the non-tumorous HSF-1184 cell line.


The crude extracts showed a slight reduction in cell viability on normal breast MCF-10A cells, 20 % approximately (Fig. 2 D), even at the higher applied doses. These results

**Table 2.** Compounds identified by ESI-MS in the *Q. sideroxylla* bark organic extract.**Tabla 2.** Identificación de compuestos por ESI-MS en extractos orgánicos de corteza de *Q. sideroxylla*.

| # Peak | RT (min) | Mass [m/z]-       | Suggested compound                              |
|--------|----------|-------------------|-------------------------------------------------|
| 6      | 2.6      | 577.1319          | Procyanidin dimer                               |
| 7      | 2.7      | 289.0702          | (+)-Catechin/(-)-Epicatechin                    |
| 10     | 3.2      | 577.1323          | Procyanidin dimer B1                            |
| 12     | 3.5      | 289.0704;579.1489 | (+)-Catechin/(-)-Epicatechin; Procyanidin dimer |
| 13     | 3.7      | 289.0702          | (+)-Catechin/(-)-Epicatechin                    |
| 14     | 3.8      | 577.1307          | Procyanidin dimer                               |
| 15     | 4.0      | 463.0873          | Quercetin 3-O-glucoside                         |
| 18     | 4.2      | 447.0920          | Ellagic acid-rhamnoside                         |
| 21     | 4.6      | 303.0498          | Dihydroquercetin                                |
| 37     | 8.2      | 331.2490          | Gallic acid hexoside                            |
| 57     | 12.5     | 311.1848          | t-caftaric acid                                 |
| 59     | 12.9     | 293.1736          | Unknow                                          |

indicate that the *P. durangensis* and *Q. sideroxylla* extracts preferentially reduced the growth of the tumoral cell while not affecting the healthy cell line. The role of phenolic compounds in the extracts is essential, since it has been observed that polyphenols such as catechins induce apoptotic cell death in MDA MB-231 cell lines but not in normal cells (MCF 10-A) (Farhan et al., 2016). The CE of both species presents catechin in their compounds, explaining the selective effect on MCF-10A.

Several investigations reported that the effects obtained on cell proliferation relate to dose and cell type, implying a different sensitivity by different cell lines (Gezer et al., 2015; Anlar et al., 2016). It has also been reported that the selective activity of cytotoxic compounds against healthy (MCF- 10A) and tumor cells is due to a broad spectrum of mechanisms responsible for the resistance. These include differences in the subcellular localization of Bik and differences in mitochondrial-mediated apoptosis that explain the sensitivity exerted by the bark extracts from *P. durangensis* and *Q. sideroxylla* in the cell lines (Studzinska-Sroka et al., 2016).



**Fig. 2.** Cytotoxic effect of *P. durangensis* and *Q. sideroxylla* bark extracts on cells. (A) HeLa cells, (B) MDA-MB-231 cells, (C) HSF-1184 cells and (D) MCF-10A. Percent cell survival in the control group (untreated cells) assumed as 100. Results are expressed as the mean  $\pm$  SD of two independent experiments (n=4). Significant difference \*p  $\leq$  0.05 versus the control group.

**Fig. 2.** Efecto citotóxico de los extractos de corteza de *P. durangensis* y *Q. sideroxylla* sobre las células. (A) Células HeLa, (B) Células MDA-MB-231, (C) Células HSF-1184 y (D) células MCF-10A. El porcentaje de supervivencia celular en el grupo de control (células no tratadas) se asumió como 100. Los resultados se expresan como la media  $\pm$  DE de dos experimentos independientes (n = 4). Diferencia significativa \* p  $\leq$  0.05 frente al grupo de control.

IC<sub>50</sub> determination provides further clarification of both species' crude and organic extracts behavior in different cell lines (Table 3). It shows that organic extracts of both species have a major cytotoxic effect on MDA-MB-231 and HeLa cells, maybe due to the concentration of metabolites exerted by solvent. Previous results report that compounds such as taxifolin and procyanidins, the most abundant present in each of the extracts, play a fundamental role in their cytotoxic activity, inducing apoptosis on upregulating the expression of the proteins in HeLa and MDA-MB-231 cells (Amalinei *et al.*, 2014).

**Table 3.** IC<sub>50</sub> for *P. durangensis* and *Q. sideroxylla* bark extracts on cancer cell lines and non-tumorigenic cells.

**Tabla 3.** IC<sub>50</sub> de los extractos de *P. durangensis* y *Q. sideroxylla* sobre la viabilidad de líneas celulares cancerosas y no tumorales.

| Extract                  | Cell lines               |                         |          |         |
|--------------------------|--------------------------|-------------------------|----------|---------|
|                          | IC <sub>50</sub> µg/mL   |                         |          |         |
|                          | HeLa                     | MDA-MB-231              | HSF-1184 | MCF-10A |
| <i>P. durangensis</i> CE | 210.30±6.75 <sup>a</sup> | 77.34±4.06 <sup>a</sup> | ND       | ND      |
| <i>P. durangensis</i> OE | 201.50±8.29 <sup>a</sup> | 37.49±0.95 <sup>c</sup> | ND       | —       |
| <i>Q. sideroxylla</i> CE | 184.03±7.83 <sup>b</sup> | 46.83±0.73 <sup>b</sup> | ND       | ND      |
| <i>Q. sideroxylla</i> OE | 103.64±3.87 <sup>c</sup> | 32.82±5.54 <sup>c</sup> | ND       | —       |

ND: not detected within the investigated concentration range. Means ± SD in each column followed by different letters are statistically different by Tukey test  $p \leq 0.05$ .

The abundance of catechins and the synergy with other metabolites within the phenolic profile of *Q. sideroxylla* could be why it showed the highest cytotoxic activity compared to *P. durangensis*. In addition, there is evidence that flavonoids that possess C2-C3 unsaturated bond and a carbonyl group at position 4 exhibit lower IC<sub>50</sub>. These two functional groups may increase the compound's activity by affording more stable flavonoid radicals via conjugation and electron delocalization (Sadeghi-Aliabadi *et al.*, 2012).

Several investigations reported cytotoxic effects of some *Pinus* and oaks species on different cell lines that were related to compounds with the biological activities (Şöhretoğlu *et al.*, 2012; Li *et al.*, 2016; Moradi *et al.*, 2016; Yang *et al.*, 2016; Amessis-Ouchemoukh *et al.*, 2017; Sarmeli *et al.*, 2017).

## CONCLUSIONS

*Quercus* and *Pinus* species studies are intense due to their antioxidant and anti-inflammatory, antimicrobial, and anticancer potential; however, studies on the phenolic profile is scarce, and is associated with the reported effect. This work demonstrated that the phenolic profile and the concentration of specific metabolites such as the procyanidin

dimer of catechin/epicatechin compounds was determinant in the cytotoxic activity evaluated in the cell lines used. *Q. sideroxylla* OE showed a potent cytotoxic activity compared to CE and OE from *P. durangensis*. This strong cytotoxic effect represents an opportunity for the valorization of the wood industry by-products, suggesting that OE of *Q. sideroxylla* may represent a good alternative in the search for new agents of natural origin to treat mammary and cervix cancer. Nevertheless, more *in vitro* and *in vivo* studies are necessary in order to understand and establish the mechanism of action.

## ACKNOWLEDGMENTS

The authors are grateful to María del Socorro González Elizondo for the identification of the study species.

## ETHICAL DISCLOSURES

The authors declare that this work did not required human or animal experiments, no patient data appear in this article, and declare no conflicts of interest.

## REFERENCES

- Abdullah, A. S. H., Mohammed A. S., Abdullah R., Mirghani M.E.S., Al-Qubaisi M. 2014. Cytotoxic effects of *Mangifera indica* L. kernel extract on human breast cancer (MCF-7 and MDA-MB-231 cell lines) and bioactive constituents in the crude extract. BMC Complementary and alternative medicine. 14(1): 1-10.
- Alzaharna, M., Alqouqa I., Cheung H. Y. 2017. Taxifolin synergizes Andrographolide-induced cell death by attenuation of autophagy and augmentation of caspase dependent and independent cell death in HeLa cells. PLoS One. 12(2): e0171325.
- Amalinei, R. L., Trifan A., Cioanca O., Miron S.D., Mihai C. T., Rotinberg P., Miron A. 2014. Polyphenol-rich extract from *Pinus sylvestris* L. bark—chemical and antitumor studies. Medical-Surgical Journal, 118(2): 551-557.
- Amessis-Ouchemoukh, N., Ouchemoukh S., Meziant N., Idiri Y., Hernanz D., Stinco C. M. 2017. Bioactive metabolites involved in the antioxidant, anticancer and anticalpain activities of *Ficus carica* L., *Ceratonia siliqua* L. and *Quercus ilex* L. extracts. Industrial Crops and Products. 95: 6-17.
- Anlar, H. G., Bacanlı M., Kutluk B., Ahmet, Başaran N. B. 2016. Cytotoxic Activity of Resveratrol in Different Cell Lines Evaluated by MTT and NRU Assays. Turkish Journal of Pharmaceutical Sciences. 13(1): 27-34.
- Bermejo, B., Pontones J. B. 1999. Los pinos mexicanos y su utilización como especies introducidas de alto potencial en varios países del mundo. Segundo Simposio sobre Avances en la Producción de Semillas Forestales en América Latina. CATIE. 8:249-253.
- Cretu, E., Karonen M., Salminen J. P., Mircea C., Trifan A., Charalambous V., Miron A. 2013. In vitro study on the antioxidant activity of a polyphenol-rich extract from *Pinus brutia* bark and its fractions. Journal of Medicinal Food. 16(11): 984-991.
- Dela Luz Cádiz-Gurrea, M., Fernández-Arroyo S., Segura-Carretero A. 2014. Pine bark and green tea concentrated extracts: antioxidant activity and comprehensive characterization of

- bioactive compounds by HPLC–ESI–QTOF–MS. International Journal of Molecular Sciences. 15(11): 20382-20402
- Evacuasiy, E., Ratnawati H., Liana L. K., Widowati W., Maesaroh M., Mozef T., Risdian C. 2014. Cytotoxic and antioxidant activities of catechins in inhibiting the malignancy of breast cancer. Oxidant and Antioxidants in Medical Science. 3(2): 141-146.
- Farhan, M., Khan H.Y., Oves M., Al-Harrasi A., Rehmani N., Arif H., Ahmad A. 2016. Cancer therapy by catechins involves redox cycling of copper ions and generation of reactive oxygen species. Toxins, 8(2): 37.
- Fernandes, A., Fernandes I., Cruz L., Mateus N., Cabral M., De Freitas V. 2009. Antioxidant and biological properties of bioactive phenolic compounds from *Quercus suber* L. Journal of Agricultural and Food Chemistry. 57(23): 11154-11160.
- Gezer, C., Yücecan S., Rattan S. I. S. 2015. Artichoke compound cynarin differentially affects the survival, growth, and stress response of normal, immortalized, and cancerous human cells. Turkish Journal of Biology. 39(2): 299-305.
- Jiménez-Sánchez, C., Lozano-Sánchez J., Gabaldón-Hernández J.A., Segura-Carretero A., Fernández-Gutiérrez A. 2015. RP-HPLC–ESI–QTOF/MS 2 based strategy for the comprehensive metabolite profiling of *Sclerocarya birrea* (marula) bark. Industrial Crops Products. 71: 214-234.
- Karonen, M., Hämäläinen M., Nieminen R., Klika K. D., Loponen J., Ovcharenko V.V., Pihlaja K. 2004. Phenolic extractives from the bark of *Pinus sylvestris* L. and their effects on inflammatory mediators nitric oxide and prostaglandin E2. Journal of Agricultural and Food Chemistry. 52(25): 7532-7540.
- Khosropanah, M. H., Dinarvand A., Nezhadhosseini A., Haghighi A., Hashemi S., Nirouza F., Dehghani H. 2016. Analysis of the antiproliferative effects of curcumin and nanocurcumin in MDA-MB231 as a breast cancer cell line. Iranian Journal of Pharmaceutical Research. 15(1), 231.
- Li, Y. Y., Feng J., Zhang X. L., Li M. Q., Cui Y. Y. 2016. Effects of *Pinus massoniana* bark extract on the invasion capability of HeLa cells. Journal of Functional. Foods. 24: 520-526.
- Lorenz, P., Heinrich M., García-Käufer M., Grunewald F., Messerschmidt V, Herrick A., Steinborn C. 2016. Constituents from oak bark (*Quercus robur* L.) inhibit degranulation and allergic mediator release from basophils and mast cells in vitro. Journal of Ethnopharmacology. 194: 642-650.
- Luna-José, A. D. L., Montalvo-Espinosa L., Rendón-Aguilar B. 2003. Los usos no leñosos de los encinos en México. Boletín de la Sociedad Botánica de México. 72: 107-117.
- Mämmelä, P., Savolainen H., Lindroos L., Kangas J., Vartiainen T. 2000. Analysis of oak tannins by liquid chromatography-electrospray ionisation mass spectrometry. Journal of Chromatography A. 891(1): 75-83.
- Marvibaigi, M., Amini N., Supriyanto E., Majid F.A.A., Jaganathan S.K., Jamil S., Nasiri R. 2016. Antioxidant Activity and ROS-Dependent Apoptotic Effect of *Scurrula ferruginea* (Jack) Danser Methanol Extract in Human Breast Cancer Cell MDA-MB-231. PLoS One. 11(7): e0158942.
- Moradi, M. T., Karimi A., Alidadi S. 2016. In vitro antiproliferative and apoptosis-inducing activities of crude ethyle alcohol extract of *Quercus brantii* L. acorn and subsequent fractions. Chinese Journal of Natural Medicines. 14(3): 196-202.
- Moreira, L., Araújo I., Costa, Correia-Branco A., Faria A., Martel F., Keating E. 2013. Quercetin and epigallocatechin gallate inhibit glucose uptake and metabolism by breast cancer cells by an estrogen receptor-independent mechanism. Experimental Cell Research. 319(12): 1784-1795.
- Moreno-Jiménez, M. R., Trujillo-Esquivel, F., Gallegos-Corona M. A., Reynoso-Camacho M. A., González-Laredo R. F., Gallegos-Infante J. A., Ramos-Gómez M. 2015. Antioxidant, anti-inflammatory and anticarcinogenic activities of edible red oak (*Quercus* spp.) infusions in rat colon carcinogenesis induced by 1, 2-dimethylhydrazine. Food Chemical Toxicology. 80: 144-153.
- Muccilli, V., Cardullo N., Spatafora C., Cunsolo V., Tringali C. 2017. α-Glucosidase inhibition and antioxidant activity of an oenological commercial tannin. Extraction, fractionation and analysis by HPLC/ESI-MS/MS and 1 H NMR. Food Chemistry. 215: 50-60.
- Pardo-García, A. I., Martínez-Gil A. M., Cadahía E., Pardo F., Alonso G. I., Salinas M. R. 2014. Oak extract application to grapevines as a plant biostimulant to increase wine polyphenols. Food Research International. 55: 150-160.
- Rameshthangam, P. y Chitra, J. P. 2017. Synergistic anticancer effect of green synthesized nickel nanoparticles and quercetin extracted from *Ocimum sanctum* leaf extract. Journal of Materials Science & Technology. 34(3): 508-522.
- Rosales-Castro, M. y González- Laredo, R. F. 2003. Comparación del contenido de compuestos fenólicos en la corteza de ocho especies de pino. Madera y Bosques, 9: 41-49.
- Rosales-Castro, M., Pérez López, M. E., Ponce Rodríguez, M. D. C. 2006. Propiedades antirradicales libres y antibacterianas de extractos de corteza de pino. Madera y Bosques, 12: 37-49.
- Rosales-Castro, M., González-Laredo, R. F., Rocha-Guzmán N. E., Gallegos-Infante J. A., Rivas-Arreola, M. J., Karchesy, J. 2012. Antioxidant activity of fractions from *Quercus sideroxyla* bark and identification of proanthocyanidins by HPLC-DAD and HPLC-MS. Holzforschung. 66: 577-584.
- Rosales-Castro, M., González-Laredo, R. F., Rivas-Arreola, M. J., Karchesy, J. 2017. Chemical analysis of polyphenols with antioxidant capacity from *Pinus durangensis* bark. Journal of Wood Chemistry and Technology. 37(5): 393-404.
- Sadeghi-Aliabadi, H., Mosavi H., Mirian M., Kakhki S., Zarghi A. 2012. The Cytotoxic and Synergistic Effects of Flavonoid Derivatives on Doxorubicin Cytotoxicity in HeLa, MDA-MB-231, and HT-29 Cancer Cells. Iranian Journal of Toxicology, 5(15): 558-564.
- Saenglee, S., Jogloy, S., Patanothai, A., Leid, M., Senawong, T. 2016. Cytotoxic effects of peanut phenolics possessing histone deacetylase inhibitory activity in breast and cervical cancer cell lines. Pharmacological Reports. 68(2): 1102-1110.
- Santos, S. A., Villaverde, J. J., Sousa, A. F., Coelho, J. F., Neto, C. P., Silvestre, A. J. 2013. Phenolic composition and antioxidant activity of industrial cork by-products. Industrial Crops and Products. 47: 262-269.
- Sarvmeili, N., Jafarian-Dehkordi, A., Zolfaghari, B. 2016. Cytotoxic effects of *Pinus eldarica* essential oil and extracts on HeLa

- and MCF-7 cell lines. *Research in Pharmaceutical Sciences*, 11(6): 476-483.
- Şöhretoğlu, D., Sabuncuoğlu, S., Harput, Ü. Ş. 2012. Evaluation of antioxidative, protective effect against H<sub>2</sub>O<sub>2</sub> induced cytotoxicity, and cytotoxic activities of three different *Quercus* species. *Food and Chemical Toxicology*. 50(2): 141-146.
- Sorice, A., Guerriero E., Volpe M.G., Capone F., La Cara F., Ciliberto G., Costantini S. 2016. Differential Response of Two Human Breast Cancer Cell Lines to the Phenolic Extract from Flaxseed Oil. *Molecules*. 21(3): 319-337.
- Soto-García, M., Rosales-Castro, M., Escalona-Cardoso, G. N., Paniagua-Castro, N. 2016. Evaluation of Hypoglycemic and Genotoxic Effect of Polyphenolic Bark Extract from *Quercus sideroxyla*. *Evidence-Based Complementary and Alternative Medicine*. 4032618.
- Soto-García, M. y Rosales-Castro M. 2016. Efecto del solvente y de la relación masa/solvente, sobre la extracción de compuestos fenólicos y la capacidad antioxidante de extractos de corteza de *Pinus durangensis* y *Quercus sideroxyla*. *Maderas. Ciencia y Tecnología*, 18: 701-714.
- Studzińska-Sroka, E., Piotrowska, H., Kucińska, M., Murias, M., Bylka, W. 2016. Cytotoxic activity of physodic acid and acetone extract from *Hypogymnia physodes* against breast cancer cell lines. *Pharmaceutical Biology*. 54(11): 2480-2485.
- Yang, I. H., Shin, J. A., Kim, L. H., Kwon, K. H., Cho, S. D. 2016. The caspase 3-dependent apoptotic effect of pycnogenol in human oral squamous cell carcinoma HSC-3 cells. *Journal of Clinical Biochemistry and Nutrition*. 15-7.