



Revista de investigación e innovación en ciencias de la salud

ISSN: 2665-2056

Fundación Universitaria María Cano

Zapata Restrepo, Nancy Julieth; Ocampo-Lopera, Diana-María; Ibarra-Rivera, David-Alejandro; Sierra-Zuluaga, Ana-Sofía; Mejía, Melisa; Espinosa-Duque, Daniel

Termination of Psychotherapy with Children and Adolescents
in the Institutional Contexts from the Parents' Perspective

Revista de investigación e innovación en ciencias de la
salud, vol. 6, no. 2, 2024, July-December, pp. 189-203

Fundación Universitaria María Cano

DOI: <https://doi.org/10.46634/riics.195>

Available in: <https://www.redalyc.org/articulo.oa?id=673278576014>

- How to cite
- Complete issue
- More information about this article
- Journal's webpage in redalyc.org

UNEM  redalyc.org

Scientific Information System Redalyc

Network of Scientific Journals from Latin America and the Caribbean, Spain and Portugal

Project academic non-profit, developed under the open access initiative

Termination of Psychotherapy with Children and Adolescents in the Institutional Contexts from the Parents' Perspective

Terminación de la psicoterapia con niños y adolescentes en los contextos institucionales desde la perspectiva de los padres

Nancy Julieth Zapata Restrepo¹  , Diana-María Ocampo-Lopera²  ,
 David-Alejandro Ibarra-Rivera¹  , Ana-Sofía Sierra-Zuluaga¹  , Melissa Mejía²  ,
 Daniel Espinosa-Duque²  

¹ School of Psychology; Universidad de San Buenaventura; Medellín; Colombia.

² School of Psychology; Universidad CES; Medellín; Colombia.



Correspondence

Nancy Julieth Zapata Restrepo.
 Email: nancy.zapata@usbmed.edu.co

How to cite

Zapata Restrepo, Nancy Julieth; Ocampo-Lopera, Diana-María; Ibarra-Rivera, David-Alejandro; Sierra-Zuluaga, Ana-Sofía; Mejía, Melissa; Espinosa-Duque, Daniel. (2024). Termination of Psychotherapy with Children and Adolescents in the Institutional Contexts from the Parents' Perspective. *Revista de Investigación e Innovación en Ciencias de la Salud*. 6(2), 189-203. <https://doi.org/10.46634/riics.195>

Received: 03/30/2023

Reviewed: 05/21/2023

Accepted: 10/13/2023

Editor

Fraidy-Alonso Alzate-Pamplona, MSc. 

Copyright

© 2024, María Cano University Foundation. The *Revista de Investigación e Innovación en Ciencias de la Salud* provides open access to all its content under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International \(CC BY-NC-ND 4.0\)](https://creativecommons.org/licenses/by-nc-nd/4.0/) license.

Declaration of interests

The authors have declared that there is no conflict of interest.

Abstract

Introduction. Knowing the forms of termination of psychotherapy and the associated factors allows us to understand this moment of treatment and to think about strategies to improve the process.

Objective. The present study seeks to identify the types of termination of psychotherapy in children and adolescents and the factors that influence them in two psychological care centers.

Methodology. A quantitative, exploratory, cross-sectional design was used, with a descriptive and association scope; 100 patients (48 children and 52 adolescents) were surveyed.

Results. Four types of termination were found: due to institutional or therapist factors, attributed to factors external to the treatment, due to user dissatisfaction, and achievement of objectives. Some associations between them and the clinical and sociodemographic characteristics of the treatment were identified.

Discussion. Institutional aspects and external conditions to the patient must be considered in constructing intervention strategies for this population. In addition, educating parents on psychological intervention and its institutional scope is required to modulate the expectations associated with the process.

Keywords

Child and adolescent; termination of psychotherapy; institutional context; psychotherapy.

Data availability

All relevant data is in the article.
For further information, contact the corresponding author.

Financing

Universidad de San Buenaventura and Universidad CES [Project Code: M3894].

Disclaimer

The content of this article is the sole responsibility of the authors and does not represent an official opinion of their institutions or of *Revista de Investigación e Innovación en Ciencias de la Salud*.

Contribution of the authors

Nancy Julieth Zapata Restrepo:

Conceptualization, formal analysis, funding acquisition, investigation, project administration, supervision, writing – review & editing.

Diana-María Ocampo-Lopera:

Conceptualization, formal analysis, funding acquisition, investigation, methodology, software, writing – original draft.

David-Alejandro Ibarra-Rivera:

Conceptualization, data curation, formal analysis, funding acquisition, investigation, methodology.

Ana-Sofía Sierra-Zuluaga:

Conceptualization, data curation, formal analysis, writing – original draft.

Melisa Mejía: Conceptualization, data curation, formal analysis, writing – original draft.

Daniel Espinosa-Duque:

Conceptualization, formal analysis, funding acquisition, investigation, methodology, project administration, software, writing – review & editing.

Resumen

Introducción. Conocer los tipos de terminación de las psicoterapias y los factores asociados, permite comprender esta dimensión del tratamiento y pensar estrategias para el mejoramiento del proceso.

Objetivo. El presente estudio busca identificar los tipos de terminación de la psicoterapia en niños y adolescentes y los factores que influyen en ellos en dos centros de atención psicológica.

Metodología. Se utilizó un diseño cuantitativo, exploratorio, transversal, con un alcance descriptivo y de asociación, fueron encuestados 100 pacientes (48 niños y 52 adolescentes).

Resultados. Se encontraron cuatro tipos de terminación: por factores institucionales o del terapeuta, atribuida a factores externos del tratamiento, por insatisfacción de los usuarios y por cumplimiento de objetivos. Se identifican algunas asociaciones entre estos y las características clínicas y sociodemográficas del tratamiento.

Discusión. Los aspectos institucionales y las condiciones externas al paciente deben ser considerados en la construcción de las estrategias de intervención con esta población. Además, se requiere un proceso de educación a padres sobre la intervención psicológica y su alcance institucional para modular las expectativas asociadas al proceso.

Palabras clave

Psicoterapia infantojuvenil; terminación de psicoterapia; ámbito institucional; psicología.

Introduction

Mental health problems in children and adolescents have been the focus of attention worldwide [1] due to its high prevalence and gradual increase in Latin America [2]. This makes it necessary to understand and optimize the interventions and treatments offered to this population [3]. It is relevant to know why this type of treatment is terminated and the elements related to such decision [4]. This is a central issue since evidence shows that there are few children and adolescents who receive specialized help for their emotional needs, and even less children who complete their treatment [5].

Studies show that there are different ways of terminating psychotherapy, understood as the termination of patient-therapist sessions [6,7]. Among them, the most referenced ways are by mutual agreement, by request of one of the participants, and by dropout. Termination of therapy does not necessarily occur because of the achievement of goals. Termination also occurs prematurely, and the factors associated with each type are different [8].

Termination by mutual agreement between therapist and patient usually occurs when the goals agreed upon in the process have been achieved [9]. In the case of child and adolescent psychotherapy, this occurs when the therapist, the child, and their parents agree to conclude the process [10]. In this agreement, parents or legal guardians have an important and active role [11] because they usually seek help, support the definition of goals, and pay for the process [12].

When a patient, therapist, or contextual factor limits the process, such termination

may occur with partial achievement of the goals [13]. Therefore, the negotiation of the therapeutic alliance and of the termination of therapy is related to the achievement of goals and to the possibilities and limitations of the participants [14].

Furthermore, unilateral termination by the therapist is associated with personal conditions (e.g., end of practice, change of job, etc.) or with the achievement of the goals established [7,15,16]. In this type of termination, the patient's clinical conditions must be considered in order to face the termination of the therapeutic relationship and the need to make referrals to continue with the process, when required [7].

Finally, dropout is understood as the unilateral termination of treatment by the patient due to lack of motivation for the process or due to some limitation that makes it impossible to continue [5,11]. In child and adolescent psychotherapy, dropout is a major challenge, as it occurs in between 20% and 60% of cases, and there are not many studies on the aspects related to this type of termination [16,5].

Research on this topic has focused on premature termination and dropout, associating it with patient factors such as personality, type of problem, severity of symptoms [17], or ethnicity [12]. Other associated factors include elements of the family context, such as the attitude of the parents or guardians towards the treatment [18], the diagnosis and their expectations [12,19]. Added to this are factors such as dissatisfaction with the process and ruptures in the therapeutic alliance [20]. Finally, some studies show that the decision to terminate treatment may be an interaction of second-order factors [13] such as financial difficulties [21] or modification of extra-therapeutic factors that changes the needs of psychotherapy [22].

In this perspective, dropout is not a sign of unachieved goals, and mutual termination does not guarantee the achievement of these goals [8]. Another point to consider is that, in child and adolescent psychotherapy, the decision to attend is often made by the caregivers or guardians. Considering this, it is important to develop motivation and to agree on therapeutic goals integrating the perspectives of all involved [10].

Additionally, the specific characteristics of the institutional context must be taken into consideration, such as the number of appointments, access to treatment, flexibility of schedules and professional rotation [23,24]. These aspects may affect the continuity of treatment [25].

Understanding the factors related to the termination of psychotherapy with children and adolescents allows therapists to identify elements connected to premature termination [5]. At the institutional level, it is essential to know the reasons associated with the termination of therapy, because this makes it possible to develop strategies to improve adherence, guide work teams, build intervention modalities, and optimize treatments for greater effectiveness.

Considering the above, the present study aims to know the forms of psychotherapy termination in child and adolescent psychotherapy and their associated factors in two university psychological care centers in Antioquia (Colombia). This makes it possible to improve interventions, identify specific elements of this context, and develop treatments closer to the patients.

Method

This study has a quantitative, exploratory, cross-sectional research design with a descriptive and association scope. The research was conducted during 2020–2021 in two psychological care centers: IPS CES Sabaneta — Universidad CES (CES) and Consultorio Psicológico Popular — Universidad de San Buenaventura (USB).

Participants

A random sampling stratified by age group was carried out. One hundred patients who had terminated therapy in the last four months (48 children and 52 adolescents) were included in the study. The children included 33 females and 15 males between the age of 5 and 12 (age $X = 9.14$; $SD = 2.2$). Among adolescents, there were 35 females and 17 males between the ages of 13 and 18 (age $X = 15.82$; $SD = 1.64$).

The inclusion criteria were: to have been in psychotherapy for at least three sessions, to be between the ages of 5 and 18, and to agree to participate by means of informed consent approved by the bioethics committees of both institutions. Patients with a diagnosis of organic brain damage or cognitive disability were excluded. The inclusion criteria of considering patients with at least three sessions was adopted because some studies show that in these sessions the preliminary evaluation and therapeutic setting are carried out and the basis of the therapeutic alliance are consolidated [26].

Variables, instruments, and data collection

A survey was conducted with questions on sociodemographic aspects of the patient, rating of psychotherapy, and reasons for terminating therapy. An expert group based on a desk review constructed the survey and defined the thematic domains and items. A pilot test of the questionnaire was conducted to determine that responses were clear and understood by the patients (see Table 1). The interviews were conducted by telephone, audio-recorded and recorded in a database for analysis.

Procedure

Initially, patients who met the study criteria were selected from institutional care records. Subsequently, parents were invited and, if they accepted the ethical conditions, the interview was carried out. The open-ended responses were then categorized, and each therapy was classified according to a main form of termination. Finally, descriptive and association analyses were performed.

Data analysis

Descriptive statistics were calculated considering frequencies, percentages, means and standard deviations according to the type of variable. Percentages were reported because standardization of the data allows for the interpretation of trends in the information. Subsequently, chi2 analysis was performed to establish the levels of association between “types of termination of therapy” and sociodemographic variables, “rating therapy” and “who decided to terminate psychotherapy”. Analyses were performed with IBM SPSS Statistics 24 software.

Results

Four types of therapy termination reported by parents were identified in the results: related to external factors ($F=25$), institutional or therapist aspects ($F=39$), attributed to patient dissatisfaction ($F=17$), and achievement of goals ($F=19$). Among these, the most frequent reasons for terminating the processes were achievement with goals ($F=19$), change of psychologist ($F=15$), and psychologist’s suggestion ($F=10$) (see Table 2).

In addition to this, 46% of the processes were terminated by parental decision (31 children and 15 adolescents), 25% by the patients (3 children and 22 adolescents), 24% by the psychologist (12 children and 12 adolescents), 4% by mutual agreement between therapist, parents, and patients (1 child and 3 adolescents), and 1% by the health insurance provider (1 child).

Table 1. Variables Included in the Analyzes.

Variable group	Variables	Values	Variable Type
Sociodemographic variables	Age group	Child-Adolescent	Dichotomous
	Gender	Male-Female	Dichotomous
	Socio-economic level	Low (1-2)	Ordinary
		Medium (3-4)	
		High (5-6)	
Therapy rating	Motivation	Low-Medium-High	Ordinary
	Therapeutic actions		
	Therapeutic relationship		
	Therapeutic progress		
	Recommend		
	Satisfaction		
Variable group	Variables	Values	Variable type
Variables of external factors that influence the completion of psychotherapy	Socioeconomic factors	Economic aspects - Academic aspects - Social aspects	Nominal
	Institutional Factors	Location - Hours and/or availability of appointments - EPS Authorizations	Nominal
End of Therapy Variables	Type of therapy termination	Open responses	Nominal
	Reason for termination		
	Who decided to terminate psychotherapy		

Regarding the external factors that influenced psychotherapy termination, 22% attributed it to economic reasons, 12% to labor or academic elements, and 3% to social aspects related to affray. In addition, institutional factors such as location (13%), schedules and appointment availability (13%), and authorizations from insurance companies (3%) were also found. As Table 3 shows, there are no descriptive differences between children and adolescents.

Furthermore, when parents were asked to rate the psychotherapy, they reported a high rating of the service in general (90%), of the therapeutic relationship (84%), of the activities carried out in therapy (76%), and 83% would return or recommend the service. The dimensions related to motivation (68%) and therapy progress (59%) were those with the lowest percentage of positive ratings. The descriptive distribution of these dimensions was similar in children and adolescents (see Table 4).

Table 2. Frequency of types and reasons for terminating therapy in children and adolescents.

Types of therapy termination	Reasons for termination	Children		Adolescents		General	
		F	%	F	%	F	%
1. Due to factors external to psychotherapy	Time incompatibility	2	4,2	2	3,8	4	4
	Economic difficulties	5	10,4			5	5
	Location	2	4,2	1	1,9	3	3
	Modification of the school's requirement	1	2,1			1	1
	Parental divorce	1	2,1			1	1
	Pandemic	1	2,1	3	5,8	4	4
	Prioritize another activity (travel, work, another process, or institution)	3	6,3	3	5,8	6	6
	Appointments in another institution	1	2,1			1	1
2. Due to institutional and/or therapist factors	Change of psychologist	6	12,5	9	17,3	15	15
	Change in institutional procedures	1	2,1	2	3,8	3	3
	Remission	2	4,2	1	1,9	3	3
	Health system processes			1	1,9	1	1
	Appointment availability	4	8,3	4	7,7	8	8
	Psychologist decision	4	8,3	5	9,6	9	9
3. Due to patient dissatisfaction	With the professional	4	8,3	3	5,8	7	7
	With the progress of therapy	3	6,3	1	1,9	4	4
	With therapy			1	1,9	1	1
	Lack of motivation			5	9,6	5	5
4. Due to achievement of goals	Achievement of goals	8	16,7	11	21,2	19	19
Total		48	100	52	100	100	100

In the analysis of variable association among the four types of psychotherapy termination identified in the study and the sociodemographic variables, a significant result was found between the termination due to goal achievement and socioeconomic level (X^2 : 6.824, DoF:1, p.0.009), with a higher percentage in middle-class patients (15%) compared to low-class (4%).

Significant associations were also found between patients who terminated therapy due to dissatisfaction and the following variables: motivation in therapy (X^2 : 31,041, DoF:2, p.0.000), actions rating (X^2 : 26.481, DoF:2, p.0.000), therapeutic relationship (X^2 : 25.597, DoF:2, p.0.000), progress perception (X^2 : 17.920, DoF:2, p.0.000), and satisfaction in therapy (X^2 :

Table 3. Factors influencing the termination of therapy.

Factors Influencing Termination	Children	Adolescent	Total
	(F - %)	(F - %)	(F)
Socioeconomic factors			
Economic aspects	12 – 25%	10 – 19%	22
Labor aspects – Academics	6 – 12,5%	6 – 11,5%	12
Social aspects (Affray)	1 – 2%	2 – 4%	3
Institutional Factors			
Location	9 – 19%	4 – 8%	13
Hours and/or availability of appointments	5 – 10,5%	8 – 15%	13
Health system authorizations	1 – 2%	3 – 6%	4

Table 4. Rating dimensions associated with psychotherapy.

Rating dimensions of therapy	Low			Medium			High			Did not report
	N	A	T	N	A	T	N	A	T	
Motivation for therapy	6	7	13	7	9	16	33	35	68	3
Therapeutic actions	8	4	12	4	5	9	36	40	76	3
Therapeutic relationship	3	4	7	4	3	7	41	43	84	2
Therapy progress	11	9	20	7	12	19	30	29	59	2
Would recommend the institution	5	3	8	2	7	9	41	42	83	0
Satisfaction with the service	2	2	4	3	2	5	42	48	90	1

Note: N=Child frequency, A=Adolescent frequency, T=Total frequency, Valuation levels = Low: Answers 1 and 2, Medium: Answer 3, and High: Answers 4 and 5.

18.844, DoF:2, p.0.000). Based on these results, the therapy rating tends to be lower when therapy ends due to dissatisfaction (See [Table 5](#)).

Additionally, a significant association was found between psychotherapies that end due to external factors and the perception of progress rating (X^2 : 11.892, DoF:2, p.0.003). Thus, a higher progress rating tends to be found in this type of termination.

Finally, when calculating the association between the types of termination and “who decided to terminate the processes,” some significant results were found. An association was found among all types of termination and parental decision: by external factors (X^2 : 8.926a, DoF:1, p.0.003); by institutional aspects and/or therapist (X^2 : 6.305, DoF:1, p.0.012); by dissatisfaction (X^2 : 4.494a, DoF:1, p.0.034); by achievement with goals (X^2 : 6.576a, DoF:1, p.0.010). Also, an association was found between the therapist’s decision and termination by institu-

Table 5. Significant associations among types of termination and sociodemographic and therapy variables.

Types of termination	Variables with significant association	Descriptive Data	χ^2	DoF	p
1. Due to factors external to psychotherapy (25)	Therapy progress rating	Low: 9	11,892	2	0,003
		High: 15			
	Caregiver decides to terminate	No: 7	9,071	1	0,003
		Yes: 18			
	Therapist decides to terminate	No: 24	7,31	1	0,007
		Yes: 1			
2. Due to institutional and/or therapist factors (39)	Therapist decides to terminate	No: 24	7,331	1	0,007
		Yes: 15			
3. Due to patient dissatisfaction (17)	Motivation for therapy	Low: 9	31,041	2	0
		Medium: 4			
		High: 4			
	Therapeutic actions rating	Low: 8	26,481	2	0
		Medium: 3			
		High: 6			
	Therapeutic relationship rating	Low: 5	25,597	2	0
		Medium: 4			
		High: 8			
	Therapy progress rating	Low: 9	17,92	2	0
		Medium: 5			
		High: 3			
	Satisfaction with the service	Low: 4	18,844	2	0
		Medium: 5			
		High: 8			
4. Due to achievement of objectives (19)	Socio-economic level	Low: 4	6,824	1	0,009
		Medium: 15			
	Caregiver decides to terminate	No: 15	5,877	1	0,015
		Yes: 4			

tional aspects (X^2 : 8,000a, DoF:1, p .0.005) and by external aspects of therapy (X^2 : 6,831, DoF:1, p .0.009). According to these analyses, in termination due to factors external to the therapy, the decision tends to be made more by the parents, and in institutional termination by the therapists. In the case of dissatisfaction there is a higher percentage of parents who decide, while in the case of achievement with goals there is less parental involvement (see [Table 5](#)). As can be seen, in children there is greater influence of parents and therapists on the termination. In adolescents, only a positive association was found between termination due to dissatisfaction and the patient's decision (X^2 : 3.890a, DoF:1, p .0.049).

Discussion

The results of this study show four types of termination of child and adolescent psychotherapy: due to institutional and/or therapist factors, attributed to external treatment factors, due to user dissatisfaction, and due to goal achievement. Within each of these types of termination, multiple reasons and associated factors are identified.

The type of termination most referred by users of child and adolescent psychotherapy was related to institutional and/or therapist factors, considering aspects such as frequent changes of psychologists, limited availability of appointments, and the decision by the psychologist to end the process. Psychotherapy requires the establishment of a therapeutic relationship, which is sensitive to professional rotation and is frequent in teaching-service institutions. Some authors have pointed out that patients treated at these centers are affected by therapist discontinuity [25] and by conclusion of academic periods for training staff [7]. This reveals the importance of improving the therapeutic frame with users at the beginning of treatment, promoting the consolidation of a patient-institution alliance that supports the user in the face of staff rotation, by creating a triage system and case assignment that takes into account the particularities of each case and by improving the process of transferring the therapy from one therapist to another.

Simultaneously, the results show how institutions and therapists have various conditions that can become barriers to treatment, such as appointment availability and timeliness in their allocation [25], as well as administrative processes within the healthcare system [23,24], which becomes a challenge for treatment continuity.

In this type of termination, it is interesting that some psychologists unilaterally decide to end the process, although some parents express that the goals were not achieved or that they did not understand the decision made in these cases. This termination may be due to the limitations that professionals have in addressing complex problems and the absence of feedback to parents [27]. Considering this, it is recommended that parents be involved in the treatment through psychoeducation processes and systematic feedback on progress. This supports their perception of the advances [28] and the acceptance of the professional's evaluation [29].

Secondly, there is terminating treatment due to external factors to the treatment, which are related to economic conditions and the prioritization of other activities. Numerous studies have shown how economic conditions are one of the factors most associated with the risk of dropout [12,21,30] and how the therapist should consider the patient's and their family's living conditions, other treatments and activities they perform, as elements that can contribute to therapy or that can affect treatment continuity [31].

This type of termination is also associated with a better perception of progress in psychotherapy, which allows us to hypothesize that, as discomfort decreases, external factors or user activities become relevant and prioritized [13,32], making the therapeutic process dispensable [30]. In this sense, it is important for therapists to anticipate that certain improvements may be associated with premature termination of the process, which should be taken into account in therapy setting and planning [22,31].

Thirdly, termination due to patient dissatisfaction, there is discontent with different aspects of therapy, such as the therapist or the perception of progress. This type of termination has been associated with therapeutic alliance failure [33,34], parental disagreement with treatment [19] and diagnosis [35]. Therefore, the importance of parental involvement in therapy is highlighted in some studies [29,36,37].

In the case of adolescents, the literature reports in the same direction as this study that, when they feel dissatisfied, it is they who decide to terminate therapy [28]. Studies have reported that when adolescents perceive collaborative sessions and less pressure to talk, they have a stronger therapeutic alliance [38,39]; therefore, it is recommended to consider the characteristics of life cycle to achieve adherence [38].

Furthermore, this type of termination highlights the importance of addressing and modulating the expectations of those seeking therapy in the therapeutic context, clarifying the reach of the treatment. It becomes important to support parents in building expectations that align with the patient's needs [28], as well as the therapist's and institution's possibilities. We recommend establishing, for the initiation of child and adolescent therapy, processes that guide caregivers regarding the reach, procedures and limitations of psychotherapy based on the evaluation of the child or adolescent.

Lastly, termination due to attainment of objectives refers to therapies in which treatment goals are achieved. At this level, parents tend to participate less in the decision to end the process. However, some studies show that this type of termination does not necessarily imply closure and can occur unilaterally when significant progress is evident [8]. This is related to some studies that explain a relationship between satisfaction with the process, symptomatic improvement, and a decrease in the need for support [6,31].

This type of termination was more common among adolescents from middle socioeconomic backgrounds, with the clarification that the treatment centers referred to in this research primarily serve low and middle socioeconomic populations. Some authors point out the positive impact that social conditions and developmental stage have on adherence and treatment outcomes [12,21]. Additionally, adolescents from middle socioeconomic backgrounds experience fewer psychosocial problems, which promotes their reflective function [40-43].

The present study has limitations, such as the small sample size, which restricts the possibility of conducting other types of analysis. However, given its exploratory and associative nature, it allows for the identification of important trends and hypotheses for future studies in this research field. Furthermore, it is crucial to achieve a broader understanding of the termination of child and adolescent treatments by conducting new studies that consider and relate the perspectives of other therapy participants, such as patients, psychotherapists, and other institutional actors.

Based on the findings of this research, it is recommended to expand the technical approach to parents or caregivers in the training of psychologists. They play a crucial role in the continuity of therapy processes, and their expectations, level of participation, and satisfaction with therapy should be taken into account. Additionally, it is important to establish an induction or framing process for therapy in institutional settings that clarifies the reach of the intervention and the level of parental participation in the process. This can also help adjust the expectations of both patients and their parents.

In addition, institutional factors should be considered as aspects of the process with children and adolescents, not only as a logistical dimension, but as an actor that, although external to the patient-parent-therapist triad, has the effect of facilitating or hindering adherence to psychotherapy. In this sense, the patient-institution relationship must be worked on, since a proper alliance with it can mitigate the effect of institutional situations, such as professional rotation.

Conclusions

The findings of this research allow us to conclude important elements for psychotherapists, mental health institutions, and formative practice processes that contribute to the reflection on child and adolescent psychotherapy. It is advisable for university psychological care centers to have triage and referral processes to evaluate the pertinence of care by a practitioner or professional, thus providing timely attention according to the needs and characteristics of patients. In this respect, it is important for institutions to be aware of the implications of the teaching-service model and its impact on the patient's process. They should also consider emphasizing the training of psychologists in culturally and contextually informed therapies, enabling them to develop tools for working with vulnerable populations and in coordination with the healthcare system.

The psychotherapist should work on developing and modulating patient and caregiver expectations to make them more realistic and help the therapist manage identified expectations within the process, ensuring they are aligned with those of the patient. This implies considering some of the barriers or temporal limitations of the process within the setting and how they can be overcome, articulating with different support networks for patients.

On the other hand, it is advisable to make use of monitoring tools for assessing patient conditions in formulating, planning, and monitoring cases, allowing necessary and timely adjustments in the process. In addition to implementing routine monitoring of interventions with process and outcome variables to enhance intervention quality and overall service.

Finally, it is an invitation to continue with this type of research, considering the limitations of this study and how they can be overcome in future research with a larger sample size and involvement of other actors in the therapeutic process within institutional contexts. Furthermore, consider incorporating more variables related to experience level, context, culture, and therapeutic process that influence therapy termination in research. This study provides initial foundations for understanding types of therapy termination in children and adolescents, as well as associated process factors within an institutional context. Continued research in this area will contribute to therapy with children and adolescents.

References

1. Fondo de las Naciones Unidas para la Infancia. Estado Mundial de la Infancia. 2021. En mi mente. Promover, proteger y cuidar la salud mental de la infancia [Internet]. Nueva York: UNICEF; 2021. 259 p. Available from: <https://www.unicef.org/es/informes/estado-mundial-de-la-infancia-2021>
2. Cuello Royert C. Mapeo en salud mental adolescente en América latina y el caribe. In: Webinar I Panorama de la Salud Mental adolescente en América Latina y el Caribe [Internet]; 2019 Oct 8: UNICEF; 2019. 18 p. Available from: <http://www.codajic.org/node/4095>.
3. Shechtman Z. Group Counseling and Psychotherapy With Children and Adolescents Theory, Research, and Practice. Theory, Research, and Practice. New York: Routledge; 2017. 306 p. doi: <https://doi.org/10.4324/9781315093369>
4. Bear HA, Dalzell K, Edbrooke-Childs J, Garland L, Wolpert M. How to manage endings in unsuccessful therapy: A qualitative comparison of youth and clinician perspectives. Psychother Res [Internet]. 2022;32(2):249-62. doi: <https://doi.org/10.1080/10503307.2021.1921304>
5. Deakin E, Gastaud M, Tiellet Nunes ML. Child psychotherapy dropout: an empirical research review. J Child Psychother [Internet]. 2012;38(2):199-209. doi: <https://doi.org/10.1080/0075417X.2012.684489>
6. Swift JK, Callahan J, Levine JC. Using Clinically Significant Change to Identify Premature Termination. Psychotherapy (Chic) [Internet]. 2009;46(3): 328-35. doi: <https://doi.org/10.1037/a0017003>
7. Vélez Gómez P, Restrepo Ochoa DA. El proceso de terminación en psicoterapia de tiempo limitado: aspectos clínicos y técnicos. CES Psicología [Internet]. 2008;1(2):58-68. Available from: <https://revistas.ces.edu.co/index.php/psicologia/article/view/80>
8. Espinosa Duque D, Ibarra DA, Ocampo Lopera DM, Montoya L, Hoyos-Pérez A, Medina LF, Tamayo S, et al. Terminación de la psicoterapia desde la perspectiva de los pacientes. CES Psicología [Internet]. 2020;13(3):201-21. doi: <https://doi.org/10.21615/cesp.13.3.12>
9. Roe D, Dekel R, Harel G, Fennig S. Client's reasons for terminating psychotherapy: a quantitative and qualitative inquiry. Psychol Psychother [Internet]. 2006;79(4):529-38. doi: <https://doi.org/10.1348/147608305X90412>
10. Vidair HB, Feyijinmi GO, Feindler EL. Termination in cognitive-behavioural therapy with children, adolescents, and parents. Psychotherapy (Chic) [Internet]. 2017;54(1):15-21. doi: <https://doi.org/10.1037/pst0000086>
11. Rojas Olivares AM, Vélez Ríos J, González García J. Alianza terapéutica: Un apoyo en la adherencia al tratamiento y al cumplimiento de los objetivos de intervención con niñas, niños y adolescentes [specialization dissertation]. [Pereira]: Universidad Católica de Pereira; 2021. 35 p. Available from: <https://repositorio.ucp.edu.co/handle/10785/8247>

12. Lambert MJ, editor. Bergin and Garfield's handbook of psychotherapy and behaviour change. 6th ed. Hoboken: John Wiley & Sons; 2013. 851 p.
13. Santangelo PR. Terminación prematura en Psicoterapia. Revista de psicoterapia [Internet]. 2020;31(115):181-95. doi: <https://doi.org/10.33898/rdp.v31i115.316>
14. Råbu M, Haavind H. Coming to an end: A case study of an ambiguous process of ending psychotherapy. Couns Psychother Res [Internet]. 2012;12(2):109-17. doi: <https://doi.org/10.1080/14733145.2011.608131>
15. Johnson E, Mellor D, Brann P. Factors associated with dropout and diagnosis in child and adolescent mental health services. Aust N Z J Psychiatry [Internet]. 2009;43(5):431-37. doi: <https://doi.org/10.1080/00048670902817687>
16. Gastaud M, Numes ML. Treatment dropout in psychoanalytical psychotherapy: searching for definition. J bras psiquiatr [Internet]. 2010;59(3):247-54. doi: <https://doi.org/10.1590/S0047-20852010000300012>
17. Reich CM, Berman JS. Are psychotherapies with more dropouts less effective? Psychother Res [Internet]. 2020;30(1):23-40. doi: <https://doi.org/10.1080/10503307.2018.1534018>
18. Hernández Castellar AK, Jiménez Araujo AM, Valencia Valencia LX. Intervención con niños y adolescentes en terapia familiar: problemáticas, teorías y técnicas de intervención. [specialization dissertation]. [Medellín]: Universidad Católica Luis Amigó; 2020. 22 p. Available from: <http://179.50.60.21:8080/xmlui/handle/ucatolicaamigo/1008>
19. Ormhaug SM, Jensen TK. Investigating treatment characteristics and first-session relationship variables as predictors of dropout in the treatment of traumatized youth. Psychother Res [Internet]. 2018;28(2):235-49. doi: <https://doi.org/10.1080/10503307.2016.1189617>
20. Nof A, Dolev T, Leibovich L, Harel J, Zilcha-Mano S. If you believe that breaking is possible, believe also that fixing is possible: a framework for ruptures and repairs in child psychotherapy. Res Psychother [Internet]. 2019;22(1):45-57. doi: <https://doi.org/10.4081/ripppo.2019.364>
21. Thompson MN, Goldberg SB, Nielsen SL. Patient financial distress and treatment outcomes in naturalistic psychotherapy. J Couns Psychol [Internet]. 2018;65(4):523-30. doi: <https://doi.org/10.1037/cou0000264>
22. Garayoa JAC, Pérez-Testor S, Magriña MA. El abandono de los tratamientos psicoterapéuticos. Clínica y salud [Internet]. 1996;7(3):271-92. Available from: <https://journals.copmadrid.org/clysa/art/b73ce398c39f506af761d2277d853a92>
23. De la Parra G, Gómez-Barris E, Zúñiga AK, Dagnino P, Valdés C. Del diván al polí-clínico: un modelo de psicoterapia para instituciones. Aprendiendo de la experiencia (empírica). Revista Argentina de Clínica Psicológica [Internet]. 2018;27(2):182-202. doi: <https://doi.org/10.24205/03276716.2018.1057>

24. Jaramillo JC, Espinosa D, Ocampo D, Arango A, Arévalo M, García L, et al. Caracterización de la atención psicoterapéutica en la IPS CES Sabaneta, Colombia, 2014-2015. CES Psicología [Internet]. 2018;11(2):97-110. doi: <https://doi.org/10.21615/cesp.11.2.9>
25. Rondón Benítez AP, Bastidas ILO, Camargo YS. Factores que influyen en la deserción terapéutica de los consultantes de un centro universitario de atención psicológica. Int J Psychol Res [Internet]. 2009;2(2):137-47. Available from: <https://revistas.usb.edu.co/index.php/IJPR/article/view/869>
26. Fernández González OM. Alianza terapéutica, acciones comunicacionales e indicadores de cambio genéricos en la fase inicial de la psicoterapia con adolescentes, y su relación con resultados y adherencia al tratamiento [Doctoral dissertation]. [Santiago de Chile]: Pontificia Universidad Católica de Chile, Universidad de Chile; 2013. 189 p. Available from: <https://repositorio.uchile.cl/handle/2250/183295>
27. Bustos Martínez DA. Psicoterapia en adolescentes: contexto, motivaciones y expectativas en la primera consulta [Master's thesis]. [Santiago de Chile]. Universidad del Desarrollo; 2017. 120 p. Available from: <https://repositorio.udd.cl/items/b1cf579d-e17d-421e-a41a-146b8af7bcf6>
28. Fernández D. Contribuciones del Contra-Edipo parental en la clínica con adolescentes. Psicol USP [Internet]. 2021;32:1-9. doi: <https://doi.org/10.1590/0103-6564e180153>
29. Bleichmar ED. Manual de psicoterapia de la relación padres e hijos. Barcelona: Paidós Ibérica; 2005. 552 p.
30. Palacín F, Silva C, Krotter L. Factores Cognitivos en la Adherencia al Tratamiento. IX Congreso Internacional De Psiquiatría. 2002. Buenos Aires: AAP.
31. León S. El psicoterapeuta como persona: apuntes personales y profesionales. Santiago de Chile: RIL Editores; 2016. 232 p.
32. Finegan M, Firth N, Wojnarowski C, Delgadillo J. Associations between socioeconomic status and psychological therapy outcomes: A systematic review and meta-analysis. Depression and Anxiety [Internet]. 2018 Apr 26;35(6):560–73. doi: <http://dx.doi.org/10.1002/da.22765>
33. Clark CM. Irreducibly human encounters: Therapeutic alliance and treatment outcome in child and adolescent psychotherapy. J Infant Child Adolesc Psychother [Internet]. 2013;12(3):228-43. doi: <https://doi.org/10.1080/15289168.2013.822751>
34. Andrade ERC, Rivera MEL, Monterrosa ANF, de Orellana AL. Satisfacción de los pacientes con el servicio de atención psicológica de una clínica universitaria. Alerta (San Salvador) [Internet]. 2021;4(2):20-7. doi: <https://doi.org/10.5377/alerta.v4i2.10889>
35. Fernández Sanz S, Fernández Gonzalez O, Krause Jacob M. Expectativas de niños y padres consultantes frente a la psicoterapia infantil. CES Psicología [Internet]. 2021;14(2):48-70. doi: <https://doi.org/10.21615/cesp.5355>

36. Jo T, Almas S. Abandono del tratamiento en una consulta externa de psiquiatría y psicología clínica. *Revista Electrónica de PortalesMédicos.com* [Internet]. 2002;2-14. Available from: https://www.portalesmedicos.com/monografias/abandono_tratamiento/index.htm
37. Londoño J. Padres, niños y psicoterapia: La inclusión parental en psicoterapia infantil desde la representación de sus participantes [Internet]. 1st ed. Medellín: Universidad de Antioquia; 2021. 146 p. Available from: <http://www.jstor.org/stable/j.ctv2rcnq0g>
38. Fernández O, Pérez C, Gloger S, Krause M. Importancia de los Cambios Iniciales en la Psicoterapia con Adolescentes. *Terapia Psicológica* [Internet]. 2015;33(3):247-55. doi: <https://doi.org/10.4067/S0718-48082015000300008>
39. Corbella S, Botella L. La alianza terapéutica: historia, investigación y evaluación. *Anales de psicología*. 2003;19(2):205-21. Available from: <https://revistas.um.es/analesps/article/view/27671>
40. Cryan G, Quiroga S. Análisis de la función reflexiva-RF en sesiones de grupo de terapia focalizada para adolescentes violentos. *Interam J Psychol* [Internet]. 2016;50(2):275-87. Available from: <https://journal.sipsych.org/index.php/IJP/article/view/125>
41. Gómez C, Lama X, Capella C. Psicoterapia y Superación de Agresiones Sexuales: Mensajes de Niños/as y Adolescentes a Pares y Psicoterapeutas. *Psykhé* [Internet]. 2021;30(2):1-15 doi: <https://doi.org/10.7764/psykhe.2019.22385>
42. Weisz J R, Kazdin AE. The present and future of evidence-based psychotherapies for children and adolescents. In: Weisz JR, Kazdin AE, editors. *Evidence-based psychotherapies for children and adolescents*. New York: Guilford Press; 2017. p. 577-95.
43. Norka M, Dangerfield M. Tratamiento basado en la mentalización para niños: Aplicaciones clínicas en el contexto actual. 1st ed. Barcelona: Herder Editorial; 2023. 496 p.