



Iberoamerican Journal of Medicine

ISSN: 2695-5075

ISSN-L: 2695-5075

eestebanz@riojasalud.es

Hospital San Pedro

España

Oliván-Gonzalvo, Gonzalo
Harlequin ichthyosis
Iberoamerican Journal of Medicine, vol. 6, núm. 1, 2024, pp. 39-41
Hospital San Pedro
España

DOI: <https://doi.org/10.53986/ibjm.2024.0005>

Disponible en: <https://www.redalyc.org/articulo.oa?id=692076381008>

- Cómo citar el artículo
- Número completo
- Más información del artículo
- Página de la revista en redalyc.org

redalyc.org

Sistema de Información Científica Redalyc
Red de revistas científicas de Acceso Abierto diamante
Infraestructura abierta no comercial propiedad de la academia



**IBEROAMERICAN
JOURNAL OF
MEDICINE**

iberoamericanjm

Journal homepage: www.iberoamjmed.com

Clinical Image

Harlequin ichthyosis

Gonzalo Oliván-Gonzalvo^{a,*} 

^a Pediatric Service, Aragon Institute of Social Services, Government of Aragon, Zaragoza, Spain

ARTICLE INFO

Article history:

Received 29 September 2023

Received in revised form 30 October 2023

Accepted 12 November 2023

Keywords:

Harlequin ichthyosis

Non-syndromic congenital ichthyosis

Autosomal recessive genetic disorder

© 2024 The Authors. Published by Iberoamerican Journal of Medicine. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Ictiosis arlequin

INFO. ARTÍCULO

Historia del artículo:

Recibido 29 Septiembre 2023

Recibido en forma revisada 30 Octubre 2023

Aceptado 12 Noviembre 2020

Palabras clave:

Ictiosis arlequin

Ictiosis congénita no sindrómica

Enfermedad autosómica recesiva

© 2024 Los Autores. Publicado por Iberoamerican Journal of Medicine. Éste es un artículo en acceso abierto bajo licencia CC BY (<http://creativecommons.org/licenses/by/4.0/>).

HOW TO CITE THIS ARTICLE: Oliván-Gonzalvo G. Harlequin ichthyosis. Iberoam J Med. 2024;6(1):39-41. doi: 10.53986/ibjm.2024.0005.

* Corresponding author.

E-mail address: golivang@aragon.es

ISSN: 2695-5075 / © 2024 The Authors. Published by Iberoamerican Journal of Medicine. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

<https://doi.org/10.53986/ibjm.2024.0005>

A 3-month-old male diagnosed with Harlequin ichthyosis was admitted to the Child Protection Center referred from a hospital due to a risky social situation and parental inability to provide the basic and medical care he needs. He is currently receiving dermatological treatment (Acitetrina 5 mg/24 hours orally, barrier and moisturizing cream in the diaper area, and sterilized vaseline all over the body every 3-4 hours), ophthalmological treatment (Thealoz eye drops, single-dose Artific, Lipolac gel and vitalizing Blefarix every 3-4 hours and Hydrosorb dressings while sleep) and daily physiotherapy. He is fed with artificial breastfeeding on demand. The skin condition has improved considerably since birth, evolving towards a phenotype similar to severe congenital ichthyosiform erythroderma, and no ocular lesions have been detected. He is the second child of consanguineous parents (first cousins) originally from Morocco. Healthy 3-year-old sister. He was born preterm

(31 weeks) with an adequate weight for gestational age (1,880 grams) after a poorly controlled pregnancy. Apgar test 8/8. In the neonatal period, he was diagnosed and successfully treated for sepsis due to coagulase-negative staphylococcus, invasive aspergillosis, necrotizing enterocolitis, and mild bronchopulmonary dysplasia. At 2 months of age, he was diagnosed with bilateral hearing loss. At the time of admission, the infant presents shiny hyperkeratotic, cracked, peeling, and slightly fissured skin, bilateral ectropion of the upper eyelids, supraciliary madarosis, eclabium, hypoplastic and folded ears, hypoplasia of the nasal bones, and hypoplasia and contracture of the fingers of the hands (Figure 1). Weight, length, and head circumference are 4SD below average for age and sex. The rest of the physical examination is apparently normal. At 4 months of age, he was referred to the emergency hospital due to acute abdomen. He is



Figure 1: A 3-month-old male affected by Harlequin Ichthyosis was treated from birth with oral retinoid, dermatological and ophthalmological care, and physiotherapy. He presents shiny hyperkeratotic, cracked, peeling, and slightly fissured skin, bilateral ectropion of the upper eyelids, supraciliary madarosis, eclabium, hypoplastic and folded ears, hypoplasia of the nasal bones, and hypoplasia and contracture of the fingers of the hands.

diagnosed with perforation of the cecum. In the immediate postoperative period, he presented oligoanuria, with progressive clinical worsening in the following 48 hours. He underwent reoperation to rule out new perforation or acute complications, revealing significant ascites and devitalized tissue. In the immediate postoperative period, he presented refractory septic shock with anuria, dying 24 hours later.

Harlequin ichthyosis is the rarest and most severe form of non-syndromic congenital ichthyosis. This autosomal recessive genetic disorder affects around 1 in 300,000 births, more likely if consanguinity is present, and there is no significant association between sex and race [1, 2]. This condition is caused by loss-of-function mutations in the ABCA12 gene, which is essential for the transportation of lipids required for the skin's barrier function [2, 3]. Newborns with harlequin ichthyosis are characterized by thickened, dry "armor-like" plaques that give an appearance of 'scales' that are separated by deep fissures covering the entire surface of the body. The eyelids may be everted (ectropion), leaving the eyes and the area around them very susceptible to infection. The lips are pulled back by the dry skin (eclabium) giving a fish mouth appearance. The nose and ears may be very poorly developed or absent entirely. Hypoplasia is sometimes found in the fingers [1-5]. There is no cure for the condition and around half of those affected die within the first few months. The most common causes of death are sepsis and acute respiratory failure due to the plaques. Other complications can include premature birth, problems with body temperature, and dehydration [1, 2, 4]. Improved neonatal intensive care, antibiotics, early treatment with oral retinoids, constant moisturizing and protection of the skin and eyes, and

coordinated multidisciplinary management may improve survival and quality of life. Early oral retinoid therapy softens scales, encourages desquamation, and improves the eclabium and ectropion in a matter of weeks [1, 4, 5]. As happened in the present case, children who survive the neonatal period usually evolve to a less severe phenotype, resembling a severe congenital ichthyosiform erythroderma [4, 5].

1. CONFLICT OF INTERESTS

The authors have no conflict of interest to declare. The authors declared that this study has received no financial support.

2. REFERENCES

1. Shibata A, Akiyama M. Epidemiology, medical genetics, diagnosis and treatment of harlequin ichthyosis in Japan. *Pediatr Int*. 2015;57(4):516-22. doi: 10.1111/ped.12638.
2. Rajpopat S, Moss C, Mellerio J, Vahlquist A, Gånemo A, Hellstrom-Pigg M, et al. Harlequin ichthyosis: a review of clinical and molecular findings in 45 cases. *Arch Dermatol*. 2011;147(6):681-6. doi: 10.1001/archdermatol.2011.9.
3. Hotz A, Kopp J, Bourrat E, Oji V, Süßmuth K, Komlosi K, et al. Mutational Spectrum of the ABCA12 Gene and Genotype-Phenotype Correlation in a Cohort of 64 Patients with Autosomal Recessive Congenital Ichthyosis. *Genes (Basel)*. 2023;14(3):717. doi: 10.3390/genes14030717.
4. Tsivilika M, Kavvadas D, Karachrysafi S, Sioga A, Papamitsou T. Management of Harlequin Ichthyosis: A Brief Review of the Recent Literature. *Children (Basel)*. 2022;9(6):893. doi: 10.3390/children9060893.
5. Glick JB, Craiglow BG, Choate KA, Kato H, Fleming RE, Siegfried E, et al. Improved Management of Harlequin Ichthyosis With Advances in Neonatal Intensive Care. *Pediatrics*. 2017;139(1):e20161003. doi: 10.1542/peds.2016-1003.