



Iberoamerican Journal of Medicine

ISSN: 2695-5075

ISSN-L: 2695-5075

eestebanz@riojasalud.es

Hospital San Pedro

España

Arias Tamayo, Sebastian Alejandro; Lugo Pinilla, María Paula; Granados Zarate, Andres; Cuastumal Martínez, José Daniel; Casabon Pastrana, Juan Felipe; Tipon Mayorga, Jesus David; Bernal Rojas, Paula Alejandra; Arango Ospina, Cesar Mauricio; Gallego Gonzalez, Edwin Alejandro; Ortega Sierra, Michael

Evidence-Based Data Regarding the Use of Antibiotic-loaded Cement in Hip Arthroplasty. A bibliometric analysis of 50 years

Iberoamerican Journal of Medicine, vol. 6, núm. 2, 2024, pp. 60-68

Hospital San Pedro

España

DOI: <https://doi.org/10.53986/ibjm.2024.0013>

Disponible en: <https://www.redalyc.org/articulo.oa?id=692077149004>

- Cómo citar el artículo
- Número completo
- Más información del artículo
- Página de la revista en [redalyc.org](https://www.redalyc.org)

[redalyc.org](https://www.redalyc.org)

Sistema de Información Científica Redalyc

Red de Revistas Científicas de América Latina y el Caribe, España y Portugal

Proyecto académico sin fines de lucro, desarrollado bajo la iniciativa de acceso abierto



**IBEROAMERICAN
JOURNAL OF
MEDICINE**

iberoamericanjm

Journal homepage: www.iberoamjmed.com

Original article

Evidence-Based Data Regarding the Use of Antibiotic-loaded Cement in Hip Arthroplasty. A bibliometric analysis of 50 years

Sebastian Alejandro Arias Tamayo ^a, María Paula Lugo Pinilla ^b, Andres Granados Zarate ^c, José Daniel Cuastumal Martínez ^d, Juan Felipe Casabon Pastrana ^e, Jesus David Tipon Mayorga ^f, Paula Alejandra Bernal Rojas ^f, Cesar Mauricio Arango Ospina ^g, Edwin Alejandro Gallego Gonzalez ^h, Michael Ortega Sierra ^{i,*}

^a Unidad Central del Valle del Cauca, School of Health Sciences, Tuluá, Colombia

^b Universidad Militar Nueva Granada, School of Medicine, Bogotá, Colombia

^c Universidad de Santander, School of Medicine, Bucaramanga, Colombia

^d Universidad del Cauca, School of Medicine, Popayán, Colombia

^e Universidad Del Sinú, School of Medicine, Cartagena, Colombia

^f Fundación Universitaria Juan N. Corpas, School of Medicine, Bogotá, Colombia

^g Universidad de Ciencias Aplicadas y Ambientales, School of Medicine, Bogotá, Colombia

^h Institución Universitaria Visión de las Américas, School of Medicine, Pereira, Colombia

ⁱ Universidad Centroccidental Lisandro Alvarado - Hospital Central Antonio María Pineda, Barquisimeto, Venezuela

ARTICLE INFO

Article history:

Received 27 December 2023

Received in revised form 22
March 2024

Accepted 31 March 2024

Keywords:

Bone cements

Anti-bacterial agents

Hip replacement arthroplasty

Bibliometrics

ABSTRACT

Introduction: The use of antibiotic-loaded cement is an intraoperative tool that has demonstrated potential benefits in hip arthroplasty. However, the global landscape of research on this topic remains unknown. The objective of this study was to analyze the scientific growth, characteristics, and metrics of global and historical research on the use of antibiotic-loaded cement in hip arthroplasty.

Material and methods: A cross-sectional bibliometric study was conducted using Scopus as the data source.

Results: A total of 523 documents published between 1973 and 2023 were selected. 89.9% (n=470) of the production consisted of original articles, with 11.85% being multi-center. 84.1% of the authors have published a single article, followed by 10.3% who have published two articles. Research has predominantly focused on adults or the elderly, with an emphasis on outcome evaluation (notably reoperation), assessing various antibiotic agents, and frequently employing retrospective designs.

Conclusions: There has been a sustained increase in research on the use of antibiotic-loaded cement in hip arthroplasty over the last 50 years. The research trend has shifted towards the evaluation of adult or elderly patients, exploration of antimicrobial agents, techniques, and health outcomes, primarily using observational and retrospective designs. An emerging research-focus is the study of hip arthroplasty in rheumatoid arthritis patients.

* Corresponding author.

E-mail address: mortegas2021@gmail.com

ISSN: 2695-5075 / © 2024 The Authors. Published by Iberoamerican Journal of Medicine. This is an open access article under the CC BY license

(<http://creativecommons.org/licenses/by/4.0/>).

<https://doi.org/10.53986/ibjm.2024.0013>

Datos basados en la evidencia sobre el uso de cemento con antibióticos en la artroplastia de cadera. Análisis bibliométrico de 50 años

INFO. ARTÍCULO

Historia del artículo:

Recibido 27 Diciembre 2023

Recibido en forma revisada

22 Marzo 2024

Aceptado 31 Marzo 2024

Palabras clave:

Cementos para huesos

Antibacterianos

Artroplastia de reemplazo de cadera

Bibliometría

RESUMEN

Introducción: El uso de cemento cargado con antibiótico, es una herramienta intraoperatoria que ha demostrado potenciales beneficios en la artroplastia de cadera. Sin embargo, se desconoce el panorama global de la investigación sobre este tópico. El objetivo de este estudio fue analizar el crecimiento científico, características y métricas de la investigación global e histórica sobre el uso de cemento cargado con antibiótico en artroplastia de cadera.

Material y métodos: Estudio bibliométrico de corte transversal, que utilizó como fuente de datos la base Scopus.

Resultados: Se seleccionaron 523 documentos publicados entre 1973 y 2023. El 89,9% (n=470) de la producción consistió en artículos originales, siendo el 11,85% multicéntricos. El 84,1% de los autores han publicado un único artículo, seguido de un 10,3% que han publicado dos artículos. La investigación se ha centrado predominantemente en adultos o ancianos, con énfasis en la evaluación de resultados (especialmente la reoperación), evaluando diversos agentes antibióticos y empleando con frecuencia diseños retrospectivos.

Conclusiones: En los últimos 50 años se ha producido un aumento sostenido de la investigación sobre el uso de cemento cargado con antibióticos en la artroplastia de cadera. La tendencia de la investigación se ha desplazado hacia la evaluación de pacientes adultos o ancianos, la exploración de los agentes antimicrobianos, las técnicas y los resultados sanitarios, utilizando principalmente diseños observacionales y retrospectivos. Un nuevo foco de investigación es el estudio de la artroplastia de cadera en pacientes con artritis reumatoide.

© 2024 Los Autores. Publicado por Iberoamerican Journal of Medicine. Éste es un artículo en acceso abierto bajo licencia CC BY (<http://creativecommons.org/licenses/by/4.0/>).

HOW TO CITE THIS ARTICLE: Arias Tamayo SA, Lugo Pinilla MP, Granados Zarate A, Cuastumal Martínez JD, Casabon Pastrana JF, Tipon Mayorga JD, Bernal Rojas PA, Arango Ospina CM, Gallego Gonzalez EA, Ortega Sierra M. Evidence-Based Data Regarding the Use of Antibiotic-loaded Cement in Hip Arthroplasty. A bibliometric analysis of 50 years. Iberoam J Med. 2024;5(2):60-68. doi: 10.53986/ibjm.2024.0013.

1. INTRODUCTION

Hip fracture is a relatively common condition in healthcare practice, especially among older adults [1]. Considering the demographic transition occurring worldwide, this pathology is increasingly observed. Depending on the personalized approach and health phenotype of the patient, the decision to undergo surgical correction is determined [2]. Adverse events and post-intervention complications are some reasons why opting for arthroplasty may be avoided, as they significantly increase the need for reintervention, morbidity, mortality, and healthcare costs [2, 3]. Among these, infectious complications are often a significant cause in practice [4].

To address this difficulty, therapeutic tools for intraoperative use in orthopedics and traumatology have been designed, such as the use of antibiotic-loaded cement [5]. Previous evidence has suggested that using this technique has a prophylactic effect against periprosthetic

joint infection, potentially improving the outcome significantly and counteracting the risk of infectious complications, even in high-risk individuals [6, 7]. Despite reports from various authors indicating its increasing use, approaches, gaps, pluralism, and research ideas associated with its use in hip arthroplasty are still unknown to date. This lack of knowledge hinders the identification of gaps and opportunities that need to be investigated, especially in regions of the world where scientific production on the topic has not been prominent.

In order to provide novel evidence on the evolution and patterns related to research in this topic, the aim of this study was to analyze the scientific growth, characteristics, and metrics of global and historical research on the use of antibiotic-loaded cement in hip arthroplasty.

2. MATERIAL AND METHODS

2.1. DESIGN AND DATA SOURCE

A cross-sectional bibliometric study was conducted using the Scopus database as the data source. It is the database with the largest amount of peer-reviewed literature globally, which is a main reason for choosing this database as previously described [8, 9].

2.2. DATA SEARCH

A search strategy based on Medical Subject Headings (MeSH) terms and synonyms was designed and structured to identify scientific articles related to the use of antibiotic-loaded cement in hip arthroplasty, regardless of the typology and research approach. Initially, a pilot test was conducted to confirm that the results obtained derived from the stated objective and strategy. After this test, the following search was employed: TITLE-ABS-KEY("Hip Replacement Arthroplasties") OR TITLE-ABS-KEY("Hip Prosthesis Implantation") OR TITLE-ABS-KEY("Hip Prosthesis Implantations") OR TITLE-ABS-KEY("Hip Replacement Arthroplasty") OR TITLE-ABS-KEY("Total Hip Replacements") OR TITLE-ABS-KEY("Total Hip Replacement") OR TITLE-ABS-KEY("Total Hip Arthroplasty") OR TITLE-ABS-KEY("Total Hip Arthroplasties") AND TITLE-ABS-KEY("Anti Bacterial Agents") OR TITLE-ABS-KEY("Antibacterial Agents") OR TITLE-ABS-KEY("Antibacterial Agent") OR TITLE-ABS-KEY("Anti-Bacterial Compounds") OR TITLE-ABS-KEY("Anti Bacterial Compounds") OR TITLE-ABS-KEY("Anti-Bacterial Agent") OR TITLE-ABS-KEY("Anti Bacterial Agent") OR TITLE-ABS-KEY("Anti-Bacterial Compound") OR TITLE-ABS-KEY("Anti Bacterial Compound") OR TITLE-ABS-KEY("Bacteriocidal Agents") OR TITLE-ABS-KEY("Bacteriocidal Agent") OR TITLE-ABS-KEY("Bacteriocide") OR TITLE-ABS-KEY("Bacteriocides") OR TITLE-ABS-KEY("Anti-Mycobacterial Agents") OR TITLE-ABS-KEY("Anti Mycobacterial Agents") OR TITLE-ABS-KEY("Anti-Mycobacterial Agent") OR TITLE-ABS-KEY("Anti Mycobacterial Agent") OR TITLE-ABS-KEY("Antimycobacterial Agent") OR TITLE-ABS-KEY("Antimycobacterial Agents") OR TITLE-ABS-KEY("Antibiotics") OR TITLE-ABS-KEY("Antibiotic") AND TITLE-ABS-KEY("cement").

As inclusion criteria, any peer-reviewed document published in a scientific journal was considered. Exclusions encompassed conference abstracts, errata, books, book chapters, among others, that did not follow the regular peer-

review process. The search was executed on December 11, 2023, with results filtered by the labels "humans" and "journals." Considering the historical nature of the analysis, no specific time frame for article inclusion was defined.

2.3. DATA STANDARDIZATION

Following the above, data related to the domains of citation, bibliographic information, abstract, and keywords were exported in .CSV format for manual review and data standardization in Microsoft Office Excel 2016. Four authors conducted the review, eliminating documents unrelated to the topic based on title, abstract, and keywords. Subsequently, these authors manually reviewed the homogeneity of variables and data to avoid discrepancies in visual and bibliometric analysis. In the case of the manuscript typology variable, these were regrouped as follows: Original Articles (including any observational or experimental design with primary data or data reanalysis), Reviews (including narrative reviews, systematic reviews with or without meta-analysis, umbrella reviews, and scoping reviews), Case Reports (including both series and individual cases), and Correspondences (including letters to the editor, comments, editorials, correspondences per se, and other short manuscript typologies sharing structure and purpose).

2.4. STATISTICAL ANALYSIS

Once the data were standardized, visual, quantitative, and bibliometric analysis was conducted. Metrics and scientometric calculations were generated to assess the evolution, general characteristics, and impact of the global evidence on the use of antibiotic-loaded cement in hip arthroplasty. The bibliometrix package in R, which allows for the calculation of quantitative scientometric indicators and visualization of results (version 4.3.1) [10], was used for this analysis. Synonyms, errors, plurals, and variants were carefully grouped using a thesaurus to homogenize the analysis. Keywords, authors, and institutions were standardized accordingly.

Descriptive analysis and characterization of the scientific production were performed. Metrics such as h-index, g-index, and m-index were used to measure the impact of authors, institutions, and countries. Additionally, the frequency of cumulative citations was measured. Definitions and specifications for the use of these metrics in bibliometric studies have been previously described [11, 12]. Frequency and percentage calculations were carried out using Microsoft Office Excel 2016.

2.5. ETHICAL STATEMENTS ANALYSIS

This study did not require approval from an ethics committee, as it did not involve research on humans, biological models, or the use of medical records.

Table 1: Overview of the global research of antibiotic-loaded cement use in hip arthroplasty (n=523)

Variable	n (%)
Article type	
Original research	470 (89.9)
Review	52 (10)
Correspondence*	1 (0.1)
Authors	
Authorships	1951
Authors of documents with single authorship	31 (3.81)
Collaboration	
Single-authored articles	38
Co-authorships per article (average)	4,69
International co-authorship	62
Keywords	2723
Journals	172
Average age of article (years)	13
Average number of citations per paper	31
Annual growth	- (3.81)

Includes letters to the editor, editorials, and comments.

3. RESULTS

Initially, 576 documents were identified, of which 523 were selected following the application of inclusion and exclusion criteria, published between 1973 and 2023, establishing a 50-year research time frame. 89.9% (n=470) of the production corresponded to original articles, with an average of approximately five authors per article, and international collaboration accounted for 11.85%. The global average of citations received per document was 31, and the annual growth was 3.8% (Table 1). A sustained growth in publications on the use of antibiotic-loaded cement in hip arthroplasty was identified, with 2022 being the year with the highest production (Figure 1-A). In contrast to citations received over time, which have been fluctuating with an average below 10 (Figure 1-B). Applying Lotka's law, it was observed that 84.1% of the authors had published only one article, followed by 10.3% with two articles.

When assessing the journals where these documents have been published, the Journal of Arthroplasty emerged as the source with the highest number of published documents (n=64) (Figure 2-A), total citations received (2637 citations) (Figure 2-B), and the highest impact obtained (both in h-index and g-index) (Figures 2 C-D). On the contrary,

concerning the m-index, the highest value (m-index = 1) is shared by four journals (Figure 2-E).

Among the most prolific authors in the studied theme, Leif Ivar Havelin (affiliated with the Universitetet i Bergen, Norway) leads the list with 11 published documents. Additionally, he is the author with the highest number of citations, h-index, and g-index obtained. In terms of prolific affiliations, Haukeland University Hospital (located in Norway) leads, having published 18 documents to date and obtaining the highest h-index (although this last value has also been achieved by the Universitetet i Bergen). Despite this, the United States stands out as the most prolific and impactful country on the topic to date, with 143 published documents and an h-index of 46 (Table 2). Interestingly, Norway is not among the top five most prolific countries.

When exploring trends and the most studied lines regarding the use of antibiotic-loaded cement in hip arthroplasty, it was observed that research has predominantly focused on adults or the elderly, with an emphasis on outcome evaluation (highlighting reoperation). Various antibiotic agents are assessed, and retrospective designs are frequently employed (Figure 3-A). Additionally, there has been a certain interest, represented by the frequency of keywords in the form of bigrams, in studies related to the evaluation of revision at two moments, outcomes in the periprosthetic joint, and cement spacer (Figure 3-B). This is evident in the co-occurrence of words in the structuring of topics through titles, where essentially three groups of networks represent the highest frequency of research in the field (Figure 3-C). An emerging theme is the inclusion of patients with rheumatoid arthritis in this research field (Figure 3-D). Regarding the international collaboration network, it was identified that the United States and the United Kingdom lead each of their networks, both strongly linked with European and Asian countries (Figure 3-E). Chile was the only Latin American country identified with participation and/or collaboration in research on the analyzed topic (Figure 3-F).

Among the articles with the highest impact obtained to date, three original studies stand out: 1) "Management of deep infection of total hip replacement" (539 citations - The Journal of Bone & Joint Surgery British Volume – 1981 – DOI:10.1302/0301-620x.63b3.7021561) [13]; 2) "Antibiotic prophylaxis in total hip arthroplasty: Effects of antibiotic prophylaxis systemically and in bone cement on the revision rate of 22,170 primary hip replacements followed 0-14 years in the Norwegian Arthroplasty Register" (323 citations - Acta Orthopaedica Scandinavica – 2003 – DOI: 10.1080/00016470310018135) [14]; and 3) "Long-term elution of antibiotics from bone-cement: An in

vivo study using the prosthesis of antibiotic-loaded acrylic cement (PROSTALAC) system" (294 citations – The Journal of Arthroplasty – 1998 – DOI: 10.1016/S0883-5403(98)90179-6) [15].

impact [16], given the metrics used in this study. Nevertheless, it is anticipated that recently launched journals, such as Antibiotics, belonging to the Infectious Diseases category, may have an m-index of 1. This is




Figure 1: Scientific evolution of research on the use of antibiotic-loaded cement in hip arthroplasty. A: Annual scientific production from 1973 to 2023. B: Average number of citations received per article per year.


4. DISCUSSION

This study provides, for the first time, an overview of the global research landscape on the use of antibiotic-loaded cement in hip arthroplasty. In general, a sustained growth in publication volume has been identified since the 1970s, albeit with a relatively low impact (indicated by the low citation rate per article). However, this finding is subjective, considering that the impact varies across different fields of knowledge [16]. It was also observed that the highest publication frequency, by source, is in high-impact journals, primarily belonging to the Orthopedics category. The observed impact could be correlated with the number of published documents and citations obtained. As it is expected that a higher number of published documents increases the probability of citation accumulation and

because the impact, while lower than that of traditional journals, aligns with the publication date of the first document related to the topic. For this reason, providing more than one metric in the analysis of the impact of medical research is valuable.


There is a relevant finding concerning the volume of publications, continuity in research, and eventual impact obtained by the various evaluated actors. For example, two of the most prolific authors are from two European countries (Norway and the Netherlands) that do not stand out among the list of the most prolific countries (in terms of total publication volume). The same applies to affiliations, where two of the five most prolific are Norwegian. Although the publication volume of the most prolific authors and affiliations has been low for the evaluated time period (50 years), compared to other biomedical areas, the impact obtained could support continuity and depth in exploring a phenomenon over time. This aligns with expectations for

well as retrospective and health outcomes studies. The above would reflect the persistence of an observational and possibly descriptive work course in different nations, probably aimed at assessing the potential efficacy, safety, and cost-effectiveness of this intervention in different age groups and clinical phenotypes of patients. Although there is an interest in researching specific objectives such as variation in intervention times, the use of certain antimicrobials (such as vancomycin or gentamicin), or the presence of complications such as septic arthritis, compared



The foregoing could be explained by the broad and upward collaboration network worldwide in the research of the topic of interest, led by the United States and the United Kingdom, which could cause heterogeneity but complementation in different ideas [19] related to the use of antibiotic-loaded cement in hip arthroplasty. In contrast, Norway reflects endogamy with nearby countries, which could correlate with a closed circle of solid and gradual growth in work. Although the most frequent research in each of these niches was not discriminated, it was identified that predominantly the research in this field has focused on assessing elderly patients, possibly due to the frailty, risk, and frequency of falls in the elderly requiring surgical hip intervention, as

to other topics, there is little evidence of trends suggesting a more experimental or translational investigative activity, with the search for biomarkers or predictors. It was observed that rheumatoid arthritis and the use of antibiotic-loaded cement for hip replacement are an emerging, interesting, and relevant research niche, given the disease burden, functional deterioration, joint involvement, and the impact on the quality of life of patients with this condition [20].



have been published more frequently in other regions of the world. However, due to the methodology and statistical analysis performed, Scopus is one of the few databases that allows these analyses by the metadata deposited. Additionally, reporting the inherent bias of what is recorded in the metadata, which does not depend on the researchers of these types of studies but is favorably modified due to the standardized process of data review and purification.

Table 2: Production and impact of authors, affiliations and most prolific countries on research on the use of antibiotic-loaded cement in hip arthroplasty

Authors	Published documents	h-index	g-index	m-index	Total citations	Current affiliation	Country
Leif Ivar Havelin	11	11	11.3	0.3	1282	Universitetet i Bergen	Norway
Javad Parvizi	9	7	10.7	0.4	483	Rothman Orthopedic Institute	USA
Clive P Duncan	9	8	10	0.3	975	The University of British Columbia	Canada
Bassam Masri	9	8	10	0.3	975	The University of British Columbia	Canada
Bernard Wim Schreurs	9	5	7.2	0.2	253	Radboud University Medical Center	Netherlands
Affiliation	Documents over time				Total documents	h-index	Country
	1973–1993	1994–2003	2004–2013	2014–2023			
Haukeland University Hospital	0	7	6	5	18	13	Norway
Universitetet i Bergen	0	8	4	3	15	13	Norway
Mayo Clinic	1	1	1	10	13	7	USA
Helios ENDO-Klinik Hamburg	3	2	1	7	13	9	Germany
Affiliation	Documents over time				Total documents	h-index	
	1973–1993	1994–2003	2004–2013	2014–2023			
USA	19	23	30	77	149	46	
United Kingdom	10	9	20	32	71	26	
Germany	11	3	8	27	49	17	
China	0	0	10	27	37	6	
Italy	1	3	11	15	30	15	

5. CONCLUSIONS

There has been a sustained increase in research on the use of antibiotic-loaded cement in hip arthroplasty over the last 50 years, primarily led by the United States and the United Kingdom. However, the most prolific contributors hail from various countries, with Norway standing out. The impact achieved by this research has been relatively low compared to other biomedical areas. It was determined that the research trend has focused on the assessment of adult or elderly patients and the exploration of antimicrobial agents, techniques, and health outcomes. This is primarily done using observational and retrospective designs, with the emerging study of rheumatoid arthritis linked to hip arthroplasty.

6. CONFLICT OF INTERESTS

The authors have no conflict of interest to declare. The authors declared that this study has received no financial support.

7. REFERENCES

- 1.Feng JN, Zhang CG, Li BH, Zhan SY, Wang SF, Song CL. Global burden of hip fracture: The Global Burden of Disease Study. *Osteoporos Int.* 2024;35(1):41-52. doi: 10.1007/s00198-023-06907-3.
- 2.Schattner A. The burden of hip fractures-why aren't we better at prevention? *QJM.* 2018;111(11):765-7. doi: 10.1093/qjmed/hcx216. PMID: 29165606.
- 3.Dong Y, Zhang Y, Song K, Kang H, Ye D, Li F. What was the Epidemiology and Global Burden of Disease of Hip Fractures From 1990 to 2019? Results From and Additional Analysis of the Global Burden of Disease Study 2019.

Clin Orthop Relat Res. 2023;481(6):1209-20. doi: 10.1097/CORR.0000000000002465.

4. Masters J, Metcalfe D, Ha JS, Judge A, Costa ML. Surgical site infection after hip fracture surgery: a systematic review and meta-analysis of studies published in the UK. *Bone Joint Res.* 2020;9(9):554-62. doi: 10.1302/2046-3758.99.BJR-2020-0023.R1.

5. Szymiski D, Walter N, Krull P, Melsheimer O, Lang S, Grimberg A et al. The Prophylactic Effect of Single vs. Dual Antibiotic-Loaded Bone Cement against Periprosthetic Joint Infection Following Hip Arthroplasty for Femoral Neck Fracture: An Analysis of the German Arthroplasty Registry. *Antibiotics (Basel).* 2023;12(4):732. doi: 10.3390/antibiotics12040732.

6. Agni NR, Costa ML, Achten J, Peckham N, Dutton SJ, Png ME, et al. High-dose dual-antibiotic loaded cement for hip hemiarthroplasty in the UK (WHiTE 8): a randomised controlled trial. *Lancet.* 2023;402(10397):196-202. doi: 10.1016/S0140-6736(23)00962-5.

7. Bingham J. When and How Should I Use Antibiotic Cement in Primary and Revision Joint Arthroplasty? *J Arthroplasty.* 2022;37(8):1435-7. doi: 10.1016/j.arth.2022.02.001.

8. Lozada-Martinez ID, Visconti-Lopez FJ, Marrugo-Ortiz AC, Ealo-Cardona CI, Camacho-Pérez D, Picón-Jaimes YA. Research and Publication Trends in Pediatric Surgery in Latin America: A Bibliometric and Visual Analysis from 2012 to 2021. *J Pediatr Surg.* 2023;58(10):2012-9. doi: 10.1016/j.jpedsurg.2023.04.003.

9. Yubo S, Ramayah T, Hongmei L, Yifan Z, Wenhui W. Analysing the current status, hotspots, and future trends of technology management: Using the WoS and scopus database. *Heliyon.* 2023;9(9):e19922. doi: 10.1016/j.heliyon.2023.e19922.

10.10. Aria M, Cuccurullo C. Bibliometrix: An R-Tool for Comprehensive Science Mapping Analysis. *J Informetr.* 2017; 11:95-975. doi: 10.1016/j.joi.2017.08.007.

11. Rousseau R, Egghe L, Guns R. *Becoming Metric-Wise: A Bibliometric Guide for Researchers.* 1st ed. Belgium: Chandos Publishing; 2018.

12. Todeschini R, Baccini A. *Handbook of Bibliometric Indicators: Quantitative Tools for Studying and Evaluating Research.* 1st ed. Italy: Wiley-VCH Verlag GmbH & Co. KGaA; 2016. doi: 10.1002/9783527681969.

13. Buchholz HW, Elson RA, Engelbrecht E, Lodenkämper H, Röttger J, Siegel A. Management of deep infection of total hip replacement. *J Bone Joint Surg Br.* 1981;63-B(3):342-53. doi: 10.1302/0301-620X.63B3.7021561.

14. Engesaeter LB, Lie SA, Espehaug B, Furnes O, Vollset SE, Havelin LI. Antibiotic prophylaxis in total hip arthroplasty: effects of antibiotic prophylaxis systemically and in bone cement on the revision rate of 22,170 primary hip replacements followed 0-14 years in the Norwegian Arthroplasty Register. *Acta Orthop Scand.* 2003;74(6):644-51. doi: 10.1080/00016470310018135.

15. Masri BA, Duncan CP, Beauchamp CP. Long-term elution of antibiotics from bone-cement: an in vivo study using the prosthesis of antibiotic-loaded acrylic cement (PROSTALAC) system. *J Arthroplasty.* 1998;13(3):331-8. doi: 10.1016/S0883-5403(98)90179-6.

16. Greenhalgh T, Raftery J, Hanney S, Glover M. Research impact: a narrative review. *BMC Med.* 2016;14:78. doi: 10.1186/s12916-016-0620-8.

17. García Ancira C. [Evaluation of Scientific Impact Assessment in Educational Researches through a Case Study]. *Rev Cuba Educ Super.* 2022;41(2):523-33.

18.18. Cañedo-Andalia R, Cruz-Font J, Nodarse-Rodríguez M. [Impact of Cuban's Health Research Published in National Journals: are there reasons for urgent intervention?]. *Rev Cuban Cardiol.* 2015;21(2):1-7.

19. Vaudano E. Research Collaborations and Quality in Research: Foes or Friends? *Handb Exp Pharmacol.* 2020;257:383-98. doi: 10.1007/164_2019_293.

20. Anaya JM. The diagnosis and clinical significance of polyautoimmunity. *Autoimmun Rev.* 2014;13(4-5):423-6. doi: 10.1016/j.autrev.2014.01.049.

21. Lozada-Martinez ID, Bolaño-Romero MP, Picón-Jaimes YA, Moscote-Salazar LR, Narvaez-Rojas AR. Quality or quantity? Questions on the growth of global scientific production. *Int J Surg.* 2022;105:106862. doi: 10.1016/j.ijsu.2022.106862.

22. Lozada-Martinez ID, Suarez-Causado A, Solana-Tinoco JB. Ethnicity, genetic variants, risk factors and cholelithiasis: The need for eco-epidemiological studies and genomic analysis in Latin American surgery. *Int J Surg.* 2022;99:106589. doi: 10.1016/j.ijsu.2022.106589.