

Relations between Spectrum Curves of Discrete Sturm-Liouville Problem with Nonlocal Boundary Conditions and Graph Theory. II

Vitkauskas, Jonas; Štikonas, Artūras

Relations between Spectrum Curves of Discrete Sturm-Liouville Problem with Nonlocal Boundary Conditions and Graph Theory. II

Lietuvos matematikos rinkinys, vol. 62 Ser. A, 2021

Vilniaus Universitetas, Lituania

Disponible en: <https://www.redalyc.org/articulo.oa?id=692674114005>

DOI: <https://doi.org/10.15388/LMR.2021.25128>



Esta obra está bajo una Licencia Creative Commons Atribución 4.0 Internacional.

Relations between Spectrum Curves of Discrete Sturm-Liouville Problem with Nonlocal Boundary Conditions and Graph Theory. II

Diskrečiojo Šturmo ir Liuvilio uždavinio su nelokaliosiomis kraštinėmis sąlygomis spektrinių kreivių ir grafų teorijos sąsajos. II

Jonas Vitkauskas vitkauskas@mif.stud.vu.lt
Vilnius University, Lithuania

 <https://orcid.org/0000-0003-1820-4557>
Artūras Štikonas arturas.stikonas@mif.vu.lt
Vilnius University, Lithuania

 <https://orcid.org/0000-0002-5872-5501>

Lietuvos matematikos rinkinys, vol. 62
Ser. A, 2021

Vilniaus Universitetas, Lithuania

Recepción: 23 Julio 2021
Publicación: 01 Diciembre 2021

DOI: <https://doi.org/10.15388/LMR.2021.25128>

Redalyc: <https://www.redalyc.org/articulo.oa?id=692674114005>

Abstract: In this paper, relations between discrete Sturm–Liouville problem with nonlocal integral boundary condition characteristics (poles, critical points, spectrum curves) and graphs characteristics (vertices, edges and faces) were found. The previous article was devoted to the Sturm–Liouville problem in the case two-points nonlocal boundary conditions.

Keywords: Sturm–Liouville problem, spectrum curves, integral boundary condition, graphs.

Summary: Šiame straipsnyje pristatomos sąsajos tarp diskrečiojo Šturmo ir Liuvilio uždavinio su nelokaliąja integraline kraštine sąlyga (poliai, kritiniai taškai ir spektrinės kreivės) bei grafų charakteristikų (viršūnės, briaunos ir veidai). Ankstesnis straipsnis buvo skirtas Šturmo ir Liuvilio uždaviniui su dvitaškinėmis nelokaliosiomis kraštinėmis sąlygomis.

Keywords: Šturmo ir Liuvilio uždavinys, spektrinės kreivės, integralinė kraštine sąlyga, grafai.

1 A discrete Sturm–Liouville Problem

In this paper, particular properties of the spectrum of a discrete Sturm–Liouville Problem (dSLP) [3, 4, 5] with Integral Boundary Condition were found using Euler’s characteristic formula [2]. In previous article [5] we have found relations between spectrum curve properties and graphs theory in the case of two-points nonlocal boundary conditions.

We introduce a uniform grid and we use notation $\bar{\omega}^h = \{t_j = jh, j = 0, \dots, n, nh = 1\}$ for $2 < n \in \mathbb{N}$ and $N^h := (0, n) \cap \mathbb{N}$, $\bar{N}^h = N^h \cup \{0, n\}$. Also, we make an assumption that ξ_1 and ξ_2 are located on the grid i.e., $\xi_1 = m_1 h = m_1/n$, $\xi_2 = m_2 h = m_2/n$, $m \in S_{\xi}^h := \{(m_1, m_2) : 0 \leq m_1 < m_2 \leq n, m_1, m_2 \in \bar{N}^h\}$

Let $\xi = m/n, \xi = (m_1/n, m_2/n), \xi = \xi_1/\xi_2 = m_+/n, \xi_- = \xi_1 - \xi_2 = m_-/n$, here $m_+ := m_1 + m_2, m_- = m_2 - m_1$. We denote $N_{odd} = \{k \in \mathbb{N} : k - odd\}, N_{even} = \{k \in \mathbb{N} : k - even\}$

Let us consider a dSLP (an approximation by Finite-Difference Scheme) [3, 5]

$$\frac{U_{i+1} - 2U_i + U_{i-1}}{h^2} = \lambda U_i, i = 1, \dots, n-1 \quad (1)$$

$\lambda \in \mathbb{C}$ with a classical discrete Dirichlet Boundary Condition (BC)

$$U_0 = 0 \quad (2)$$

and Integral Boundary Condition (approximated by trapezoidal formula):

$$U_n = \gamma h \left(\frac{U_{m_1} + U_{m_2}}{2} + \sum_{k=m_1+1}^{m_2-1} U_k \right) \quad (3)$$

Let us consider a bijection (see [1])

$$\lambda = \lambda^h(q) := \frac{4}{h^2} \sin^2 \frac{\pi q h}{2} \quad (4)$$

between $C_\lambda := \mathbb{C}$ and $C_q^h := R_q^h \cup C_q^{h-}, R_q^h := R_y^- \cup \{0\} \cup R_x^h \cup \{n\} \cup R_y^{h+}, R_y^- := \{q = iy : y > 0\}, R_x^h := \{q = x : 0 < x < n\}, R_q^{h+} := \{q = n + iy : y > 0\}, C_q^{h+} := \{q = x + iy : 0 < x < n, y > 0\}, C_q^{h-} := \{q = x + iy : 0 < x < n, y < 0\}.$

The general solution U_j for a discrete equation (1) is equal to:

$$U_j = C_1 \sin(\pi q t_j) (1 - hq)^{-1\pi - 1q - 1} + C_2 \cos(\pi q t_j)$$

Then by using BCs (2) and (3) we get an equation:

$$Z^h(q) = \gamma P_\xi^h(q), q \in C_q^h$$

where functions $Z^h(q)$ and $P_\xi^h(q)$ are as follows:

$$Z^h(q) = \frac{\sin(\pi q)}{\pi q} \cdot \frac{\sin(\frac{\pi}{2} q h)}{\pi q h \cos(\frac{\pi}{2} q h)}, \left(q \right) = \frac{\sin(\frac{\pi}{2} q (\xi_2 - \xi_1))}{\pi q} \cdot \frac{\sin(\frac{\pi}{2} q (\xi_2 + \xi_1))}{\pi q}$$

Constant Eigenvalues. For any constant eigenvalue $\lambda \in \mathbb{C}$ there exists the Constant Eigenvalue Point (CEP) $q \in \mathbb{C}$. CEP are roots of the system [1]:


$$Z^h(q) = 0, P_{\xi}^h(q) = 0$$

For every CEP c_j we define nonregular Spectrum Curve $N_j = \{c_j\}$


Nonconstant eigenvalues. Let us consider Complex Characteristic Function:

$$\gamma_c(q, \xi) := \frac{Z^n(q)}{P_{\xi}^h(q)}, q \in \mathbb{C}_q. \quad (5)$$


All nonconstant eigenvalues (which depend on the parameter γ) are γ -points of (Complex-Real) Characteristic Function (CF)[6]. CF $\gamma(q)$ is the restriction of Complex CF $\gamma_c(q)$ on a set $D_{\xi} := \{q \in \mathbb{C}_q^h : \text{Im} \text{ (see more in [5])}. We call such$



(a) $m = (0, 2)$



(b) $m = (1, 2)$



(c) $m = (0, 1)$

Fig. 1.
Spectrum Curves for $n = 2$ [4]




Fig. 2.
Spectrum Curves for $n = 3$ [4].




Fig. 3.
Spectrum Curves for $n = 4$ [4].




Fig. 4.
Spectrum Curves for $n = 5$ [4].

There exist Pole Points (PP) of the second order. They are described as follows:

$$p_k^{12} = 2nk / \gcd(m_+, m_-, k = 1, \dots, [n / p_1^{12}]).$$

All PP of the second order are in R_x^h . Note that there could be a PP of second order in Ramification Point $q = n$, so they become simple PP of the first order.

Two or more Spectrum Curves may intersect at CP. We denote a set $CP\beta = \{b_i, i = 1, \dots, n_b\}$, where n_b is the number of CPs at C_q^h . The number of CPs at R_q^h and C_q^{h+} we denote as n_{cr} and n_{cr}^+ , respectively. Note that the part of the spectrum domain in set C_q^{h+} is symmetric to the part in set C_q^{h-} . So, $n_b = n_{cr} + 2n_{cr}^+$. If $b \in \beta$ then $\deg^+(b)$ is one unit larger than the order of this CP.

The pole at $q = \infty$ is of

$$n_\infty = n - m_2 \quad (6)$$

order. If $m_2 < n$, $q = \infty$ is a PP. For $m_2 = n$, $q = \infty$ is a Removable Singularity Point.

For poles and CP $\deg^+(q)$, $q \in P \cup B \cup \{\infty\}$, corresponds to the number of outgoing Spectrum Curves at that point. Note that incoming Spectrum Curves alternate with outgoing, so $\deg^+(q) = \deg^-(q)$

2 Relations between dSLP and graphs properties in the case of integral BC

It is possible to define relations between properties of dSLP and graph theory. Poles or CPs refer to vertices of a certain graph and parts of Spectrum Curves could be interpreted as edges. In our case, we have a simple balanced weakly connected digraph. Definitions and notations in graphs theory are described in [5].




Fig. 5.
Spectrum Curves for $n = 6$ [4].

2.1 Properties of Spectrum Curves

Poles, CPs, regular and nonregular Spectrum Curves, CEPs were found by A. Skučaitė [4].

There is $n - 1$ Spectrum Curves for every $n \in \mathbb{N}$, $n \geq 2$. Nonregular Spectrum Curves are CEPs and belong to $R_x^h = (0, n)$. The number of such Spectrum Curves is equal to

$$n_{ce} = \left[\frac{n-1}{2n} dbd(2n, m_+) \right] + \left[\frac{n-1}{2n} dbd(2n, m_-) \right] - \left[\frac{n-2}{2n} dbd(2n, m_+, m_-) \right] \quad (7)$$

Number of regular Spectrum Curves $n_{nce} = n - 1 - n_{ce}$. The poles of CF belong to $R_x^h \cup \{0\} \cup \{n\} \cup \{\infty\}$ and $n_p + n_\infty = n_{nce}$. So, we have formula

$$n_p + n_{ce} = m_2 - 1 \quad (8)$$

Let us denote

$$\deg_r^+ = \sum_{b \in B \cap R_q^h} \deg^+(b), \deg_c^+ = \sum_{b \in B \cap C_q^{h\pm}} \deg^+(b) = 2 \sum_{b \in C_q^{h\pm}} \deg^+(b).$$

Let n_c is the number of Spectrum Curves parts in $C_q^{h\pm}$ between two CP (including $q = 0$ and $q = n$).

2.2 Spectrum domain as a graph

We consider Spectrum domain as graph on sphere (Riemann sphere C^-) because Ch S2. The poles and CPs of the CF are the vertices of this graph. The point is the pole or CP.

Lemma 1. The number of vertices is

$$\begin{aligned} u &= n_p - n_{2p} + n_b + 1 - [m_2 \text{ } n] \\ &= n_p - n_{2p} + n_{cr} + 2n_{cr}^+ + 1 - [m_2 \text{ } n] \\ &= m_2 - n_{ce} - n_{2p} + n_{cr} + 2n_{cr}^+ - [m_2 \text{ } \emptyset] \end{aligned}$$

Lemma 2. The number of edges is

$$e = n_p + \deg_r^+ + \deg_c^+ + n_\infty. \quad (10)$$

Lemma 3. The number of faces is

$$(11)$$

This lemma is valid for $n_c = n_{cr}^+ = 0$. Each part of spectrum curve between two CPs $b_1, b_2 \in R_q^h$ increases the number of faces by one. So, this formula is valid for the case $n_{cr}^+ = 0$. Each additional CP $CPb \in C_q^{h\pm}$ increases the number of faces by $2(\deg^+(b) - 1)$ and number parts of Spectrum Curves between this CP and other CPs by $2\deg^+(b) - 1$.

Numbers of spectrum vertices, edges and faces, expressed by the formulas above, inserted to the Euler's characteristic's formula of sphere $u - e + f = 2$ give new relation.

Theorem 1. The Euler's characteristic's formula is equivalent to

$$\begin{aligned} \sum_{b \in B} \deg^+(b) &= \deg_c^+ + \deg_r^+ = n_\infty + n_{2p} + 2n_c + n_{cr} - 1 + [m_2 / n] \\ &= n - m_2 + n_{2n} + 2n_c + n_{cr} - 1 + [m_2 \text{ } n] \end{aligned} \quad (12)$$

This formula was derived in [4] when are not CPs in $C_q^{h\pm}$ ($\deg_c^+ = 0$), all CPs are of the first order ($\deg_r^+ = 2n_{cr}$) and $n_c = 0$. Then it can be rewritten as

$$n_{cr} = n_{2p} + n_{\infty} - 1 + \lfloor m_2/n \rfloor = n_{2n} + n - m_2 - 1 + \lfloor m_2/n \rfloor$$

Corollary 1. The number of edges is

$$\begin{aligned} e &= 2n_{\infty} + n_p + n_{2p} + n_{cr} + 2n_c - 1 + \lfloor m_2/n \rfloor \\ &= 2n - m_2 - 2 - n_{ce} + n_{2p} + n_{cr} + 2n_c + \lfloor m_2/n \rfloor \end{aligned} \quad (13)$$

Remark 1. In the case $m_2 < n$ the formulas (9)–(13) are

$$\begin{aligned} u &= m_2 - n_{ce} - n_{2p} + n_{cr} + 2n_c^+, \\ e &= 2n - m_2 - 2 - n_{ce} + n_{2p} + n_{cr} + 2n_c \\ f &= 2(n - m_2 + n_{2p} + n_c - n_{cr}^+), \deg_r^+ + \deg_c^+ = n - m_2 - 1 + n_{2p} + 2n_c + n_{cr}, \end{aligned}$$

where n_{ce} is defined by (7).

In the case $m_2 = n$ we have $n_{\infty} = 0$, $n_p = n - 1 - n_{ce}$ and $n_{2p} = 0$, $n_{cr} = 0$ (so, $\deg_r^+ = 0$) (see [4]). In this case we get $\deg_c^+ = 2n_c = 0$ and $n_{cr}^+ = 0$. Thus, for are valid. Note, that in this case we have Cycle digraph $m_2 = n$ the following formulas

$$\begin{aligned} u &= n - n_{ce} \\ e &= n - n_{ce} \\ f &= 2 \end{aligned}$$

are valid. Note, that in this case we have Cycle digraph

Remark 2. If $n_{cr}^+ = 0$ ($\deg_c^+ = 0$) then $\deg_r^+ = n_{cr} = 2n_c + n - m_2 + n_{2p} - 1 > n_{cr}$ shows that there are exist CPs in R_q^h of the second or the higher order.

References

1. K. Bingele, A. Bankauskiene, A. Štikonas. Spectrum curves for a discrete Sturm–Liouville problem with one integral boundary condition. *Nonlinear Anal. Model. Control*, **24**(5):755–774, 2019. <https://doi.org/10.15388/NA.2019.5.5>.
2. E. Manstavičius. *Analizė ir tikimybinė kombinatorika*. TEV, Vilnius, 2007.
3. M. Sapagovas. *Diferencialinių lygčių kraštiniai uždaviniai su nelokaliosiomis sąlygomis*. Mokslo aidai, Vilnius, 2007.
4. A. Skučaitė. *Investigation of the spectrum for Sturm–Liouville problem with a nonlocal integral boundary condition*. PhD thesis, Vilnius University, 2016.
5. J. Vitkauskas, A. Štikonas. Relations between spectrum curves and discrete Sturm–Liouville problem with nonlocal boundary conditions and graph theory. *Proc. of the Lithuanian Mathematical Society*, **61**:1–6, 2020. <https://doi.org/10.15388/LMR.2020.22474>.
6. A. Štikonas, O. Štikonienė. Characteristic functions for Sturm–Liouville problems with nonlocal boundary conditions. *Math. Model. Anal.*, **14**(2):229–246, 2009. <https://doi.org/10.3846/1392-6292.2009.14.229-246>.