



Nonlinear Analysis: Modelling and Control
ISSN: 1392-5113
ISSN: 2335-8963
nonlinear@mii.vu.lt
Vilniaus Universitetas
Lituania

Sign-changing solutions for Kirchhoff-type problems involving variable-order fractional Laplacian and critical exponents

Lianga, Sihui; Molica Bisci, Giovanni; Zhang, Binlin

Sign-changing solutions for Kirchhoff-type problems involving variable-order fractional Laplacian and critical exponents

Nonlinear Analysis: Modelling and Control, vol. 27, núm. 3, 2022

Vilniaus Universitetas, Lituania

Disponible en: <https://www.redalyc.org/articulo.oa?id=694173185009>


DOI: <https://doi.org/10.15388/namc.2022.27.26575>




Esta obra está bajo una Licencia Creative Commons Atribución 4.0 Internacional.

Sign-changing solutions for Kirchhoff-type problems involving variable-order fractional Laplacian and critical exponents


Sihu Lianga ^{a 1} liangsihua@163.com
Changchun Normal University, China

 <https://orcid.org/0000-0002-4260-7762>

Giovanni Molica Bisci ^b giovanni.molicabisci@uniurb.it
Università degli Studi di Urbino Carlo Bo, Italia

 <https://orcid.org/0000-0003-4802-1727>

Binlin Zhang ^{c 2 3} zhangbinlin2012@163.com
University of Science and Technology, China

 <https://orcid.org/0000-0003-1481-7313>

Nonlinear Analysis: Modelling and Control, vol. 27, núm. 3, 2022

Vilniaus Universitetas, Lituania

Recepción: 12 Abril 2021
Revisado: 27 Noviembre 2021
Publicación: 28 Marzo 2022

DOI: <https://doi.org/10.15388/namc.2022.27.26575>

Redalyc: <https://www.redalyc.org/articulo.oa?id=694173185009>

Abstract: In this paper, we are concerned with the Kirchhoff-type variable-order fractional Laplacian problem with critical variable exponent. By using constraint variational method and quantitative deformation lemma we show the existence of one least energy solution, which is strictly larger than twice of that of any ground state solution.

Keywords: Kirchhoff-type problem, variable-order fractional Laplacian, variational method, signchanging solution.

1 Introduction and main results

In this paper, we are interested in the existence of least energy nodal solutions for the following Kirchhoff-type variable-order fractional Laplacian problems with critical variable growth:

$$\begin{aligned} (a + b[u]_{s(\cdot)}^2)(-\Delta)^{s(\cdot)}u &= |u|^{q(x)-2}u + \lambda f(x, u) \quad \text{in } \Omega, \\ u &= 0 \quad \text{in } \mathbb{R}^N \setminus \Omega, \end{aligned} \quad (1)$$

where

$$[u]_{s(\cdot)} := \left(\iint_{\mathbb{R}^{2N}} \frac{|u(x) - u(y)|^2}{|x - y|^{N+2s(x,y)}} dx dy \right)^{1/2},$$

$a, b \geq 0, s(\cdot) : \mathbb{R}^N \times \mathbb{R}^N \rightarrow (0, 1)$ is a continuous function, Ω is a bounded domain in \mathbb{R}^N with Lipschitz boundary, $\lambda > 0$ is a parameter, $N > 2s(x, y)$ for all $(x, y) \in \Omega \times \Omega$, $(-\Delta)^{s(\cdot)}$ is the variable-order fractional Laplace operator, $4 < q(x) \leq 2^*(x) := 2N/(N - 2s(x, x))$ for all $x \in \Omega$. The variable-order fractional Laplace operator $(-\Delta)^{s(\cdot)}$ is defined as follows:

$$(-\Delta)^{s(\cdot)} \varphi(x) = 2PV \int_{\mathbb{R}^N} \frac{\varphi(x) - \varphi(y)}{|x - y|^{N+2s(x,y)}} dy$$

along any $\varphi \in C_0^\infty(\Omega)$, where PV denotes the Cauchy principle value. As $s(\cdot) \equiv \text{const}$, the variable-order fractional Laplace operator $(-\Delta)^{s(\cdot)}$ reduces to the usual fractional Laplace operator; see [14,15] for the concise introduction to the fractional Laplace operator and related variational results.

We now impose the assumptions on the functions $s(\cdot)$ and f that will in full force throughout the paper. Firstly, we suppose that $s : \mathbb{R}^N \times \mathbb{R}^N \rightarrow (0, 1)$ is a continuous function satisfying the following assumptions:

$$(s1) \quad 0 < s_- := \min_{(x,y) \in \mathbb{R}^N \times \mathbb{R}^N} s(x, y) \leq s_+ := \max_{(x,y) \in \mathbb{R}^N \times \mathbb{R}^N} s(x, y) < 1;$$

$$(s2) \quad s(\cdot) \text{ is symmetric, that is, } s(x, y) = s(y, x) \text{ for all } (x, y) \in \mathbb{R}^N \times \mathbb{R}^N.$$

From now on, for the variable exponents m , we set

$$\underline{m} = \text{ess inf}_{x \in \Omega} m(x), \quad \overline{m} = \text{ess sup}_{x \in \Omega} m(x).$$

Moreover, we suppose that $f \in C^1(\mathbb{R}, \mathbb{R})$ satisfies the following conditions:

$$(f1) \quad \lim_{t \rightarrow 0} f(x, t)/|t|^3 = 0;$$

$$(f2) \quad \text{there exist } \theta(x) \in (4, 2^*(x)) \text{ and } C > 0 \text{ such that } |f(x, t)| \leq C(1 + |t|^{\theta(x)-1})$$

for all $t \in \mathbb{R}$ and all $x \in \Omega$; for all $\cdot \in \mathbb{R}$ and all $\cdot \in \cdot$;

$$(f3) \quad f(x, t)/|t|^3 \text{ is a strictly increasing function of } t \in \mathbb{R} \setminus \{0\}.$$

A typical example of function fulfilling hypotheses (f1)–(f3) is as follows:

$$f(x, t) = |t|^{\theta(x)-2}t, \text{ where } t \in \mathbb{R} \text{ and } x \in \Omega.$$

The main driving force for studying problem (1) includes two aspects. On the one hand, when $s(\cdot) = 1$ Eq. (1) reduces to the general Kirchhoff-type model. Recently, some researchers also explored such equations in the study of nonlinear vibrations theoretically or experimentally. For example, Carrier [1] used a more rigorous method to deduce a more general Kirchhoff model. Moreover, the nonlocal Kirchhoff problems of parabolic type can model several biological systems such as population density; see, for instance, [4]. In fact, the energy functionals of (1) have obviously different properties from the case $b = 0$, and thus, several mathematical difficulties arise naturally in the study of the case $b \neq 0$ by variational and topological methods. It is worth mentioning that Fiscella and Valdinoci [9] deduced a new Kirchhoff model involving the

fractional Laplacian by considering the nonlocal aspect of the tension arising from nonlocal measurements of the fractional length of the string; see [9, App. A] for more details.

In recent years, finding sign-changing solutions to the Kirchhoff-type problems has been an attractive subject, and many interesting results have been obtained. In the following, let us sketch some advances related to the subject of our paper. Concerning the advances of Kirchhoff-type problems in the bounded domains, Zhang and Perera in [26] applied the method of invariant sets of descent flow to investigate the existence of signchanging solutions for Kirchhoff-type problems; see also Mao and Zhang in [12] for more related results via similar approaches. Using the constraint variational methods, Shuai in [18] obtained that Kirchhoff-type problems has one least energy sign-changing solution u_b and the energy of u_b strictly larger than the ground state energy. After that, with the help of non-Nehari manifold method, Tang and Cheng in [19] generalized some results obtained in [18]; see also [2] for more general Kirchhoff-type function in this direction. In [20], Wang obtained the following results for Kirchhoff-type equation with critical growth by employing the constraint variational method and the quantitative deformation lemma: the existence of least energy sign-changing solutions u , and the energy of u is strictly larger than twice that of the ground state solutions. Concerning the advances in the abstract Kirchhoff framework, here we just review two papers as follows: by using the minimization argument and a quantitative deformation lemma, Figueiredo et al. in [7] investigated the existence of a sign-changing solution for the following Kirchhoff-type equation:

$$-M\left(\int_{\Omega} |\nabla u|^2 dx\right) \Delta u = g(u) \quad \text{in } \Omega, \quad u = 0 \quad \text{on } \partial\Omega,$$

where Ω is a bounded domain in

where Ω is a bounded domain in \mathbb{R}^3 , $M: \mathbb{R}_0^+ \rightarrow \mathbb{R}$ is a continuous function with some appropriate assumptions, and g is a superlinear C^1 class function with subcritical growth. In unbounded domains, Figueiredo and Santos Júnior in [8] obtained a least energy signchanging solution to a class of nonlocal Schrödinger–Kirchhof problems involving only continuous functions by using a minimization argument and a quantitative deformation lemma. Moreover, the authors also proved that the problem has infinitely many nontrivial solutions when it presents symmetry. In [3], Cheng and Gao studied the following Kirchhoff-type problem, which involves a fractional Laplacian operator:

$$\left(a + b \iint_{\mathbb{R}^{2N}} \frac{|u(x) - u(y)|^2}{|x - y|^{N+2s}} dx dy\right) (-\Delta)^s u + V(x)u = f(x, u) \quad \text{in } \mathbb{R}^N,$$

where $0 < s < 1$ is a constant, f satisfies subcritical growth. The authors proved the existence of least energy sign-changing solutions for this

problem by using the constraint variation method and quantitative deformation lemma.

On the other hand, variable-order fractional Laplacian problems was introduced by Xiang et al. in [25]. They studied the following variable-order fractional Laplacian problems involving variable exponents:

$$\begin{aligned} (-\Delta)^{s(\cdot)} u + \lambda V(x)u &= \alpha |u|^{p(x)-2} u + \beta |u|^{q(x)-2} u \quad \text{in } \Omega, \\ u &= 0 \quad \text{in } \mathbb{R}^N \setminus \Omega. \end{aligned} \quad (2)$$

Under some suitable assumptions, they showed that problem (2) admits at least two distinct solutions by applying the mountain pass theorem and Ekeland's variational principle. Subsequently, Wang and Zhang in [21] proved the existence of infinitely many solutions for possibly degenerate Kirchhoff-type variable-order fractional Laplacian problems by using the new version of Clark's theorem due to Liu and Wang in [11]. Very recently, Xiang et al. in [24] obtained the existence of two solutions for a class of degenerate Kirchhoff-type variable-order fractional Laplacian problems by employing the Nehari manifold approach.

However, regarding the existence of sign-changing solutions for Kirchhoff-type variable-order fractional Laplacian problems involving variable exponents, there has been no paper in the literature as far as we know. Hence, a natural question is whether or not there exists sign-changing solutions of problem (1)? Another interesting question is whether or not the same conclusion still holds for critical exponent $q(x) = 2^*(x) := 2N/(N - 2s(x))$? The goal of the present paper is to give an affirmative answer.

The corresponding energy functional $\mathcal{I}_{b,\lambda} : H_0^{s(\cdot)}(\Omega) \rightarrow \mathbb{R}$ to problem (1) is defined by

$$\mathcal{I}_{b,\lambda}(u) = \frac{a}{2} [u]_{s(\cdot)}^2 + \frac{b}{4} [u]_{s(\cdot)}^4 - \lambda \int_{\Omega} F(x, u) \, dx - \int_{\Omega} \frac{1}{q(x)} |u|^{q(x)} \, dx. \quad (3)$$

It is standard to verify that $\mathcal{I}_{b,\lambda}$ belongs to $C^1(H_0^{s(\cdot)}(\Omega), \mathbb{R})$, and the critical points of $\mathcal{I}_{b,\lambda}$ are the solutions of problem (1). Furthermore, if we write $u^+(x) = \max\{u(x), 0\}$ and $u^-(x) = \min\{u(x), 0\}$ for $u \in H_0^{s(\cdot)}(\Omega)$, then every solution $u \in H_0^{s(\cdot)}(\Omega)$ problem (1) with the property that $u^{\pm} \neq 0$ is a sign-changing solution of problem (1).

Now, we give the following first main result.

Theorem 1. *Assume that (s1).(s2) and (f1).(f3) hold. Then there exists $\lambda^* > 0$ such that for all $\lambda \geq \lambda^*$, problem (1) has a least energy sign-changing solution u_b*

Another objective of this paper is to establish the so-called energy doubling property (cf. [22]), i.e., the energy of any sign-changing solution of problem (1) is strictly bigger than twice that of the ground state solution. We have the following result.

Theorem 2. *Assume that (s1).(s2) and (f1).(f3) hold. Then there exists $\lambda^{**} > 0$ such that for all $\lambda \geq \lambda^{**}$, $c^* := \inf_{u \in \mathcal{N}_{b,\lambda}} \mathcal{I}_{b,\lambda}(u) > 0$ is achieved and $\mathcal{I}_{b,\lambda}(u) > 2c^*$.*

where $\mathcal{N}_{b,\lambda} = \{u \in H_0^{s(\cdot)}(\Omega) \setminus \{0\} : \langle (\mathcal{I}_{b,\lambda})'(u), u \rangle = 0\}$, and u is the least energy sign-changing solution obtained in Theorem 1. In particular, $c^* > 0$ is achieved either by a positive or a negative function.

Remark 1. As an application of Theorem 2, problem (1) with $s = 9/10$, $2^*(x) = 5$, and $\theta(x) = 9/2$,

$$(a + b[u]_{9/10}^2)(-\Delta)^{9/10}u = |u|^3u + \lambda|u|^{5/2}u \quad \text{in } \Omega, \\ u = 0 \quad \text{in } \mathbb{R}^3 \setminus \Omega$$

has a least energy sign-changing solution u with energy doubling property.

2 Preliminaries

In this section, we first recall some definitions and results of variable exponent Lebesgue spaces (see [5,6,16]), which will be used later.

Let $N \geq 1$ and $\Omega \subset \mathbb{R}^N$ be a nonempty open set. A measurable function $p : \overline{\Omega} \rightarrow [1, \infty)$ is named a variable exponent.

The variable exponent Lebesgue space is

$$L^{p(\cdot)}(\Omega) = \left\{ u : \Omega \rightarrow \mathbb{R} \text{ is a measurable function: } \eta_{p(\cdot)}(u) = \int_{\Omega} |u(x)|^{p(x)} dx < \infty \right\}$$

with the Luxemburg norm

$$\|u\|_{L^{p(\cdot)}(\Omega)} = \inf \left\{ \varrho > 0 : \eta_{p(\cdot)}(\varrho^{-1}u) \leq 1 \right\},$$

then $L^{p(\cdot)}(\Omega)$ is a Banach space, and when p is bounded, we have the following relations:

$$\min \left\{ \|u\|_{L^{p(\cdot)}(\Omega)}^p, \|u\|_{L^{p(\cdot)}(\Omega)}^{\bar{p}} \right\} \leq \eta_{p(\cdot)}(u) \leq \max \left\{ \|u\|_{L^{p(\cdot)}(\Omega)}^p, \|u\|_{L^{p(\cdot)}(\Omega)}^{\bar{p}} \right\}.$$

That is, if p is bounded, then norm convergence is equivalent to convergence with respect to the modular $\eta_{p(\cdot)}$. For the bounded exponent, the dual space $(L^{p(\cdot)}(\Omega))'$ can be identified with $L^{p'(\cdot)}(\Omega)$ where the conjugate exponent p' is defined by $p' = p/(p(x)-1)$. If $1 < \underline{p} \leq \bar{p} < \infty$, then the variable exponent Lebesgue space $L^{p(\cdot)}(\Omega)$ is separable and reflexive. So we can see that Hölder's inequality is still valid in the variable exponent Lebesgue space. For all $\varphi \in L^{p(\cdot)}(\Omega)$, $\phi \in L^{p'(\cdot)}(\Omega)$ with $p(x) \in (1, \infty)$, the following inequality holds:

$$\int_{\Omega} |\varphi\phi| dx \leq \left(\frac{1}{\underline{p}} + \frac{1}{\underline{p}'} \right) \|\varphi\|_{L^{p(\cdot)}(\Omega)} \|\phi\|_{L^{p'(\cdot)}(\Omega)} \leq 2 \|\varphi\|_{L^{p(\cdot)}(\Omega)} \|\phi\|_{L^{p'(\cdot)}(\Omega)}.$$

Next, we give some definitions and results of variable-order fractional Sobolev spaces.

Define $H^{s(\cdot)}(\Omega)$ as the linear space of Lebesgue measurable functions from \mathbb{R}^N to \mathbb{R} such that any function $u = 0$ in $\mathbb{R}^N \setminus \Omega$ belongs to $L^2(\Omega)$ and

$$[u]_{s(\cdot)} := \left(\iint_{\mathbb{R}^{2N}} \frac{|u(x) - u(y)|^2}{|x - y|^{N+2s(x,y)}} dx dy \right)^{1/2} < \infty.$$

Equip $H^{s(\cdot)}(\Omega)$ with the norm

$$\|u\|_{H^{s(\cdot)}(\Omega)} = \left(\|u\|_{L^2(\Omega)}^2 + [u]_{s(\cdot)}^2 \right)^{1/2}.$$

Similar to the proof of Lemma 7 in [17], we can show that $(H_0^{s(\cdot)}(\Omega), [\cdot]_{s(\cdot)})$ is a Hilbert space. In this paper, we used norm $\|\cdot\| = [\cdot]_{s(\cdot)}$ to study problem (1).

Lemma 1. (See [25, Lemma 2.1].) The embeddings $H_0^{s_2}(\Omega) \hookrightarrow H_0^{s(\cdot)}(\Omega) \hookrightarrow H_0^{s_1}(\Omega)$ are continuous. Moreover, if $N > 2s_1$, for any fixed constant exponent $t \in [1, 2N/(N - 2s_1)]$, $H_0^{s(\cdot)}(\Omega)$ can be continuously embedded into $L^t(\Omega)$.

The following embedding theorem shows that the variable-order fractional Sobolev space is related with the variable exponent Sobolev spaces.

Lemma 2. (See [25, Thm. 2.1].) Let $\Omega \subset \mathbb{R}^N$ be a smooth bounded domain. Assume that $s : \mathbb{R}^N \times \mathbb{R}^N \rightarrow (0, 1)$ and $p : \overline{\Omega} \rightarrow (1, \infty)$ are two continuous functions satisfying (s1)–(s2) and $2 < p(x) < 2N/(N - 2s(x, x))$, respectively. Then there exists $C_p = C(N, \overline{p}, \overline{s}, \underline{s}) > 0$ such that for any $u \in H_0^{s(\cdot)}(\Omega)$, it holds that

$$\text{If } u \in H_0^{\overline{s}(\cdot)}(\Omega),$$

That is, the embedding $H_0^{s(\cdot)}(\Omega) \hookrightarrow L^{p(x)}(\Omega)$ is continuous. Furthermore, this embedding is compact. If $u \in H_0^{s(\cdot)}(\Omega)$, then there exists $C_p = C(N, \overline{p}, \overline{s}, \underline{s}) > 0$ such that

$$\|u\|_{L^{p(x)}(\Omega)} \leq C_p [u]_{s(\cdot)}.$$

3 Some technical lemmas

Now, for fixed $u \in H_0^{s(\cdot)}(\Omega)$ with $u^\pm \neq 0$, we define function $\sigma_u : \mathbb{R}_0^+ \times \mathbb{R}_0^+ \rightarrow \mathbb{R}$ and mapping $T_u : \mathbb{R}_0^+ \times \mathbb{R}_0^+ \rightarrow \mathbb{R}^2$ by

$$\sigma_u(\alpha, \beta) = \mathcal{I}_{b,\lambda}(\alpha u^+ + \beta u^-)$$

and

$$T_u(\alpha, \beta) = (\langle (\mathcal{I}_{b,\lambda})'(\alpha u^+ + \beta u^-), \alpha u^+ \rangle, \langle (\mathcal{I}_{b,\lambda})'(\alpha u^+ + \beta u^-), \beta u^- \rangle). \quad (4)$$

Lemma 3. Assume that (s1).(s2) and (f1).(f3) hold. If $u \in H_0^{s(\cdot)}(\Omega)$ with $u^\pm \neq 0$, then σ_u has the following properties:

(i) The pair (α, β) is a critical point σ_u of with $\alpha, \beta > 0$ if and only if $\alpha u^+ + \beta u^- \in \mathcal{M}_{b,\lambda}$;

(ii) The function σ_u has a unique critical point (α_u, β_u) on $(0, \infty) \times (0, \infty)$, which is also the unique maximum point of σ_u on $[0, \infty) \times [0, \infty)$. Furthermore, if $\langle (\mathcal{I}_{b,\lambda})'(u), u^\pm \rangle \leq 0$, then $0 < \alpha_u, \beta_u < 1$.

Proof. (i) By definition of σ_u we have that

$$\begin{aligned} \nabla \sigma_u(\alpha, \beta) &= (\langle (\mathcal{I}_{b,\lambda})'(\alpha u^+ + \beta u^-), u^+ \rangle, \langle (\mathcal{I}_{b,\lambda})'(\alpha u^+ + \beta u^-), u^- \rangle) \\ &= \left(\frac{1}{\alpha} \langle (\mathcal{I}_{b,\lambda})'(\alpha u^+ + \beta u^-), \alpha u^+ \rangle, \frac{1}{\beta} \langle (\mathcal{I}_{b,\lambda})'(\alpha u^+ + \beta u^-), \beta u^- \rangle \right). \end{aligned}$$

Thus, item (i) holds

(ii) For any $u \in H_0^{s(\cdot)}(\Omega)$ $u^\pm \neq 0$, α_u, β_u .

From (f1) and (f2), for any $\varepsilon > 0$, there is $C_\varepsilon > 0$ such that

$$|f(x, t)| \leq \varepsilon |t| + C_\varepsilon |t|^{\theta(x)-1} \quad \text{for all } t \in \mathbb{R}. \quad (5)$$

Then by the Sobolev embedding theorem we have

$$\begin{aligned} &\langle (\mathcal{I}_{b,\lambda})'(\alpha u^+ + \beta u^-), \alpha u^+ \rangle \\ &\geq (a - \lambda \varepsilon C_2) \alpha^2 \|u^+\|^2 + b \alpha^4 \|u^+\|^4 - (\alpha^q + \alpha^{\bar{q}}) C_1 \max\{\|u^+\|^{\bar{q}}, \|u^+\|^q\} \\ &\quad - \lambda C_\varepsilon C_3 (\alpha^{\bar{\theta}} + \alpha^{\bar{\theta}}) \max\{\|u^+\|^{\bar{\theta}}, \|u^+\|^{\theta}\}. \end{aligned}$$

Choose $\varepsilon > 0$, such that $(a - \lambda \varepsilon C_2) > 0$. Since $q, \bar{q} > 4$, we have that $\langle (\mathcal{I}_{b,\lambda})'(\alpha u^+ + \beta u^-), \alpha u^+ \rangle > 0$ for α small enough and all $\beta \geq 0$. Similarly, we are also able to prove that $\langle (\mathcal{I}_{b,\lambda})'(\alpha u^+ + \beta u^-), \beta u^- \rangle > 0$ for β small enough and all $\alpha \geq 0$. Therefore, there exists $\delta_1 > 0$ such that

$$\langle (\mathcal{I}_{b,\lambda})'(\delta_1 u^+ + \beta u^-), \delta_1 u^+ \rangle > 0, \quad \langle (\mathcal{I}_{b,\lambda})'(\alpha u^+ + \delta_1 u^-), \delta_1 u^- \rangle > 0. \quad (6)$$

On the other hand, by (f2) and (f3) we claim

$$f(x, t)t > 0, \quad t \neq 0; \quad F(x, t) \geq 0, \quad t \in \mathbb{R}. \quad (7)$$

Therefore, choose $\alpha = \delta_2^* > \delta_1$. If $\beta \in [\delta_1, \delta_2^*]$ δ_2^* is large enough, it follows that

$$\begin{aligned} &\langle (\mathcal{I}_{b,\lambda})'(\delta_2^* u^+ + \beta u^-), \delta_2^* u^+ \rangle \\ &\leq a(\delta_2^*)^2 \|u^+\|^2 + b(\delta_2^*)^4 \|u^+\|^4 + b(\delta_2^*)^4 \|u^+\|^2 \|u^-\|^2 \\ &\quad - (\delta_2^*)^q \int_{\Omega} |u^+|^{q(x)} dx \leq 0. \end{aligned}$$

Similarly, we have $\langle (\mathcal{I}_{b,\lambda})'(\alpha u^+ + \delta_2^* u^-), \delta_2^* u^- \rangle \leq 0$. Let $\delta_2 > \delta_2^*$ be large enough, we obtain

$$\langle (\mathcal{I}_{b,\lambda})'(\delta_2^* u^+ + \beta u^-), \delta_2^* u^+ \rangle < 0 \quad \text{and} \quad \langle (\mathcal{I}_{b,\lambda})'(\alpha u^+ + \delta_2^* u^-), \delta_2^* u^- \rangle < 0 \quad (8)$$

for all $\alpha, \beta \in [\delta_1, \delta_2]$. Combining (6) and (8) with Miranda's theorem [13], there exists $(\alpha_u, \beta_u) \in \mathbb{R}^+ \times \mathbb{R}^+$ such that $T_{u(\alpha, \beta)} = (0, 0)$, i.e., $\alpha u^+ + \beta u^- \in \mathcal{M}_{\lambda, \cdot}$.

According to the proof in [20], we can prove the uniqueness of the pair (α_u, β_u) .

Lastly, we prove that $0 < \alpha_u, \beta_u \leq 1$ if $\langle (\mathcal{I}_{b,\lambda})'(u), u^\pm \rangle \leq 0$.

Suppose $\alpha_u \geq \beta_u > 0$. By $\alpha_u u^+ + \beta_u u^- \in \mathcal{M}_{b,\lambda}$ we have

$$\begin{aligned} & a\alpha_u^2 \|u^+\|^2 + b\alpha_u^4 \|u^+\|^4 + b\alpha_u^4 \|u^+\|^2 \|u^-\|^2 \\ & \geq a\alpha_u^2 \|u^+\|^2 + b\alpha_u^4 \|u^+\|^4 + b\alpha_u^2 \beta_u^2 \|u^+\|^2 \|u^-\|^2 \\ & = \lambda \int_{\Omega} f(x, \alpha_u u^+) \alpha_u u^+ dx + \int_{\Omega} |\alpha_u u^+|^{q(x)} dx. \end{aligned} \quad (9)$$

On the other hand, by $\langle (\mathcal{I}_{b,\lambda})'(u), u^+ \rangle \leq 0$ we have

$$a\|u^+\|^2 + b\|u^+\|^4 + b\|u^+\|^2 \|u^-\|^2 \leq \lambda \int_{\Omega} f(x, u^+) u^+ dx + \int_{\Omega} |u^+|^{q(x)} dx. \quad (10)$$

So, according to (9) and (10), we obtain

$$\begin{aligned} a\left(\frac{1}{\alpha_u^2} - 1\right) \|u^+\|^2 & \geq \lambda \int_{\Omega} \left[\frac{f(x, \alpha_u u^+)}{(\alpha_u u^+)^3} - \frac{f(x, u^+)}{(u^+)^3} \right] (u^+)^4 dx \\ & + \int_{\Omega} (\alpha_u^{(q(x)-2)} - 1) |u^+|^{q(x)} dx. \end{aligned}$$

In view of (f3), we conclude that $\alpha_u \leq 1$. Thus, we have that $0 < \alpha_u, \beta_u \leq 1$.

Lemma 4. Let $c_{b,\lambda} = \inf_{u \in \mathcal{M}_{b,\lambda}} \mathcal{I}_{b,\lambda}(u)$. Then we have that $\lim_{\lambda \rightarrow \infty} c_{b,\lambda} = 0$.

Proof. For any $u \in \mathcal{M}_{b,\lambda}$, we have

$$a\|u^\pm\|^2 + b\|u^\pm\|^4 + b\|u^+\|^2 \|u^-\|^2 = \lambda \int_{\Omega} f(x, u^\pm) u^\pm dx + \int_{\Omega} |u^\pm|^{q(x)} dx.$$

Then by (5) and Sobolev inequalities we get

$$\begin{aligned} a\|u^\pm\|^2 &\leq \lambda \int_{\Omega} f(x, u^\pm) u^\pm dx + \int_{\Omega} |u^\pm|^{q(x)} dx \\ &\leq \lambda \varepsilon C_1 \|u^\pm\|^2 + \lambda C_\varepsilon \min\{\|u^\pm\|^\theta, \|u^\pm\|^{\bar{\theta}}\} \\ &\quad + C \min\{\|u^\pm\|^{\frac{q}{2}}, \|u^\pm\|^{\bar{q}}\}. \end{aligned}$$

Thus, we obtain

$$(a - \lambda \varepsilon C_1) \|u^\pm\|^2 \leq \lambda C_\varepsilon \min\{\|u^\pm\|^\theta, \|u^\pm\|^{\bar{\theta}}\} + C \min\{\|u^\pm\|^{\frac{q}{2}}, \|u^\pm\|^{\bar{q}}\}.$$

Choosing ε small enough such that $a - \lambda \varepsilon C_1 > 0$, since $\frac{\theta}{2}, \bar{\theta}, \frac{q}{2}, \bar{q} > 4$, there exists $\rho > 0$ such that

$$\|u^\pm\| \geq \rho \quad \text{for all } u \in \mathcal{M}_{b,\lambda}. \quad (11)$$

On the other hand, for any $u \in \mathcal{M}_{b,\lambda}$, it is obvious that $\langle (\mathcal{I}_{b,\lambda})'(u), u \rangle = 0$. Thanks to (f2) and (f3), we obtain that

$$f'(x, t)t - 3f(x, t) > (<) 0 \quad \text{for all } t > 0,$$

This fact implies that

$$\Theta(x, t) := f(x, t)t - 4F(x, t) \geq 0 \quad (12)$$

is increasing when $t > 0$ and decreasing when $t < 0$ for almost every $x \in \Omega$. Then we have

$$\mathcal{I}_{b,\lambda}(u) = \mathcal{I}_{b,\lambda}(u) - \frac{1}{4} \langle (\mathcal{I}_{b,\lambda})'(u), u \rangle \geq \frac{a}{4} \|u\|^2.$$

From above discussions we have that $\mathcal{I}_{b,\lambda}(u) > 0$ for all $u \in \mathcal{M}_{b,\lambda}$. Therefore, $\mathcal{I}_{b,\lambda}$ is bounded below on $\mathcal{M}_{b,\lambda}$, that is, $c_{b,\lambda} = \inf_{u \in \mathcal{M}_{b,\lambda}} \mathcal{I}_{b,\lambda}(u)$ is well defined.

Let $u \in H_0^{s(\cdot)}(\Omega)$ with $u^\pm \neq 0$ be fixed. By Lemma 3, for each $\lambda > 0$, there exist $\alpha_\lambda, \beta_\lambda > 0$ such that $\alpha_\lambda u^+ + \beta_\lambda u^- \in \mathcal{M}_{b,\lambda}$. By Lemma 3 we have

$$\begin{aligned} 0 \leq c_{b,\lambda} &= \inf_{u \in \mathcal{M}_{b,\lambda}} \mathcal{I}_{b,\lambda}(u) \leq \mathcal{I}_{b,\lambda}(\alpha_\lambda u^+ + \beta_\lambda u^-) \\ &\leq \frac{a}{2} \|\alpha_\lambda u^+ + \beta_\lambda u^-\|^2 + \frac{b}{4} \|\alpha_\lambda u^+ + \beta_\lambda u^-\|^4 \\ &\leq a\alpha_\lambda^2 \|u^+\|^2 + a\beta_\lambda^2 \|u^-\|^2 + 2b\alpha_\lambda^4 \|u^+\|^4 + 2b\beta_\lambda^4 \|u^-\|^4. \end{aligned}$$

For our purpose, we just prove that $\alpha_\lambda \rightarrow 0$ and $\beta_\lambda \rightarrow 0$ as $\lambda \rightarrow \infty$.

Let

$$\mathcal{T}_u = \{(\alpha_\lambda, \beta_\lambda) \in \mathbb{R}_0^+ \times \mathbb{R}_0^+ : T_u(\alpha_\lambda, \beta_\lambda) = (0, 0), \lambda > 0\},$$

where T_u is defined as (4). By (5) there holds

$$\begin{aligned} & \min\{\alpha_\lambda^{\bar{q}}, \alpha_\lambda^q\} \int_{\Omega} |u^+|^{q(x)} dx + \min\{\beta_\lambda^{\bar{q}}, \beta_\lambda^q\} \int_{\Omega} |u^-|^{q(x)} dx \\ & \leq \int_{\Omega} \alpha_\lambda^{q(x)} |u^+|^{q(x)} dx + \int_{\Omega} \alpha_\beta^{q(x)} |u^-|^{q(x)} dx \\ & \quad + \lambda \int_{\Omega} f(x, \alpha_\lambda u^+) \alpha_\lambda u^+ dx + \lambda \int_{\Omega} f(x, \beta_\lambda u^-) \beta_\lambda u^- dx \\ & = a \|\alpha_\lambda u^+ + \beta_\lambda u^-\|^2 + b \|\alpha_\lambda u^+ + \beta_\lambda u^-\|^4 \\ & \leq 2a\alpha_\lambda^2 \|u^+\|^2 + 2a\beta_\lambda^2 \|u^-\|^2 + 4b\alpha_\lambda^4 \|u^+\|^4 + 4b\beta_\lambda^4 \|u^-\|^4. \end{aligned}$$

Hence, T_u is bounded. Let $\{\lambda_n\} \subset \mathbb{R}^+$ be such that $\lambda_n \rightarrow \infty$ as $n \rightarrow \infty$. Then there exist α_0 and β_0 such that $(\alpha_{\lambda_n}, \beta_{\lambda_n}) \rightarrow (\alpha_0, \beta_0)$ as $n \rightarrow \infty$.

Now, we claim $\alpha_0 = \beta_0 = 0$. Suppose, by contradiction, that $\alpha_0 > 0$ or $\beta_0 > 0$. By $\alpha_{\lambda_n} u^+ + \beta_{\lambda_n} u^- \in \mathcal{M}_{b, \lambda_n}$ for any $n \in \mathbb{N}$, we have

$$\begin{aligned} & a \|\alpha_{\lambda_n} u^+ + \beta_{\lambda_n} u^-\|^2 + b \|\alpha_{\lambda_n} u^+ + \beta_{\lambda_n} u^-\|^4 \\ & = \lambda_n \int_{\Omega} f(x, \alpha_{\lambda_n} u^+ + \beta_{\lambda_n} u^-) (\alpha_{\lambda_n} u^+ + \beta_{\lambda_n} u^-) dx \\ & \quad + \int_{\Omega} |\alpha_{\lambda_n} u^+ + \beta_{\lambda_n} u^-|^{q(x)} dx. \end{aligned} \tag{13}$$

Thanks to $\alpha_{\lambda_n} u^+ \rightarrow \alpha_0 u^+$ and $\beta_{\lambda_n} u^- \rightarrow \beta_0 u^-$ in $H_0^{s(\cdot)}(\Omega)$,

$$\begin{aligned} & \int_{\Omega} f(x, \alpha_{\lambda_n} u^+ + \beta_{\lambda_n} u^-) (\alpha_{\lambda_n} u^+ + \beta_{\lambda_n} u^-) dx \\ & \rightarrow \int_{\Omega} f(x, \alpha_0 u^+ + \beta_0 u^-) (\alpha_0 u^+ + \beta_0 u^-) dx > 0 \end{aligned}$$

as $n \rightarrow \infty$. It follows a contradiction with equality (13) from two facts: $\lambda_n \rightarrow \infty$ as $n \rightarrow \infty$. and $\{\alpha_{\lambda_n} u^+ + \beta_{\lambda_n} u^-\}$ is bounded in $H_0^{s(\cdot)}(\Omega)$. Hence, $\alpha_0 = \beta_0 = 0$.

Therefore, we conclude that $\lim_{\lambda \rightarrow \infty} c_{b, \lambda} = 0$.

Lemma 5. *There exists $\lambda^* > 0$ such that for all $\lambda \geq \lambda^*$, the infimum $c_{b, \lambda}$ is achieved.*

Proof. By the definition of $c_{b,\lambda}$ there exists a sequence $\{u_n\} \subset \mathcal{M}_{b,\lambda}$ such that $\lim_{n \rightarrow \infty} \mathcal{I}_{b,\lambda}(u_n) = c_{b,\lambda}$. Obviously, $\{u_n\}$ is bounded in $H_0^{s(\cdot)}(\Omega)$. then, up to a subsequence, still denoted by $\{u_n\}$, there exists $u \in H_0^{s(\cdot)}(\Omega)$ such that $u_n \rightharpoonup u$. Since the embedding $H_0^{s(\cdot)}(\Omega) \hookrightarrow L^t(\Omega)$ is compact, for all $t \in (2, 2^*(x))$, we have

$$u_n \rightarrow u \quad \text{in } L^t(\Omega), \quad u_n \rightarrow u \quad \text{a.e. } x \in \Omega.$$

Hence, $u_n^\pm \rightarrow u^\pm$, a.e. $x \in \Omega$, and

$$u_n^\pm \rightharpoonup u^\pm \quad \text{in } H_0^{s(\cdot)}(\Omega), \quad u_n^\pm \rightarrow u^\pm \quad \text{in } L^t(\Omega).$$

By Lemma 3 we have

$$I_{b,\lambda}(\alpha u_n^+ + \beta u_n^-) \leq I_{b,\lambda}(u_n) \quad \text{for all } \alpha, \beta \geq 0.$$

Then by Brézis–Lieb lemma and Fatou’s lemma we get

$$\begin{aligned} & \liminf_{n \rightarrow \infty} \mathcal{I}_{b,\lambda}(\alpha u_n^+ + \beta u_n^-) \\ & \geq I_{b,\lambda}(\alpha u^+ + \beta u^-) + \frac{a\alpha^2}{2} \lim_{n \rightarrow \infty} \|u_n^+ - u^+\|^2 + \frac{a\beta^2}{2} \lim_{n \rightarrow \infty} \|u_n^- - u^-\|^2 \\ & \quad + \frac{b\alpha^4}{2} \lim_{n \rightarrow \infty} \|u_n^+ - u^+\|^2 \|u^+\|^2 + \frac{b\beta^4}{2} \lim_{n \rightarrow \infty} \|u_n^- - u^-\|^2 \|u^-\|^2 \\ & \quad + \frac{b\alpha^4}{4} \left(\lim_{n \rightarrow \infty} \|u_n^+ - u^+\|^2 \right)^2 + \frac{b\beta^4}{4} \left(\lim_{n \rightarrow \infty} \|u_n^- - u^-\|^2 \right)^2 \\ & \quad - \int_{\Omega} \frac{\alpha^{q(x)}}{q(x)} |u_n^+ - u^+|^{q(x)} dx - \int_{\Omega} \frac{\beta^{q(x)}}{q(x)} |u_n^- - u^-|^{q(x)} dx \\ & \geq I_{b,\lambda}(\alpha u^+ + \beta u^-) + \frac{a\alpha^2}{2} A_1 + \frac{b\alpha^4}{2} A_1 \|u^+\|^2 + \frac{b\alpha^4}{4} A_1^2 - \frac{\max\{\alpha^{\underline{q}}, \alpha^{\bar{q}}\}}{\underline{q}} B_1 \\ & \quad + \frac{a\beta^2}{2} A_2 + \frac{b\beta^4}{2} A_2 \|u^-\|^2 + \frac{b\beta^4}{4} A_2^2 - \frac{\max\{\alpha^{\underline{q}}, \alpha^{\bar{q}}\}}{\underline{q}} B_2, \end{aligned}$$

where

$$\begin{aligned} A_1 &= \lim_{n \rightarrow \infty} \|u_n^+ - u^+\|^2, & A_2 &= \lim_{n \rightarrow \infty} \|u_n^- - u^-\|^2, \\ B_1 &= \lim_{n \rightarrow \infty} |u_n^+ - u^+|_{q(x)}^{q(x)}, & B_2 &= \lim_{n \rightarrow \infty} |u_n^- - u^-|_{q(x)}^{q(x)}. \end{aligned}$$

That is, one has

$$\begin{aligned} \mathcal{I}_{b,\lambda}(\alpha u^+ + \beta u^-) &+ \frac{a\alpha^2}{2}A_1 + \frac{b\alpha^4}{2}A_1\|u^+\|^2 + \frac{b\alpha^4}{4}A_1^2 - \frac{\max\{\alpha^{\underline{q}}, \alpha^{\bar{q}}\}}{\underline{q}}B_1 \\ &+ \frac{a\beta^2}{2}A_2 + \frac{b\beta^4}{2}A_2\|u^-\|^2 + \frac{b\beta^4}{4}A_2^2 - \frac{\max\{\beta^{\underline{q}}, \beta^{\bar{q}}\}}{\underline{q}}B_2 \leq c_{b,\lambda} \end{aligned} \quad (14)$$

for all $\alpha \geq 0$ and all $\beta \geq 0$.

Now, we claim that $u^\pm \neq 0$. In fact, since the situation $u^- \neq 0$ is analogous, we just prove $u^+ \neq 0$. By contradiction we suppose $u^+ \equiv 0$. Hence, let $\beta = 0$ in (14), and we have

$$\frac{a\alpha^2}{2}A_1 + \frac{b\alpha^4}{4}A_1^2 - \frac{\max\{\alpha^{\underline{q}}, \alpha^{\bar{q}}\}}{\underline{q}}B_1 \leq c_{b,\lambda} \quad \text{for all } \alpha \geq 0. \quad (15)$$

Case 1: $B_1 = 0$.

If $A_1 = 0$, that is, $u_n^+ \rightarrow u^+$ in $H_0^{s(\cdot)}(\Omega)$. From Lemma (11) we obtain $\|u^+\| > 0$, which contradicts our supposition. If $A_1 > 0$, by (14) and Lemma 4 we have

$$0 < \frac{a\alpha^2}{2}A_1 + \frac{b\alpha^4}{4}A_1^2 \leq c_{b,\lambda} \rightarrow 0 \quad \text{for all } \alpha \geq 0 \text{ and } \lambda \rightarrow +\infty.$$

The inequality is absurd. Anyway, we have a contradiction.

Case 2: $B_1 > 0$.

On the one hand, by Lemma 4 there exists $\lambda^* > 0$ such that

$$c_{b,\lambda} < \Lambda := \min\{C_q^{-\underline{q}}, C_q^{-\bar{q}}\}a^{\underline{q}/(\underline{q}-2)} \quad \text{for all } \lambda \geq \lambda^*.$$

On the other hand, since $B_1 > 0$, we obtain $A_1 > 0$. Hence, by means of (15) we have

$$\begin{aligned} \Lambda &\leq \left[\frac{(aA_1)^{\underline{q}/2}}{B_1} \right]^{2/(\underline{q}-2)} \leq \max_{0 \leq \alpha \leq 1} \left\{ \frac{a\alpha^2}{2}A_1 - \frac{\alpha^{\underline{q}}}{\underline{q}}B_1 \right\} \\ &\leq \max_{0 \leq \alpha \leq 1} \left\{ \frac{a\alpha^2}{2}A_1 + \frac{b\alpha^4}{4}A_1^2 - \frac{\alpha^{\underline{q}}}{\underline{q}}B_1 \right\} \leq c_{b,\lambda}, \end{aligned}$$

which is a contradiction. Hence, we deduce that $\|u^+\| > 0$.

Second, we prove $B_1 = B_2 = 0$.

Since the situation $B_2 = 0$ is analogous, we only prove $B_1 = 0$. By contradiction we suppose that $B_1 > 0$.

Case 1: $B_2 > 0$.

According to $B_1, B_2 > 0$ and Sobolev embedding, we obtain that $A_1, A_2 > 0$. Let

$$\varphi(\alpha) = \frac{a\alpha^2}{2}A_1 + \frac{b\alpha^4}{4}A_1^2 - \frac{\max\{\alpha^{\underline{q}}, \alpha^{\bar{q}}\}}{\underline{q}}B_1 \quad \text{for all } \alpha \geq 0.$$

It is easy to see that $\varphi(\alpha) > 0$ for $\alpha > 0$ small enough and $\varphi(\alpha) < 0$ for $\alpha < 0$ large enough. Hence, by the continuity of $\varphi(\alpha)$ there exists $\tilde{\alpha} > 0$ such that

$$\begin{aligned} & \frac{a\tilde{\alpha}^2}{2}A_1 + \frac{b\tilde{\alpha}^4}{4}A_1^2 - \frac{\max\{\tilde{\alpha}^q, \tilde{\alpha}^{\bar{q}}\}}{q}B_1 \\ &= \max_{s \geq 0} \left\{ \frac{a\alpha^2}{2}A_1 + \frac{b\alpha^4}{4}A_1^2 - \frac{\max\{\alpha^q, \alpha^{\bar{q}}\}}{q}B_1 \right\}. \end{aligned}$$

Similarly, there exists $\tilde{\beta} > 0$ such that

$$\begin{aligned} & \frac{a\tilde{\beta}^2}{2}A_2 + \frac{b\tilde{\beta}^4}{4}A_2^2 - \frac{\max\{\tilde{\beta}^q, \tilde{\beta}^{\bar{q}}\}}{q}B_2 \\ &= \max_{s \geq 0} \left\{ \frac{a\beta^2}{2}A_2 + \frac{b\beta^4}{4}A_2^2 - \frac{\max\{\beta^q, \beta^{\bar{q}}\}}{q}B_2 \right\}. \end{aligned}$$

Since $[0, \tilde{\alpha}] \times [0, \tilde{\beta}]$ is compact and σ is continuous, there exists $(\alpha_u, \beta_u) \in (0, \tilde{\alpha}) \times (0, \tilde{\beta})$ such that

$$\sigma(\alpha_u, \beta_u) = \max_{(\alpha, \beta) \in [0, \tilde{\alpha}] \times [0, \tilde{\beta}]} \sigma(\alpha, \beta).$$

Now, we prove that $(\alpha_u, \beta_u) \in (0, \tilde{\alpha}) \times (0, \tilde{\beta})$. Note that if β is small enough, we obtain

$$\begin{aligned} \sigma(\alpha, 0) &= \mathcal{I}_{b, \lambda}(\alpha u^+) < \mathcal{I}_{b, \lambda}(\alpha u^+) + \mathcal{I}_{b, \lambda}(\beta u^-) \\ &\leq \mathcal{I}_{b, \lambda}(\alpha u^+ + \beta u^-) = \sigma(\alpha, \beta) \end{aligned}$$

for all $\alpha \in [0, \tilde{\alpha}]$. Hence, there exists $\beta_0 \in [0, \tilde{\beta}]$ such that

$$\sigma(\alpha, 0) \leq \sigma(\alpha, \beta_0) \quad \text{for all } \alpha \in [0, \tilde{\alpha}].$$

That is, any point of $(\alpha, 0)$ with $0 \leq \alpha \leq \tilde{\alpha}$ is not the maximizer of σ . This yields that $(\alpha_u, \beta_u) \notin [0, \tilde{\alpha}] \times \{0\}$. Similarly, we obtain $(\alpha_u, \beta_u) \notin \{0\} \times [0, \tilde{\alpha}]$. On the other hand, it is easy to see that

$$\frac{a\alpha^2}{2}A_1 + \frac{b\alpha^4}{2}A_1\|u^+\|^2 + \frac{b\alpha^4}{4}A_1^2 - \frac{\max\{\alpha^q, \alpha^{\bar{q}}\}}{q}B_1 > 0 \quad (16)$$

and

$$\frac{a\beta^2}{2}A_2 + \frac{b\beta^4}{2}A_2\|u^-\|^2 + \frac{b\beta^4}{4}A_2^2 - \frac{\max\{\beta^q, \beta^{\bar{q}}\}}{q}B_2 > 0 \quad (17)$$

for $\alpha \in (0, \tilde{\alpha}]$, $\beta \in (0, \tilde{\beta}]$.

Then we have

$$\begin{aligned} \Lambda \leq & \frac{a\tilde{\alpha}^2}{2}A_1 + \frac{b\tilde{\alpha}^4}{4}A_1^2 - \frac{\max\{\alpha^q, \alpha^{\bar{q}}\}}{q}B_1 + \frac{b\tilde{\alpha}^4}{2}A_1\|u^+\|^2 \\ & + \frac{a\beta^2}{2}A_2 + \frac{b\beta^4}{2}A_2\|u^-\|^2 + \frac{b\beta^4}{4}A_2^2 - \frac{\max\{\beta^q, \beta^{\bar{q}}\}}{q}B_2 \end{aligned}$$

and

$$\begin{aligned} \Lambda \leq & \frac{a\tilde{\beta}^2}{2}A_2 + \frac{b\tilde{\beta}^4}{4}A_2^2 - \frac{\max\{\beta^q, \beta^{\bar{q}}\}}{q}B_2 + \frac{b\tilde{\beta}^4}{2}A_2\|u^-\|^2 \\ & + \frac{a\alpha^2}{2}A_1 + \frac{b\alpha^4}{2}A_1\|u^+\|^2 + \frac{b\alpha^4}{4}A_1^2 - \frac{\max\{\alpha^q, \alpha^{\bar{q}}\}}{q}B_1 \end{aligned}$$

for all $\alpha \in [0, \tilde{\alpha}]$ and all $\beta \in [0, \tilde{\beta}]$. Therefore, according to (14), we conclude $\sigma(\alpha, \tilde{\beta}) \leq 0$, $\sigma(\tilde{\alpha}, \beta) \leq 0$ for all $\alpha \in [0, \tilde{\alpha}]$ and all $\beta \in [0, \tilde{\beta}]$. Thus, $(\alpha_u, \beta_u) \notin \{\tilde{\alpha}\} \times [0, \tilde{\beta}]$ and $(\alpha_u, \beta_u) \notin [0, \tilde{\alpha}] \times \{\tilde{\beta}\}$. Finally, we get that $(\alpha_u, \beta_u) \in (0, \tilde{\alpha}) \times (0, \tilde{\beta})$. Hence, it follows that (α_u, β_u) is a critical point of σ . This implies that $\alpha_u u^+ + \beta_u u^- \in \mathcal{M}_{b,\lambda}$. From (14), (16), and (17) we have

$$\begin{aligned} c_{b,\lambda} & \geq \mathcal{I}_{b,\lambda}(\alpha_u u^+ + \beta_u u^-) \\ & + \frac{a\alpha_u^2}{2}A_1 + \frac{b\alpha_u^4}{2}A_1\|u^+\|^2 + \frac{b\alpha_u^4}{4}A_1^2 - \frac{\max\{\alpha^q, \alpha^{\bar{q}}\}}{q}B_1 \\ & + \frac{a\beta_u^2}{2}A_2 + \frac{b\beta_u^4}{2}A_2\|u^-\|^2 + \frac{b\beta_u^4}{4}A_2^2 - \frac{\max\{\beta^q, \beta^{\bar{q}}\}}{q}B_2 \\ & > \mathcal{I}_{b,\lambda}(\alpha_u u^+ + \beta_u u^-) \geq c_{b,\lambda}, \end{aligned}$$

which is a contradiction.

Case 2: $B_2 = 0$.

In this case, we can maximize in $[0, \tilde{\alpha}] \times \mathbb{R}_0^+$. Indeed, it is possible to show that there exist $\beta_0 \in \mathbb{R}_0^+$ such that

$$\mathcal{I}_{b,\lambda}(\alpha_u u^+ + \beta_u u^-) \leq 0 \quad \text{for all } (\alpha, \beta) \in [0, \tilde{\alpha}] \times [\beta_0, \infty).$$

Hence, there is $(\alpha_u, \beta_u) \in [0, \tilde{\alpha}] \times [0, \infty)$ such that

$$\sigma(\alpha_u, \beta_u) = \max_{(\alpha, \beta) \in [0, \tilde{\alpha}] \times [0, \infty)} \sigma(\alpha, \beta).$$

In the following, we prove that $(\alpha_u, \beta_u) \in (0, \tilde{\alpha}) \times \mathbb{R}^+$. It is noted that $\sigma(\alpha, 0) < \sigma(\alpha, \beta)$ for $\alpha \in [0, \tilde{\alpha}]$ and β small enough, so we have $(\alpha_u, \beta_u) \notin [0, \tilde{\alpha}] \times \{0\}$. Meanwhile, $\sigma(0, \beta) < \sigma(\alpha, \beta)$ for $\beta \in [0, \infty)$ and α small enough, then we have $(\alpha_u, \beta_u) \notin \{0\} \times \mathbb{R}_0^+$.

On the other hand, it is obvious that

$$\begin{aligned} A \leq & \frac{a\tilde{\alpha}^2}{2}A_1 + \frac{b\tilde{\alpha}^4}{4}A_1^2 - \frac{\max\{\alpha^q, \alpha^{\bar{q}}\}}{q}B_1 + \frac{b\tilde{\alpha}^4}{2}A_2\|u^+\|^2 \\ & + \frac{a\beta^2}{2}A_2 + \frac{b\beta^4}{2}A_2\|u^-\|^2 + \frac{b\beta^4}{4}A_2^2 \quad \text{for all } \beta \in [0, \infty). \end{aligned}$$

Hence, we obtain that $\sigma(\tilde{\alpha}, \beta) \leq 0$ for all $\beta \in \mathbb{R}_0^+$. Thus, $(\alpha_u, \beta_u) \notin \{\tilde{\alpha}\} \times \mathbb{R}_0^+$. Hence, $(\alpha_u, \beta_u) \in (0, \tilde{\alpha}) \times \mathbb{R}^+$. That is, (α_u, β_u) is an inner maximizer of σ in $[0, \tilde{\alpha}] \times \mathbb{R}_0^+$. Hence, $\alpha_u u^+ + \beta_u u^- \in \mathcal{M}_{b, \lambda}$. Then it follows from (16) that

$$\begin{aligned} c_{b, \lambda} & \geq \mathcal{I}_{b, \lambda}(\alpha_u u^+ + \beta_u u^-) \\ & + \frac{a\alpha_u^2}{2}A_1 + \frac{b\alpha_u^4}{2}A_1\|u^+\|^2 + \frac{b\alpha_u^4}{4}A_1^2 - \frac{\max\{\alpha^q, \alpha^{\bar{q}}\}}{q}B_1 \\ & + \frac{a\beta_u^2}{2}A_2 + \frac{b\beta_u^4}{2}A_2\|u^-\|^2 + \frac{b\beta_u^4}{4}A_2^2 \\ & > \mathcal{I}_{b, \lambda}(\alpha_u u^+ + \beta_u u^-) \geq c_{b, \lambda}, \end{aligned}$$

which is absurd.

Therefore, from the above arguments we have that $B_1 = B_2 = 0$.

Finally, we prove that $c_{b, \lambda}$ is achieved. Since $u^\pm \neq 0$, by Lemma 1, there exist $\alpha_u, \beta_u > 0$ such that

$$\hat{u} := \alpha_u u^+ + \beta_u u^- \in \mathcal{M}_{b, \lambda}.$$

Furthermore, it is easy to see that $\langle (\mathcal{I}_{b, \lambda})'(u), u^\pm \rangle \leq 0$. By Lemma 3 we obtain $0 < \alpha_u, \beta_u < 1$. Since $u_n \in \mathcal{M}_{b, \lambda}$, according to Lemma 4, we get

$$\mathcal{I}_{b, \lambda}(\alpha_u u_n^+ + \beta_u u_n^-) \leq \mathcal{I}_{b, \lambda}(u_n^+ + u_n^-) = \mathcal{I}_{b, \lambda}(u_n).$$

Thanks to (f3), $B_1 = B_2 = 0$. and the norm in $H_0^{s(\varrho)}$ is lower semicontinuous, we have

$$\begin{aligned}
 c_{b,\lambda} &\leq \mathcal{I}_{b,\lambda}(\widehat{u}) - \frac{1}{4} \langle (\mathcal{I}_{b,\lambda})'(\widehat{u}), \widehat{u} \rangle \\
 &\leq \frac{a}{4} \|\widehat{u}\|^2 + \int_{\Omega} \left(\frac{1}{4} - \frac{1}{q(x)} \right) |\widehat{u}|^{q(x)} dx + \frac{\lambda}{4} \int_{\Omega} [f(x, \widehat{u})\widehat{u} - 4F(x, \widehat{u})] dx \\
 &\leq \frac{a}{4} \|u\|^2 + \int_{\Omega} \left(\frac{1}{4} - \frac{1}{q(x)} \right) |u|^{q(x)} dx + \frac{\lambda}{4} \int_{\Omega} [f(x, u)u - 4F(x, u)] dx \\
 &\leq \liminf_{n \rightarrow \infty} \left[\mathcal{I}_{b,\lambda}(u_n) - \frac{1}{4} \langle (\mathcal{I}_{b,\lambda})'(u_n), u_n \rangle \right] \leq c_{b,\lambda}.
 \end{aligned}$$

Therefore, $\alpha_u = \beta_u = 1$, and $c_{b,\lambda}$ is achieved by $u_b := u^+ + u^- \in \mathcal{M}_{b,\lambda}$.

4 Proof of main results

In this section, we prove our main results. First, we prove Theorem 1. In fact, thanks to Lemma 5, we just prove that the minimizer u_b for $c_{b,\lambda}$ is indeed a sign-changing solution of problem (1).

Proof of Theorem 1. Since $u_b \in \mathcal{M}_{b,\lambda}$, we have

$$\langle (\mathcal{I}_{b,\lambda})'(u_b), u_b^+ \rangle = \langle (\mathcal{I}_{b,\lambda})'(u_b), u_b^- \rangle = 0.$$

By Lemma 3 and Lemma 4, for $(\alpha, \beta) \in (\mathbb{R}_+ \times \mathbb{R}_+) \setminus (1, 1)$, we have

$$\mathcal{I}_{b,\lambda}(\alpha u_b^+ + \beta u_b^-) < \mathcal{I}_{b,\lambda}(u_b^+ + u_b^-) = c_{b,\lambda}. \quad (18)$$

If $(\mathcal{I}_{b,\lambda})'(u_b) \neq 0$, then there exist $\delta > 0$ and $\theta > 0$ such that

$$\|(\mathcal{I}_{b,\lambda})'(v)\| \geq \theta \quad \text{for all } \|v - u_b\| \geq 3\delta.$$

Choose $\tau \in (0, \min\{1/2, \delta/(\sqrt{2}\|u_b\|)\})$. Let

$$D := (1 - \tau, 1 + \tau) \times (1 - \tau, 1 + \tau)$$

and

$$g(\alpha, \beta) = \alpha u_b^+ + \beta u_b^- \quad \text{for all } (\alpha, \beta) \in D.$$

In view of (18), it is easy to see that

$$\bar{c}_\lambda := \max_{\partial\Omega} \mathcal{I}_{b,\lambda} \circ g < c_{b,\lambda}.$$

Let $\varepsilon := \min\{(c_{b,\lambda} - \bar{c}_\lambda)/2, \theta\delta/8\}$ and $S_\delta := B(u_b, \delta)$, by Lemma 2.3 in [23] there exists a deformation $\eta \in C([0, 1] \times D, D)$ such that

- a) $\eta(1, v) = v$ if $v \notin (\mathcal{I}_{b,\lambda})^{-1}([c_{b,\lambda} - 2\varepsilon, c_{b,\lambda} + 2\varepsilon] \cap S_\delta)$;
- b) $\eta(1, (\mathcal{I}_{b,\lambda})^{c_{b,\lambda}+\varepsilon} \cap S_\delta) \subset (\mathcal{I}_{b,\lambda})^{c_{b,\lambda}-\varepsilon}$;
- c) $\mathcal{I}_{b,\lambda}(\eta(1, v)) \leq \mathcal{I}_{b,\lambda}(v)$ for all $v \in H_0^{s(\cdot)}(\Omega)$.

First, we need to prove that

$$\max_{(\alpha, \beta) \in \bar{D}} \mathcal{I}_{b,\lambda}(\eta(1, g(\alpha, \beta))) < c_{b,\lambda}. \quad (19)$$

In fact, it follows from Lemma 1 that $\mathcal{I}_{b,\lambda}(g(\alpha, \beta)) \leq c_{b,\lambda} < c_{b,\lambda} + \varepsilon$. That is,

$$g(\alpha, \beta) \in (\mathcal{I}_{b,\lambda})^{c_{b,\lambda}+\varepsilon}.$$

On the other hand, we have

$$\begin{aligned} \|g(\alpha, \beta) - u_b\|^2 &= \|(\alpha - 1)u_b^+ + (\beta - 1)u_b^-\|^2 \\ &\leq 2((\alpha - 1)^2\|u_b^+\|^2 + (\beta - 1)^2\|u_b^-\|^2) \\ &\leq 2\tau\|u_b\|^2 < \delta^2, \end{aligned}$$

which shows that $g(\alpha, \beta) \in S_\delta$ for all $(\alpha, \beta) \in D$.

Therefore, by (b) we have $\mathcal{I}_{b,\lambda}(\eta(1, g(s, t))) < c_{b,\lambda} - \varepsilon$. Hence, (19) holds.

In the following, we prove that $\eta(1, g(D)) \cap \mathcal{M}_{b,\lambda} \neq \emptyset$, which contradicts the definition of $c_{b,\lambda}$.

Let $h(\alpha, \beta) := \eta(1, g(\alpha, \beta))$.

$$\begin{aligned} \Psi_0(\alpha, \beta) &:= (\langle (\mathcal{I}_{b,\lambda})'(g(\alpha, \beta)), u_b^+ \rangle, \langle (\mathcal{I}_{b,\lambda})'(g(\alpha, \beta)), u_b^- \rangle) \\ &= (\langle (\mathcal{I}_{b,\lambda})'(\alpha u_b^+ + \beta u_b^-), u_b^+ \rangle, \langle (\mathcal{I}_{b,\lambda})'(\alpha u_b^+ + \beta u_b^-), u_b^- \rangle) \\ &:= (\varphi_u^1(\alpha, \beta), \varphi_u^2(\alpha, \beta)), \end{aligned}$$

and

$$\Psi_1(\alpha, \beta) := \left(\frac{1}{\alpha} \langle (\mathcal{I}_{b,\lambda})'(h(\alpha, \beta)), (h(\alpha, \beta))^+ \rangle, \frac{1}{\beta} \langle (\mathcal{I}_{b,\lambda})'(h(\alpha, \beta)), (h(\alpha, \beta))^- \rangle \right).$$

By the direct calculation we have

$$\begin{aligned} \left. \frac{\varphi_u^1(\alpha, \beta)}{\partial \alpha} \right|_{(1,1)} &= a \|u_b^+\|^2 + 3b \|u_b^+\|^4 + b \|u_b^+\|^2 \|u_b^-\|^2 \\ &\quad - \int_{\Omega} (q(x) - 1) |u_b^+|^{q(x)} dx - \lambda \int_{\Omega} \partial_{\alpha} f(x, u_b^+) (u_b^+)^2 dx, \\ \left. \frac{\varphi_u^1(\alpha, \beta)}{\partial \beta} \right|_{(1,1)} &= 2b \|u_b^+\|^2 \|u_b^-\|^2, \quad \left. \frac{\varphi_u^2(\alpha, \beta)}{\partial \alpha} \right|_{(1,1)} = 2b \|u_b^+\|^2 \|u_b^-\|^2, \\ \left. \frac{\varphi_u^2(\alpha, \beta)}{\partial \beta} \right|_{(1,1)} &= a \|u_b^-\|^2 + 3b \|u_b^-\|^4 + b \|u_b^+\|^2 \|u_b^-\|^2 \\ &\quad - \int_{\Omega} (q(x) - 1) |u_b^-|^{q(x)} dx - \lambda \int_{\Omega} \partial_{\beta} f(x, u_b^-) (u_b^-)^2 dx. \end{aligned}$$

Let

$$M = \begin{bmatrix} \left. \frac{\varphi_u^1(\alpha, \beta)}{\partial \alpha} \right|_{(1,1)} & \left. \frac{\varphi_u^2(\alpha, \beta)}{\partial \alpha} \right|_{(1,1)} \\ \left. \frac{\varphi_u^1(\alpha, \beta)}{\partial \beta} \right|_{(1,1)} & \left. \frac{\varphi_u^2(\alpha, \beta)}{\partial \beta} \right|_{(1,1)} \end{bmatrix}.$$

By (f3), for $t \neq 0$, we have

$$\partial_t f(x, t) t^2 - 3f(x, t) t > 0 \quad \text{for a.e. } x \in \Omega.$$

Then, since $u_b \in \mathcal{M}_{b,\lambda}$, we have

$$\det M = \left. \frac{\varphi_u^1(\alpha, \beta)}{\partial \alpha} \right|_{(1,1)} \cdot \left. \frac{\varphi_u^2(\alpha, \beta)}{\partial \beta} \right|_{(1,1)} - \left. \frac{\varphi_u^1(\alpha, \beta)}{\partial \beta} \right|_{(1,1)} \cdot \left. \frac{\varphi_u^2(\alpha, \beta)}{\partial \alpha} \right|_{(1,1)} > 0.$$

Since $\Psi_0(\alpha, \beta)$ is C^1 a function and $(1,1)$ is the unique isolated zero point of Ψ_0 , by using the degree theory we deduce that $\deg(\Psi_1, D, 0) = 1$.

Hence, combining (19) with (a), we obtain

$$g(\alpha, \beta) = h(\alpha, \beta) \quad \text{on } \partial D.$$

Consequently, we obtain $\deg(\Psi_1, D, 0) = 1$. Therefore, $\Psi_1(\alpha_0, \beta_0) = 0$ for some $(\alpha_0, \beta_0) \in D$ so that

$$\eta(1, g(\alpha_0, \beta_0)) = h(\alpha_0, \beta_0) \in \mathcal{M}_{b,\lambda},$$

which contradicts (19).

From the above discussions we deduce that u is a sign-changing solution for problem (1).

Finally, we prove that u has exactly two nodal domains. To this end, we assume by contradiction that

$$u_b = u_1 + u_2 + u_3 \quad \text{with } u_i \neq 0, \quad u_1 \geq 0, \quad u_2 \leq 0$$

and

$$\text{suppt}(u_i) \cap \text{suppt}(u_j) = \emptyset \quad \text{for } i \neq j, \quad i, j = 1, 2, 3,$$

and

$$\langle (\mathcal{I}_{b,\lambda})'(u), u_i \rangle = 0 \quad \text{for } i = 1, 2, 3.$$

Setting $v := u_1 + u_2$, we see that $v^+ = u_1$ and $v^- = u_2$, i.e., $v^\pm \neq 0$. Then there exist a unique pair (α_v, β_v) of positive numbers such that $\alpha_v u_1 + \beta_v u_2 \in \mathcal{M}_{b,\lambda}$. Hence,

$$\mathcal{I}_{b,\lambda}(\alpha_v u_1 + \beta_v u_2) \geq c_b^\lambda.$$

Moreover, using the fact that $\langle (\mathcal{I}_{b,\lambda})'(u), u_i \rangle = 0$, we obtain $\langle (\mathcal{I}_{b,\lambda})'(v), v^\pm \rangle < 0$. From Lemma 3(ii) we have that

$$(\alpha_v, \beta_v) \in (0, 1] \times (0, 1].$$

On the other hand, we have

$$0 = \frac{1}{4} \langle (\mathcal{I}_{b,\lambda})'(u), u_3 \rangle < \mathcal{I}_{b,\lambda}(u_3) + \frac{b}{4} \|u_1\|^2 \|u_3\|^2 + \frac{b}{4} \|u_2\|^2 \|u_3\|^2.$$

Hence, by (12) we obtain

$$\begin{aligned}
 c_{b,\lambda} &\leq \mathcal{I}_{b,\lambda}(\alpha_v u_1 + \beta_v u_2) \\
 &= \mathcal{I}_{b,\lambda}(\alpha_v u_1 + \beta_v u_2) - \frac{1}{4} \langle (\mathcal{I}_{b,\lambda})'(\alpha_v u_1 + \beta_v u_2), (\alpha_v u_1 + \beta_v u_2) \rangle \\
 &= \frac{a}{4} (\|\alpha_v u_1\|^2 + \|\beta_v u_2\|^2) + \frac{\lambda}{4} \int_{\Omega} [f(x, \alpha_v u_1)(\alpha_v u_1) - 4F(x, \alpha_v u_1)] dx \\
 &\quad + \frac{\lambda}{4} \int_{\Omega} [f(x, \beta_v u_2)(\beta_v u_2) - 4F(x, \beta_v u_2)] dx \\
 &\quad + \int_{\Omega} \left(\frac{1}{4} - \frac{1}{q(x)} \right) \alpha_v^{q(x)} |u_1|^{q(x)} dx + \int_{\Omega} \left(\frac{1}{4} - \frac{1}{q(x)} \right) \beta_v^{q(x)} |u_2|^{q(x)} dx \\
 &\leq \mathcal{I}_{b,\lambda}(u_1 + u_2) - \frac{1}{4} \langle (\mathcal{I}_{b,\lambda})'(u_1 + u_2), (u_1 + u_2) \rangle \\
 &= \mathcal{I}_{b,\lambda}(u_1 + u_2) + \frac{1}{4} \langle (\mathcal{I}_{b,\lambda})'(u), u_3 \rangle + \frac{b}{4} \|u_1\|^2 \|u_3\|^2 + \frac{b}{4} \|u_2\|^2 \|u_3\|^2 \\
 &< \mathcal{I}_{b,\lambda}(u_1) + \mathcal{I}_{b,\lambda}(u_2) + \mathcal{I}_{b,\lambda}(u_3) + \frac{b}{4} (\|u_2\|^2 + \|u_3\|^2) \|u_1\|^2 \\
 &\quad + \frac{b}{4} (\|u_1\|^2 + \|u_3\|^2) \|u_2\|^2 + \frac{b}{4} (\|u_1\|^2 + \|u_2\|^2) \|u_3\|^2 \\
 &= \mathcal{I}_{b,\lambda}(u) = c_{b,\lambda},
 \end{aligned}$$

which is a contradiction, that is, $u_3 = 0$ and u_b has exactly two nodal domains.

By Theorem 1 we obtain a least energy sign-changing solution u_b of problem (1). Next, we prove that the energy of u_b is strictly larger than two times the ground state energy.

Proof of Theorem 2. Similar to the proof of Lemma 5, there exists $\lambda_1^* > 0$ such that for all $\lambda \geq \lambda_1^*$ and for each $b > 0$, there exists $v_b \in \mathcal{N}_{b,\lambda}$ such that $\mathcal{I}_{b,\lambda}(v_b) = c^* > 0$. By standard arguments (see [10, Cor. 2.13]) the critical points of the functional $\mathcal{I}_{b,\lambda}$ on $\mathcal{N}_{b,\lambda}$ are critical points of $\mathcal{I}_{b,\lambda}$ in $H_0^{s(\cdot)}(\Omega)$, and we obtain $(\mathcal{I}_{b,\lambda})'(v_b) = 0$. That is, u_b is a ground state solution of (1).

According to Theorem 1, we know that problem (1) has a least energy sign-changing solution u_b , which changes sign only once when $\lambda \geq \lambda^*$.

Let

Let $\lambda^{**} = \max\{\lambda^*, \lambda_1^*\}$. Suppose that $u_b = u_b^+ + u_b^-$. As in the proof of Lemma 3, there exist $\alpha_{u_b^+} > 0$ and $\beta_{u_b^-} > 0$ such that $\alpha_{u_b^+} u_b^+ \in \mathcal{N}_{b,\lambda}$, $\beta_{u_b^-} u_b^- \in \mathcal{N}_{b,\lambda}$. Furthermore, Lemma 3 implies that $\alpha_{u_b^+}, \beta_{u_b^-} \in (0, 1)$. Therefore, in view of Lemma 3, we have

$$\begin{aligned}
 2c^* &\leq \mathcal{I}_{b,\lambda}(\alpha_{u_b^+} u_b^+) + \mathcal{I}_{b,\lambda}(\beta_{u_b^-} u_b^-) \leq \mathcal{I}_{b,\lambda}(\alpha_{u_b^+} u_b^+ + \beta_{u_b^-} u_b^-) \\
 &< \mathcal{I}_{b,\lambda}(u_b^+ + u_b^-) = c_{b,\lambda}.
 \end{aligned}$$

Hence, it follows that $c^* > 0$ cannot be achieved by a sign-changing function.

References

1. G.F. Carrier, On the nonlinear vibration problem of the elastic string, *Q. Appl. Math.*, 3(2):157– 165, 1945, <https://doi.org/10.1090/qam/12351>.

2. B. Cheng, J. Chen, B. Zhang, Least energy nodal solutions for Kirchhoff-type Laplacian problems, *Math. Methods Appl. Sci.*, 43(6):3827–3849, 2020, <https://doi.org/10.1002/mma.6157>.
3. K. Cheng, Q. Gao, Sign-changing solutions for the stationary Kirchhoff problems, involving the fractional Laplacian in \mathbb{R} , *Acta Math. Sci., Ser. B, Engl. Ed.*, 38(6):1712–1730, 2018, [https://doi.org/10.1016/S0252-9602\(18\)30841-5](https://doi.org/10.1016/S0252-9602(18)30841-5).
4. M. Chipot, B. Lovat, Some remarks on nonlocal elliptic and parabolic problems, *Nonlinear Anal., Theory Methods Appl.*, 30(7):4619–4627, 1997, [https://doi.org/10.1016/S0362-546X\(97\)00169-7](https://doi.org/10.1016/S0362-546X(97)00169-7).
5. L. Diening, P. Harjulehto, P. Hästö, M. Ružička, *Lebesgue and Sobolev spaces with variable exponents*, Lect. Notes Math., Vol. 2017, Springer, Berlin, Heidelberg, 2011, <https://doi.org/10.1007/978-3-642-18363-8>.
6. X. Fan, D. Zhao, On the spaces $L_p(x)(\Omega)$ and $W_{m,p}(x)(\Omega)$, *J. Math. Anal. Appl.*, 263(2): 424–446, 2001, <https://doi.org/10.1006/jmaa.2000.7617>.
7. G.M. Figueiredo, N. Ikoma, J.R. Santos Jr., Existence and concentration result for the Kirchhoff type equations with general nonlinearities, *Arch. Ration. Mech. Anal.*, 213(3):931–979, 2015, <https://doi.org/10.1007/s00205-014-0747-8>.
8. G.M. Figueiredo, J.R. Santos Jr., Existence of a least energy nodal solution for a Schrödinger– Kirchhoff equation with potential vanishing at infinity, *J. Math. Phys.*, 56(5):051506, 2015, <https://doi.org/10.1063/1.4921639>.
9. A. Fiscella, E. Valdinoci, A critical Kirchhoff type problem involving a nonlocal operator, *Nonlinear Anal., Theory Methods Appl.*, 94:156–170, 2014, <https://doi.org/10.1016/j.na.2013.08.011>.
10. X. He, W. Zou, Ground states for nonlinear Kirchhoff equations with critical growth, *Ann. Mat. Pura Appl. (4)*, 193(2):473–500, 2014, <https://doi.org/10.1007/s10231012-0286-6>.
11. Z. Liu, Z.-Q. Wang, On Clark’s theorem and its applications to partially sublinear problems, *Ann. Inst. Henri Poincaré, Anal. Non Linéaire*, 32(5):1015–1037, 2015, <https://doi.org/10.1016/j.anihpc.2014.05.002>.
12. A. Mao, Z. Zhang, Sign-changing and multiple solutions of Kirchhoff type problems without the P.S. condition, *Nonlinear Anal., Theory Methods Appl.*, 70(3):1275–1287, 2009, <https://doi.org/10.1016/j.na.2008.02.011>.
13. C. Miranda, Un’osservazione su un teorema di Brouwer, *Boll. Un. Mat. Ital.* (2), 3:5–7, 1940.
14. G. Molica Bisci, V.D. Radulescu, R. Servadei, *Variational methods for nonlocal fractional problems*, Encycl. Math. Appl., Vol. 162, Cambridge Univ. Press, Cambridge, 2016, <https://doi.org/10.1017/CBO9781316282397>.
15. E. Di Nezza, G. Palatucci, E. Valdinoci, Hitchhiker’s guide to the fractional Sobolev spaces, *Bull. Sci. Math.*, 136(5):521–573, 2012, <https://doi.org/10.1016/j.bulsci.2011.12.004>.
16. V.D. Radulescu, D.D. Repovš, *Partial Differential Equations with Variable Exponents. Variational Methods and Qualitative Analysis*, Monogr. Res. Notes Math., Chapman and Hall/CRC, New York, 2015, <https://doi.org/10.1201/b18601>.

17. R. Servadei, E. Valdinoci, Mountain pass solutions for non-local elliptic operators, *J. Math. Anal. Appl.*, 389(2):887–898, 2012, <https://doi.org/10.1016/j.jmaa.2011.12.032>.
18. W. Shuai, Sign-changing solutions for a class of Kirchhoff-type problem in bounded domains, *J. Differ. Equations*, 259(4):1256–1274, 2015, <https://doi.org/10.1016/j.jde.2015.02.040>.
19. X. Tang, B. Cheng, Ground state sign-changing solutions for Kirchhoff type problems in bounded domains, *J. Differ. Equations*, 261(4):2384–2402, 2016, <https://doi.org/10.1016/j.jde.2016.04.032>.
20. D. Wang, Least energy sign-changing solutions of Kirchhoff-type equation with critical growth, *J. Math. Phys.*, 61(1):011501, 2020, <https://doi.org/10.1063/1.5074163>.
21. L. Wang, B. Zhang, Infinitely many solutions for Kirchhoff-type variable-order fractional Laplacian problems involving variable exponents, *Appl. Anal.*, 100(11):2418–2435, 2021, <https://doi.org/10.1080/00036811.2019.1688790>.
22. T. Weth, Energy bounds for entire nodal solutions of autonomous superlinear equations, *Calc. Var. Partial Differ. Equ.*, 27(4):421–437, 2006, <https://doi.org/10.1007/s00526006-0015-3>.
23. M. Willem, *Minimax Theorems*, Prog. Nonlinear Differ. Equ. Appl., Vol. 24, Birkhäuser, Boston, MA, 1996, <https://doi.org/10.1007/978-1-4612-4146-1>.
24. M. Xiang, D. Hu, B. Zhang, Y. Wang, Multiplicity of solutions for variable-order fractional Kirchhoff equations with nonstandard growth, *J. Math. Anal. Appl.*, 501(1):124269, 2021, <https://doi.org/10.1016/j.jmaa.2020.124269>.
25. M. Xiang, B. Zhang, D. Yang, Multiplicity results for variable-order fractional Laplacian equations with variable growth, *Nonlinear Anal.*, 178:190–204, 2019, <https://doi.org/10.1016/j.na.2018.07.016>.
26. Z. Zhang, K. Perera, Sign changing solutions of Kirchhoff type problems via invariant sets of descent flow, *J. Math. Anal. Appl.*, 317(2):456–463, 2006, <https://doi.org/10.1016/j.jmaa.2005.06.102>.

Notas de autor

- a College of Mathematics, Changchun Normal University, Changchun, 130032, China liangsihua@163.com
- 1 The author was supported by the Foundation for China Postdoctoral Science Foundation (grant No. 2019M662220), Scientific research projects for Department of Education of

Jilin Province, China (JJKH20210874KJ), and Natural Science Foundation of Jilin Province.

- b Dipartimento di Scienze Pure e Applicate (DiSPeA), Università degli Studi di Urbino Carlo Bo, Urbino, 61029, Italy giovanni.molicabisci@uniurb.it
 - c College of Mathematics and Systems Science, Shandong University of Science and Technology, Qingdao, 266590, China zhangbinlin2012@163.com
- 2 The author was supported by National Natural Science Foundation of China (11871199 and 12171152), Shandong Provincial Natural Science Foundation, PR China (ZR2020MA006), and Cultivation Project of Young and Innovative Talents in Universities of Shandong Province.
 - 3 Corresponding author.