



Nonlinear Analysis: Modelling and Control
ISSN: 1392-5113
ISSN: 2335-8963
nonlinear@mii.vu.lt
Vilniaus Universitetas
Lituania

Stability of solutions of Caputo fractional stochastic differential equations*

Xiao, Guanli; Wang, JinRong

Stability of solutions of Caputo fractional stochastic differential equations*

Nonlinear Analysis: Modelling and Control, vol. 26, núm. 4, 2021

Vilniaus Universitetas, Lituania

Disponible en: <https://www.redalyc.org/articulo.oa?id=694173200002>

DOI: <https://doi.org/10.15388/namc.2021.26.22421>



Esta obra está bajo una Licencia Creative Commons Atribución 4.0 Internacional.


Stability of solutions of Caputo fractional stochastic differential equations*

Guanli Xiao glxiaomath@126.com

Guizhou University, China

JinRong Wang sci.jrwang@gzu.edu.cn

Qufu Normal University, China

 <https://orcid.org/0000-0002-6642-1946>

Abstract: In this paper, we study the stability of Caputo-type fractional stochastic differential equations. Stochastic stability and stochastic asymptotical stability are shown by stopping time technique. Almost surely exponential stability and p -th moment exponentially stability are derived by a new established Itô's formula of Caputo version. Numerical examples are given to illustrate the main results.

Keywords: Caputo fractional derivative, stochastic differential equations, stability.

Nonlinear Analysis: Modelling and Control, vol. 26, núm. 4, 2021

Vilniaus Universitetas, Lithuania

Recepción: 21 Marzo 2020

Revisado: 20 Agosto 2020

Publicación: 01 Julio 2021

DOI: <https://doi.org/10.15388/namc.2021.26.22421>

Redalyc: <https://www.redalyc.org/articulo.oa?id=694173200002>

1 Introduction

Recently, fractional derivative is used to study the properties of memory and genetic for complex systems in different fields of application; see [2, 6]. Wu et al. [23] introduced a new result for Mittag-Leffler stability analysis of fractional discrete-time neural networks via fixed point technique. Huang et al. [10] gave the analysis for variable order fractional discrete-time recurrent neural networks. For more details on fractional differential equations, we refer to the monographs [12, 25].

Stochastic is one of the essential properties of the world, and the stability is a top priority to the system application in practice. Stability analysis of stochastic systems is very necessary. Lyapunov direct method and frequency analysis method have been popular among scholars for their intuitive concept, general method, clear physical meaning and rigorous theory and have become the main tools to study the stability of differential systems. At the same time, Burton et al. [3,4,5] presented the application of fixed point technology to stability analysis and fractional differential equations. Existence, uniqueness and stability of solutions to stochastic partial differential equations have been studied by many researchers; see [2, 8, 15, 17, 20]. Many new achievements have been used in dealing with stochastic integral and differential equations; see [1, 9, 11, 18, 19, 21, 24]. For more details regarding functional analysis and mathematical analysis, we refer to [13, 14, 22].

Recently, Doan et al. [8] studied asymptotic separation between solutions of the following Caputo fractional stochastic differential equations:

$$\begin{aligned} {}^C D_{0+}^{\alpha} X(t) &= f(t, X(t)) + g(t, X(t)) \frac{dW(t)}{dt}, \quad t > 0, \alpha \in (0, 1], \\ X(0) &= X_0, \end{aligned} \quad (1)$$

where $CD\alpha$ denotes the Caputo fractional derivative, $f, g : [0, +\infty) \times \mathbb{R} \rightarrow \mathbb{R}$ are measurable functions, and $\{W(t), t \in [0, +\infty)\}$ is a standard scalar Brownian motion on an underlying complete filtered probability space $(\Omega, \mathcal{F}, \mathcal{F} := \{\mathcal{F}_t\}_{t \in [0, +\infty)}, P)$.

For each $t \in [0, +\infty)$, $X_t := L^2(\Omega, \mathcal{F}_t, P)$ denotes the space of all t -measurable, mean-squared integrable functions $u : \Omega \rightarrow \mathbb{R}$ with $\|u\|_2^2 = E |u|^2$. A process $X : [0, +\infty) \rightarrow L^2(\Omega, \mathcal{F}, P)$ is called \mathcal{F} -adapted if $X(t) \in \mathcal{F}_t$, $t \in [0, +\infty)$.

Doan et al. [8, Thm. 1] applied contraction mapping principle to derive the existence and uniqueness result of (1) by imposing the Lipschitz condition on f and g .

Note that asymptotic behavior and exponential stability of fractional stochastic differential equations in the sense of expected have been studied in [6, 16]. However, fractional Itô formula of Caputo version, stochastic stability and stochastic asymptotic stability of (1) in a probabilistic sense have not been established.

In this paper, we present stochastic stability, stochastic asymptotic stability, almost surely exponential stability and p th moment exponential stability for (1). In Section 3, we establish a new established fractional Itô's formula of Caputo version, and in Section 4 we present the main results. Numerical simulation illustrates our theoretical results in the final section.

2 Preliminaries

Definition 1. (See [12].) Let $\alpha \in (0, 1]$, $T > 0$, $f : [0, T] \rightarrow \mathbb{R}$ be a differentiable function. The Caputo fractional derivative of f is defined as

$${}^C D_{0+}^{\alpha} f(t) = \frac{1}{\Gamma(1-\alpha)} \int_0^t \frac{f'(\tau)}{(t-\tau)^{\alpha}} d\tau, \quad t \in [0, T],$$

where $\Gamma(\alpha) := \int_0^{\infty} \tau^{\alpha-1} e^{-\tau} d\tau$ is Gamma function.

Definition 2. (See [12].) Let $\alpha \in (0, 1]$, $f : [0, +\infty) \rightarrow \mathbb{R}$. The Caputo fractional integral of f is defined as

$$I_{0+}^{\alpha} f(t) = \frac{1}{\Gamma(\alpha)} \int_0^t (t-\tau)^{\alpha-1} f(\tau) d\tau, \quad t \in [0, T].$$

Definition 3. (See [8].) For each $X_0 \in X_0$, a F -adapted X is called a solution of (1) if the following holds for $t \in [0, +\infty)$:

$$\begin{aligned} X(t) &:= X(t; X_0) \\ &= X_0 + \frac{1}{\Gamma(\alpha)} \left(\int_0^t (t-\tau)^{\alpha-1} f(\tau, X(\tau)) d\tau \right. \\ &\quad \left. + \int_0^t (t-\tau)^{\alpha-1} g(\tau, X(\tau)) dW(\tau) \right). \end{aligned} \quad (2)$$

we introduce the following assumptions:

(H1) There exists a constant $L > 0$ such that for all $X, X^* \in \mathbb{R}$, $t \in [0, +\infty)$,

$$|f(t, X) - f(t, X^*)| + |g(t, X) - g(t, X^*)| \leq L|X - X^*|.$$

(H2) $g(\cdot, 0)$ is essentially bounded, i.e.,

$$\|g(t, 0)\|_\infty := \operatorname{ess\,sup}_{t \in [0, +\infty)} |g(t, 0)| < +\infty,$$

and $g(\cdot, 0)$ is L^2 integrable, i.e., $\int_0^{+\infty} |g(t, 0)|^2 dt < +\infty$.

Lemma 1. (See [8, Thm. 1].) Suppose that (H1) and (H2) hold. Then for $\alpha \in (1/2, 1)$, (1) has a unique solution $X(\cdot) \in X_t := L^2(\Omega, \mathcal{F}_t, P)$ given by (2).

Definition 4. (See [15, Def. 2.1].)

(i) The trivial solution of (1) is said to be stochastically stable or stable in probability if for every pair of $\varepsilon \in (0, 1)$ and $r > 0$, there exists a $\delta = \delta(\varepsilon, r, 0) > 0$ such that $P\{|X(t)| < r\} \geq 1 - \varepsilon$, $t \geq 0$, whenever $|X_0| < \delta$. Otherwise, it is said to be stochastically unstable.

(ii) The trivial solution of (1) is said to be stochastically asymptotically stable if it is stochastically stable and, moreover, for every $\varepsilon \in (0, 1)$, there exists a $\delta_0 = \delta_0(\varepsilon) > 0$ such that $P\{\lim_{t \rightarrow +\infty} X(t) = 0\} \geq 1 - \varepsilon$ whenever $|X_0| < \delta_0$ and 0 denotes n -dimensional zero vector.

Definition 5. (See [15, Def. 3.1].) The trivial solution of (1) is said to be almost surely exponentially stable if

$$\lim_{t \rightarrow +\infty} \sup \frac{1}{t} \ln |X(t)| < 0 \quad \text{a.s.}$$

for all $X_0 \in \mathbb{R}^n$, where “a.s.” means “almost surely”.

Definition 6. (See [15, Def. 4.1].) The trivial solution of (1) is said to be p th moment exponentially stable if there is a pair of positive constants

λ and C such that $E(\|X(t)\|^p) \leq C \|X_0\|^p e^{-\lambda t}$ for all $t \geq 0$, $X_0 \in \mathbb{R}$. If $p = 2$, then it reduces to exponentially stable in mean square.

Lemma 2. (See [15, Lemma 2.4].) Let \mathcal{F} is a σ -algebra, and let $\{A_k\} \subset \mathcal{F}$ and $\sum_{k=1}^{\infty} P(A_k) < +\infty$. Then $P(\lim_{k \rightarrow \infty} \sup A_k) = 0$.

3 Itô's formula of Caputo version

In this part, we introduce the Itô's formula of Caputo fractional version. It pointed out the rules for differentiating a function of Caputo fractional stochastic process.

Let $W(t)$, $t \geq 0$, be a standard scalar Brownian motion, and let $Y \in C^{2,1}(\mathbb{R} \times \mathbb{R}_+, \mathbb{R})$ denotes the family of all real-valued functions $Y(z, t)$ defined on $\mathbb{R} \times \mathbb{R}_+$ such that they are continuously twice differentiable in z and once in t .

Let $Z(t)$, $t \geq 0$, be an Itô process for $dZ(t) = \tilde{f}(t)dt + \tilde{g}(t)dW(t)$, where $\tilde{f} \in L^1(\mathbb{R}_+, \mathbb{R})$ and $\tilde{g} \in L^2(\mathbb{R}_+, \mathbb{R})$.

We recall the standard one-dimensional Itô formula.

Lemma 3. (See [15, p. 32, Thm. 6.2].) Let $Y(\cdot) := Y(Z(\cdot), \cdot) \in C^{2,1}(\mathbb{R} \times \mathbb{R}_+, \mathbb{R})$. Then $Y(t)$, $t \geq 0$ is an Itô process given by

$$dY(t) = \left[Y_t(Z(t), t) + Y_Z(Z(t), t)\tilde{f}(t) + \frac{1}{2}Y_{ZZ}(Z(t), t)\tilde{g}^2(t) \right] dt + Y_Z(Z(t), t)\tilde{g}(t)dW(t) \quad a.s.$$

Now, set $T > 0$. Suppose that $X(t)$ is an Itô process for

$${}^C D_{0+}^\alpha \tilde{X}(t) = f(t) + g(t) \frac{dW(t)}{dt}, \quad t \in [0, T], \quad \alpha \in \left(\frac{1}{2}, 1 \right), \quad (3)$$

with the initial condition $X(0) = X_0$.

From Lemma 1, (3) has the following unique solution for $t \in [0, T]$:

$$\tilde{X}(t) = X_0 + \frac{1}{\Gamma(\alpha)} \left(\int_0^t (t-\tau)^{\alpha-1} f(\tau) d\tau + \int_0^t (t-\tau)^{\alpha-1} g(\tau) dW(\tau) \right).$$

Note that when $t \in [0, T]$, (3) has an equivalent form

$$\begin{aligned} d\tilde{X}(t) &= \tilde{X}'(t)dt \\ &= \frac{\alpha-1}{\Gamma(\alpha)} \left(\int_0^t (t-\tau)^{\alpha-2} f(\tau) d\tau + \int_0^t (t-\tau)^{\alpha-2} g(\tau) dW(\tau) \right) dt, \end{aligned} \quad (4)$$

where $f(\cdot)(t-\cdot)^{\alpha-2} \in L^1[0, T]$ and $g(\cdot)(t-\cdot)^{\alpha-2} \in L^2[0, T]$.

Now, we are ready to introduce the following fractional Itô's formula of the Caputo version.

Theorem 1. Let $Y(\cdot) := Y(X(\cdot), \cdot) \in C^{2,1}(\mathbb{R} \times \mathbb{R}_+, \mathbb{R})$. Then $Y(\cdot)$ is an Itô process given by

$$\begin{aligned} dY(\tilde{X}(t), t) &= Y_t(\tilde{X}(t), t) dt \\ &+ \frac{\alpha-1}{\Gamma(\alpha)} Y_{\tilde{X}}(\tilde{X}(t), t) \int_0^t (t-\tau)^{\alpha-2} f(\tau) d\tau dt \\ &+ \frac{\alpha-1}{\Gamma(\alpha)} Y_{\tilde{X}}(\tilde{X}(t), t) \int_0^t (t-\tau)^{\alpha-2} g(\tau) dW(\tau) dt. \end{aligned}$$

Proof. From Lemma 3 via (4) we can derive that

$$\begin{aligned} dY(\tilde{X}(t), t) &= \frac{\partial Y(\tilde{X}(t), t)}{\partial t} dt + \frac{\partial Y(\tilde{X}(t), t)}{\partial \tilde{X}} d\tilde{X}(t) + \frac{1}{2} \frac{\partial^2 Y(\tilde{X}(t), t)}{\partial \tilde{X}^2} (d\tilde{X}(t))^2 \\ &= Y_t(\tilde{X}(t), t) dt + \frac{\alpha-1}{\Gamma(\alpha)} Y_{\tilde{X}}(\tilde{X}(t), t) \int_0^t (t-\tau)^{\alpha-2} f(\tau) d\tau dt \\ &+ \frac{\alpha-1}{\Gamma(\alpha)} Y_{\tilde{X}}(\tilde{X}(t), t) \int_0^t (t-\tau)^{\alpha-2} g(\tau) dW(\tau) dt. \end{aligned}$$

The proof is completed.

4 Stability results

Let $k > 0$ be arbitrary, denote $S_k := \{X(\cdot) \in \mathbb{R} : |X(\cdot)| < k\}$, $a \wedge b$ denotes the minimum of a and b , $a \vee b$ denotes the maximum of a and b , $I\{\cdot\}$ is indicative function.

(V1) There exists a positive definite function $V \in C^{2,1}(S_k \times [0, +\infty); \mathbb{R}_+)$ such that for all $(X(t), t) \in (S_k \times [0, +\infty))$, $\alpha \in (1/2, 1)$,

$$\begin{aligned} L^\alpha V(X(t), t) &:= V_t(X(t), t) \\ &+ \frac{\alpha-1}{\Gamma(\alpha)} V_X(X(t), t) \int_0^t f(X(\tau), \tau) (t-\tau)^{\alpha-2} d\tau \\ &\leq 0. \end{aligned} \tag{5}$$

From (V1), $V(0, t) \equiv 0$, and there is a continuous nondecreasing function μ such that $V(X(t), t) \geq \mu(|X(t)|)$ for all $(X(t), t) \in (S_k \times [0, +\infty))$.

The following result is motivated by [15, p. 111, Thm. 2.2].

Theorem 2. Assume (H1), (H2) and (V1) hold, $\alpha \neq (1/2, 1)$. Then the trivial solution of (1) is stochastically stable.

Proof. Since the proof is similar to [15, p. 111, Thm. 2.2], we only give the sketch of proofs.

Let $\varepsilon \neq (0, 1)$ and $r > 0$ be arbitrary and assume that $r < k$. By the continuity of V and $V(0, 0) = 0$ one can find $\delta = \delta(\varepsilon, r) > 0$ such that

$$\frac{1}{\varepsilon} \sup_{X \in S_\delta} V(X(t), t) \leq \mu(r). \quad (6)$$

Obviously, $\delta < r$. Fix $X_0 \notin S_\delta$ and let η be the first exit time of $X(t)$ from S_r , i.e., $\eta = \inf\{t > 0: X(t) \notin S_r\}$.

By Theorem 1, for any $t > 0$, we can get

$$\begin{aligned} & V(X(\eta \wedge t), \eta \wedge t) \\ &= V(X_0, 0) + \int_0^{\eta \wedge t} V_s(X(s), s) \, ds \\ & \quad + \frac{\alpha - 1}{\Gamma(\alpha)} \int_0^{\eta \wedge t} V_X(X(s), s) \int_0^s (s - \tau)^{\alpha-2} f(X(\tau), \tau) \, d\tau \, ds \\ & \quad + \frac{\alpha - 1}{\Gamma(\alpha)} \int_0^{\eta \wedge t} V_X(X(s), s) \int_0^s (s - \tau)^{\alpha-2} g(X(\tau), \tau) \, dW(\tau) \, ds \\ &= V(X_0, 0) + \int_0^{\eta \wedge t} L^\alpha V(X(\tau), \tau) \, d\tau \\ & \quad + \frac{\alpha - 1}{\Gamma(\alpha)} \int_0^{\eta \wedge t} V_X(X(s), s) \int_0^s (s - \tau)^{\alpha-2} g(X(\tau), \tau) \, dW(\tau) \, ds. \end{aligned} \quad (7)$$

Take the expectation on (7) and note $L^\alpha V \leq 0$. By modulus inequality, for any $t > 0$, $1/2 < \alpha < 1$, we can get

$$\begin{aligned} & \frac{\alpha - 1}{\Gamma(\alpha)} \left| \mathbf{E} \left(\int_0^{\eta \wedge t} V_X(X(s), s) \int_0^s (s - \tau)^{\alpha-2} g(X(\tau), \tau) \, dW(\tau) \, ds \right) \right| \\ & \leq \frac{\alpha - 1}{\Gamma(\alpha)} \mathbf{E} \left| \int_0^{\eta \wedge t} V_X(X(s), s) \left(\int_0^s (s - \tau)^{\alpha-2} g(X(\tau), \tau) \, dW(\tau) \right) \, ds \right| \\ & \leq 0. \end{aligned}$$

Following the same procedure in the proof of [15, p. 111, Thm. 2.2] via (6), one has

$$P\{\eta \leq t\} \leq \varepsilon. \text{ Let } t \rightarrow +\infty, \text{ i.e., } P\{\eta < +\infty\} \leq \varepsilon.$$

Then we have $P\{X(t) \leq r\} \geq 1 - \varepsilon$ for all $t \geq 0$. By Definition 4(i) the trivial solution of (1) is stochastically stable.

(V2) The positive definite decreasing function $V \in C^{2,1}(S_k \times [0, +\infty); \mathbb{R}^+)$ such that $L^\alpha V < 0$, $\alpha \in (1/2, 1)$, where $L^\alpha V$ is defined in (5).

From (V2), $V(0, t) = 0$, and there exist continuous nondecreasing functions μ_1, μ_2, μ_3 such that

$$\mu_1(|X(t)|) \leq V(X(t), t) \leq \mu_2(|X(t)|), \quad L^\alpha V(X(t), t) \leq -\mu_3(|X(t)|)$$

for all $(X(t), t) \in (S_k \times [0, +\infty))$.

The following result is motivated by [15, p. 112, Thm. 2.3].

Theorem 3. Assume (H1), (H2) and (V2) hold. Then the trivial solution of (1) is stochastically asymptotically stable.

Proof. From Theorem 2 the trivial solution of (1) is stochastically stable. Following the same procedure in the proof of [15, p. 112, Thm. 2.3] and using Theorem 1, one can show that there exists $\delta_0 = \delta_0(\varepsilon) > 0$ such that $P\{\lim_{t \rightarrow +\infty} X(t) = 0\} \geq 1 - \varepsilon$ for $|X_0| < \delta_0$, $\varepsilon \in (0, 1)$.

Based on Definition 4(ii), the trivial solution of (1) is stochastically asymptotically stable.

(V3) $V \in C^{2,1}(\mathbb{R}^n \times [0, +\infty); \mathbb{R}^+)$, and there exist constants $c_1 > 1$, $c_2 \in \mathbb{R}$, $c_3 \geq 0$ such that

- (i) $c_1|X(t)| \leq V(X(t), t)$,
- (ii) $L^\alpha V(X(t), t) \leq c_2 V(X(t), t)$,
- (iii) $|V_x(X(t), t)|^2 \int_0^t |g(X(t), t)(s - \tau)^{\alpha-2}|^2 d\tau \geq c_3 V^2(X(t), t)$ for all $X(t) \neq 0$, $\alpha \in (1/2, 1)$ and $t \geq 0$.

The following result is motivated by [15, p. 121, Thm. 3.3].

Theorem 4. Assume (H1), (H2) and (V3) hold. Then

$$\limsup_{t \rightarrow +\infty} \frac{1}{t} \ln |X(t)| \leq -\frac{1}{\ln c_1} \frac{1 - \alpha}{\Gamma(\alpha)} (c_2 + c_3) \quad a.s. \quad (8)$$

In particular, if $c_2 + c_3 > 0$, the trivial solution of (1) is almost surely exponentially stable.

Proof. Fix any $X_0 \neq 0$. From Theorem 1 and (V3)(ii), (iii), for $\alpha \in (1/2, 1)$, we have

$$\begin{aligned}\ln V(X(t), t) &= \ln V(X_0, 0) + \int_0^t \frac{V_s(X(s), s)}{V(X(s), s)} ds \\ &\quad + \frac{\alpha - 1}{\Gamma(\alpha)} \int_0^t \frac{V_X(X(s), s) \int_0^s f(X(\tau), \tau)(s - \tau)^{\alpha-2} d\tau}{V(X(s), s)} ds \\ &\quad + \frac{\alpha - 1}{\Gamma(\alpha)} \int_0^t \frac{V_X(X(s), s) \int_0^s g(X(\tau), \tau)(s - \tau)^{\alpha-2} dW(\tau)}{V(X(s), s)} ds \\ &\leq \ln V(X_0, 0) + \int_0^t \frac{L^\alpha V(X(s), s)}{V(X(s), s)} ds \\ &\quad + \frac{\alpha - 1}{\Gamma(\alpha)} \int_0^t \frac{V_X(X(s), s) \int_0^s g(X(\tau), \tau)(s - \tau)^{\alpha-2} dW(\tau)}{V(X(s), s)} ds.\end{aligned}$$

Set $M(t) = \int_0^t (V_X(X(s), s) \int_0^s g(X(\tau), \tau)(s - \tau)^{\alpha-2} dW(\tau) / V(X(s), s)) ds$. Then let $n = 1, 2, \dots$. For arbitrary $\varepsilon \in (0, 1)$, by (V3)(iii) we can get

$$\mathbf{P} \left\{ \sup_{0 \leq t \leq n} \left| M(t) + \varepsilon \int_0^t \frac{V_X^2(X(s), s) \int_0^s |g(X(\tau), \tau)(s - \tau)^{\alpha-2}|^2 d\tau}{V^2(X(s), s)} ds \right| \geq c_3 t \right\} \leq \varepsilon.$$

Using Lemma 2, we get that almost surely

$$\begin{aligned}M(t) &\leq c_3 t - \varepsilon \int_0^t \frac{V_X^2(X(s), s) \int_0^s |g(X(\tau), \tau)(s - \tau)^{\alpha-2}|^2 d\tau}{V^2(X(s), s)} ds \\ &= (1 - \varepsilon)c_3 t.\end{aligned}\tag{9}$$

Thus, by (V3)(iii) and (9) we have

$$\ln V(X(t), t) \leq \ln V(X_0, 0) - \frac{1 - \alpha}{\Gamma(\alpha)} [c_2 + (1 - \varepsilon)c_3] t.$$

Then we can get

$$\frac{1}{t} \ln V(X(t), t) \leq -\frac{1 - \alpha}{\Gamma(\alpha)} [c_2 + (1 - \varepsilon)c_3] + \frac{\ln V(X_0, 0)}{t}.$$

Thus

$$\lim_{t \rightarrow +\infty} \sup \frac{1}{t} \ln V(X(t), t) \leq -\frac{1 - \alpha}{\Gamma(\alpha)} [c_2 + (1 - \varepsilon)c_3].$$

Using (V3)(i),

$$\lim_{t \rightarrow +\infty} \sup \frac{1}{t} \ln c_1 |X(t)| \leq \lim_{t \rightarrow \infty} \frac{1}{t} \ln V(X(t), t) \leq -\frac{1-\alpha}{\Gamma(\alpha)} [c_2 + (1-\varepsilon)c_3].$$

Finally, we get

$$\lim_{t \rightarrow +\infty} \sup \frac{1}{t} \ln |X(t)| \leq -\frac{1}{\ln c_1} \frac{1-\alpha}{\Gamma(\alpha)} [c_2 + (1-\varepsilon)c_3].$$

Since ε is arbitrarily, we have (8).

Note that $c_1 > 1$, then, if $c_2 + c_3 > 0$, we have $-(1/\ln c_1)((1-\alpha)/\Gamma(\alpha))(c_2 + c_3) < 0$. By Definition 5 the trivial solution of (1) is almost surely exponentially stable.

(V4) $V \in C^{2,1}(\mathbb{R}^n \times [0, +\infty); \mathbb{R}_+)$, and there exist constants $c_1 > 1$, $c_2 \in \mathbb{R}$, $c_3 \geq 0$ such that

- (i) $c_1 |X(t)| \geq V(X(t), t) > 0$,
- (ii) $L_\alpha V(X(t), t) \geq c_2 V(X(t), t)$,
- (iii) $|V_x(X(t), t)|^2 \int_0^t |g(X(t), t)(s-\tau)^{\rho-1}|^2 d\tau \leq c V^2(X(t), t)$ for all $X(t) \neq 0$ and $t \geq 0$, $1/2 < \alpha < 1$.

The following result is motivated by [15, p. 123, Thm. 3.5].

Remark 1. Assume (H1), (H2) and (V4) hold. Then

$$\lim_{t \rightarrow +\infty} \inf \frac{1}{t} \ln |X(t)| \geq -\frac{1}{\ln c_1} \frac{1-\alpha}{\Gamma(\alpha)} (c_2 + c_3) \quad \text{a.s.} \quad (10)$$

If $c_2 + c_3 < 0$, then $-(1/\ln c_1)((1-\alpha)/\Gamma(\alpha))(c_2 + c_3) > 0$. Thus, almost all the sample paths of $X(\cdot)$ will tend to infinity (see (10)), i.e., the trivial solution of (1) is almost surely exponentially unstable.

(H3) The following inequality holds for $K > 0$:

$$\int_0^t f(X(\tau), \tau) (t-\tau)^{\alpha-2} d\tau \leq K |X(t)|, \quad (X, t) \in \mathbb{R} \times [0, +\infty). \quad (11)$$

The following result is motivated by [15, p. 128, Thm. 4.2].

Theorem 5. Assume (H1), (H2) and (H3) hold. Then the trivial solution of (1) is pth moment exponentially stable (also almost surely exponentially stable).

Proof. Let $n = 1, 2, \dots$. By Theorem 1 and (11), for any $n-1 \leq t \leq n$,

$$\begin{aligned}
 |X(t)|^p &= |X_0|^p + \frac{\alpha-1}{\Gamma(\alpha)} \int_0^t p |X(s)|^{p-1} \int_0^s f(X(\tau), \tau) (s-\tau)^{\alpha-2} d\tau ds \\
 &\quad + \frac{\alpha-1}{\Gamma(\alpha)} \int_0^t p |X(s)|^{p-1} \int_0^s g(X(\tau), \tau) (s-\tau)^{\alpha-2} dW(\tau) ds \\
 &\leq |X_0|^p + pK \frac{\alpha-1}{\Gamma(\alpha)} \int_0^t |X(s)|^p ds \\
 &\quad + \frac{\alpha-1}{\Gamma(\alpha)} \int_0^t p |X(s)|^{p-1} \int_0^s g(X(\tau), \tau) (s-\tau)^{\alpha-2} dW(\tau) ds.
 \end{aligned}$$

Since $1/2 < \alpha < 1$, by modulus inequality we obtain

$$\begin{aligned}
 &\frac{\alpha-1}{\Gamma(\alpha)} \left| \mathbb{E} \left(\int_0^t p |X(s)|^{p-1} \int_0^s g(X(\tau), \tau) (s-\tau)^{\alpha-2} dW(\tau) ds \right) \right| \\
 &\leq \frac{p(\alpha-1)}{\Gamma(\alpha)} \mathbb{E} \left| \int_0^t |X(s)|^{p-1} \left(\int_0^s g(X(\tau), \tau) (s-\tau)^{\alpha-2} dW(\tau) \right) ds \right| \\
 &\leq 0.
 \end{aligned}$$

Hence,

$$\mathbb{E} \left(\sup_{0 \leq t \leq n} |X(t)|^p \right) \leq |X_0|^p + pK \frac{\alpha-1}{\Gamma(\alpha)} \int_0^t \mathbb{E} \left(\sup_{0 \leq s \leq n} |X(s)|^p \right) ds.$$

Using Gronwall's inequality,

$$\mathbb{E} \left(\sup_{0 \leq t \leq n} |X(t)|^p \right) \leq |X_0|^p e^{-\lambda t}, \quad \lambda = \frac{1}{\Gamma(\alpha)} pK(1-\alpha) > 0.$$

By Definition 6 the trivial solution of (1) is pth moment exponentially stable. Now, let $\varepsilon \in (0, 1)$ be arbitrary, then

$$\begin{aligned}
 \mathbb{P} \left\{ \sup_{0 \leq t \leq n} |X(t)|^p > e^{-(\lambda-\varepsilon)t} \right\} &\leq e^{(\lambda-\varepsilon)t} \cdot \mathbb{E} \left(\sup_{0 \leq t \leq n} |X(t)|^p \right) \\
 &\leq |X_0|^p e^{-\varepsilon t}.
 \end{aligned}$$

By Lemma 2 we get

$$\sup_{n-1 \leq t \leq n} |X(t)|^p \leq e^{-(\lambda-\varepsilon)t} \quad \text{a.s.}$$

Consequently, for almost all $\omega \in \Omega$, $n-1 \leq t \leq n$,

$$\frac{1}{t} \ln |X(t)| \leq \frac{1}{pt} \ln |X(t)|^p \leq -\frac{1}{pt} (\lambda - \varepsilon)t \leq -\frac{(\lambda - \varepsilon)t}{pn}.$$

Hence,

$$\lim_{t \rightarrow +\infty} \sup \frac{1}{t} \ln |X(t)| \leq -\frac{\lambda - \varepsilon}{p} \quad \text{a.s.}$$

Since $\varepsilon > 0$ is arbitrary, we obtain

$$\lim_{t \rightarrow +\infty} \sup \frac{1}{t} \ln |X(t)| \leq -\frac{\lambda}{p} \quad \text{a.s.}$$

By Definition 5 the trivial solution of (1) is almost surely exponential stability.

5 Examples

The following result is motivated by [15, p. 121, Thm. 3.3].

Example 1. Consider the stochastic differential equation on

$$\begin{aligned} {}^C D_{0+}^{\alpha} X(t) &= f(X(t), t) + g(X(t), t) \frac{dW(t)}{dt}, \quad t \geq 0, \quad \alpha \in \left(\frac{1}{2}, 1\right), \\ X(0) &= X_0, \end{aligned} \quad (12)$$

where $X_0 \in \mathbb{R}$ and $X_0 \neq \infty$, $f: \mathbb{R} \times \mathbb{R}_+ \rightarrow \mathbb{R}$, $g: \mathbb{R} \times \mathbb{R}_+ \rightarrow \mathbb{R}$.

Let the functions f, g satisfy assumptions (H1) and (H2), and for any $t \geq 0$,

$$\int_0^t f(X(\tau), \tau) (t - \tau)^{\alpha-2} d\tau < +\infty. \quad (13)$$

Then for any $t \geq 0$, there is a pair of positive constants θ and K such that

$$-K \leq \frac{\alpha-1}{\Gamma(\alpha)} \int_0^t f(X(\tau), \tau) (t - \tau)^{\alpha-2} d\tau + \theta \leq K. \quad (14)$$

Next, let $t \geq 0$, we define

$$V(X(t), t) = e^{|X(t)|} \exp \left[-\frac{\alpha-1}{\Gamma(\alpha)} \int_0^t f(X(\tau), \tau)(t-\tau)^{\alpha-2} d\tau + \theta \right].$$

By (14), for $t \geq 0$, we can get

$$e^{|X(t)|} e^{-\varepsilon K} \leq V(X(t), t) \leq e^{|X(t)|} e^{\varepsilon K}.$$

Hence, V is positive definite. Next, by (13), for $t \geq 0$, we have

$$\begin{aligned} L^\alpha V(X(t), t) &= e^{|X(t)|} \exp \left[-\frac{\alpha-1}{\Gamma(\alpha)} \int_0^t f(X(\tau), \tau)(t-\tau)^{\alpha-2} d\tau + \theta \right] \\ &\quad \times \left(\frac{\alpha-1}{\Gamma(\alpha)} \int_0^t f(X(\tau), \tau)(t-\tau)^{\alpha-2} d\tau + \theta \right) \\ &\quad + e^{|X(t)|} \exp \left[-\frac{\alpha-1}{\Gamma(\alpha)} \int_0^t f(X(\tau), \tau)(t-\tau)^{\alpha-2} d\tau + \theta \right] \\ &\quad \times \frac{\alpha-1}{\Gamma(\alpha)} \int_0^t f(X(\tau), \tau)(t-\tau)^{\alpha-2} d\tau \\ &\leq -\theta e^{-K} e^{|X(t)|} < 0. \end{aligned}$$




Figure 1
Brownian motion $W(t)$ and white noise $dW(t)/dt$, $t \in [0, 500]$.




Figure 2
Stochastic stability and instability when $\alpha = 1$.

From above (V2) holds. By Theorem 2 we can conclude that under (13) and (14), the trivial solution of (12) is stochastically stable.

Let r and β be constants when $\alpha = 1$, $f(X, t) = rX$ and $g(X, t) = \beta X$, the solution of (12) can be seen in [15, Ex. 5.5]. Our chosen Brownian motion and white noise are shown in Fig. 1. Using MATLAB software, trajectories of stochastically stable on $r < \beta^2/2$ and $r > \beta^2/2$ and corresponding unstable results are shown in Fig. 2.

Example 2. We consider the one-dimensional fractional Langevin equation on $t \geq 0$

$${}^C D_{0+}^{\alpha} X(t) = -rX(t) + \sigma \frac{dW(t)}{dt}, \quad \alpha \in \left(\frac{1}{2}, 1\right), \quad r > 0, \\ X(0) = X_0. \quad (15)$$

The solution of (15) can be expressed as

$$X(t) = X_0 t^{\alpha-1} E_{\alpha, \alpha}(-rt^{\alpha}) + \sigma \int_0^t (t-\tau)^{\alpha-1} E_{\alpha, \alpha}(-r(t-\tau)^{\alpha}) dW(\tau), \quad t \geq 0.$$




Figure 3
Almost surely exponential stability of fractional Langevin equation when $\alpha = 0.6$ and 0.7 .




Figure 4

Almost surely exponential stability of fractional Langevin equation when $\alpha = 0.8$ and 0.9 .

If $X_0 > 0$ and $X_0 \neq \infty$, according to the results for asymptotic behavior of Mittag-Leffler functions in [7], the solution of (15) is exponentially decaying under the disturbance of white noise.

On the other hand, from Lemma 1 $X(t) \in L^2(\Omega, \mathcal{F}_t, P)$. Since $r > 0$, we can get

$$\int_0^t -rX(\tau)(t-\tau)^{\alpha-2} d\tau < +\infty, \quad t \geq 0.$$

Obviously, there are a pair of positive constants θ and K such that

$$-\overline{K} \leq \frac{\alpha-1}{\Gamma(\alpha)} \int_0^t -rX(\tau)(t-\tau)^{\alpha-2} d\tau + \overline{\theta} \leq \overline{K}, \quad t \geq 0.$$

From Example 1 the solution of (15) is stochastically stable. Using MATLAB software, the solution trajectories for $\alpha = 0.6, 0.7, 0.8, 0.9$ are shown in Figs. 3 and 4.

When




Figure 5

Almost surely exponential stability of fractional Langevin equation when $\alpha = 1$ and white noise vanishes in (15) when $\alpha = 0.6$.

$\alpha = 1$, the solution of (15) can be directly expressed as

$$X(t) = X_0 e^{-rt} + \sigma \int_0^t e^{-r(t-\tau)} dW(\tau), \quad t \geq 0,$$

and its trajectory is shown in Fig. 5.

6 Conclusion

This paper studies the stability of Caputo-type fractional stochastic differential equations. The Itô's formula of Caputo version is established and used to prove the stochastic stability, stochastic asymptotical stability, almost surely exponential stability and pth moment exponential stability. Numerical examples are given to illustrate the stability results.

Acknowledgments

The authors are grateful to the referees for their careful reading of the manuscript and their valuable comments. We thank the editor also.

References

- 1 M.A. Abdullin, S.V. Meleshko, F.S. Nasyrov, A new approach to the group analysis of onedimensional stochastic differential equations, *J. Appl. Mech. Tech. Phys.*, **55**(2):191–198, 2014, <https://doi.org/10.1134/S0021894414020011>
- 2 B. Bayour, D.F.M. Torres, Existence of solution to a local fractional nonlinear differential equation, *J. Comput. Appl. Math.*, **312**:127–133, 2017, <https://doi.org/10.1016/j.cam.2016.01.014>
- 3 T. Burton, T. Furumochi, Asymptotic behavior of solutions of functional differential equations by fixed point theorems, *Dyn. Syst. Appl.*, **11**(4):499–519, 2002.
- 4 T. Burton, T. Furumochi, Krasnoselskii's fixed point theorem and stability, *Nonlinear Anal., Theory Methods Appl.*, **49**:445–454, 2002, [https://doi.org/10.1016/S0362-546X\(01\)00111-0](https://doi.org/10.1016/S0362-546X(01)00111-0)
- 5 T. Burton, B. Zhang, Fixed points and fractional differential equations: Examples, *Fixed Point Theory*, **14**:313–325, 2013.
- 6 A. Chadha, S.N. Bora, Existence and exponential stability for neutral stochastic fractional differential equations with impulses driven by Poisson jumps, *Stochastics*, **90**(5):663–681, 2018, <https://doi.org/10.1080/17442508.2017.1402899>
- 7 N.D. Cong, T.S. Doan, S. Siegmund, H.T. Tuan, On stable manifolds for planar fractional differential equations, *Appl. Math. Comput.*, **226**:157–168, 2014, <https://doi.org/10.1016/j.amc.2013.10.010>
- 8 D.T. Dr. Son, P.T. Huong, P.E. Kloeden, H.T. Tuan, Asymptotic separation between solutions of Caputo fractional stochastic differential equations,

- Stochastic Anal. Appl.*, **36**(4):654–664, 2018, <https://doi.org/10.1080/07362994.2018.1440243>
- 9 M. Hu, S. Ji, S. Peng, Y. Song, Backward stochastic differential equations driven by α -Brownian motion, *Stochastic Processes Appl.*, **124**(1):759–784, 2014, <https://doi.org/10.1016/j.spa.2013.09.010>
- 10 L.-L. Huang, J.H. Park, G.-C. Wu, Z.-W. Mo, Variable-order fractional discrete-time recurrent neural networks, *J. Comput. Appl. Math.*, **370**:112633, 2020, <https://doi.org/10.1016/j.cam.2019.112633>
- 11 S. Jin, K. Kobayashi, Strong approximation of stochastic differential equations driven by a time-changed Brownian motion with time-space-dependent coefficients, *J. Math. Anal. Appl.*, **476**(2):619–636, 2019, <https://doi.org/10.1016/j.jmaa.2019.04.001>
- 12 A.A. Kilbas, H.M. Srivastava, J.J. Trujillo, *Theory and Applications of Fractional Differential Equations*, Elsevier, Amsterdam, 2006.
- 13 R. Klen, V. Manojlovic', S. Simic', M. Vuorinen, Bernoulli inequality and hypergeometric functions, *Proc. Am. Math. Soc.*, **142**(2):559–573, 2014, <https://doi.org/10.1090/S0002-9939-2013-11781-8>
- 14 E. Malkowski, V. Rakoc'evic', *Advanced Functional Analysis*, CRS Press, Boca Raton, FL, 2019, <https://doi.org/10.1201/9780429442599>
- 15 X. Mao, *Stochastic Differential Equations and Application*, 2nd ed., Horwood Publishing, Philadelphia, 2007.
- 16 D. Nguye, Asymptotic behavior of linear fractional stochastic differential equations with timevarying delays, *Commun. Nonlinear Sci. Numer. Simul.*, **19**(1):1–7, 2014, <https://doi.org/10.1016/j.cnsns.2013.06.004>
- 17 B. Øksendal, *Stochastic Differential Equations an Introduction with Application*, 5th ed., Springer, Berlin, 2000, <https://doi.org/10.1007/978-3-662-03185-8>
- 18 B. Pei, Y. Xu, J. L. Wu, Stochastic averaging for stochastic differential equations driven by fractional Brownian motion and standard Brownian motion, *Appl. Math. Lett.*, **100**:106–006, 2020, <https://doi.org/10.1016/j.aml.2019.106006>
- 19 S. Peng, Multi-dimensional α -Brownian motion and related stochastic calculus under α -expectation, *Stochastic Processes Appl.*, **118**(12):2223–2253, 2008, <https://doi.org/10.1016/j.spa.2007.10.015>
- 20 R. Sakthivel, P. Revathi, Y. Ren, Existence of solutions for nonlinear fractional stochastic differential equations, *Nonlinear Anal., Theory Methods Appl.*, **81**:70–86, 2013, <https://doi.org/10.1016/j.na.2012.10.009>
- 21 Y. Song, Properties of α -martingales with finite variation and the application to α -Sobolev spaces, *Stochastic Processes Appl.*, **129**(6):2066–2085, 2019, <https://doi.org/10.1016/j.spa.2018.07.002>
- 22 V. Todorc'evic', *Harmonic Quasiconformal Mappings and Hyperbolic Type Metrics*, Springer, Cham, 2019, <https://doi.org/10.1007/978-3-030-22591-9>
- 23 G.-C. Wu, T. Abdeljawad, J. Liu, D. Baleanu, K.-T. Wu, Mittag-Leffler stability analysis of fractional discrete-time neural networks via fixed point technique, *Nonlinear Anal. Model. Control*, **24**(6):919–936, 2019, <https://doi.org/10.15388/NA.2019.6.5>

- 24 X. Zhang, H. Yi, H. Shu, Nonparametric estimation of the trend for stochastic differential equations driven by small α -stable noises, *Stat. Probab. Lett.*, **151**:8–16, 2019, <https://doi.org/10.1016/j.spl.2019.03.012>
- 25 Y. Zhou, J. Wang, L. Zhang, *Basic Theory of Fractional Differential Equations*, World Scientific, Singapore, 2016.

Notes

- * This work is partially supported by the National Natural Science Foundation of China (11661016), Training Object of High Level and Innovative Talents of Guizhou Province ((2016)4006), Department of Science and Technology of Guizhou Province (Fundamental Research Program [2018]1118) and Guizhou Data Driven Modeling Learning and Optimization Innovation Team ([2020]5016).

Notas de autor

Department of Mathematics, Guizhou University, Guiyang 550025, Guizhou, China