



Revista Colombiana de Biotecnología

ISSN: 0123-3475

ISSN: 1909-8758

Instituto de Biotecnología, Universidad Nacional de Colombia

Herrera-Cruz, Edwin; Vásquez-Machado, Gersson; Estrada-Posada, Ana;
Pardo-Camacho, Kamylo; Atencio-García, Víctor; Yepes-Blandón, Jonny
Use of probiotics in the catfish *Sorubim cuspicaudus* larviculture
Revista Colombiana de Biotecnología, vol. XXV, no. 2, 2023, July-December, pp. 50-61
Instituto de Biotecnología, Universidad Nacional de Colombia

DOI: <https://doi.org/10.15446/rev.colomb.biote.v25n2.110786>

Available in: <https://www.redalyc.org/articulo.oa?id=77677359005>

- How to cite
- Complete issue
- More information about this article
- Journal's webpage in redalyc.org

redalyc.org

Scientific Information System Redalyc

Network of Scientific Journals from Latin America and the Caribbean, Spain and Portugal

Project academic non-profit, developed under the open access initiative

Use of probiotics in the catfish *Sorubim cuspicaudus* larviculture

Uso de probióticos en la larvicultura del bagre *Sorubim cuspicaudus*

Edwin Herrera-Cruz^{*}; Gersson Vásquez-Machado^{**}; Ana Estrada-Posada^{***};
Kamylo Pardo-Camacho^{****}; Víctor Atencio-García^{*****}; Jonny Yepes-Blandón^{*****}

DOI: 10.15446/rev.colomb.biote.v25n2.110786

RESUMEN

Sorubim cuspicaudus, un bagre migratorio distribuido en las cuencas de los ríos Magdalena, Sinú y Catatumbo, se encuentra catalogado como una especie vulnerable a la extinción. La producción de alevinos en ambientes controlados es considerada como una estrategia crucial para su conservación, y la larvicultura es una fase crítica en la cría de esta especie. Para mejorar esta fase crítica de producción de alevinos se utilizan probióticos. El estudio tuvo como objetivo evaluar el uso de probióticos (*Bacillus subtilis* y *Bacillus licheniformis*) durante la fase de larvicultura de *S. cuspicaudus*. Las larvas, a las 42 horas post eclosión (1,5±0,1mg, longitud total 5,7±0,4mm) fueron tratadas con cuatro niveles de inclusión de probióticos en el agua: 0, 5, 10 y 20ppm durante 22 días. La calidad del agua se mantuvo dentro de los rangos adecuados para la larvicultura de especies neotropicales de bagre y los parámetros evaluados fueron el aumento de peso (Gw), el aumento de longitud (GL), la tasa específica de crecimiento (G), la tasa de supervivencia (S), la resistencia al estrés (Sr), la longitud del pliegue intestinal (LF) y el recuento de unidades formadoras de colonias (UFC). Los resultados mostraron mayores GL (22,23±3,5 mm), Gw (40,0±12,6 mg), G (14,9±1,5%/día), LP (205±72,7 µm) y UFC (118,7±80,9) a 20 ppm (p<0,05). Sin embargo, S y Sr no mostraron diferencias significativas entre tratamientos (p>0,05). Los resultados de este estudio sugieren que los probióticos (*Bacillus subtilis* y *Bacillus licheniformis*) podrían utilizarse como alternativa para avanzar en la larvicultura de *S. cuspicaudus*.

Palabras clave: *Bacillus*, crecimiento, pliegues intestinales, peces migratorios, pimelodidae.

ABSTRACT

Sorubim cuspicaudus, a migratory catfish distributed in the Magdalena, Sinú, and Catatumbo river basins, is categorized as vulnerable to extinction. Production of fingerlings in controlled environments stands as a strategic conservation approach, and larviculture is a critical phase in rearing this species. Probiotics are used for improvement in the critical stages of fingerling production. The study aimed to evaluate the use of probiotics (*Bacillus subtilis* and *Bacillus licheniformis*) during the larviculture phase of *S. cuspicaudus*.

* GIOANE Research group on native and exotic aquatic organisms. Facultad de Ciencias Agrarias, Universidad de Antioquia, Medellín, Colombia. ORCID: <https://orcid.org/0000-0002-5527-2375>
 ** HISTOLAB, Bogotá, Colombia. ORCID: <https://orcid.org/0000-0002-4737-7038>
 *** ISAGEN S.A. E.S.P, Medellín, Colombia. ORCID: <https://orcid.org/0000-0003-3585-3719>
 **** GIOANE Research group on native and exotic aquatic organisms. Facultad de Ciencias Agrarias, Universidad de Antioquia, Medellín, Colombia. ORCID: <https://orcid.org/0009-0003-3838-5989>
 ***** FMVZ/DCA/CINPIC, Universidad de Córdoba, Montería, Colombia. ORCID: <https://orcid.org/0000-0002-2533-1995>
 ***** GIOANE Research group on native and exotic aquatic organisms. Facultad de Ciencias Agrarias, Universidad de Antioquia, Medellín, Colombia. ORCID: <https://orcid.org/0000-0001-6276-5488> (Corresponding autor), jonny.yepes@udea.edu.co

cus. Larvae at 42 hours post-hatching (1.5 ± 0.1 mg, total length 5.7 ± 0.4 mm) were treated with four levels of probiotic inclusion in the water: 0, 5, 10, and 20 ppm for 22 days. Water quality remained within suitable ranges for neotropical catfish species larviculture and the parameters assessed were weight gain (Gw), length gain (Gl), specific growth rate (G), survival rate (S), stress resistance (Sr), intestinal fold length (LF), and colony-forming units (CFU) count. Results showed higher Gl (22.23 ± 3.5 mm), Gw (40.0 ± 12.6 mg), G ($14.9 \pm 1.5\%$ /day), LP ($205 \pm 72.7 \mu\text{m}$), and CFU (118.7 ± 80.9) were found at 20 ppm ($p < 0.05$). However, S and Sr exhibited no significant differences among treatments ($p > 0.05$). The findings of this study suggest that probiotics (*Bacillus subtilis* and *Bacillus licheniformis*) could be used as an alternative to advance in the *S. cuspidatus* larviculture.

Key words: *Bacillus*, growth, intestinal folds, migratory fish, pimelodidae.

Recibido: Mayo 9 de 2023 **Aprobado:** Noviembre 14 de 2023

INTRODUCTION

The Trans-Andean shovelnose catfish, *Sorubim cuspidatus*, is a migratory species that inhabits the Magdalena, Cauca, Sinú, and Catatumbo basins. It is categorized as vulnerable due to a gradual decline in its capture volumes in recent years (Mojica *et al.*, 2012). In 2006, the Magdalena river's basin reported a capture volume of 416.5 tons, which drastically decreased to 65.8 tons in 2017, marking a reduction of 84% (INCODER, 2006; De la Hoz Maestre *et al.*, 2017). Over the past decades, the Magdalena basin has experienced substantial environmental changes resulting from mining, oil extraction, agriculture, damming, and other anthropogenic activities. These alterations have led to habitat loss and degradation of water quality, consequently affecting native fish populations (Mojica *et al.*, 2012; Lozano *et al.*, 2017).

Ex situ conservation strategies such as captive reproduction allow the massive production of fingerlings of fish species with some vulnerability (Prieto-Guevara *et al.*, 2015). Atencio *et al.*, (2010), state that three phases can be identified in the production of fingerlings: management of broodstock, induced reproduction, and larviculture. Considerable progress has been made in the initial two phases (Prieto-Guevara *et al.*, 2015). However, the most notable challenges have been observed in larviculture, particularly during the initial feeding phase. During this period, mortality attributed to cannibalism is documented (Atencio *et al.*, 2010) along with elevated stress levels that disrupt the fish's physiological equilibrium (homeostasis). This disruption leads to immunosuppression, increased diseases susceptibility, and decreased survival rates (Aerts *et al.*, 2018). Consequently, it is imperative to explore alternatives or strategies to enhance the performance of larviculture for this catfish. Among the successful options used in other fishes, probiotics stands out. These microorganisms play a pivotal role in maintaining the intestinal microbial balance, providing nutritional benefits, offering protection against pathogens, and contributing to both development and immune responses (Pérez-Sánchez *et al.*, 2018; Paixão *et al.*, 2020).

Probiotics are used in domestic animal production systems and have been introduced in recent years in commercial aquaculture, both in larviculture and in grow-out cultures (Martínez Cruz *et al.*, 2012; Newaj-Fyzul *et al.*, 2014; Pérez-Sánchez *et al.*, 2014; Ribas & Piferrer, 2014; Kumar *et al.*, 2016; Hoseinifar *et al.*, 2016; Cai *et al.*, 2018). The beneficial effect of probiotics on aquaculture production is attributed to their ability to improve water quality, reduce accumulated organic matter, inhibit pathogens through various pathways, enhance the immune system, and provide nutritional benefits by influencing digestibility and feed utilization (Carnevali *et al.*, 2006; Nayak, 2010; Cai *et al.*, 2018). Additionally, probiotics presents functions similar to antibiotics without generating drug resistance (Thurlow *et al.*, 2019).

Nevertheless, it is imperative to consider factors such as the selection of strains capable of giving benefits to the host and maintaining stability at sufficient densities over time (Balcázar *et al.*, 2006; Begley *et al.*, 2006; Interaminense *et al.*, 2018). Similarly, the host species, developmental phase (larvae, fingerlings, grow-out), environmental culture conditions, and administration route must be considered (Soltani *et al.*, 2019; Ringø *et al.*, 2020). Probiotics can be applied orally through food, primarily influencing the intestine, or indirectly by administration into the culture water to enhance its quality (Qi *et al.*, 2009).

The studies reveal that the use of mixtures of probiotic microorganisms is more effective than the independent strains in the control of pathogens, in the improvement of productive performance parameters and in the establishment of microorganism populations, observing synergistic processes (Van Doan *et al.*, 2018; Soltani *et al.*, 2019; Kuebutornye *et al.*, 2020). Among the bacteria with probiotic capacities is the genus *Bacillus*; one of the most used in aquaculture (Ringø *et al.*, 2020). *Bacillus* is spore-forming bacteria highly resistant to aggressive physical and chemical conditions, surviving in various environments, including freshwater, marine sediments, sand, thermal waters, arctic soils, and the gastrointestinal tract of fish, crustaceans, and mollusks; providing a wide range of beneficial effects (Soltani *et al.*, 2019).

Bacillus species can play an important role in removing waste products for more sustainable aquaculture, reducing stress in fish, leading to better immunophysiological balance, better growth and higher survival in aquatic animals (Soltani *et al.*, 2019). In addition, *Bacillus* stimulates the presence of beneficial bacteria and decreases the load of pathogens in ponds. Most species in this genus are producers of various secondary metabolites containing antibiotics, helpful chemicals, and enzymes (Zokaeifar *et al.*, 2012; Cai *et al.*, 2018; Ringø *et al.*, 2020).

The objective of the study was to evaluate the effect of *Bacillus subtilis* and *Bacillus licheniformis* as a probiotic at different levels of inclusion in the water on the growth and gastrointestinal development of the catfish *Sorubim cuspicaudus* larvae.

METHODOLOGY

Biological material and treatments. The research was conducted at San Silvestre S.A. Fishculture Station (Barrancabermeja, Santander, Colombia), situated at geographic coordinates 7°06'31" north latitude and 73° 51'25" west longitude, at an elevation of 80 m.a.s.l. and average annual temperature of 28.4°C. Larvae were obtained through the hormonal induction of *S. cuspicaudus* broodstock originating from the middle basin of the Magdalena River. These broodstock had been adapted to captive conditions for over a year, being fed *ad libitum* with forage fish and supplemented daily with a diet of 34% crude protein based with a 1% fish biomass equivalent ratio.

The females and the males were induced at 10 µg/Kg of GnRH analog of live weight (Pardo-Carrasco *et al.*, 2015). The embryos were placed in incubators with up-flow water (4L/min), and newly hatching larvae were transferred to circular pools with constant aeration at a density of 80 larvae/L. 2720 larvae of 42 hours post hatching (HPE) were placed in 16 aquariums of 10L. Dissolved oxygen, pH, temperature, electrical conductivity, and dissolved solids were measured daily with a multiparameter probe (YSI, Pro Plus, USA), while hardness and alkalinity by titration (HACH, 1690001, USA), and turbidity with a turbidimeter (LaMate, 2020we, USA) at the beginning, in the middle and at the end of the experiment. For 22 days, four probiotic inclusion levels were evaluated in the water containing 50% *Bacillus subtilis* and 50% *Bacillus licheniformis*: 0 (control), 5, 10, and 20 ppm each with four replicates per treatment. In each aquarium, 170 larvae were stocked, fed twice a day (morning and afternoon), with recently hatched brine shrimp nauplii (instar I) as follows: days 1 to 5, 10 nauplii/ml were

offered; from 6 to 14 20 nauplii/ml were offered and from 15 to 22 40 nauplii/ml were offered.

Growth and survival. At the end of the experiment, 15 larvae were taken per experimental unit for growth evaluation. The larvae were weighed on an analytical balance (Chyo, JS-110, Japan) and the total length was measured with an image analyzer (ImageJ, USA). The following parameters were estimated:

Weight gain (Gw) = $W_f - W_o$; where W_f and W_o correspond to the final and initial weight of the larvae. Length gain (Gl) = $L_f - L_o$; where L_f and L_o correspond to the final and initial total length of the larvae. Specific growth rate (G) = $100 \times [\ln(W_f) - \ln(W_o)] / t$; where t is the culture time expressed in days and \ln is the natural logarithm. The final survival (S) is estimated by dividing the final number of larvae by the total number of initial larvae in each experimental unit.

Stress resistance test. 10 larvae were taken per replicate and placed out of the water on absorbent paper for eight minutes, then transferred to a container with water from the same replicate for 15 minutes, and survival was estimated based on the method used by Atencio Garcia *et al.*, (2003).

Microbiological analysis of water. On days 0 and 22 of the experiment, bacterial count was conducted using nutrient agar plates. For this purpose, 0.1 ml of water from the aquariums was extracted, diluted to 1/1000 with a 0.9% saline solution, and then incubated at 30±2°C for 48 hours. Counts ranging from 30 to 300 colony-forming units (CFUs) were considered for enumeration (CFU/ml × 10³).

Histological analysis. Fifteen larvae per replicate were collected at the initial and final of the experiment. These larvae were anesthetized using eugenol (40 ppm), fixed in 10% buffered formalin, embedded in paraffin, and subjected to histotechnical processing. Subsequently, 5 µm histological sections were generated employing a rotary microtome (Leica, RM2125, Germany) and stained with hematoxylin-eosin. A histological description of the digestive tract was performed, and the length of the intestinal folds in the anterior, middle, and posterior regions was measured (n=3 per region). This measurement was conducted utilizing an image analyzer (Image J, USA) employing the methodology suggested by (Mokhtar *et al.*, 2015).

Statistical analysis. A completely randomized design was employed, and all evaluated parameters were subjected to normality tests (Shapiro-Wilk test) and homo-

scedasticity (Levene test). Subsequently, they were subjected to ANOVA, followed by a Tukey Multiple Range test. In all instances, a significance level of $p < 0.05$ was adopted to indicate statistical differences among treatments. The analyzed parameters were presented as mean \pm standard deviation (SD). Statistical analysis was executed using the SAS program (version 9.1, 2004, USA).

All animals handling procedures followed the standards for using laboratory animals outlined by the Committee on the Care and Use of Laboratory Animal Resources of the National Research Council (National Academies, USA). This research had a permit issued by the Colombian National Aquaculture and Fisheries Authority - AUNAP (Resolution 0955 of May 27, 2020).

RESULTS

Growth performance. The growth, survival, and stress resistance outcomes are presented in Table 1. Larvae treated to a probiotic inclusion level of 20 ppm exhibited higher Gw, Gl, and G, displaying a statistically significant difference from the other treatments ($p < 0.05$). In contrast, the control group showed lower values. Survival rates (between 31.0% to 23.1%) and stress resistance (between 87.5% to 97.5%) displayed no significant statistical difference among the treatments ($p < 0.05$).

Digestive tract and intestinal folds. Figure 1 shows the histological features of the intestine in *S. cuspidatus* larvae at the beginning of the experiment. In the control treatment, more pronounced development was evident in the intestine middle and posterior sections than in the anterior section. Among larvae receiving probiotic supplementation (5, 10, and 20 ppm), the most caudal portion of the esophagus exhibited finger-like folds oriented both caudally and, to a lesser extent, cranially. These folds were covered by a simple cylindrical epithelium supported by the basal lamina, and, under this, a thin layer of loose connective tissue known as the lamina propria. Moreover, enhanced fold development was observed in the foregut, demonstrating a nearly similar relationship with the middle and posterior regions, when contrasted with the larvae in the control group.

At the end of the experiment, less development of the esophagus, stomach and intestine was observed in the control treatment concerning the treatments including probiotics, especially at 20 ppm (figure 2). In addition, differences in size, length, width, and development of intestinal folds of the larvae treated with the probiotic and the control group were observed. Also, in the treatments with the inclusion of probiotics, it was observed

that in the middle portion of the intestine, there was a greater presence of goblet cells in the epithelium and, in the posterior portion, the presence of multiple epithelial cells with small intracytoplasmic vacuoles.

Figure 3 shows the measurement of the intestinal folds at the experiment's beginning and end. Initially, no significant difference was observed between treatments ($p < 0.05$), which allows us to infer that the catfishes presented the same initial conditions. At the end of the experiment, a significantly greater length of intestinal folds ($p < 0.05$) was found in larvae with the inclusion of 20 ppm of probiotics ($205.3 \pm 72.7 \mu\text{m}$), compared to the control ($119.7 \pm 31.4 \mu\text{m}$); but without statistical difference with the larvae maintained at 5 ppm ($162.2 \pm 34.5 \mu\text{m}$) and 10 ppm ($179.0 \pm 39.9 \mu\text{m}$).

Physicochemical parameters of water. The average values of the water quality parameters are presented in table 2. The higher temperature occurred in the 5 ppm treatment ($26.8 \pm 0.6^\circ\text{C}$) and the lower in the control group ($26.6 \pm 0.5^\circ\text{C}$), observing statistical difference between these values ($p < 0.05$). The higher dissolved oxygen concentration was recorded in the control group ($7.1 \pm 0.4 \text{ mg/l}$) and the lower in 5 ppm ($6.7 \pm 1.0 \text{ mg/l}$), showing significant differences ($p < 0.05$). For the parameters pH, electrical conductivity, dissolved solids, hardness and total alkalinity, the higher averages occurred at 20 ppm treatment, while the lower were at 5 ppm (Table 2). The most elevated turbidity ($7.7 \pm 0.2 \text{ NTU}$) was recorded in the 20 ppm treatment and the lowest ($0.6 \pm 0.25 \text{ NTU}$) in the control group, observing a statistical difference between these values ($p < 0.05$).

Microbiological analysis of water. Figure 4 presents the counts of colony-forming units (CFU) in the culture water in the different treatments. At the beginning of the experiment, a higher number of CFU was observed at 20 ppm ($138.7 \pm 81.9 \text{ CFU/ml} \times 10^3$), being statistically different from the control group and at 5 ppm ($p < 0.05$). At the end of the experiment (day 22), a higher CFU concentration was recorded in the treatment with 20 ppm inclusion ($83.5 \pm 64 \text{ CFU/ml} \times 10^3$) without observing a significant statistical difference when the probiotics were included at 5 ppm ($36.5 \pm 16.0 \text{ UFC}$) and 10 ppm ($55.0 \pm 29.5 \text{ UFC}$), but with the control group ($5.3 \pm 1.2 \text{ UFC}$) ($p < 0.05$).

DISCUSSION

The results of this study suggest that the mixture of probiotics (50% *Bacillus subtilis* - 50% *Bacillus licheniformis*) supplied in the culture water benefited effect the catfish larviculture. The larvae with 20 ppm inclusion of probiot-




Figure 1. Photomicrographs of intestinal folds of catfish *Sorubim cuspicaudus* larvae at the beginning of the experiment treated to different inclusions of probiotics in the water. 0 ppm, control (A), 5 ppm (B), 10 ppm (C), 20 ppm (D). EC: epithelial cells; EN: stratified flat epithelium nuclei; IF: intestinal folds; L: lumen; MT: muscular tunic; TA: mucosal tunic; PL: propia layer; SCE: simple cylindrical epithelium; ST: serosa tunic (H&E, 40X).

Table 1. Growth performance of catfish *Sorubim cuspicaudus* larvae, treated to different levels of inclusion of probiotics in the water. G, specific growth rate; Gl, length gain; Gw, weight gain. Different letters in the column indicate a significant difference ($p < 0.05$).

Treatments	Gl (mm)	Gw (mg)	G (%/día)	Survival (%)	Stress resistance (%)
0 ppm	14.9±3.0 ^c	18.0±5.9 ^c	11.4±1.4 ^c	27.1±6.7	90.9±15.7
5 ppm	19.30±4.5 ^b	27.0±14.8 ^b	12.8±2.4 ^b	31.0±8.5	92.5±9.6
10 ppm	15.13±5.8 ^c	25.0±10.8 ^b	12.7±1.8 ^b	30.2±7.2	97.5±5.0
20 ppm	22.23±3.5 ^a	40.0±12.6 ^a	14.9±1.5 ^a	23.1±7.5	87.5±10.2

Mean values followed by different letters in the same column differ significantly ($P < 0.05$).




Figure 2. Photomicrographs of intestinal folds of catfish *Sorubim cuspidus* larvae at the end of the experiment treated to different probiotic inclusions in the water. 0 ppm, control (A), 5 ppm (B), 10 ppm (C), 20 ppm (D). AS: artemia cyst; EC: epithelial cells; IF: intestinal folds; K: kidney; L: lumen; PL: propia layer; MT: medular tissue; NE: nuclei of the stratified squamous epithelium; SM: skeletal muscle; ST: serous tunic; MT: muscular tunic (H&E, 20X).




Figure 3. Length of intestinal folds of catfish *Sorubim cuspidus* larviculture treated to different levels of inclusion of probiotics. Different letter indicate a significant difference ($p < 0.05$).




Figure 4. Colony-forming units (CFU) in the water of catfish *Sorubim cuspicaudus* larviculture treated to different levels of inclusion of probiotics. Different letters indicate a significant difference ($p < 0.05$).

Table 2. Physicochemical parameters of water during larviculture of catfish *Sorubim cuspicaudus* treated to different levels of probiotic inclusion.

Parameters	Treatments			
	0 ppm	5 ppm	10 ppm	20 ppm
Temperature (°C)	26.6±0.5 ^a	26.8±0.6 ^b	26.7±0.6 ^b	26.7±0.6 ^{ab}
Dissolved oxygen (mg/l)	7.1±0.4 ^c	6.7±1.0 ^a	6.8±0.5 ^{ab}	7.0±0.5 ^{bc}
pH	7.6±0.2 ^a	7.5±0.2 ^a	7.6±0.1 ^b	7.7±0.1 ^c
Conductivity(µS/cm)	189.9±33.9 ^a	181.1±23.6 ^a	210.4±37.2 ^b	226.2±37.1 ^c
Dissolved solids (mg/l)	120.6±20.8 ^b	113.9±15.2 ^a	133.1±24.0 ^c	142.6±23.5 ^d
Turbidity (NTU)	0.6±0.25 ^a	1.6±0.6 ^a	4.99±1.3 ^b	7.1±0.2 ^c
Hardness (mg CaCO ₃ /l)	10.4±1.2 ^a	10.4±1.1 ^a	13.7±1.6 ^{ab}	16.1±3.7 ^b
Alkalinity (mg CaCO ₃ /l)	53.3±1.50 ^a	52.9±1.3 ^a	56.2±3.7 ^{ab}	60.0±3.7 ^b

Different letters in the same column indicate significant difference ($p < 0.05$).

ics showed better growth (Gw, GL, G) than the larvae of the control group and the other treatments (5 and 10 ppm). Some authors maintain that probiotics promote digestive activity early, through the release of enzymes (proteases and amylases), and facilitate the production of vitamins, prevent intestinal disorders, and promote and improve the digestibility of food (Zhang *et al.*, 2010; Avella *et al.*, 2012; Carnevali *et al.*, 2017; Interaminense *et al.*, 2018; Van Doan *et al.*, 2018).

This action of the probiotics possibly accelerated the growth of *S. cuspidatus* larvae, especially in the treatment with 20 ppm inclusion. In other studies, similar effects have been observed in the parameters of productive performance as in the herbivorous carp *Ctenopharyngodon idellus* when 2.4×10^7 CFU/g of *B. subtilis* were added for 42 days (Tang *et al.*, 2019). In *Oreochromis niloticus*, the inclusion of three probiotic bacteria in a 1:1:1 ratio with 1×10^8 CFU/ml of *Bacillus velezensis* TPS3N, *B. subtilis* and *B. amyloliquefaciens* improved final weight, weight gain and feed conversion ratio. In addition, the probiotics separately also improved the final weight and weight gain compared to the control (Kuebutornye *et al.*, 2020). A better productive performance has also been reported in crustaceans such as *M. rosenbergii*, when 9.35×10^8 CFU/g of *B. licheniformis* MTCC 429 was added for 90 days (Sudha *et al.*, 2019) and in *L. vannamei* with the addition of 10^9 CFU per kg/56 days of *B. subtilis* (Tsai *et al.*, 2019).

It is possible that, in the present study, the interaction of *B. subtilis* and *B. licheniformis* in the larviculture of *S. cuspidatus* has potentiated the individual effect of each one of the probiotics, and thus explains the better productive performance obtained in the treatments with greater inclusion of probiotics (10 and 20 ppm).

Among the beneficial effects attributed to probiotics of the genus *Bacillus* is the generation of bacteriocins, which can exert a favorable impact on the growth and resilience to pathogenic microorganisms in aquatic organisms (Soltani *et al.*, 2019; Ringø *et al.*, 2020) which could be considered among the factors that improved the growth of the catfish larvae treated with probiotics.

In this study, introducing *Bacillus subtilis* and *Bacillus licheniformis* into the culture water prompted the proliferation of intestinal folds. This expansion likely augmented the nutrient absorption region and presumably influenced the larval intestinal well-being. The histological analyses of the catfish's digestive tract indicated an increasing trend in the length of the intestinal folds as the inclusion of *Bacillus* increased. A potential advantage of extending the length of the intestinal folds is the likely

enlargement of the intestinal absorptive surface (Cao *et al.*, 2019; Sewaka *et al.*, 2019; Kong *et al.*, 2021). In the catfish larvae, the length of the intestinal folds increased, with notable structural advancement in treatments supplemented with probiotics (5-20 ppm) regarding the control group. Comparable findings have been noted in *Channa argus* larvae when their diet was enriched with *Lactococcus lactis*, *Enterococcus faecalis*, or their combination (1×10^8 CFU/g); these larvae exhibited a significant increase in intestinal fold height compared to the group without probiotics (Kong *et al.*, 2021). The intestine holds vital significance in nutrient digestion and absorption in fish, and the integrity of its tissue structure significantly impacts animal growth (Nicholson *et al.*, 2012; Kong *et al.*, 2021). Parameters such as muscle thickness, height, and fold width indicate intestinal health in aquatic organisms (Kong *et al.*, 2021).

In the present study, in larvae treated with probiotics, a greater presence of goblet cells was recorded, particularly in the middle portion of the intestine and the presence of multiple epithelial cells with small intracytoplasmic vacuoles, as well as greater development of the esophagus and stomach. Similar results were obtained in tilapia (*Oreochromis niloticus*) fed with diets supplemented with enzymes and probiotics (20 ppm) containing *Bacillus subtilis*, *Bacillus licheniformis* and *Bacillus pumilus*, finding a greater number of goblet cells and a greater diameter of intestinal folds (Adeoye *et al.*, 2016).

Even though studies affirm that applying probiotics, including bacteria of the *Bacillus* genus, is a beneficial practice for aquaculture since it improves animal welfare and the immune system, increasing resistance to diseases (Interaminense *et al.*, 2018). There are also reports of adverse effects, such as those in Cerezuela *et al.*, (2012) in sea-bream *Sparus aurata* (50 g of weight), who reported signs of edema, intestinal inflammation and decreased height of the intestinal folds when they were treated with *Bacillus subtilis* (1×10^7 CFU/g) in the food; which suggests that the effect of probiotics is species-specific and therefore it is necessary to evaluate the probiotic, the dosage and route of application for each species.

In the present study, there was no significant difference in survival and resistance to stress in the different treatments, possibly because the culture conditions were adequate in all the evaluated treatments. Similar results have been found in other studies. For example, with juveniles of *Amphiprion ocellaris*, Paixão *et al.*, (2020) found that at the highest concentration of *Lactobacillus plantarum* (1×10^8 CFU/g), the productive performance parameters improved without affecting survival. The larval phase of aquaculture species is critical and with high

susceptibility to diseases given the poor development of the immune system (Prieto-Guevara *et al.*, 2015; Aerts *et al.*, 2018); therefore, increasing disease resistance in larviculture and enhancing feed efficiency has led to the use of probiotics as a prophylactic alternative (Sorieul *et al.*, 2018; Ringø *et al.*, 2020).

However, an adequate dosage of probiotics improves animal welfare conditions and water quality and allows for obtaining animals of excellent quality and resistance (Midhun *et al.*, 2019; Mukherjee *et al.*, 2019; Paixão *et al.*, 2020). It has been reported that adding 2×10^8 CFU/g of *B. subtilis* for 56 days increased resistance to stress to disease caused by *A. hydrophila* in the sturgeon *Acipenser dabryanus* (Di *et al.*, 2019).

Cai *et al.*, (2018) found that the probiotics *B. licheniformis* and *B. flexus* improved water quality in *L. vannamei* postlarvae cultures. Ringø *et al.*, (2020) stated that spore-producing bacteria such as *Bacillus* can be supplied directly in the culture water because it is an efficient and economical transport route and improves water quality. In the present study, the electrical conductivity, dissolved solids, and turbidity increased significantly in the treatments with 10 and 20 ppm inclusion of probiotics, possibly caused by metabolites or excess of microorganisms.

In addition, the application of probiotics influenced the number of microorganisms present in the culture water, being higher ($p < 0.05$) when it was included at 20 ppm compared to the control group. Nevertheless, although the CFU showed higher values when the probiotics were included at 20 ppm, these values did not show a statistical difference to those registered when the probiotic was included at 5 and 10 ppm ($p > 0.05$). These results suggest that the inclusion of probiotics influenced the concentration of microorganisms (UFC/ml), which affected the increase in the length of intestinal folds. Other studies have reported similar results (Gisbert *et al.*, 2013; Carnevali *et al.*, 2017; Sewaka *et al.*, 2019; Kong *et al.*, 2021).

CONCLUSIONS

Based on the findings of this study, it can be concluded that a combination of *Bacillus subtilis* and *Bacillus licheniformis* in a 1:1 ratio holds promise as a beneficial supplementation strategy in the larviculture of *S. cuspidatus*. This supplementation positively impacts on development, growth, and overall productive performance. Notably, it contributed to the enhancement of microbiota, gastrointestinal tract development, and particularly the elongation of intestinal folds.

Ethics statement: The authors declare that they agree with this publication and have made contributions that justify their authorship, that there is no conflict of interest of any kind, and that they have complied with all relevant ethical and legal requirements and procedures. All funding sources are fully detailed in the acknowledgments section. The respective signed legal document is in the journal's archives. They are entirely described in the acknowledgments section.

ACKNOWLEDGMENTS

We thank the company ISAGEN S.A. (Colombia), for financing the study, the company Piscícola San Silvestre S.A. (Barrancabermeja, Colombia) for the logistical support and the company OCEANOS S.A. (Colombia) for the supply of probiotics.

REFERENCES

- Adeoye, A. A., Yomla, R., Jaramillo-Torres, A., Rodiles, A., Merrifield, D. L., & Davies, S. J. (2016). Combined effects of exogenous enzymes and probiotic on Nile tilapia (*Oreochromis niloticus*) growth, intestinal morphology and microbiome. *Aquaculture*, 463, 61–70. <https://doi.org/10.1016/j.aquaculture.2016.05.028>
- Aerts, J., Schaeck, M., De Swaef, E., Ampe, B., & Decostere, A. (2018). *Vibrio lentus* as a probiotic candidate lowers glucocorticoid levels in gnotobiotic sea bass larvae. *Aquaculture*, 492, 40–45. <https://doi.org/10.1016/j.aquaculture.2018.03.059>
- Atencio Garcia, V., Kerguelén, E., Lina, W., & Ana, N. (2003). Manejo de la primera alimentación del bocachico (*Prochilodus magdalenae*). *Revista Mvz Córdoba*, 8. <https://doi.org/10.21897/rmvz.1049>
- Atencio, V., Buelvas, V. M. P., Espitia, F. P., Mestra, R. O., & Carrasco, S. C. P. (2010). Manejo de la primera alimentación de dorada *Brycon sinuensis* ofreciendo larvas de bocachico *Prochilodus magdalenae*. *Revista Colombiana de Ciencias Pecuarias*, 23(3), Article 3. <https://doi.org/10.17533/udea.rccp.324593>
- Avella, M. A., Place, A., Du, S.-J., Williams, E., Silvi, S., Zohar, Y., & Carnevali, O. (2012). *Lactobacillus rhamnosus* Accelerates Zebrafish Backbone Calcification and Gonadal Differentiation through Effects on the GnRH and IGF Systems. *PLOS ONE*, 7(9), e45572. <https://doi.org/10.1371/journal.pone.0045572>

- Balcázar, J., Decamp, O., Vendrell, D., De Blas, I., & Ruiz-Zarzuela, I. (2006). Health and nutritional properties of probiotics in fish and shellfish. *Microbial Ecology in Health and Disease*, 18(2), 65–70. <https://doi.org/10.1080/08910600600799497>
- Begley, M., Hill, C., & Gahan, C. G. M. (2006). Bile Salt Hydrolase Activity in Probiotics. *Applied and Environmental Microbiology*, 72(3), 1729–1738. <https://doi.org/10.1128/AEM.72.3.1729-1738.2006>
- Cai, Y., Yuan, W., Shifeng, W., Guo, W., Li, A., Wu, Y., Chen, X., Ren, Z., & Zhou, Y. (2018). In vitro screening of putative probiotics and their dual beneficial effects: To white shrimp (*Litopenaeus vannamei*) postlarvae and to the rearing water. *Aquaculture*, 498. <https://doi.org/10.1016/j.aquaculture.2018.08.024>
- Cao, H., Yu, R., Zhang, Y., Hu, B., Jian, S., Wen, C., Kajbaf, K., Kumar, V., & Yang, G. (2019). Effects of dietary supplementation with β -glucan and *Bacillus subtilis* on growth, fillet quality, immune capacity, and antioxidant status of Pengze crucian carp (*Carassius auratus* var. Pengze). *Aquaculture*, 508, 106–112. <https://doi.org/10.1016/j.aquaculture.2019.04.064>
- Carnevali, O., de Vivo, L., Sulpizio, R., Gioacchini, G., Olivotto, I., Silvi, S., & Cresci, A. (2006). Growth improvement by probiotic in European sea bass juveniles (*Dicentrarchus labrax*, L.), with particular attention to IGF-1, myostatin and cortisol gene expression. *Aquaculture*, 258(1), 430–438. <https://doi.org/10.1016/j.aquaculture.2006.04.025>
- Carnevali, O., Maradonna, F., & Gioacchini, G. (2017). Integrated control of fish metabolism, wellbeing and reproduction: The role of probiotic. *Aquaculture*, 472, 144–155. <https://doi.org/10.1016/j.aquaculture.2016.03.037>
- Cerezuela, R., Fumanal, M., Tapia-Paniagua, S. T., Meseguer, J., Moriñigo, M. A., & Esteban, M. A. (2012). Histological alterations and microbial ecology of the intestine in gilthead seabream (*Sparus aurata* L.) fed dietary probiotics and microalgae. *Cell and Tissue Research*, 350(3), 477–489. <https://doi.org/10.1007/s00441-012-1495-4>
- De la Hoz Maestre, J., Duarte, L. O., & Manjarrés-Martínez, L. (2017). *Estadísticas de desembarco y esfuerzo de las pesquerías artesanales e industriales de Colombia entre marzo y diciembre de 2017*. <https://doi.org/10.13140/RG.2.2.14291.68645>
- Di, J., Chu, Z., Zhang, S., Huang, J., Du, H., & Wei, Q. (2019). Evaluation of the potential probiotic *Bacillus subtilis* isolated from two ancient sturgeons on growth performance, serum immunity and disease resistance of *Acipenser dabryanus*. *Fish & Shellfish Immunology*, 93, 711–719. <https://doi.org/10.1016/j.fsi.2019.08.020>
- Gisbert, E., Castillo, M., Skalli, A., Andree, K. B., & Badiola, I. (2013). *Bacillus cereus* var. Toyoi promotes growth, affects the histological organization and microbiota of the intestinal mucosa in rainbow trout fingerlings. *Journal of Animal Science*, 91(6), 2766–2774. <https://doi.org/10.2527/jas.2012-5414>
- Hoseinifar, S. H., Sun, Y., & Caipang, C. M. (2016). Short-chain fatty acids as feed supplements for sustainable aquaculture: An updated view. *Aquaculture Research*, 48. <https://doi.org/10.1111/are.13239>
- INCODER. (2006). *PESCA Y ACUICULTURA COLOMBIA 2006*.
- Interaminense, J. A., Vogeley, J. L., Gouveia, C. K., Portela, R. W. S., Oliveira, J. P., Andrade, H. A., Peixoto, S. M., Soares, R. B., Buarque, D. S., & Bezerra, R. S. (2018). In vitro and in vivo potential probiotic activity of *Bacillus subtilis* and *Shewanella* algae for use in *Litopenaeus vannamei* rearing. *Aquaculture*, 488, 114–122. <https://doi.org/10.1016/j.aquaculture.2018.01.027>
- Kong, Y., Li, M., Chu, G., Liu, H., Shan, X., Wang, G., & Han, G. (2021). The positive effects of single or conjoint administration of lactic acid bacteria on *Channa argus*: Digestive enzyme activity, antioxidant capacity, intestinal microbiota and morphology. *Aquaculture*, 531, 735852. <https://doi.org/10.1016/j.aquaculture.2020.735852>
- Kuebutornye, F. K. A., Tang, J., Cai, J., Yu, H., Wang, Z., Abarike, E. D., Lu, Y., Li, Y., & Afriyie, G. (2020). In vivo assessment of the probiotic potentials of three host-associated *Bacillus* species on growth perfor-

- mance, health status and disease resistance of *Oreochromis niloticus* against *Streptococcus agalactiae*. *Aquaculture*, 527, 735440. <https://doi.org/10.1016/j.aquaculture.2020.735440>
- Kumar, V., Roy, S., Meena, D. K., & Sarkar, U. K. (2016). Application of Probiotics in Shrimp Aquaculture: Importance, Mechanisms of Action, and Methods of Administration. *Reviews in Fisheries Science & Aquaculture*, 24(4), 342–368. <https://doi.org/10.1080/23308249.2016.1193841>
- Lozano, G., Hernández, D., Chaves, N., Valderrama, M., Mojica, J., & Gómez, F. (2017). Characterization of skin patterns in *Pseudoplatystoma Magdaleniatum*. *2017 Sustainable Internet and ICT for Sustainability (SustainIT)*, 1–3. <https://doi.org/10.23919/SustainIT.2017.8379806>
- Martínez Cruz, P., Ibáñez, A. L., Monroy Hermosillo, O. A., & Ramírez Saad, H. C. (2012). Use of Probiotics in Aquaculture. *International Scholarly Research Notices*, 2012, e916845. <https://doi.org/10.5402/2012/916845>
- Midhun, S. J., Neethu, S., Arun, D., Vysakh, A., Divya, L., Radhakrishnan, E. K., & Jyothis, M. (2019). Dietary supplementation of *Bacillus licheniformis* HGA8B improves growth parameters, enzymatic profile and gene expression of *Oreochromis niloticus*. *Aquaculture*, 505, 289–296. <https://doi.org/10.1016/j.aquaculture.2019.02.064>
- Mojica, J. I., Usma Oviedo, J. S., Alvarez León, R., & Lasso, C. A. (Eds.). (2012). *Libro rojo de peces dulceacuícolas de Colombia (2012)*. Instituto de Investigación de Recursos Biológicos Alexander von Humboldt.
- Mokhtar, D., Abd-Elhafez, E., & Hassan, A. (2015). Light and Scanning Electron Microscopic Studies on the Intestine of Grass Carp (*Ctenopharyngodon idella*): I- Anterior Intestine. *Journal of Aquaculture Research and Development*, 6. <https://doi.org/10.4172/2155-9546.1000374>
- Mukherjee, A., Chandra, G., & Ghosh, K. (2019). Single or conjoint application of autochthonous *Bacillus* strains as potential probiotics: Effects on growth, feed utilization, immunity and disease resistance in Rohu, *Labeo rohita* (Hamilton). *Aquaculture*, 512, 734302. <https://doi.org/10.1016/j.aquaculture.2019.734302>
- Nayak, S. K. (2010). Probiotics and immunity: A fish perspective. *Fish & Shellfish Immunology*, 29(1), 2–14. <https://doi.org/10.1016/j.fsi.2010.02.017>
- Newaj-Fyzul, A., Al-Harbi, A. H., & Austin, B. (2014). Review: Developments in the use of probiotics for disease control in aquaculture. *Aquaculture*, 431, 1–11. <https://doi.org/10.1016/j.aquaculture.2013.08.026>
- Nicholson, J. K., Holmes, E., Kinross, J., Burcelin, R., Gibson, G., Jia, W., & Pettersson, S. (2012). Host-gut microbiota metabolic interactions. *Science (New York, N.Y.)*, 336(6086), 1262–1267. <https://doi.org/10.1126/science.1223813>
- Paixão, P. E. G., do Couto, M. V. S., da Costa Sousa, N., Abe, H. A., Reis, R. G. A., Dias, J. A. R., Meneses, J. O., Cunha, F. S., Santos, T. B. R., da Silva, I. C. A., Medeiros, E. dos S., & Fujimoto, R. Y. (2020). Autochthonous bacterium *Lactobacillus plantarum* as probiotic supplementation for productive performance and sanitary improvements on clownfish *Amphiprion ocellaris*. *Aquaculture*, 526, 735395. <https://doi.org/10.1016/j.aquaculture.2020.735395>
- Pardo-Carrasco, S., Salas Villalva, J., Reza Gaviria, L., Espinosa Araujo, J., & Atencio-García, V. (2015). Cryopreservation of Trans-Andean shovelnose catfish (*Sorubim cuspicaudus*) semen using dimethylacetamide. *CES Medicina Veterinaria y Zootecnia*, 10(2), 122–131.
- Pérez-Sánchez, T., Mora-Sánchez, B., & Balcázar, J. L. (2018). Biological Approaches for Disease Control in Aquaculture: Advantages, Limitations and Challenges. *Trends in Microbiology*, 26(11), 896–903. <https://doi.org/10.1016/j.tim.2018.05.002>
- Pérez-Sánchez, T., Ruiz-Zarzuela, I., de Blas, I., & Balcázar, J. L. (2014). Probiotics in aquaculture: A current assessment. *Reviews in Aquaculture*, 6(3), 133–146. <https://doi.org/10.1111/raq.12033>
- Prieto-Guevara, M., Atencio Garcia, V., & Pardo Carrasco, S. (2015). *El bagre blanco Sorubim cuspicaudus y su potencial en acuicultura*.
- Qi, Z., Zhang, X.-H., Boon, N., & Bossier, P. (2009). Probiotics in aquaculture of China—Current state, prob-

- lems and prospect. *Aquaculture*, 290(1), 15–21. <https://doi.org/10.1016/j.aquaculture.2009.02.012>
- Ribas, L., & Piferrer, F. (2014). The zebrafish (*Danio rerio*) as a model organism, with emphasis on applications for finfish aquaculture research. *Reviews in Aquaculture*, 6(4), 209–240. <https://doi.org/10.1111/raq.12041>
- Ringø, E., Van Doan, H., Lee, S. h., Soltani, M., Hoseinifar, S. h., Harikrishnan, R., & Song, S. k. (2020). Probiotics, lactic acid bacteria and bacilli: Interesting supplementation for aquaculture. *Journal of Applied Microbiology*, 129(1), 116–136. <https://doi.org/10.1111/jam.14628>
- Sewaka, M., Trullas, C., Chotiko, A., Rodkhum, C., Chansue, N., Boonanuntanasarn, S., & Pirarat, N. (2019). Efficacy of synbiotic Jerusalem artichoke and *Lactobacillus rhamnosus* GG-supplemented diets on growth performance, serum biochemical parameters, intestinal morphology, immune parameters and protection against *Aeromonas veronii* in juvenile red tilapia (*Oreochromis spp.*). *Fish & Shellfish Immunology*, 86, 260–268. <https://doi.org/10.1016/j.fsi.2018.11.026>
- Soltani, M., Ghosh, K., Hoseinifar, S. H., Kumar, V., Lymbery, A. J., Roy, S., & Ringø, E. (2019). Genus bacillus, promising probiotics in aquaculture: Aquatic animal origin, bio-active components, bioremediation and efficacy in fish and shellfish. *Reviews in Fisheries Science & Aquaculture*, 27(3), 331–379. <https://doi.org/10.1080/23308249.2019.1597010>
- Sorieul, L., Wabete, N., Ansquer, D., Mailliez, J.-R., Pallud, M., Zhang, C., Lindivat, M., Boulo, V., & Pham, D. (2018). Survival improvement conferred by the *Pseudoalteromonas* sp. NC201 probiotic in *Litopenaeus stylirostris* exposed to *Vibrio nigripulchritudo* infection and salinity stress. *Aquaculture*, 495, 888–898. <https://doi.org/10.1016/j.aquaculture.2018.06.058>
- Sudha, A., Saravana Bhavan, T. Manjula, R. Kalpana, & M. Karthik. (2019). BACILLUS LICHENIFORMIS AS A PROBIOTIC BACTERIUM FOR CULTURE OF THE PRAWN MACROBRACHIUM ROSENBERGII. *Research Journal of Life Sciences, Bioinformatics, Pharmaceutical and Chemical Sciences*, 05(04). <https://doi.org/10.26479/2019.0504.05>
- Thurlow, C. M., Williams, M. A., Carrias, A., Ran, C., Newman, M., Tweedie, J., Allison, E., Jescovitch, L. N., Wilson, A. E., Terhune, J. S., & Liles, M. R. (2019). *Bacillus velezensis* AP193 exerts probiotic effects in channel catfish (*Ictalurus punctatus*) and reduces aquaculture pond eutrophication. *Aquaculture*, 503, 347–356. <https://doi.org/10.1016/j.aquaculture.2018.11.051>
- Tsai, C.-Y., Chi, C.-C., & Liu, C.-H. (2019). The growth and apparent digestibility of white shrimp, *Litopenaeus vannamei*, are increased with the probiotic, *Bacillus subtilis*. *Aquaculture Research*, 50(5), 1475–1481. <https://doi.org/10.1111/are.14022>
- Van Doan, H., Hoseinifar, S. H., Khanongnuch, C., Kanpiengjai, A., Unban, K., Van Kim, V., & Srichaiyo, S. (2018). Host-associated probiotics boosted mucosal and serum immunity, disease resistance and growth performance of Nile tilapia (*Oreochromis niloticus*). *Aquaculture*, 491, 94–100. <https://doi.org/10.1016/j.aquaculture.2018.03.019>
- Zhang, Q., Ma, H., Mai, K., Zhang, W., Liufu, Z., & Xu, W. (2010). Interaction of dietary *Bacillus subtilis* and fructooligosaccharide on the growth performance, non-specific immunity of sea cucumber, *Apostichopus japonicus*. *Fish & Shellfish Immunology*, 29(2), 204–211. <https://doi.org/10.1016/j.fsi.2010.03.009>
- Zokaeifar, H., Balcázar, J. L., Saad, C. R., Kamarudin, M. S., Sijam, K., Arshad, A., & Nejat, N. (2012). Effects of *Bacillus subtilis* on the growth performance, digestive enzymes, immune gene expression and disease resistance of white shrimp, *Litopenaeus vannamei*. *Fish & Shellfish Immunology*, 33(4), 683–689. <https://doi.org/10.1016/j.fsi.2012.05.027>