



Revista Salud Uninorte

ISSN: 0120-5552

ISSN: 2011-7531

Fundación Universidad del Norte, División de Ciencias de la

SOTO LESMES, VIRGINIA; PARRADO LOZANO, YANETH; GÓMEZ RAMÍREZ, OLGA; GÓMEZ RAMÍREZ, ARLEN; RODRÍGUEZ, MARTHA FABIOLA
Multidrug-resistant Bacteria in Child Care Centers
Revista Salud Uninorte, vol. 36, núm. 3, 2020, Septiembre-Diciembre, pp. 532-544
Fundación Universidad del Norte, División de Ciencias de la

DOI: <https://doi.org/10.14482/sun.36.3.616.92>

Disponible en: <https://www.redalyc.org/articulo.oa?id=81769338003>

- ▶ [Cómo citar el artículo](#)
- ▶ [Número completo](#)
- ▶ [Más información del artículo](#)
- ▶ [Página de la revista en redalyc.org](#)



Sistema de Información Científica Redalyc

Red de Revistas Científicas de América Latina y el Caribe, España y Portugal
Proyecto académico sin fines de lucro, desarrollado bajo la iniciativa de acceso abierto

ARTÍCULO ORIGINAL

DOI: <http://dx.doi.org/10.14482/sun.36.3.616.92>

Multidrug-resistant Bacteria in Child Care Centers

Bacterias multirresistentes en Centros de Cuidado de niños

VIRGINIA SOTO LESMES¹, YANETH PARRADO LOZANO², OLGA GÓMEZ RAMÍREZ³,
ARLEN GOMEZ RAMÍREZ⁴, MARTHA FABIOLA RODRÍGUEZ⁵

¹ Nurse, M.Sc., Ph.D. Profesora titular. Facultad de Enfermería, Universidad Nacional de Colombia. visotol@unal.edu.co. Orcid: orcid.org/0000-0002-3296-8812. CVLAC: https://scienti.colciencias.gov.co/cvlac/visualizador/generarCurriculoCv.do?cod_rh=0000161012. ID: 41773751

² Nurse, M.Sc., Ph.D.. Profesora titular. Facultad de Enfermería, Universidad Nacional de Colombia. ymparradol@unal.edu.co. Orcid: orcid.org/0000-0003-4665-5355. CVLAC: https://scienti.colciencias.gov.co/cvlac/visualizador/generarCurriculoCv.do?cod_rh=0000213942. ID: 41788675

³ Nurse, M.Sc., Ph.D. Profesora asociada. Facultad de Enfermería, Universidad Nacional de Colombia. ojgomezr@unal.edu.co. Orcid: orcid.org/0000-0002-9160-4170. CVLAC: https://scienti.colciencias.gov.co/cvlac/visualizador/generarCurriculoCv.do?cod_rh=0000664200. ID: 52504659

⁴ Veterinary, Ph.D. Profesora asistente. Facultad de Medicina Veterinaria y de Zootecnia, Universidad Nacional de Colombia. apgomezr@unal.edu.co. Orcid: orcid.org/0000-0001-6460-5162. CVLAC: https://scienti.colciencias.gov.co/cvlac/visualizador/generarCurriculoCv.do?cod_rh=0000520829. ID: 52775892

⁵ Bact., M.Sc. Profesora asistente. Facultad de Ciencias de la Salud, Universidad de La Salle. mafarodriguez@unisalle.edu.co. Orcid: orcid.org/0000-0003-1237-9305. CVLAC: https://scienti.colciencias.gov.co/cvlac/visualizador/generarCurriculoCv.do?cod_rh=0000508578. ID: 51.922.969

Institución donde se llevó a cabo la investigación: Universidad de La Salle, sede norte (Av. Carrera 7 No 179-03 - Tel: (571) 677 26 99). Universidad de Nacional de Colombia, Cra 30 # 45-03, Edificio 228, (+57 1) 316 5000 Ext. 17006, 17020. Bogotá D.C., Colombia.

■ ABSTRACT

Purpose: To describe the antibiotic resistance of isolated bacteria, on the surfaces of child care centers.

Methods: Swabs were used to sample the surfaces in 266 childcare centers in Bogotá (Colombia). Bacterial characterization and susceptibility patterns were verified, using a computerized system.

Results: A total of 151 different bacteria were obtained, primarily in the kitchens (36%). *Staphylococcus* was a frequent find, especially *S. hominis*, *S. saprophyticus*, and *S. epidermidis*; 2.7% were resistant to methicillin. *Klebsiella oxytoca* was the most isolated bacteria in kitchens, with high resistance to antibiotics.

Conclusion: This investigation shows the importance of identifying the bacteria present in different environments to which children are exposed, to continuously supervise hygiene habits in child day care centers.

Key Words: Environment; Bacteria; Anti-Bacterial Agents; Child; Child Day Care Centers.

■ RESUMEN

Objetivo: Describir la resistencia a los antibióticos de las bacterias aisladas en las superficies de los centros de cuidado infantil.

Materiales y métodos: Se utilizaron hisopos para tomar muestras de las superficies en 266 centros de cuidado infantil en Bogotá (Colombia). La caracterización bacteriana y los patrones de susceptibilidad se verificaron utilizando un sistema computarizado.

Resultados: Se obtuvieron un total de 151 bacterias diferentes, principalmente en las cocinas (36%). *Staphylococcus* fue un hallazgo frecuente, especialmente: *S. hominis*, *S. saprophyticus* y *S. epidermidis*; El 2,7% era resistente a la meticilina. *Klebsiella oxytoca* fue la bacteria más aislada en las cocinas, con alta resistencia a los antibióticos.

Conclusión: Esta investigación muestra la importancia de identificar las bacterias presentes en diferentes entornos a los que los niños están expuestos para supervisar continuamente los hábitos de higiene en los centros de cuidado diurno infantil.

Palabras claves: Ambiente; bacterias; Antibacterianos; niño; Guarderías Infantiles.

INTRODUCTION

The World Health Organization report that, in 2016, children under the age of five mainly died of diarrhea, acute respiratory infections, and malaria (1). These types of diseases represent a greater burden for lower-income countries, due to flaws in food sanitation, hygiene, and safety, which facilitates pathogen propagation, especially enteric ones (2). Among infectious diseases, those of bacterial etiology are being associated with environmental disinfection and its health conditions, more specifically, surface contamination facilitates the propagation of pathogens among children and family members (3,4).

It is known that the health impact of environmental microbiological exposure is greater in children under the age of five, due to physiological immaturity, and deficient hygiene habits, such as hand washing and adequate toilet use. Similarly, child care centers contain greater indoor concentration (3870 UFC/mm³) of bacteria, when compared to schools and older adult care centers (5) and how to effectively prevent these, remains. This study aimed (i. The air in these environments presents a wide range of bacteria, such as *Bacteroidetes*, *Bacillus*, *Ruminococcus*, *Coprococcus*, *Enterobacter*, *Flavobacterium*, *Staphylococcus*, *Micrococcus*, and *Corynebacterium* (6).

One study identified that *Bacillus*, *Staphylococcus*, *Brevibacillus*, *Pseudomona*, *Moraxella*, *Enterococcus*, *Acinetobacter*, and *Microbacterium* isolate on the toys and furniture of day care centers (7). Furthermore, *Pseudomonas*, *Pantoea*, *Bacillus*, *Staphylococcus*, and *Streptococcus* have been identified in home refrigerators and bathrooms (8) whereas in classrooms, *Sphingomonas*, *Staphylococcus*, and *Enterobacteria* prevailed (9).

Even though most of these bacteria are not primary pathogens, some can survive and be transmitted through the surfaces of indoor environments, and cause community-acquired infections. These bacteria may be nosocomial, and present multidrug resistance, as is the case of methicillin-resistant *Staphylococcus aureus* (MRSA) (10). However, genetic studies with community-acquired MRSA strains have revealed differences in relation to those of nosocomial origins, demonstrating that they are susceptible to many other antibiotics, such as tobramycin, gentamycin, lincomycin, cloranfenicol, vancomycin (11). Studies have shown an increase of community-acquired MRSA among children. These strains are not multi-resistant, but can present additional resistance to erythromycin (12). Other community-acquired pathogens that are multi-resistant to antimicro-

bials are *E.coli* and *Klebsiella spp.* These are common causing agents of urinary infections, with a high percentage of extended-spectrum beta-lactamase producing bacteria (13)Rio Grande do Norte State capital, northeastern Brazil, from 2007 to 2010. A total of 1,082 positive samples were evaluated; *E. coli* was the most prevalent pathogen (60.4%. Possible factors associated with the prevalence of antimicrobial resistance in the community are the inappropriate prescription of antibiotics, the genetic variability of bacteria, and the genetic expression of inducible resistance; situations that require further study (10).

Considering the influence of the environment in transmitting infectious diseases among children, the problem of bacterial resistance, and its relevance to public health, the objective of this study was to characterize the pattern of antibiotic resistance of the bacterial biota isolated from the surfaces of child care centers.

MATERIALS AND METHODS

Study design and area: This was a cross-sectional descriptive study, conducted in 2016, in 230 government-funded child care centers located in the urban perimeter of Bogotá, Colombia. The city had 897 public child care centers, and a population of 11.973 children, between the ages of 2 and 5, at the time of data collection. Sample size was calculated using Win Episcopo 2.0® software (University of Edinburgh, 2000), set at 266 child care centers by assuming a 50% prevalence of infectious agents, confidence level of 95%, and an expected margin of error of 5%. Probabilistic sampling was used, and informed consent was given by all 266 centers, ensuring voluntary participation. This study was approved by the Research Ethics Committee of the Universidad Nacional de Colombia, and was developed according to the principles enshrined in the Declaration of Helsinki.

Sample collection: First, the researchers verified that the facilities contained the following items: at least one bathroom, one kitchen, and a play and eating area for the children. Samples were taken in the first hours of the morning, before the arrival of the children, and then, following the cleaning and disinfection protocol indicated by the person in charge of each center. Samples were taken from surfaces in the kitchens, bathrooms, and places to which children were directly exposed, spent most of their time in, and carried out their play activities (chairs, tables, mattresses, toys, and colored pencils). Swab samples were collecting using the COPAN Venturi Transystem® (Innovation, Italy) (STUART sterile transport swab system), which were then trans-

ported and processed up to eight hours after collection in the Laboratory of Integrative Health of the Universidad de La Salle, where all microbiological analyses took place. The personnel in charge of sample collection were previously trained, and they wore all the necessary personal protective equipment to avoid cross-contamination (disposable caps, gloves, mouth masks, and aprons). The protocols were conducted according to the Standard Operation Procedures (SOP) validated in the lab in previous studies.

Antimicrobial susceptibility profile: The samples were seeded in blood and MacConkey agar, and incubated at 37° C for 24 hours. Bacterial colonies were classified as either Gram-positive or Gram-negative, using Gram staining, and were isolated in trypticase soy agar, under the same incubation conditions. Microbiological typing of the isolates and antibiotic susceptibility was carried out using the Vitek® 2 compact computerized system (Biomérieux, Marcy l'Etoile, France) with Vitek 2-GP cards® (Reference 22218 for Gram-positive), Vitek® 2- GN cards (Reference 414163 for Gram negative), Vitek® 2 AST-P577 (Gram-positive susceptibility), and Vitek® 2 AST-NO82 (Gram-negative susceptibility), according to the protocol described by Herold et al. 1998 (14) To analyze antimicrobial susceptibility, the software used cutoff points established in the Clinical and Laboratory Standards Institute (15). Moreover, as recommended by the manufacturer for antimicrobial identification and susceptibility, the following ATCC strains were used for quality control: *Klebsiella oxytoca* ATCC® 700324, and *Staphylococcus aureus* ATCC®29213. The results were analyzed using descriptive statistics, calculated in terms of frequencies and percentages.

RESULTS

The pattern of antimicrobial susceptibility of the 151 bacteria, isolated from the environments of 266 childcare centers, was analyzed (72 Gram-positive and 79 Gram-negative bacteria). The greatest number of bacteria were found on kitchen surfaces (n=58), followed by bathrooms (n=38), classrooms (n=35) and mattresses, toys, and colored pencils (n=20).

The most common genus of bacteria was *Staphylococcus spp* (72 isolates). Furthermore, 94.4% of these isolates (68/72) were coagulase-negative *Staphylococcus* (CNS), and the most representative species was *S. epidermidis*, with 23 isolates, followed by *S. haemolyticus* (14/72). Additionally, 5.6% were identified as *S. aureus* (4/72). Table 1 shows the distribution of these bacteria among the studied surfaces. [Table 1].

The most common Gram-negative bacilli were *Klebsiella* (*K. oxytoca* and *K. pneumoniae*), *Enterobacter* (*E. cloacae*, *E. aerogenes*, *E. amnigenus* and *E. asburiae*), *Citrobacter* (*C. freundii*, *C. koseri* and *C. amalonaticus*), and *Raoultella* (*R. ornithinolytica* and *R. planticola*) (Table 1).

Table 1. Frequency of isolates by surface analyzed in the early childhood day care centers

Genus	Kitchen	Bathroom	Classroom	Toys/Mattresses	Total (%)
<i>Staphylococcus</i>	8	20	27	17	72 (47.7)
<i>Klebsiella</i>	22	2	4	0	28 (18.5)
<i>Enterobacter</i>	12	1	1	0	14 (9.3)
<i>Citrobacter</i>	4	4	1	0	9 (6.0)
<i>Raoultella</i>	5	3	0	1	9 (6.0)
<i>Serratia</i>	3	3	0	0	6 (4.0)
<i>Escherichia</i>	1	2	1	1	5 (3.3)
<i>Kluyvera</i>	2	0	1	0	3 (2.0)
<i>Acinetobacter</i>	1	1	0	0	2 (1.3)
<i>Leclercia</i>	0	0	0	1	1 (0.7)
<i>Proteus</i>	0	1	0	0	1 (0.7)
<i>Pseudomona</i>	0	1	0	0	1 (0.7)
Total	58	38	35	20	151 (100)

Source: elaboration of the authors.

Regarding the analysis of antimicrobial resistance patterns, species belonging to the *Staphylococcus* genus were resistant to some of the analyzed antibiotics, except for *S. simulans*. The highest percentages of resistance were found against erythromycin (48.6%), oxacillin and tetracycline (26.4%), cephoxytine (23.6%), and clindamycin (20.8%). None of the strains were quinolone-resistant (Table 2). Two *S. hominis* strains (2.7%) presented Methicillin-resistant phenotypes (MRS), which originated from bathrooms and toys/mattresses. Seventeen strains of *Staphylococcus* (23.6%), isolated, primarily, from toys/mattresses and classrooms, presented multidrug resistance, especially to beta-lactamases, macrolides, and clindamycin (Table 3).

Table 2. Number of antibiotic-resistant strains of *Staphylococcus*, isolated from surfaces in early childhood day care centers

Genus	N	Antibiotic														
		Pattern	V	T	C	O	Te	Mi	G	E	Cl	Q/D	Li	N	R	T/S
<i>S. epidermidis</i>	23	R/I	1/0	1/1	6/0	6/0	7/0	0/0	0/0	11/0	4/0	1/0	0/0	0/0	1/0	2/0
<i>S. haemolyticus</i>	14	R/I	2/0	3/0	2/0	2/0	3/0	0/0	0/0	6/0	3/0	2/0	0/0	1/1	2/0	0/0
<i>S. warneri</i>	8	R/I	1/0	1/0	2/0	2/0	3/0	0/0	0/0	7/0	1/2	0/0	0/0	0/0	0/0	0/0
<i>S. saprophyticus</i>	6	R/I	0/0	0/0	1/0	2/0	3/0	0/0	0/0	4/0	1/0	0/0	0/0	0/0	0/0	0/0
<i>S. capitis</i>	6	R/I	0/0	0/0	2/0	1/0	0/0	0/0	1/0	1/0	0/1	0/0	0/0	0/0	0/0	0/0
<i>S. hominis</i>	5	R/I	0/0	0/0	3/0	3/0	2/0	0/0	0/0	4/0	1/0	0/0	0/0	0/0	0/1	0/0
<i>S. aureus</i>	4	R/I	0/1	0/0	0/0	0/0	0/0	0/0	0/0	0/0	0/0	0/0	0/0	0/0	0/0	0/0
Other CNS	6	R/I	0/0	0/0	1/0	3/0	1/0	0/0	0/0	1/1	1/0	0/0	0/0	0/0	0/0	0/0
Total	72	R+I	5	6	17	19	19	0	1	35	15	3	0	4	4	2
		%	6.9	8.3	23.6	26.4	26.4	0	1.4	48.6	20.8	4.2	0	5.6	5.6	2.8

R/I: Resistant/Intermediate. V: Vancomycin, T: Teicoplanin, C: Cefoxitin, O: Oxacillin, Te: Tetracycline, Mi: Minocycline, G:Gentamicin, E: Erythromycin, Cl: Clindamycin, Q/D: Quinupristin/Dalfoprisitn, Li: Linezolid, N:Nitrofurantoin, , R: Rifampicin, T/S: Trimethoprim/Sulfamethoxazole.

Source: elaboration of the authors.

Table 3. Patterns of resistance of *Staphylococcus* strains isolated in this study

Species	Resistance patterns	Isolates (n)
<i>S. hominis</i>	C,O,E,Cl,T/S	1
	C,O,E	1
	C,O,Te,E	1
<i>S. epidermidis</i>	V,T,C,O,E,Cl,Q/D,R	1
	C,O,E,C,l	1
	Te,E,T/S	2
	C,O,Te,E	1
	C,O,Cl,N	2
	T,E,Cl	1
<i>S. haemolyticus</i>	V,T,C,O,E,Cl,Q/D,N,R	2
	C,O,Te,E,Cl	1
<i>S. warneri</i>	V,T, E,Cl	1
	C,O,Te,E.	2
<i>S. saprophyticus</i>	C,O,Te,E.	2
<i>S. cohnii</i>	C,O,E,Cl,	1

V: Vancomycin, T: Teicoplanin, C: Cefoxitin, O: Oxacillin, Te: Tetracycline, E: Erythromycin, Cl: Clindamycin, Q/D: Quinupristin/Dalfoprisitn, N:Nitrofurantoin, R: Rifampicin, T/S: Trimethoprim/Sulfamethoxazole.

Source: elaboration of the authors.

The Gram-negative bacilli identified in the study were most resistant to ampicillin (57.0%) and cephalosporins (cephalothin - 49.4%, Cefuroxime Axetil - 34.2% and Cephuroxime - 30.4%). *Enterobacteriaceae* isolates did not display multidrug resistance; however, most were resistant to cephalosporin. Of the non-fermenting Gram-negative bacilli, *Acinetobacter* was resistance to beta-lactamases and nitrofurantoin. The only isolate of *P. aureginosa* in this study was multidrug resistant (beta-lactamases, nitrofurantoin, and Trimethoprim/Sulfamethoxazole) (Table 4).

Table 4. Number of antimicrobial resistant strains of Gram-negative bacilli isolated from surfaces in early childhood day care centers

Genus	n	Pattern	Antibiotic													
			A	AS	C1	C2	C2-A	C3	C3*	C3p	C4	Am	G	NIT	T/S	
<i>Citrobacter</i>	9	R/I	-/0	-/0	7/0	4/0	4/0	0/0	0/0	0/0	0/0	0/0	0/0	0/1	0/1	2/0
<i>Enterobacter</i>	14	R/I	-/0	-/0	14/0	8/0	10/0	0/0	0/0	0/0	0/0	0/0	0/0	0/0	1/2	2/0
<i>Escherichia</i>	5	R/I	2/0	1/0	1/3	0/0	0/0	0/0	0/0	0/0	0/0	0/0	0/0	1/0	1/0	
<i>Klebsiella</i>	28	R/I	23/0	2/0	1/1	1/0	1/0	0/0	0/0	0/0	0/0	0/0	0/0	1/3	1/0	
<i>Kluyvera</i>	3	R/I	3/0	2/0	3/0	1/0	2/0	0/0	0/0	0/0	0/0	0/0	0/0	0/0	0/0	
<i>Leclercia</i>	1	R/I	0/0	0/0	0/0	0/0	0/0	0/0	0/0	0/0	0/0	0/0	0/0	0/0	0/0	
<i>Proteus</i>	1	R/I	0/0	0/0	0/0	0/0	0/0	0/0	1/0	0/0	0/0	0/0	0/0	1/0	0/0	
<i>Raoultella</i>	9	R/I	9/0	0/0	2/0	0/1	0/1	0/0	0/0	0/0	0/0	0/0	0/0	0/0	1/0	
<i>Serratia</i>	6	R/I	5/0	0/0	6/0	6/0	6/0	0/0	0/0	0/0	0/0	0/0	0/0	6/0	0/0	
<i>Acinetobacter</i>	2	R/I	2/0	1/0	-/0	2/0	2/0	0/0	0/0	0/1	-/0	-/0	0/0	2/0	0/0	
<i>Pseudomona</i>	1	R/I	1/0	1/0	1/0	1/0	1/0	1/0	0/0	1/0	0/0	0/0	0/0	1/0	1/0	
Total	79	R	45	7	39	24	27	1	1	2	0	0	1	19	8	
		%	57.0	8.9	49.4	30.4	34.2	1.3	1.3	2.5	0	0	1.3	24.1	10.1	

R/I: Resistant/Intermediate. A: Ampicillin, AS: Ampicillin/sulbactam, C1: Cephalothin, C2: Cefuroxime, C2-A: Cefuroxime Axetil, C3: Cefotaxime, C3*: Ceftazidime, C3p: Ceftriaxone, C4: Cefepime, Am: Amikacin, G: Gentamicin, Nitrofurantoin, T/S: Trimethoprim/Sulfamethoxazole. In the table: (-) not assessed.

Source: elaboration of the authors.

DISCUSSION

Child care centers can represent transmission sites for infectious diseases, especially because of the exposure of children to surfaces considered reservoirs for microorganisms. In the present study, the most common isolates found in classrooms, bathrooms, mattresses, and toys was *Sta-*

phylococcus spp, especially CNS. This finding is consistent with that of previous studies, which have shown that these environments contain high levels and diversity of microorganisms, with a predominance of phyla *Firmicutes* and *Proteobacteria*, and genera *Bacillus* and *Staphylococcus* (6,7).

CNS have been isolated from the hands of children (16), and more generally, from skin and mucous membrane microbiota, reaching densities greater than 10³ strains/cm² (17), enough for contamination through direct contact or flaking skin. Normal skin flora is considered one of the main origins of bacteria in indoor environments (18,19), especially in bathrooms (8) and other surfaces of the home (9), similar to the findings of the present study. CNS are normal commensal bacteria present on healthy human skin and mucous membranes; however, approximately 10% are multidrug resistant (17).

A study found that 42% of the isolates of microorganisms found on skin were resistant to erythromycin, 24.3% to tetracycline, and 13.2% to oxacillin (19), similar to the findings presented here. However, in the present study, CNS presented greater resistance, as the automated methodologies revealed the presence of the MRS phenotype in two *S. hominis* strains, and 17 strains resistant to beta-lactamases, macrolides, and clindamycin (*S. hominis*, *S. haemolyticus*, *S. cohnii*, *S. saprophyticus*, *S. epidermidis* and *S. warneri*). The most commonly isolated phenotype profile of CNS was resistant to ampicillin, penicillin, oxacillin, and trimetroprim-sulfametoxazol, either with or without susceptibility to first-generation cephalosporins, clindamycin, erythromycin, gentamicin, and tetracycline. The present study corroborated the emergence of oxacillin-resistant and multidrug resistant CNS, which has been reported as a matter of public health concern in recent years (16).

In the present study, four strains of *S. aureus* were isolated from classrooms, mattresses, toys, and kitchens were sensitive to methicillin (MSSA) and the other analyzed antibiotics. Only one strain presented intermediate resistance to vancomycin. The results corroborate those of a study conducted by Scott et al. (8), which reported methicillin-sensitive *S. aureus* isolated from toys; however, the bacteria identified in bathrooms and kitchens were found to be resistant (MRSA). This finding is noteworthy, considering that children can acquire these bacteria through surface contact, enhancing their transmission and generating an increase in community-acquired MRSA infections.

Another relevant finding in the present study was the isolation of a multidrug-resistant strain of *P. aeruginosa* from bathroom surfaces. This bacterium has been reported to be a multidrug-re-

sistant nosocomial pathogen that affects respiratory and urinary tracts, causing outbreaks in hospitals (20). However, it has also been associated with infections in people, including children, who participate in recreational activities involving exposure to sources of water in which this microorganism can survive. *P. aeruginosa* can be present in the water supply and sanitation systems, forming biofilm, its intrinsic resistance to various antibiotics, and high capacity to acquire genes that inhibit the action mechanisms of new antimicrobials, can incur in complications for the treatment of infections (21).

Pepperell et al., have reported that enterobacteria, especially *Citrobacter* and *Enterobacter*, presented multidrug resistance associated with the genetic recombination processes triggered by the action of an integrase that enables integrons to adhere to resistance genes. The importance of *Citrobacter* and similar species stems from its capacity to survive long periods because of its low level of virulence; thus, resistance genes accumulate, which can be transmitted to other more highly virulent organisms (22). Considering these studies, special mention goes to the risks found in the present investigation, such as bacteria belonging to the *Enterobacteriaceae* (*Citrobacter*, *Serratia*, *Enterobacter*, and *Kluyveria*) family that were 100% resistant against cephalosporins, specifically first-generation ones. This can be explained by the constant and prolonged use of antibiotics and high production of beta-lactamases.

The results of the present study did not show multidrug resistance in *Enterobacteriaceae*. These findings are consistent with those reported by Odhiambo et al. (23), in which most of the enteric microorganisms did not present multidrug resistance against commonly used antimicrobials, such as ampicillin, tetracycline, and co-trimoxazole. However, the authors also informed resistance against these drugs by *Salmonella spp*, *Shigella flexineri*, and *Shigella spp* isolates.

In the present study, bacteria such as *Citrobacter*, *E. coli*, *Kuyvera*, *Pantoea* and *Enterobacter* were resistant to more than three antibiotics, whereas *Klebsiella* and *Serratia* were resistant to four others. These profiles are significant, considering reports of the presence of multidrug-resistant genes in strains isolated from non-hospitalized children, indicating the transmission of these microorganisms in the community (24).

Contributions of the present research include the characterization of microorganisms and antibiotic susceptibility patterns of bacterial biota, isolated from different early childhood day care

centers in Bogota. These microorganisms are communicable and generate risk of infection among children. This study points to the need to conduct interventions relative to hand hygiene and carrying out surface cleaning and disinfection protocols in child care centers. Such practices can reduce exposure of children to potentially-infectious microorganisms. Efforts must be employed to ensure healthy environments, and reduce the unnecessary prescription of antibiotics, which is not only related to the issue of multidrug resistance, but also generates burdens for health services and the resources of families.

Conflicts of interest: None

Acknowledgments: To Colciencias, Universidad Nacional de Colombia and Universidad de La Salle for the support in the development of research.

REFERENCES

1. World Health Organization. World Health statistics 2018. Monitoring health for the SDGs, sustainable development goals. WHO, editor. Geneva; 2018. 100 p.
2. BD 2017 Risk Factor Collaborators. Global , regional , and national comparative risk assessment of 84 behavioural , environmental and occupational , and metabolic risks or clusters of risks for 195 countries and territories , 1990 – 2017 : a systematic analysis for the Global Burden of Dis. *Lancet*. 2018;392(10):1923–94.
3. Torner N, Soldevila N, Garcia JJ, Launes C, Godoy P, Castilla J. Effectiveness of non-pharmaceutical measures in preventing pediatric influenza : a case – control study. *BMC Public Health*. 2015;1–8.
4. Acharya D, Singh JK, Adhikari M, Gautam S, Pandey P, Dayal V. Journal of Infection and Public Health Association of water handling and child feeding practice with childhood diarrhoea in rural community of Southern Nepal. *J Infect Public Health*. 2018;11(1):69–74.
5. Madureira J, Paciência I, Cavaleiro J, Costa C, Teixeira JP, De Oliveira E. Assessment and determinants of airborne bacterial and fungal concentrations in different indoor environments: Homes, child day-care centres, primary schools and elderly care centres. *Atmos Environ*. 2015;109:139–46.
6. Deng W, Chai Y, Lin H, So WWM, Ho KWK, Tsui AKY, et al. Distribution of bacteria in inhalable particles and its implications for health risks in kindergarten children in Hong Kong. *Atmos Environ*. 2016;128:268–75.

7. Prussin AJ, Vikram A, Bibby KJ, Marr LC. Seasonal Dynamics of the Airborne Bacterial Community and Selected Viruses in a Children's Daycare Center. Yao M, editor. *PLoS One* [Internet]. 2016 Mar 4 [cited 2020 Sep 6];11(3):e0151004. Available from: <https://dx.plos.org/10.1371/journal.pone.0151004>
8. Jeon Y-SS, Chun J, Kim B-SS. Identification of household bacterial community and analysis of species shared with human microbiome. *Curr Microbiol*. 2013 Nov;67(5):557–63.
9. Kelley ST, Gilbert JA. Studying the microbiology of the indoor environment. 2013;1–9.
10. Vindel A, Cercenado E. [Methicillin-resistant *Staphylococcus aureus* carrying the *mecC* gene: an emerging problem?]. Vol. 34, *Enfermedades infecciosas y microbiología clínica*. Spain; 2016. p. 277–9.
11. Zhang C, Guo L, Chu X, Shen L, Guo Y, Dong H, et al. Presence of the Panton-Valentine Leukocidin Genes in Methicillin-Resistant *Staphylococcus aureus* Is Associated with Severity and Clinical Outcome of Hospital-Acquired Pneumonia in a Single Center Study in China. Otto M, editor. *PLoS One* [Internet]. 2016 Jun 1 [cited 2020 Sep 6];11(6):e0156704. Available from: <https://dx.plos.org/10.1371/journal.pone.0156704>
12. Miller MB, Weber DJ, Goodrich JS, Popowitch EB, Poe MD, Nyugen V, et al. Prevalence and Risk Factor Analysis for Methicillin-Resistant *Staphylococcus aureus* Nasal Colonization in Children attending child care center. *J Clin Microbiol*. 2011;49(3):1041–7.
13. Cunha MA, Assunção GLM, Medeiros IM, Freitas MR. Antibiotic resistance patterns of urinary tract infections in a northeastern Brazilian capital. *Rev Inst Med. Sao Paulo* [Internet]. 2016 [cited 2020 Sep 6];58. Available from: [/pmc/articles/PMC4793943/?report=abstract](https://pubmed.ncbi.nlm.nih.gov/2793943/)
14. Herold BC, Immergluck LC, Maranan MC, Lauderdale DS, Gaskin RE, Boyle-Vavra S, et al. Community-acquired methicillin-resistant *Staphylococcus aureus* in children with no identified predisposing risk. *JAMA*. 1998;279(8):593–8.
15. CLSI. Performance Standards for Antimicrobial Susceptibility Testing; Twenty-Third Informational Supplement. 2013.
16. Candido De Almeida MC, Corrêa I. Bacterias presentes en las manos de los niños en edad escolar en la Unidad de Internación Pediátrica. *Investig y Educ en Enfermería*. 2012;2:240–4.

17. May L, Klein EY, Rothman RE, Laxminarayan R. Trends in antibiotic resistance in coagulase-negative staphylococci in the United States, 1999 to 2012. *Antimicrob Agents Chemother* [Internet]. 2014 Mar 1 [cited 2020 Sep 6];58(3):1404–9. Available from: <http://dx.doi.org/10.1128>
18. Hospodsky D, Qian J, Nazaroff WW, Yamamoto N, Bibby K, Rismani-Yazdi H, et al. Human occupancy as a source of indoor airborne bacteria. *PLoS One*. 2012;7(4).
19. Freeman S, Okoroafor NO, Gast CM, Koval M, Nowowiejski D, O'Connor E, et al. Crowdsourced Data Indicate Widespread Multidrug Resistance in Skin Flora of Healthy Young Adults. *J Microbiol Biol Educ*. 2016;17(1):172–82.
20. Cholley P, Gbaguidi-Haore H, Bertrand X, Thouverez M, Plésiat P, Hocquet D, et al. Molecular epidemiology of multidrug-resistant *Pseudomonas aeruginosa* in a French university hospital. *J Hosp Infect*. 2010 Dec;76(4):316–9.
21. Hocquet D, Muller A, Bertrand X. What happens in hospitals does not stay in hospitals: antibiotic-resistant bacteria in hospital wastewater systems. *J Hosp Infect*. 2016 Aug;93(4):395–402.
22. Liu L, Chen D, Liu L, Lan R, Hao S, Jin W, et al. Genetic Diversity, Multidrug Resistance, and Virulence of *Citrobacter freundii* From Diarrheal Patients and Healthy Individuals. *Front Cell Infect Microbiol* [Internet]. 2018 Jul 10 [cited 2020 Sep 6];8(Jul):233. Available from: <https://www.frontiersin.org/article/10.3389/fcimb.2018.00233/full>
23. Odhiambo F, Galgalo T, Wences A, Maina Muchemi O, E WK, Chepkemoi Tonui J, et al. Antimicrobial resistance: capacity and practices among clinical laboratories in Kenya, 2013. *Pan Afr Med J*. 2014;19:1–8.
24. Hijazi SM, Fawzi MA, Ali FM, Abd El Galil KH. Multidrug-resistant ESBL-producing Enterobacteriaceae and associated risk factors in community infants in Lebanon. *J Infect Dev Ctries*. 2016 Sep;10(9):947–55.