



Revista Salud Uninorte

ISSN: 0120-5552

ISSN: 2011-7531

Fundación Universidad del Norte, División de Ciencias de la

ÁLVAREZ, FRANCISCA KEITH; SANDOVAL, JAVIERA SCHEU;
NAVARRO, DANTON MORENO; DÍAZ-NARVÁEZ, VÍCTOR
Frequency of pediatric dentomaxilar anomalies in
Orthodontics Barros Luco Trudeau healthcare complex, Chile
Revista Salud Uninorte, vol. 37, no. 3, 2021, September-December, pp. 683-695
Fundación Universidad del Norte, División de Ciencias de la

DOI: <https://doi.org/10.14482/sun.37.3.617.645>

Available in: <https://www.redalyc.org/articulo.oa?id=81771260011>


- How to cite
- Complete issue
- More information about this article
- Journal's webpage in redalyc.org

UNEN
redalyc.org

Scientific Information System Redalyc

Network of Scientific Journals from Latin America and the Caribbean, Spain and Portugal

Project academic non-profit, developed under the open access initiative



Fecha de recepción: abril 13 de 2020
Fecha de aceptación: julio 23 de 2021

ARTÍCULO ORIGINAL

<https://dx.doi.org/10.14482/sun.37.3.617.645>

Frequency of pediatric dentomaxilar anomalies in Orthodontics Barros Luco Trudeau healthcare complex, Chile

Frecuencia de anomalías dentomaxilares pediátricas en ortodoncia perteneciente al complejo asistencial Barros Luco Trudeau, Chile

FRANCISCA KEITH ÁLVAREZ¹, JAVIERA SCHEU SANDOVAL², DANTÓN MORENO NAVARRO³, Víctor Díaz-Narváez⁴

¹ Academic titles: DDS Universidad San Sebastian. Complejo asistencial Hospital Barros Luco Trudeau. Gran Avenida Jose Miguel Carrera 3204, San Miguel, Región Metropolitana de Chile. +56966091146. francisca.keith@gmail.com. Av.Suecia 1818, depto 604, Providencia, Santiago de Chile. 7500000 Orcid: <http://orcid.org/0000-0002-3566-737X>.

Contribution in the article: Research idea, sample examiner, writing of the manuscript, Criticism, final approval of the manuscript and author of the correspondence.

² DDS Universidad Mayor. Complejo asistencial Hospital Barros Luco Trudeau. Gran Avenida Jose Miguel Carrera 3204, San Miguel, Región Metropolitana Santiago de Chile. +56982337346. javiera.scheu@gmail.com. Orcid: <http://orcid.org/0000-0002-3977-7736>.

Contribution in the article: Research idea, Sample examiner, writing of the manuscript, translation of the manuscript and final approval of the manuscript.

³ DDS Universidad de Chile, Orthodontist CONACEO. Complejo asistencial Hospital Barros Luco Trudeau. Gran Avenida Jose Miguel Carrera 3204, San Miguel, Región Metropolitana Santiago de Chile. +56996790250. dantonmoreno@yahoo.es Orcid: <http://orcid.org/0000-0003-2989-9657>.

Contribution in the article: Research idea, Sample examiner, writing of the manuscript, provides the subject of the manuscript and final approval of the manuscript.

⁴ Ph.D Universidad Andres Bello. School of Dentistry. Universidad Andres Bello. Av. República 239, Santiago, Región metropolitana de Chile. +56991614015. victor.diaz@unab.cl. Orcid: <http://orcid.org/0000-0002-5486-0415>.

Contribution in the article: Criticism, writing of the manuscript, correction and final approval of the manuscript

Corresponding author: Francisca Keith Álvarez. francisca.keith@gmail.com . None of the mentioned authors has a conflict of interest. Self-funded research.

RESUMEN

Introducción: Las anomalías dentomaxilares (ADM) son la tercera patología bucal más frecuente en Chile.

Objetivo: Determinar la distribución de frecuencia de anomalías dentomaxilares observadas en los pacientes atendidos por el Departamento de Ortodoncia dentro del Servicio de Especialidades Odontológicas (SEO) perteneciente al Complejo Asistencial Barros Luco Trudeau, Santiago de Chile. La población pertenece a pacientes pediátricos entre 5 y 15 años de edad que presentan ADM, atendidos dentro del servicio de salud metropolitano sur.

Método: La muestra de 250 pacientes con ADM fué recolectada a partir del mes de abril a septiembre del año 2019, por lo tanto, es una muestra por conveniencia, los datos fueron extraídos de las fichas clínicas de los pacientes. La clasificación de las anomalías estudiadas fue realizada mediante la clasificación biogenética de Bonn modificada por la cátedra de la Universidad de Chile.

Resultado: Se observó que la ADM intermaxilar más frecuente de la muestra es en el sentido transversal y corresponde en el espacio lateral a la mordida cruzada unilateral, por otra parte, la ADM intramaxilar más frecuente es la alteración de tamaño dentoalveolar por diente grande. Todas las ADM intermaxilares coincidieron en ser más frecuentes a la edad de 9 años.

Conclusión: Se logró determinar la distribución de las frecuencias de ADM en pacientes atendidos dentro del SEO, siendo esta en el sentido transversal, correspondiente a mordida cruzada unilateral. Con el presente estudio se pretende hacer una contribución para futuros estudios que utilicen metodologías similares.

Palabras clave: Maloclusión; Ortodoncia; Epidemiología; Anomalías dentales.

ABSTRACT

Introduction: Dentomaxilar anomalies (DMA) are the third most common oral pathology in Chile.

Objective: To determine the frequent of dentomaxilar anomalies observed in patients treated in the Department of Orthodontic of Dental Specialties Service (DSS) belonging to the Barros Luco Trudeau Healthcare Complex, Santiago, Chile.

The population belongs to pediatric patients between 5 and 15 years, who have DMA, treated in the southern metropolitan health service.

Method: The sample of 250 patients with DMA was chosen from April to September of the year 2019, therefore, the data was extracted from the clinical records of the patients. The classification of the anomalies was carried out using the Bonn's biogenetic method, modified by the University of Chile.

Result: It was observed that the most frequent intermaxillary DMA of the sample is in the transverse direction and corresponds to the unilateral crossbite, on the other hand, the most frequent intramaxillary DMA is the alteration of dentoalveolar per large tooth. All intermaxillary DMA agreed to be more frequent at the age of 9 years.

Conclusion: It was possible to define the frequency of the DMA in patients treated in the DSS, this being in the transverse direction, corresponding to unilateral crossbite. With the present study it is intended to make a contribution for future studies that use similar methodologies.

Key words: Malocclusion; Orthodontics; Epidemiology; Dental anomalies, Pediatrics, Interceptive.

INTRODUCTION

Dentomaxilar anomalies (DMA) constitute a public health problem because it is one of the most prevalent oral pathologies in Chile and in the rest of the world from the epidemiological point of view.

DMA correspond to the third most prevalent oral pathology in the Chilean population after dental caries and gingival or periodontal diseases. (1-8)

DMA correspond to a group of alterations in the development of the jaws that frequently appear with altered dental position that affect the shape, function and aesthetics of the stomatognathic

system, whose symptoms include changes in the numbers, shapes and structures of the teeth. (3-4,9-11)

The etiology is multifactorial; therefore, DMA may be the result of one or more alterations, which include hereditary factors, alterations during the development stage, alteration of eruption and exfoliation patterns of developing teeth, environmental factors, among others. In the environmental factors are stomatognathic dysfunctions or parafunctions, also called Non-physiological oral habits. (4,6,8,9,12)

These anomalies affect the aesthetics and functionality of the stomatognathic system and constitute the immediate cause of the loss of the harmonic relationship between the jaws and teeth, many of these anomalies are progressive and can transcend to the adult life in the interpersonal relationships, they can develop a feeling of shame about their dental appearance, feeling shy in social situations, even losing job opportunities, which translates into discontent, deterioration in the quality of life, social and functional limitations. (2,9,10,13)

Orthodontics is the branch of dentistry that is responsible for preventing, diagnosing, intercepting, and treating dental malpositions and/or maxillofacial disorders. When appear the first signs of the disease we begin to interpose barriers to prevent its unfavorable evolution, all of them are interceptive orthodontic treatment. Is estimated that the need for posterior orthodontic treatment is reduced, therefore, early treatment is recommended to achieve the greatest amount of benefits for the patient, including better use of growth potential, less need for extractions, more limited fixed appliances treatments, leading to better and more stable over time, even orthognathic surgery could be avoid

From an epidemiological point of view, interceptive orthodontics turns out to be the most optimal treatment option, since it allows to control and treat these anomalies at their least serious moment, enhancing the success of the treatment. (5,14-18)

The Bonns school postulated a classification where DMA are grouped into six main groups: three groups of intermaxillary DMA and three groups of intramaxillary DMA. In Chile, the biogenetic classification has been modified by the Department of Dentomaxilar Orthopedics of the University of Chile since 1965, which is based on the Bonn classification. The classification gains high value for making diagnoses of epidemiological utility. (3,4,12):

Intermaxillary anomalies: Vertical direction: open bite and overbite; Transverse direction: bis to bis bite, unilateral cross bite and bilateral cross bite; Sagittal: inverted Simple bite, accommodation inverted bite without secondary alteration, accommodation inverted bite with secondary alteration and distocclusions. Intramaxillary anomalies: Alterations of dental size, alterations in the number of pieces and alterations due to loss of dental tissue.

The success of the treatments also depends on the resources, the management and the ability to optimize both aspects. Knowledge the number of patients to be treated and, above all, the type of the anomalies should be treated would be important for planning and management resources. It is extremely important to make an epidemiological diagnosis of DMA in a specific population. (7,19)

In our country, there is no national study that allows establishing real data on the prevalence of DMA. The review of the available studies, most of them carried out in the Metropolitan Region, provides only local information, which cannot be extrapolated to the entire country due to the great diversity of these studies designs, the great variability between the age groups, the size and selection of the samples, etc. (20)

The objective of this work is to determine the frequency of DMA present in patients treated in the Orthodontic Specialty of the Dental Specialties Service (DSS) of the Barros Luco Trudeau Healthcare Complex in Santiago, Chile. Applying a standardized measurement methodology to be able to compare with other studied populations.

MATERIALS AND METHODS

Research Design: exploratory, descriptive, and cross-sectional.

Location: it was carried out in the Dental Specialties Service (DSS) in the Orthodontic area, in the Barros Luco Trudeau Healthcare Complex, Santiago, Metropolitan Region, Chile.

This complex serves several districts that belong to the South Metropolitan area. The study was carried out between the months of April and September of the year 2019.

Procedure: Two professional dentist examiners were chosen, who were calibrated through a specific course for this purpose. A pilot sample consisting of 30 patients, randomly selected with

properly diagnosed DMAs. These samples were examined by both dentists independently and for their respective classification (there was no contact between them during this procedure). The samples were coded and randomized to each of the calibrators.

Population: patients in orthodontics treatment in the DSS. Inclusion criteria: Pediatric patients in orthodontic treatment in DSS, patients with DMA in one or more senses of space. Exclusion criteria: Patients with bad behavior, syndromic patients.

Sample: The total sample consisted of 250 pediatric patients with DMA in an age range of 5 to 15 years, 124 female patients and 126 male patients.

Data collection: the same clinical file that the patients had realized during their orthodontic treatment, the realization of this file consisted of an anamnesis of the patients that records dental information, including the patient oral habits, followed by a clinical examination.

The collected data were tabulated in the Excel® program. The classification used in this study is the Bonns biogenetics, modified by the department of dentomaxilar orthopedics at the Faculty of Dentistry of the University of Chile (3,4,12)

Statistical tests: calibration between subjects was performed using the Kappa test.

The frequency of cases was estimated as the division between a specific type of DMA over the total DMA (20). The level of significance used was $\alpha \leq 0.06$.

Ethical aspects: It was approved by the Barros Luco Trudeau Healthcare Complex ethics committee with resolution number 3972 dated August 26, 2019.

RESULTS

The total sample consisted of 250 pediatric patients with DMA of which 49,6% was female and 50,4% male. The distribution of the sample by age denotes that most of the children were 9 years old, represented by 28,4%, followed by the age of 10 years corresponding to 20,4% and, thirdly the age of 8 years corresponding to 19,2%.

Analyzing the vertical anomalies, 24,8% presented an open bite and 15,5% an overbite. The most frequent vertical DMA in both genders is the open bite where the female gender predominates over the male. (Table 1)

Table 1. Distribution of vertical intermaxillary DMA according to sex

	Type of DMA	female	male	total
	n	n	n	%
Overbite	19	20	39	15,5
Open bite	34	28	62	24,8
No alteration	71	78	149	59,6

In relation to the transverse anomalies, it was observed that 56,4% correspond to the anterior area with dental crowding and 21,2% with protrusion. DMA with dental crowding is the most frequent in both genders, but it is more prevalent in the female gender. In the lateral area, 17,2% correspond to unilateral crossbite, 6,8% bilateral crossbite. The unilateral crossbite DMA is the most frequent in both genders, with the masculine gender prevailing. (Table 2)

Table 2. Distribution of cross-sectional intermaxillary DMA according to sex

	Type of DMA	female	male	total
	n	n	n	%
Anterior zone				
- With protrusion	23	30	53	21,2
- With crowding	75	66	141	56,4
- No alteration	26	30	56	22,4
Lateral zone				
- Cross				
Unilateral	18	25	43	17,2
Bilateral	7	10	17	6,8
- Bis a bis	0	0	0	0
- No alteration	99	91	190	76

The intermaxillary anomalies 34,8% presented inverted simple bite, 14,4% presented distocclusions, the most frequent DMA in both sexes is the inverted simple bite where the male gender predominates over the female. (Table 3)

Table 3. Distribution of sagittal intermaxillary DMA according to sex

	Type of DMA	female	male	total
	n	n	n	%
Distocclusions	19	17	36	14,4
Simple inverted bite	34	53	87	34,8
Inverted bite by accommodation without secondary modification	0	0	0	0
Inverted bite by accommodation with secondary modifications	0	0	0	0
No alteration	71	56	127	50,8

With respect to intramaxillary DMA in relation to the dentomaxilar size, 68,0% corresponds to a large tooth and 3,2% to a small tooth.

According to the alteration by number of teeth, it was found 3,2% corresponds to supernumeraries and 1,6% to agenesis. The loss of dental tissue corresponds to 15,6%. The most frequent DMA is the alteration of dentoalveolar size by large tooth and its distribution by genders is symmetrical. (Table 4)

Table 4. Distribution of intramaxillary DMA according to sex

	Type of DMA	female	male	total
	n	n	n	%
Dentomaxilar size				
- Big tooth	85	85	170	68,0
- Little tooth	5	3	8	3,2
- No alteration	34	38	72	28,8
Number of dental pieces				
- Supernumerary	1	7	8	3,2
- Agenesis	2	2	4	1,6
- No alteration	121	117	238	95,2
Tissue loss				
- Dental (various)	19	20	39	15,6
- No alteration	105	106	211	84,4

Finally, remembering that for each examined can be more than one DMA, it is observed that of all the intermaxillary DMAs in the three directions of space, the most frequent is in the transverse direction with 79,9%, followed by 49,2 % in the sagittal direction and finally 40,4% in the vertical direction it is also evident that at the age of 9 years it is the most frequent with respect to the total of the DMA in the three directions of space. In the vertical and sagittal sense, the masculine gender predominates over the female and in the transversal sense, the female gender predominates over the masculine. (Table 5)

Table 5. Distribution of intermaxillary DMA according to sense of space, age and sex

Age	Vertical DMA			Sagittal DMA			Transversal DMA		
(Year)	Female	Male	%	Female	Male	%	Female	Male	%
5	1	0	0,4	1	1	0,8	2	0	0,8
6	1	3	1,6	5	3	3,2	4	3	2,8
7	3	5	3,2	7	7	5,6	14	9	9,2
8	9	11	8,0	9	13	8,8	15	20	14,0
9	17	19	14,4	11	20	12,4	33	29	24,8
10	12	8	7,2	12	12	9,6	16	27	17,2
11	6	2	3,2	4	8	4,8	7	7	5,6
12	2	2	1,6	1	1	0,8	7	3	4,0
13	2	0	0,8	2	2	1,6	4	1	2,0
14	0	0	0,0	0	2	0,8	0	2	0,8
15	0	0	0,0	1	1	0,8	1	1	0,8
Total	53	48	40,4%	53	70	49,2%	103	102	79%

DISCUSSION

There are no previous studies corresponding to the sample observed in this work, it is pertinent to analyze the national studies in the Metropolitan Region that resemble ours.

In 1972 Richards et al already revealed that 65% to 72,6% of Chilean children between 5 and 14 years old had some type of DMA. This study is similar to the age range of our study however, he maintains a different methodology in the rest of its research compared to ours. (21)

In 2005 in the southern area of the Metropolitan Region, Carreño et al found that 29,3% of the children between the ages of 2 and 6 had interceptable DMA. The most prevalent DMA were open bite with 17,1% and compressions with 6,65%. Our study also measures patients in the southern area of the metropolitan region, among our results in patients with DMA, the most frequent is in the transverse direction due to compressions with 79%, causing cross bite and crowding in the anterior sector and in the lateral sector, this is followed by sagittal DMA with 49,2% where the most frequent corresponding to the inverted simple bite. (22)

Scott and Calderón, in 2009, carried out a study at the primary level of health care within the southeast metropolitan service, where they measure the implementation of the “norms for the prevention and interception of dentomaxilar anomalies” in 1998, the study determined that only 9,3% of the population enrolled in the primary health service in the South had access to interceptive orthodontic treatment. This data is relevant when considering that the present study measures the frequency of DMA in the population extrapolated to the current southern metropolitan health area, specifically within the Barros Luco Trudeau Hospital. (23)

Pardo et al carried a study in 2018 in children from 3 and 10 years of age, with a total of 106 patients attended in the clinic of the child and adolescent of 4th year of the faculty of dentistry of the University of Chile, which yielded a 45,1% prevalence of DMA in any of the 3 senses of space. In the case of the present study, children between 5 and 15 years old, represented in a total of 250 patients, had 100% DMA due to being in orthodontic treatment, so our results distribute the frequencies of DMA according to the direction of the space they affect. Both studies used modified Bonn's Biogenetic classification and a more similar methodology compared to the rest of the studies. (12)

As can be seen in this study, DMA affects a high percentage of the population under 15 years old, with a different degree of commitment, since there is not enough infrastructure or human resources in the country to provide them with an adequate solution. (20)

In conclusion, with the results, it was possible to analyze and compare the different frequencies of DMA, present in all the patients in the sample, with respect to the type of DMA, age and gender, mainly. However, we consider ourselves in debt for not being able to measure the real prevalence of DMA, since it requires access to a universal sample that includes even several undiagnosed patients. It is worth mentioning that this debt is extrapolated at the national level since there is no

updated literature on the prevalence of DMA through Chile, which is important to determine the need for treatment and thus be able to provide adequate coverage.

REFERENCES

- MINSAL, Departamento de Salud Bucal División de Prevención y Control de Enfermedades. Análisis de situación de salud bucal en Chile, 2010.
- Cueto, A., Skog, F., Muñoz, M., Espinoza, S., Muñoz, D., & Martínez, D. Prevalencia de Anomalías Dentomaxilares y Necesidad de Tratamiento en Adolescentes. *International journal of odontostomatology*, 2017; 11(3): 333-338. Doi: 10.4067/S0718-381X2017000300333
- González A. Frecuencia de pacientes con mordida cruzada derivados al Programa conducente al título profesional de especialista en Ortodoncia y Ortopedia Dento Máxilo Facial de la Facultad de Odontología de la Universidad de Chile entre los años 2013-2015 [Internet]. Santiago, Chile: Universidad de Chile - Facultad de Odontología; 2016 [citado: 2020, febrero]. Disponible en: <http://repositorio.uchile.cl/handle/2250/147230>.
- Martens Landeta C. Prevalencia de mordida abierta y malos hábitos bucales en niños de 3 y 7 años de sectores vulnerables de Santiago. *Tesis Doctoral. Universidad Andrés Bello*, 2015 (Chile).
- Burgos, D. Prevalencia de maloclusiones en niños y adolescentes de 6 a 15 años en Frutillar, Chile. *International journal of odontostomatology*, 2014; 8(1): 13-19. Doi: 10.4067/S0718-381X2014000100002
- Wauters, M., Vergara, D., Vergara, P., Pérez, D., Parada, J., & Aguilera, J. Prevalencia de anomalías dentomaxilares y articulatorias en escolares, Cochamó 2014. *Revista Chilena de Ortodoncia*, 2015; 32 (2): 74-81.
- Calderón, N., Cerna, N., Escudero, D., Guzmán, C., Leppe, J., & Álvarez, E. Descripción de la implementación de las normas de prevención e intercepción de anomalías dentomaxilares a nivel primario del Servicio de Salud Metropolitano Oriente. *Rev Chil Ortod*, 2009; 26(2): 54-62.
- Muente, R., Aburto, R., & Tapia, P. Prevalencia de maloclusiones y necesidad de tratamiento de ortodoncia en niños de octavo básico de la comuna de Viña del Mar. *Rev Chil Ortod*, 2015; 32(2): 82-88.
- Singh, R. N., Shahi, A. K., Ramesh, V., Sharma, S., Kumar, S., & Chandra, S. Prevalence of malocclusion and orthodontic treatment needs among 12-15 years old school children in Patna, Eastern India. *J Family Med Prim Care*, 2019; 8(9): 2983. Doi: 10.4103/jfmpc.jfmpc_681_19

- Bilgic, F., Gelgor, I. E., & Celebi, A. A. Malocclusion prevalence and orthodontic treatment need in central Anatolian adolescents compared to European and other nations' adolescents. *Dental Press J Orthod*, 2015; 20(6): 75-81. Doi: 10.1590/2177-6709.20.6.075-081.oar
- Baron, C., Houchmand-Cuny, M., Enkel, B., & Lopez-Cazaux, S. Prevalence of dental anomalies in French orthodontic patients: A retrospective study. *Archives de Pédiatrie*, 2018; 25(7): 426-430. Doi: 10.1016/j.arcped.2018.07.002
- Pardo, M. C., Araya, F. M., Prado, A. G., & Miranda, M. A. C. Prevalencia de anomalías dentomaxilares en dentición primaria y mixta primera fase en la Facultad de Odontología de la Universidad de Chile, 2018. *Odontología sanmarquina*, 2019; 22(3): 181-185. Doi: 10.15381/os.v22i3.16707
- Yao, J., Li, D. D., Yang, Y. Q., McGrath, C. P. J., & Mattheos, N. What are patients' expectations of orthodontic treatment: a systematic review. *BMC oral health*, 2016; 16(1): 19.
- Burhan, A. S., & Nawaya, F. R. Preventive and interceptive orthodontic needs among Syrian children. *J Egypt Public Health Assoc*, 2016; 91(2): 90-94. Doi: 10.1097/01.EPX.0000483238.83501.60
- Ministerio de Salud Chile. Guía Clínica Salud Oral integral para niños y niñas de 6 años. Santiago: MINSAL; 2013.
- Sandoval, P., & Bizcar, B. Beneficios de la implementación de ortodoncia interceptiva en la clínica infantil. *International journal of odontostomatology*, 2013; 7(2): 253-265. Doi: 10.4067/S0718-381X2013000200016
- Carrasco-Sierra, M., Mendoza-Castro, A. M., & Andrade-Vera, F. M. Implementación de la ortodoncia interceptiva. *Dominio de las Ciencias*, 2018; 4(1): 332-340. Doi: 10.23857/dc.v4i1.745
- Santiesteban-Ponciano F, Alvarado-Torres E. Ortodoncia interceptiva - revisión bibliográfica. *Rev Latinoamericana de ortodoncia y odontopediatria*. 2015. Doi: 10.23857/dc.v4i1.745
- Sollenius, O., Petrén, S., Björnsson, L., Norlund, A., & Bondemark, L. Health economic evaluations in orthodontics: a systematic review. *Eur J Orthod*, 2016; 38(3): 259-265. Doi: 10.1093/ejo/cjv040
- Normas en la Prevención e Intercepción de Anomalías Dentomaxilares". División de salud de las personas, Departamento odontológico, MINSAL, 1998.
- Richards C., Gantz R., Cabrera J., Ayala J. (1972). "Las anomalías dentomaxilares, daño poblacional y realidad social". *Revista Dental de Chile*, 62: 47-49.

Carreño S., Calderón N. (2005). "Prevelencia de anomalías dentomaxilares interceptables y malos hábitos bucales en niños con dentición temporal completa del área sur de la región Metropolitana". *Revista Chilena Ortodoncia* 22:18-25.

Scott K. Descripción de la implementación de las normas de prevención e intercepción de anomalías dentomaxilares a nivel primario del Servicio de Salud Metropolitano Sur Oriente [Internet]. Santiago, Chile: Universidad de Chile - Facultad de Odontología; 2009 [citado: 2020, marzo]. Disponible en: <http://repositorio.uchile.cl/handle/2250/134985>